



The Case for Guided Self Help for People with intellectual disabilities

Journal:	<i>Advances in Mental Health and Intellectual Disabilities</i>
Manuscript ID	AMHID-10-2016-0030.R1
Manuscript Type:	Personal View
Keywords:	Intellectual Disability, Guided self-help, Mental Health, Psychological treatments, Depression, anxiety

SCHOLARONE™
Manuscripts

The Case for Guided Self Help for People with Intellectual Disabilities

Abstract

Purpose

This article examines Guided Self-Help (GSH), and some of the barriers as to why it is not routinely available for people with intellectual disabilities (ID).

Design

This article offers an overview of GSH and the potential benefits of it as an intervention for people with ID with mild depression and/or anxiety.

Findings

The current literature reports the successful use and effectiveness of GSH in the general population. However despite this there is little evidence that it is being used in practice for people with ID.

Originality

This paper offers an overview of GSH and advocates for its increasing use for people with ID to help bring about equality in mental health care.

Introduction

It is accepted that people with intellectual disability (ID) have higher rates of mental illness. Indeed it is estimated that between 20.1% -22.41% of adults will experience mental illness (excluding challenging behavior (CB) (Cooper, Smiley, Morrison, et al, 2007). This compares to an estimate of 16% in the general population (Department of Health, 2003). This article looks at Guided Self-Help (GSH) and examines why it is not being made routinely available for people with ID, in spite of evidence of its efficacy in the general population. GSH is fast becoming an important intervention for the management of common mental health problems such as depression and anxiety (National Collaborating Centre for Mental Health, 2010). However there little evidence to suggest this treatment is available to or being developed for people with ID is Mild depression and anxiety are the targets of GSH, however often these conditions can be overlooked, both people with ID and in the general population, where it is estimated that a third of people with depression and half of those with anxiety are undiagnosed and therefore not treated. This has a financial and human cost with milder forms of depression and anxiety being associated with increased risk of mortality (Russ, Stamatakis, Hamer, et al, 2012).

National Policy such as the Green Light Tool Kit (National Development Team for Inclusion, 2013) has attempted to improve mental healthcare for people with ID and drive the mainstream agenda of equitable access to mental healthcare. For many there is still difficulty accessing mental health services and those that do are less likely to receive psychological treatments (Michaels, 2008). Paradoxically the availability of psychological interventions for people with ID with a range of less intrusive person centered treatment options being available. Indeed only a decade ago ID was an exclusion criterion in studies evaluating psychological treatments. Mason, (2007) puts forward five factors that are believed to influence psychological therapy outcomes:

- the perceived effectiveness of clinicians
- individual clinician competence
- how well the service is resourced in terms of the number of clinicians
- the level of the client's disability
- the presence of diagnostic overshadowing bias.

1
2
3 However in spite of a growth in the availability of psychological therapies for people
4 with ID, a reliance on proxy based reporting has meant that self-report options have
5 traditionally been ignored (Fujiura, 2012).
6
7
8
9

10 11 **Barriers to psychological treatments and GSH**

12
13 There is little in terms of self-help materials available to people with ID outside the
14 context of individual psychological therapy. Internet GSH, may not be an option for
15 some as they cannot be easily accessed; other GSH programmes may require high
16 levels of health literacy about a condition e.g., depression, which could exclude
17 people with ID. People with ID may lack opportunities to engage in or to enjoy
18 activities that enhance or protect their mental wellbeing. Given a lack of accessible
19 GSH materials there are a number of things to be aware of when providing treatment
20 such as difficulty in comprehension, being able to understand their feelings and
21 emotions or conceptualise. e.g. guilt, low self-esteem or self-worth. The inability to
22 articulate or the clinician's style of questioning or basic awareness of ID can mean
23 these complex emotions are missed and therefore not considered.
24
25
26
27
28
29
30
31

32 For those who do access treatment a lack of knowledge of the needs of and how to
33 support people with ID can lead to treatment failure and fuel the idea that these
34 types of treatments do not work and are of little use. Reasonable adjustments are
35 required to enable equitable access to healthcare and health outcomes. These
36 factors should not be a barrier to treatment, but something that needs to be
37 considered when planning treatments. For example a clinician might ask a patient
38 with ID to complete an online measure of depression as part of their commitment to
39 offering equal access to services. However if they cannot read or can't understand
40 the questions they will be unable to participate. Adding a reasonable adjustment
41 such as a voiceover or access to someone who can support the activity is likely to
42 contribute to a better health outcome and ensure equity.
43
44
45
46
47
48
49
50
51

52 53 **What is GSH**

54
55 The NICE Guidelines (GG90) (National Collaborating Centre for Mental Health,
56 2010, p182) define GSH as,
57
58
59
60

1
2
3 “... a self-administered intervention designed to treat depression, which makes use
4 of a range of books or other self-help manuals derived from an evidence-based
5 intervention and designed specifically for the purpose. A healthcare professional (or
6 paraprofessional, for example, graduate and low-intensity workers in mental health)
7 facilitates the use of this material by introducing, monitoring and reviewing the
8 outcome of such treatment.
9
10

11
12
13 The use of low-intensity interventions such as GSH to treat mild depression and
14 promote positive mental health, mean that many people do not need or go on to
15 receive more intrusive treatments, that may produce unwanted side effects and that
16 are less well tolerated. GSH allows the involvement of others who are important to
17 the person to support them either formally or informally in line with Recovery
18 principles (Lovell & Richards, 2008), allowing self-management of symptoms where
19 possible to reduce dependence on services (Lovell, Bower, Richards, *et al*, 2008).
20 There have been attempts to standardise the implementation of GSH within the
21 United Kingdom, although there is as yet no consensus. NICE (2010) and the
22 Scottish Executive (2006) both identified four essential components of GSH:
23
24
25
26
27
28
29
30

NICE (2010)	Scottish Executive (2006)
<ul style="list-style-type: none"> • frequent support 	<ul style="list-style-type: none"> • information on common mental health problems
<ul style="list-style-type: none"> • minimum support, 	<ul style="list-style-type: none"> • advice and coping
<ul style="list-style-type: none"> • group psychoeducation 	<ul style="list-style-type: none"> • self-directed structured plan
<ul style="list-style-type: none"> • support by mail 	<ul style="list-style-type: none"> • supported self-help

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 According to NICE guidelines, individual GSH programmes based on Cognitive
49 Behavioural Therapy should:

- 50
51
52
53
54
55
56
57
58
59
60
- Include the provision of written materials of an appropriate reading age (or alternative media to support access)
 - Be supported by a trained practitioner, who typically facilitates the self-help programme and reviews progress and outcome

- Consist of up to six to eight sessions (face-to-face and via telephone) normally taking place over 9 to 12 weeks, including follow-up.

(National Collaborating Centre for Mental Health, 2010, pp., 13-214).

Evidence for GSH as an effective Mental Health Intervention

Research into GSH in ID is poor. There are case studies that have focused on self- and the use of techniques and strategies involving diaries, self-monitoring and relaxation exercises, which are completed as homework, following individual or group sessions. These studies have demonstrated that people with ID are able to use and benefit from self-help techniques central to GSH. Taylor (2002) reported twelve studies (1986–2002) the majority of which focused upon skills training within a cognitive behavioural framework, using self-instruction and interpersonal problem solving aimed at addressing cognitive deficits, rather than to modify cognitive content and distortions. These studies measured a number of variables, including anger, aggression, self-esteem and depression. Although not all participants showed a decrease in these behaviours, there was evidence of measurable improvements for participants.

In the general population, studies of GSH using CBT techniques have reported positive outcomes and have been endorsed by the Government as an effective means of combating depression (Department of Health, 2001a). The evidence base for GSH has been subject to a number of reviews. The NICE guidelines (National Collaborating Centre for Mental Health, 2010, pp. 184-187) reviewed 18 book based GSH using Randomised Controlled Trials (RCT) methodology. Two of the studies reported a beneficial effect for those with mild depression and sub threshold depression, trending towards statistical significance. The five studies characterised by frequent support with minimum duration reported a large effect when reporting reduction of depressive symptoms with waiting list controls. The other studies considered had insufficient data and wide confidence intervals, to be able to reach any meaningful conclusions. Of the eighteen RCTs reported in the NICE guidelines that met inclusion criteria, two examined individual GSH with guided support, ten individual with minimal support, three groups GSH (psycho education) and three GSH by mail.

1
2
3 Gellatly, Bower, Hennessy, *et al* (2007), examined the role of moderators on
4 effectiveness of GSH e.g., patient populations, study design, intervention content
5 and compared RCTs versus controls in the treatment of depressive symptoms.
6
7

8 Thirty-four studies were identified which included 39 comparisons. Greater
9 effectiveness was associated with a number of factors including: recruitment outside
10 of clinical settings, those with a diagnosis of depression rather than people at risk of
11 depression and use of CBT techniques. In terms of delivery, Lovell, Bower,
12 Richards, *et al* (2008a) found no evidence that the number of sessions or how GSH
13 was delivered e.g., mail, computer face to face was related to outcomes. However
14 outcomes improved when GSH was based on CBT and those with mild to moderate
15 depression were found to do better than those with a more severe clinical
16 presentation.
17
18
19
20
21
22

23 Since the NICE guidelines on depression were originally published both GSH and
24 self-help have received greater recognition and acceptance as a legitimate treatment
25 modality for depression (IAPTs, 2010). There has been further guidance published to
26 help those facilitating the treatment and to distinguish it from other interventions.
27 IAPTs (2010) has given guidance on developing self-help materials, which includes
28 that they are technically accurate, engage with the person and maintain that
29 engagement, the sessions reflect the persons own life and provide a structure so
30 that they can see change brought about by the treatment. For this to happen it is
31 necessary that appropriate materials are available to support the person during the
32 intervention.
33
34
35
36
37
38
39
40
41
42

43 **Psychological input and GSH for people with ID**

44
45 There is evidence that people with ID not only can benefit from, but can be
46 instrumental in developing new approaches with clinicians e.g. The SAINT (Chaplin
47 *et al*, 2014, 2017). The SAINT is a GSH resource for people with ID, presented in an
48 easy read format, designed to encourage people with an ID to recognise and
49 identify their feelings, particularly those that may cause or lead to distress and
50 impact on the person's daily lives and mental health. The person is encouraged with
51 or without support to identify their feelings and following this is asked to select an
52 appropriate coping strategy or intervention from a checklist to use. The person also
53
54
55
56
57
58
59
60

has a diary which can be used to record not only their feelings and what strategies they used to cope, but also the things they may have done well or enjoyed during the day. This provides an insight into a person's mental health over time, and can assist families, carers and professionals to not only monitor threats to mental wellbeing, but also to see the effects of any coping strategies used. Currently the SAINT is being tried in clinical practice and we are to await the report (Russ, 2016). In addition the author has received reports of its use in local services not only to provide GSH but to act as a structure for nurses to base 1:1 or group sessions they have with clients with ID and to assist treatment goals such as problem solving and teaching good mental health strategies. In terms of feedback from those receiving the SAINT, below is a selection of comments

"I like the coping strategies"

"I just used the book any time I get depressed"

"I use it during the day if I got staff I can talk to. Some staff can be sort of thing, not very understanding"

"I have found it very good and found it very helpful a lot of people have written down feeling diaries sad and helpless sought of thing and has helped me with my moods as well"

Conclusion

There is a considerable burden of mental health for people with ID and it is only in recent history that psychological treatment options such as GSH have been explored. Self-help and GSH have been found to be effective in the wider population and the absence of specific evidence of its effectiveness relating to people with ID should not preclude its use, providing reasonable adjustments are made.

People with ID are more at risk to mental health problems. These conditions can be difficult to detect and therefore can go unnoticed. There are a number of reports of the successful use of various self-monitoring and stress management techniques including symptom diaries and the use of strategies aimed at reducing distress that are consistent with GSH approaches. There are few GSH materials available for people with ID. However early evidence of the SAINT demonstrate that people with ID can engage and use this intervention to improve their health and mental

1
2
3 wellbeing. The use of low level treatments such as GSH should be made more
4 widely available for people with ID particularly given the evidence for its
5 effectiveness in the general population.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Bhaumik, S., Gangadharan, S., Hiremath, A., et al (2011) Psychological treatments in intellectual disability: the challenges of building a good evidence base. *British Journal of Psychiatry*, **198**, 428-430.

Chaplin, E.; Craig, T.; McCarthy, J.; & Bouras, N. (2017) The SAINT a guided self-help intervention for people with intellectual disabilities, *Journal of Mental Health Research in Intellectual Disabilities* Accepted 10.01.2017

Chaplin, E, McCarthy, J, Hardy, S, Underwood, L, Spain, D, Cronin, P, Peyton, L and Henry, J (2014) The SAINT a Guided Self Help approach for people with intellectual disability, Pavilion Brighton.

Cooper, S. A., Smiley, E., Morrison, J., et al (2007) Mental ill-health in adults with intellectual disabilities: prevalence and associated factors. *The British Journal of Psychiatry*, **190**, 27-35.

Department of Health (2003) Better or worse: a longitudinal study of the mental health of adults living in private households in Great Britain. London: The Stationery Office.

Fujiura, G. T. (2012) Self-reported health of people with intellectual disability. *Intellectual and Developmental Disabilities*, **50**, 352-369.

Gellatly, J., Bower, P., Hennessy, S., et al (2007) What makes self-help interventions effective in the management of depressive symptoms? Meta-analysis and meta-regression. *Psychological Medicine*, **37**.

IAPTs (2010) Good practice guidance on the use of self-help materials within IAPT services Available from <http://www.iapt.nhs.uk/silo/files/good-practice-guidance-on-the-use-of-selfhelp-materials-within-iapt-services.pdf> accessed 20th October 2016.

Lovell, K., Bower, P., Richards, D., et al (2008) Developing guided self-help for depression using the Medical Research Council complex interventions framework: a description of the modelling phase and results of an exploratory randomised controlled trial. *BMC Psychiatry*, **8**, 91.

Lovell, K. & Richards, D. (2008) A recovery programme for depression. The University of Manchester, The University of York,: Rethink.

Mason, J., (2007). The provision of psychological therapy to people with intellectual disabilities: an investigation into some of the relevant factors. *Journal of Intellectual Disability Research*, **51**(3), pp.244-249.

1
2
3 **Michaels, J. (2008)** Healthcare for all: Report of the Independent Inquiry Into access
4 to healthcare for people with learning disabilities. London: Aldridge Press

5
6 **National Collaborating Centre for Mental Health (2004)** Depression: management
7 of depression in primary and secondary care - NICE guidance, London.

8
9 ---- **(2010)** Depression: The NICE Guidelines on the treatment and management of
10 depression in adults (Updated Edition). The British Psychological Society and The
11 Royal College of Psychiatrists.

12
13 **National Development Team for Inclusion (2013)** A guide to auditing and
14 improving your mental health services so that it is effective in supporting
15 people with autism and people with learning disabilities, Bath, National
16 Development Team for Inclusion

17
18
19 **Russ, M (2016)**, Implementing the SAINT in Clinical Practice, Poster presentation,
20 22nd International Network for Psychiatric Nursing Research (NPNR) Conference
21 2016 15 Sep 2016 - 16 Sep 2016, Nottingham Conference Centre, Burton Street,
22 Nottingham, NG1 4BU

23
24 **Russ, T. C., Stamatakis, E., Hamer, M., et al (2012)** Association between
25 psychological distress and mortality: individual participant pooled analysis of 10
26 prospective cohort studies. *British Medical Journal*, **345**, 4933-4933.

27
28 **Scottish Executive (2006)** National Evaluation Of The 'Doing Well By People With
29 Depression' Programme, A report of the three year programme to enhance services
30 in primary care for people with mild to moderate depression. Edinburgh, The Scottish
31 Government.

32
33
34 **Taylor, J. L. (2002)** A review of the assessment and treatment of anger and
35 aggression in offenders with intellectual disability. *Journal of Intellectual Disability*
36 *Research*, **46 Suppl 1**, 57-73.

37
38 **Thompson Prout, H. & Nowak-Drabik, K. (2003)** Psychotherapy With Persons
39 Who Have Mental Retardation: An Evaluation of Effectiveness. *American Journal of*
40 *Mental Retardation*, **108**, 82-93.

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60