

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/124107>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

OPEN

Transplantation Publish Ahead of Print

DOI: 10.1097/TP.0000000000002931

Recipient Comorbidity and Survival Outcomes after Kidney Transplantation: a UK-wide Prospective Cohort Study

Diana A Wu (MChB)^{1*}, Matthew L Robb (PhD)^{2*}, John L R Forsythe (MD)^{1,2}, Clare Bradley (PhD)³, John Cairns (MPhil)⁴, Heather Draper (PhD)⁵, Christopher Dudley (MD)⁶, Rachel J Johnson (MSc)², Wendy Metcalfe (MD)¹, Rommel Ramanan (MD)⁶, Paul Roderick (MD)⁷, Charles R V Tomson (DM (Oxon))⁸, Christopher J E Watson (MD)⁹, J Andrew Bradley (PhD)⁹, Gabriel C Oniscu (MD)¹

*Joint first authors

¹Transplant Unit, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, EH16 4SA, UK

²NHS Blood and Transplant, Fox Den Road, Stoke Gifford, Bristol, BS34 8RR, UK

³Health Psychology Research Unit, Orchard Building, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK

⁴London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK

⁵Health Sciences, University of Warwick, Coventry, UK

⁶Department of Renal Medicine, Southmead Hospital, Southmead Road, Bristol, BS10 5NB, UK

⁷Academic Unit of Primary Care and Population Sciences, Faculty of Medicine, University of Southampton, Tremona Road, Southampton, SO16 6YD, UK

⁸Department of Renal Medicine, Freeman Hospital, Newcastle upon Tyne, NE12 9NY, UK

⁹Department of Surgery, University of Cambridge and the NIHR Cambridge Biomedical Research Centre, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 0QQ, UK

Correspondence to: Gabriel C. Oniscu, Consultant Transplant Surgeon and Honorary Reader, Transplant Unit, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, EH16 4SA, UK. Tel: 0044-1312421715, gabriel.oniscu@ed.ac.uk

Authorship

All authors contributed to the study design. DAW conducted the literature review. DAW and MLR conducted the data analysis. DAW and GCO drafted the manuscript. All authors interpreted the data, revised the drafts and approved the final version.

Disclosure

All authors declared no conflicts of interests.

Funding

This work is part of the Access to Transplantation and Transplant Outcome Measures (ATTOM) research programme funded by the National Institute for Health Research (grant number RP-PG-0109-10116). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. The funding body had no role in study design, data collection, data analysis, data interpretation, writing of the manuscript or decision to submit for publication.

Abbreviations

ARD, adjusted risk difference

ATTOM, Access to Transplantation and Transplant Outcome Measures

BMI, body mass index

CI, confidence interval

CIT, cold ischaemia time

CLD, chronic liver disease

cRF, calculated reaction frequency

CVD, cerebrovascular disease

DDKT, deceased-donor kidney transplant

DGF, delayed graft function

ERA-EDTA, European Renal Association – European Dialysis and Transplant Association

ESRD, end-stage renal disease

HF, heart failure

HLA, human leukocyte antigen

HR, hazard ratio

IHD, ischaemic heart disease

LDKT, living-donor kidney transplant

MM, mismatches

PVD, peripheral vascular disease

SE, standard error

UK, United Kingdom

WHO, World Health Organisation

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Comorbidity is increasingly common in kidney transplant recipients, yet the implications for transplant outcomes are not fully understood. We analysed the relationship between recipient comorbidity and survival outcomes in a UK-wide prospective cohort study – ATTOM.

Methods: 2100 adult kidney transplant recipients were recruited from all 23 UK transplant centers between 2011-2013. Data on 15 comorbidities were collected at the time of transplantation. Multivariable Cox regression models were used to analyse the relationship between comorbidity and 2-year graft survival, patient survival and transplant survival (earliest of graft failure or patient death) for deceased-donor kidney transplant (DDKT) recipients (n=1288) and living-donor kidney transplant (LDKT) recipients (n=812).

Results: For DDKT recipients, peripheral vascular disease (HR 3.04, 95%CI 1.37, 6.74, p=0.006) and obesity (HR 2.27, 95%CI 1.27, 4.06, p=0.006) were independent risk factors for graft loss, while heart failure (HR 3.77, 95%CI 1.79, 7.95, p=0.0005), cerebrovascular disease (HR 3.45, 95%CI 1.72, 6.92, p=0.0005) and chronic liver disease (HR 4.36, 95%CI 1.29, 14.71, p=0.018) were associated with an increased risk of mortality. For LDKT recipients, heart failure (HR 3.83, 95%CI 1.15, 12.81, p=0.029) and diabetes (HR 2.23, 95%CI 1.03, 4.81, p=0.042) were associated with poorer transplant survival.

Conclusion: The key comorbidities that predict poorer 2-year survival outcomes after kidney transplantation have been identified in this large prospective cohort study. The findings will facilitate assessment of individual patient risks and evidence-based decision making.

Introduction

Kidney transplantation is widely regarded as the treatment of choice for end-stage renal disease (ESRD). However, outcomes after transplantation vary considerably between patients and prediction of individual risk is challenging due to the increasing prevalence of complex comorbidity among the ESRD population. Conditions such as diabetes, hypertension and obesity which contribute to the development of ESRD are on the rise,¹ while ESRD itself is an important risk factor for other comorbidities such as cardiovascular disease.^{2, 3} Over the past decade, the proportion of deceased-donor kidney transplant (DDKT) recipients older than 60 years of age has increased from 17% to 29% in the UK,⁴ and the burden of comorbidity among patients undergoing kidney transplantation has also risen significantly.⁵⁻⁷

Despite this, there are limited data on the impact of comorbidity on transplant outcomes. A small number of studies have demonstrated the overall detrimental effect of comorbidity on transplant outcomes using various comorbidity indices.^{5, 8-10} However, this does not allow characterisation of the risks associated with specific comorbid conditions. Retrospective registry analyses have identified several comorbidities as risk factors for transplant outcomes, but the results show considerable heterogeneity and are limited by the reliability of the data.¹¹⁻¹³ Up-to-date and reliable evidence is essential to enable clinicians to fully inform patients of their individual risks and likely outcomes, thereby facilitating shared decision-making and informed consent.

We conducted a national prospective cohort study to investigate the impact of a wide range of baseline comorbid conditions on survival outcomes following kidney transplantation. We report the two-year survival outcomes of the study which was conducted as part of the Access to Transplantation and Transplant Outcome Measures (ATTOM) research programme.

Materials and Methods

Study design and participants

ATTOM is a national prospective cohort study investigating the factors that influence access to and outcomes from renal transplantation in the UK. A full description of the ATTOM protocol has been reported previously.¹⁴ A cohort of 2262 incident kidney transplant recipients were recruited to ATTOM at the time of transplantation, from all 23 UK renal transplant centers. In each center recruitment took place over a 12-month period between 1st November 2011 and 31st March 2013. Patients aged 18-75 years were eligible for recruitment. For the purposes of this analysis, multi-organ transplant recipients (n=162) were excluded. The final study sample (n=2100) represented 73% of eligible study participants from the national kidney-only transplant population (Figure 1). Patients were followed up for two years from the date of transplant. DDKT recipients (n=1288) and living-donor kidney transplant (LDKT) recipients (n=812) were analysed separately.

Data variables

The variables of interest were recipient comorbidities at the time of transplantation comprising diabetes, ischaemic heart disease (IHD), heart failure (HF), atrial fibrillation, cardiac valve replacement, pacemaker, cerebrovascular disease (CVD), peripheral vascular disease (PVD), abdominal aortic aneurysm, chronic respiratory disease, chronic liver disease (CLD), blood borne viruses, malignancy, mental illness (definitions given in Table S1, SDC, <http://links.lww.com/TP/B802>) and body mass index (BMI).

The primary outcome measures were graft survival, patient survival and transplant survival. Graft survival was defined as the time from transplantation to graft failure (earliest of return to dialysis or retransplantation), with censoring for death with a functioning graft, at last follow-up or at 2 years. Patient survival was defined as the time from transplantation to patient death, with censoring at last follow-up or at two years. Transplant survival is a

composite outcome defined as the time from transplantation to the earliest of graft failure or patient death, with censoring at last follow-up or at two years.

Potential confounders considered in multivariable analyses included (a) recipient variables: age, gender, ethnicity, primary renal disease (as classified by ERA-EDTA codes¹⁵), time on dialysis, previous transplantation, sensitisation level, smoking status; (b) donor variables: age, gender, ethnicity, BMI; (c) transplant variables: human leukocyte antigen (HLA) mismatches (MM), cold ischaemia time (CIT), delayed graft function (DGF). Ethnicity was coded as White, Black, Asian and Other (including Chinese and mixed origin). Recipient calculated reaction frequency (cRF) $\geq 85\%$ was used to define highly sensitised recipients. The cRF is the percentage of a pool of 10,000 UK donors to whom the recipient has unacceptable HLA antibodies. HLA mismatches were classified into 4 levels as defined by the current UK deceased-donor kidney allocation scheme: level 1 (000 HLA-A, B, DR MM), level 2 (0DR + 0/1B MM), level 3 (0DR + 2B MM) or (1DR + 0/1B MM) and level 4 (1DR + 2B MM) or (2DR MM).¹⁶

Data collection

Baseline recipient variables (including comorbidity) were collected by trained research nurses at the time of transplantation from patient interviews, case notes, local electronic patient information systems and/or confirmed with the patient's named consultant nephrologist.

Independent validation of 5% of data entries in all research sites confirmed $>98\%$ concordance for all data fields.¹⁴ Donor and transplant variables and 2-year graft and patient survival data were obtained through linkage with the UK Transplant Registry.

Statistical methods

Baseline characteristics were compared with chi-squared tests for categorical data and Mann-Whitney U tests for nonparametric continuous data. The impact of comorbidity on two-year survival outcomes was examined using Kaplan-Meier estimates and Cox proportional hazards

regression models. DDKT and LDKT recipients were analysed separately. As there were no significant differences in outcomes between recipients of donors after circulatory death and donors after brain death, all DDKT recipients were analysed together. For DDKT recipients, separate multivariable models were built for the three different outcomes of transplant, graft and patient survival. For LDKT recipients, modelling was only possible for transplant survival, as the lower number of graft failures and patient deaths prevented modelling of graft and patient survival separately. All comorbidities were considered for inclusion in the multivariable models, and those leading to a significant ($p < 0.05$) change in log likelihood were retained using a manual backward elimination method. Models were adjusted for statistically significant variables as well as variables selected *a priori* on the basis of clinical relevance. Continuous variables were explored as linear, fractional polynomials and categorical variables. In all models, the effect of the time on dialysis variable was only found to be significant after 3 years, and thus it was converted to a binary variable (<3 years versus ≥ 3 years) as this provided the best fit in each model. The relationship between recipient BMI and graft survival was also found to be better represented by converting BMI to a categorical variable, in accordance with the World Health Organisation (WHO) BMI classifications.¹⁷ Potential interactions between all variables were tested, none were significant. The proportional hazards assumption was found to be satisfied for all variables after checking log cumulative hazards plots and Schoenfeld residuals. Frailty models were used to check for inter-center variation by using the likelihood ratio test to assess the change in -2LogL after inclusion of transplant center as a random effect. The adjusted risk difference (ARD) was calculated using methods described by Laubender et al.¹⁸ The ARD describes the absolute effect of the comorbidity risk factor on survival probabilities after adjustment for covariates in the multivariable model. Standard errors of the ARD were derived from bootstrap methods using 1000 resamples of the data. Patients with missing data were excluded, the extent of

missing data is shown in Table S2 (SDC, <http://links.lww.com/TP/B802>). Sensitivity analyses were conducted to test robustness of the results; each model was adjusted for a risk score developed from UK Transplant Registry data for kidney transplants performed in the 5 years prior to the study recruitment period (2006 - 2011), rather than adjusting for individual confounding factors. This minimised the number of degrees of freedom in the models, and enabled checking for any missed comorbidity effects. All analyses were conducted using SAS® 9.4 (SAS Institute Inc, Cary, USA).

Ethics approval

East of England Research Ethics Committee (reference number 11/EE/0120).

Results

Baseline characteristics

Characteristics of the DDKT (n=1288) and LDKT (n=812) recipients, donors and transplants are shown in Table 1. These were consistent with UK Transplant Registry data for the study recruitment period.^{19, 20} The demographics of recruited versus excluded patients were compared (Table S3, SDC, <http://links.lww.com/TP/B802>). There was a higher proportion of White patients in the recruited group compared with the excluded group, however there were no significant differences in age group, gender or type of transplant. Table 2 shows the prevalence of comorbidity in the study cohort at the time of transplantation. DDKT recipients had significantly higher rates of diabetes (16.0% vs 10.3%, p=0.0002), IHD (9.8% vs 7.0%, p=0.029), HF (3.1% vs 1.6%, p=0.033), CVD (5.8% vs 3.1%, p=0.004) and PVD (3.3% vs 1.7%, p=0.027) compared with LDKT recipients.

DDKT recipients

Transplant survival

Overall, there were 134 “transplant failures” (85 graft failures and 49 patient deaths). The Kaplan-Meier estimate for two-year transplant survival was 89.4% (95% confidence interval

[CI] 87.6, 91.0). After adjustment for relevant factors in the multivariable Cox regression model, HF (HR 2.39, 95% CI 1.30, 4.37, $p=0.005$) and CVD (HR 2.33, 95% CI 1.40, 3.88, $p=0.001$) were associated with a significant increase in the risk of transplant failure (Table 3). There was no significant inter-center variation in transplant survival when including transplant center as a random effect in the model (difference in $-2\text{LogL}=0.02$, degrees of freedom [df]=1, $p=0.885$). For HF, the ARD was 0.117 (standard error [SE] 0.052) (i.e. patients with heart failure had an 11.7% increased risk of transplant failure within 2 years compared to those without heart failure, after adjustment for all other factors in the multivariable model). For CVD, the ARD was 0.101 (SE 0.043). The effect of adding DGF to the final model is shown in Table S4 (SDC, <http://links.lww.com/TP/B802>).

a) Graft survival

At two years, there were 85 graft failures, and the Kaplan-Meier estimate of graft survival was 93.2% (95% CI 91.7, 94.5). Multivariable Cox regression modelling showed PVD (HR 3.04, 95% CI 1.37, 6.74, $p=0.006$) and obesity (BMI ≥ 30.0) (HR 2.27, 95% CI 1.27, 4.06, $p=0.006$, compared with normal BMI 18.5 – 24.9) to be independent risk factors for graft loss (Table 3). The obesity variable was explored further in the model by dividing it into class I and class II and above (BMI 30.0 – 34.9 and ≥ 35.0 respectively) (Table S5, SDC, <http://links.lww.com/TP/B802>). There were too few patients with obesity class III (BMI ≥ 40.0) ($n=7$) to include this as a separate category. There was no significant variation in the risk of graft failure for the different classes of obesity, therefore the broader category of obesity (BMI ≥ 30.0) was retained in the main model (Table 3). No center effect on graft survival was found when modelling center as a random effect (difference in $-2\text{LogL}=0.23$, $df=1$, $p=0.632$). Among patients with PVD, the risk of graft failure was highest in the first ten days following transplantation, as demonstrated by the initial steep drop in the survival curve before the more gradual decline (Figure 2A); 85.7% graft failures in the PVD group occurred

during this early postoperative period, compared with 26.9% among patients without PVD. In contrast, the impact of obesity on graft survival followed a more gradual decline over the two-year period (Figure 2B). Unadjusted two-year graft survival estimates for patients with and without PVD and obesity are shown in Table 4. The ARD for PVD was 0.104 (SE 0.058) and for obesity was 0.060 (SE 0.029). The incidence of delayed graft function was 31.1% for all patients, 48.7% for patients with PVD and 39.1% for patients with obesity. Adding DGF to the final model resulted in a reduction in the effect of PVD (Table S4, SDC, <http://links.lww.com/TP/B802>). The cause of graft failure for all patients as well as patients with PVD and obesity in the DDKT cohort is shown in Table 5.

b) Patient survival

There were 56 patient deaths, of which 49 were deaths with a functioning graft. The two-year Kaplan-Meier survival estimate was 95.4% (95% CI 94.1, 96.5). The comorbidities significantly associated with inferior patient survival in the multivariable model included HF (HR 3.77, 95% CI 1.79, 7.95, $p=0.0005$), CVD (HR 3.45, 95% CI 1.72, 6.92, $p=0.0005$) and CLD (HR 4.36, 95% CI 1.29, 14.71, $p=0.018$) (Table 3). There were no significant center differences in patient survival (difference in $-2\text{LogL}=0.01$, $df=1$, $p=0.925$). Among patients with HF and CVD, just over half of patient deaths occurred in the second year after transplantation (55.6% and 58.3% respectively), while 100% of deaths among patients with CLD occurred within the first year post transplantation. This is demonstrated by the survival curves in Figures 3A, 3B and 3C. Unadjusted 2-year patient survival estimates for patients with and without HF, CVD and CLD are shown in Table 6. For HF, CVD and CLD the ARD was 0.159 (SE 0.057), 0.041 (SE 0.027) and 0.056 (SE 0.091) respectively. The effect of adding DGF to the final model is shown in Table S4 (SDC, <http://links.lww.com/TP/B802>).

LDKT recipients

In the LDKT cohort it was only possible to model transplant survival, as the smaller number of recipients and outcome events prevented meaningful analysis of separate graft and patient survival models. There were 42 “transplant failures” (26 graft failures and 16 patient deaths). The Kaplan-Meier estimate for transplant survival at 2 years was 94.7% (95% CI 92.9, 96.0). The multivariable model demonstrated significantly higher risk of transplant failure for HF (HR 3.83, 95% CI 1.15, 12.81, $p=0.029$) and diabetes (HR 2.23, 95% CI 1.03, 4.81, $p=0.042$) (Table 7). There was no significant center effect on LDKT transplant survival (difference in $-2\text{LogL}=0.11$, $df=1$, $p=0.741$). The ARD for HF was 0.121 (SE 0.099) and for diabetes was 0.056 (SE 0.036).

Sensitivity analyses

Each multivariable model was checked by adjusting for a risk score (Boxes S1, S2, S3 and S4) rather than entering the confounding factors individually into the model (Tables S6, S7, S8 and S9, SDC, <http://links.lww.com/TP/B802>). No additional comorbidities were identified as significant, and hazard ratios were very similar to the original models, confirming the reliability of the results.

Discussion

In this national observational study, we have collected data prospectively on a wide range of comorbid conditions and identified those that predict poorer survival outcomes after kidney transplantation. Among DDKT recipients, PVD and obesity were associated with a two- to three-fold increased risk of graft failure within two years of transplantation, while the risk of death was three- to four-fold higher with HF, CVD and CLD. For LDKT recipients, HF and diabetes were associated with significant detrimental effects on overall transplant survival, but longer follow up is required to determine the separate effects on graft and patient survival.

Among DDKT recipients, a history of PVD increased the risk of graft failure by 10.4% after adjusting for confounding factors, with the majority of graft failures occurring in the early postoperative period. PVD is typically diagnosed clinically by measuring the ankle-brachial pressure index (ABPI), and our results are in agreement with a US study of 819 patients which reported a 2.77 times increased risk of graft failure for patients with a low ABPI (<0.9).²¹ Preexisting PVD affecting the aorta or iliac arteries may complicate implantation of the kidney graft, resulting in difficult anastomoses, cholesterol emboli or hypoperfusion of the graft, and subsequent failure in the early postoperative period.^{22, 23} Our data showed a high incidence of technical operative issues as the cause of graft failure among PVD patients (42.9%). We also found that the addition of DGF to the regression model for DDKT graft survival reduced the effect of PVD and is thus a potential mediator of this effect. Despite being a high risk group, patients with PVD still derive a significant survival benefit from transplantation compared with dialysis.^{24, 25} As such, a history of PVD should not preclude transplantation, but given the high risk of early complications, appropriate preoperative planning and informed consent of patients is crucial.

Obesity is an ongoing topic of controversy with regard to patient suitability for kidney transplantation. Some centers do not exclude patients with obesity, while others restrict access to the waiting list at specific BMI thresholds, which may differ considerably between centers, and even between clinicians within the same center.²⁶ Despite conflicting outcomes from early single-center studies, more recent meta-analyses have confirmed the detrimental effect of obesity on graft survival.²⁷⁻³⁰ Our results are in keeping with this evidence; with obesity conferring a 6% increased risk of graft failure among DDKT recipients. The mechanisms for this are unclear. There was a high incidence of acute rejection as a cause of graft failure among obese patients (44%) and this could be a potential cause for the higher risk of graft failure associated with obesity. Difficulties in achieving and maintaining the

narrow therapeutic target concentrations of immunosuppressive drugs in obese patients have previously been reported.³¹

We found that HF was associated with a 15.9% higher risk of mortality after DDKT and 12.1% higher risk of transplant failure after LDKT. We acknowledge that in patients on dialysis, it can be difficult to make a clear distinction between HF and fluid overload; however, our findings demonstrate that a diagnosis of heart failure in the patient's record predicts poorer survival, irrespective of how the diagnosis was made or the exact pathophysiology. It is also noteworthy that although HF was identified as a significant risk factor, no effect was observed for IHD. Our findings concur with the results of a US study which found that pretransplant impaired left ventricular systolic function (on single photon emission computed tomography (SPECT)) was associated with a significantly higher risk of both cardiac mortality and all-cause mortality after kidney transplantation, while cardiac ischaemia (on SPECT) was not.³² Our findings suggest that either IHD does not increase the risk of death within two years post transplantation, or that current risk stratification of patients with IHD in the UK is effective.

CVD was associated with a 4.1% elevated risk of death among DDKT recipients. It is known that patients with ESRD have more severe carotid atherosclerosis than the general population and are at substantially greater risk of stroke.³³⁻³⁵ A large US registry analysis demonstrated that transplantation reduced the risk of cerebrovascular events from 11.8% to 6.8% compared to patients remaining on the waiting list.³⁶ However, previous CVD remains a strong risk factor for further post transplantation events and mortality.^{35, 37, 38} Post transplantation cerebrovascular events are associated with high mortality,³⁹ which is worse for haemorrhagic strokes (48%) compared with ischaemic strokes (6%).³⁸ In a prospective randomised controlled trial including 1652 kidney transplant recipients (ALERT trial), the use of Fluvastatin did not reduce the incidence of cerebrovascular events or mortality.³⁸ Further

trials are needed to assess the ability of therapies to reduce the risk of further cerebrovascular events and mortality in this high risk population.

CLD was independently associated with 5.6% increased risk of mortality within two years of DDKT. There is a paucity of published research regarding CLD in kidney transplant outcomes. Previous studies have focussed on the role of hepatitis B and C related liver disease as predictors of increased mortality after kidney transplantation.⁴⁰⁻⁴² The present study is the first to demonstrate that CLD of any aetiology leads to reduced survival after DDKT. Further research is required to understand the underlying mechanisms.

Interestingly, a diagnosis of diabetes was identified as a risk factor for transplant failure among LDKT recipients, but not for DDKT recipients. The reason for this finding is unclear. Diabetes is a well-recognised risk factor for mortality after transplantation, primarily due to elevated cardiovascular risk.⁴³ It may be that this cardiovascular risk was actually accounted for by other comorbidity variables in the models for DDKT recipients, while in the LDKT cohort with a significantly lower prevalence of other comorbidities, diabetes may have served as more general marker of poorer outcomes. A recent large population cohort study in Australia and New Zealand demonstrated that patients with Type 2 diabetes had significantly poorer survival after kidney transplantation, with the highest risk being among younger patients under the age of 40 years.⁴⁴ In our study the LDKT population was significantly younger than the DDKT population and this may explain why diabetes was a significant risk factor in this population. The 5.6% higher risk of transplant failure among patients with diabetes (and 12% higher risk for patients with heart failure discussed previously) must be given due consideration in the context of LDKT, given the potential implications for both the recipient as well as the live donor.

A major strength of the present study is that it is a prospective and comprehensive analysis of a large cohort of transplant recipients from all UK transplant centers. The cohort included a large proportion of the national adult transplant population with a minimal amount of missing data, which adds to the reliability of the study. There are a number of limitations to this study. First, for practical reasons we used relatively broad definitions for each comorbidity and were unable to distinguish between differing levels of severity or duration for each condition. All comorbidity data were collected at the time of transplantation when patients were recruited to the study. Therefore, we were unable to assess the progression or improvement of each condition after transplantation, and whether this impacted on outcomes. Secondly, it should be noted that the study population is largely of white ethnicity and thus conclusions with respect to other ethnic groups may be less certain. Thirdly, due to the favourable survival outcomes of LDKT recipients, we were only able to analyse the composite outcome of transplant survival in this cohort, as there were too few events for separate analysis of graft and patient survival. Transplant survival (also known as graft survival not censored for death) is a commonly analysed end-point in the transplant literature, as it demonstrates the overall success of a transplant.^{45, 46} However, in the DDKT analysis we found that this method masked the importance of several comorbidity risk factors that were found to be significant when analysing graft and patient survival separately. Therefore, it is important that we carry out separate graft and patient survival analyses in the LDKT cohort after longer follow-up time. Finally, the results from this study describe associations and no causation can be inferred.

This study quantifies the risks associated with specific comorbid conditions in the context of kidney transplantation. The findings can be utilised in everyday clinical practice to fully inform patients of their individual risks and outcomes, to inform future wait-listing and

allocation policy and also to guide further research into improving the outcomes of patients with specific comorbidities.

Acknowledgments

The authors are very grateful to the research nurses involved in the data collection, to the patient representatives involved in the study development and to the patients who participated in the study.

ACCEPTED

References

1. Hsu CY, Iribarren C, McCulloch CE, et al. Risk factors for end-stage renal disease: 25-year follow-up. *Arch Intern Med.* 2009;169(4):342-350.
2. Longenecker JC, Coresh J, Powe NR, et al. Traditional cardiovascular disease risk factors in dialysis patients compared with the general population: the CHOICE Study. *J Am Soc Nephrol.* 2002;13(7):1918-1927.
3. Alani H, Tamimi A, Tamimi N. Cardiovascular co-morbidity in chronic kidney disease: Current knowledge and future research needs. *World J Nephrol.* 2014;3(4):156-168.
4. NHS Blood and Transplant. UK Transplant Registry. Organ Donation and Transplantation Annual Activity Report. Available at <https://www.odt.nhs.uk/statistics-and-reports/annual-activity-report>. Accessed March 7, 2019.
5. Wu C, Evans I, Joseph R, et al. Comorbid conditions in kidney transplantation: association with graft and patient survival. *J Am Soc Nephrol.* 2005;16(11):3437-3444.
6. Chang SH, Russ GR, Chadban SJ, et al. Trends in kidney transplantation in Australia and New Zealand, 1993-2004. *Transplantation.* 2007;84(5):611-618.
7. Weinhandl ED, Snyder JJ, Israni AK, et al. Effect of Comorbidity Adjustment on CMS Criteria for Kidney Transplant Center Performance. *Am J Transplant.* 2009;9(3):506-516.
8. Jassal SV, Schaubel DE, Fenton SSA. Baseline comorbidity in kidney transplant recipients: a comparison of comorbidity indices. *Am J Kidney Dis.* 2005;46(1):136-142.

9. Fabbian F, De Giorgi A, Manfredini F, et al. Impact of comorbidity on outcome in kidney transplant recipients: a retrospective study in Italy. *Intern Emerg Med*. 2016;11(6):825-832.
10. Grosso G, Corona D, Mistretta A, et al. Predictive value of the Charlson comorbidity index in kidney transplantation. *Transplant Proc*. 2012;44(7):1859-1863.
11. Kotwal S, Webster AC, Cass A, et al. Comorbidity recording and predictive power of comorbidities in the Australia and New Zealand dialysis and transplant registry compared with administrative data: 2000-2010. *Nephrology (Carlton)*. 2016;21(11):930-937.
12. Machnicki G, Pinsky B, Takemoto S, et al. Predictive Ability of Pretransplant Comorbidities to Predict Long-Term Graft Loss and Death. *Am J Transplant*. 2009;9(3):494-505.
13. Farrugia D, Cheshire J, Begaj I, et al. Death within the first year after kidney transplantation--an observational cohort study. *Transpl Int*. 2014;27(3):262-270.
14. Oniscu GC, Ramanan R, Wu D, et al. Access to Transplantation and Transplant Outcome Measures (ATTOM): study protocol of a UK wide, in-depth, prospective cohort analysis. *BMJ Open*. 2016;6:e010377.
15. Venkat-Raman G, Tomson CR, Gao Y, et al. New primary renal diagnosis codes for the ERA-EDTA. *Nephrol Dial Transplant*. 2012;27(12):4414-4419.
16. Johnson RJ, Fuggle SV, O'Neill J, et al. Factors influencing outcome after deceased heart beating donor kidney transplantation in the United Kingdom: an evidence base for a new national kidney allocation policy. *Transplantation*. 2010;89(4):379-386.
17. World Health Organization. BMI Classification. Available at <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>. Accessed August 7, 2019.

18. Laubender RP, Bender R. Estimating adjusted risk difference (RD) and number needed to treat (NNT) measures in the Cox regression model. *Stat Med.* 2010;29(7-8):851-859.
19. NHS Blood and Transplant. Organ Donation and Transplantation Activity Report 2012/13. Available at https://nhsbtmediaservices.blob.core.windows.net/organ-donation-assets/pdfs/activity_report_2012_13.pdf. Accessed January 30, 2018.
20. NHS Blood and Transplant. Organ Donation and Transplantation Activity Report 2011/2012. Available at https://nhsbtmediaservices.blob.core.windows.net/organ-donation-assets/pdfs/activity_report_2011_12.pdf. Accessed January 30, 2018.
21. Patel SI, Chakkera HA, Wennberg PW, et al. Peripheral arterial disease preoperatively may predict graft failure and mortality in kidney transplant recipients. *Vasc Med.* 2017;22(3):225-230.
22. Shishehbor MH, Aksut B, Poggio E, et al. Presence of peripheral artery disease in renal transplant outcomes - Don't throw the baby out with the bath water. *Vasc Med.* 2017;22(3):231-233.
23. Droupy S, Eschwège P, Hammoudi Y, et al. Consequences of iliac arterial atheroma on renal transplantation. *J Urol.* 2006;175(3 Pt 1):1036-1039.
24. Cassuto J, Babu S, Laskowski I. The survival benefit of kidney transplantation in the setting of combined peripheral arterial disease and end-stage renal failure. *Clin Transplant.* 2016;30(5):545-555.
25. Gill JS, Tonelli M, Johnson N, et al. The impact of waiting time and comorbid conditions on the survival benefit of kidney transplantation. *Kidney Int.* 2005;68(5):2345-2351.
26. Akolekar D, Oniscu GC, Forsythe JL. Variations in the assessment practice for renal transplantation across the United Kingdom. *Transplantation.* 2008;85(3):407-410.

27. Gore JL, Pham PT, Danovitch GM, et al. Obesity and outcome following renal transplantation. *Am J Transplant*. 2006;6(2):357-363.
28. Meier-Kriesche HU, Arndorfer JA, Kaplan B. The impact of body mass index on renal transplant outcomes: a significant independent risk factor for graft failure and patient death. *Transplantation*. 2002;73(1):70-74.
29. Hill CJ, Courtney AE, Cardwell CR, et al. Recipient obesity and outcomes after kidney transplantation: a systematic review and meta-analysis. *Nephrol Dial Transplant*. 2015;30(8):1403-1411.
30. Lafranca JA, IJermans JN, Betjes MG, et al. Body mass index and outcome in renal transplant recipients: a systematic review and meta-analysis. *BMC Med*. 2015;13:111.
31. Hortal L, Fernández A, Losada A, et al. Study of the cyclosporine concentration at 2 hours in stable renal transplant patients and relation to body mass index. *Transplant Proc*. 2001;33(7-8):3110-3111.
32. Siedlecki A, Foushee M, Curtis JJ, et al. The impact of left ventricular systolic dysfunction on survival after renal transplantation. *Transplantation*. 2007;84(12):1610-1617.
33. Seliger SL, Gillen DL, Longstreth WT, Jr., et al. Elevated risk of stroke among patients with end-stage renal disease. *Kidney Int*. 2003;64(2):603-609.
34. Kennedy R, Case C, Fathi R, et al. Does renal failure cause an atherosclerotic milieu in patients with end-stage renal disease? *Am J Med*. 2001;110(3):198-204.
35. Findlay MD, Thomson PC, MacIsaac R, et al. Risk factors and outcome of stroke in renal transplant recipients. *Clin Transplant*. 2016;30(8):918-924.
36. Lentine KL, Rey LAR, Kolli S, et al. Variations in the Risk for Cerebrovascular Events after Kidney Transplant Compared with Experience on the Waiting List and after Graft Failure. *Clin J Am Soc Nephrol*. 2008;3(4):1090-1101.

37. de Mattos AM, Prather J, Olyaei AJ, et al. Cardiovascular events following renal transplantation: role of traditional and transplant-specific risk factors. *Kidney Int.* 2006;70(4):757-764.
38. Abedini S, Holme I, Fellström B, et al. Cerebrovascular events in renal transplant recipients. *Transplantation.* 2009;87(1):112-117.
39. Oliveras A, Roquer J, Puig JM, et al. Stroke in renal transplant recipients: epidemiology, predictive risk factors and outcome. *Clin Transplant.* 2003;17(1):1-8.
40. Yu T-M, Lin C-C, Shu K-H, et al. Increased risk of hepatic complications in kidney transplantation with chronic virus hepatitis infection: A nationwide population-based cohort study. *Sci Rep.* 2016;6:21312.
41. Fabrizi F, Martin P, Dixit V, et al. Hepatitis C virus antibody status and survival after renal transplantation: meta-analysis of observational studies. *Am J Transplant.* 2005;5(6):1452-1461.
42. Fabrizi F, Dixit V, Martin P, et al. Hepatitis B and survival after renal transplant: meta-analysis of observational studies. *J Viral Hepat.* 2014;21(8):542-550.
43. Cosio FG, Hickson LJ, Griffin MD, et al. Patient survival and cardiovascular risk after kidney transplantation: the challenge of diabetes. *Am J Transplant.* 2008;8(3):593-599.
44. Lim WH, Wong G, Pilmore HL, et al. Long-term outcomes of kidney transplantation in people with type 2 diabetes: a population cohort study. *Lancet Diabetes Endocrinol.* 2017;5(1):26-33.
45. EBPG Expert Group on Renal Transplantation. European Best Practice Guidelines for Renal Transplantation. Section IV: Long-term management of the transplant recipient. IV.13 Analysis of patient and graft survival. *Nephrol Dial Transplant.* 2002;17 Suppl 4:60-67.

46. OPTN/SRTR 2016 Annual Data Report: Preface. *Am J Transplant*. 2018;18 Suppl 1:1-9.

ACCEPTED

Figures

Figure 1. Study population and analyses. Patients were recruited from all 23 UK renal transplant centers. Recruitment took place over a 12-month period in each center, between 1st November 2011 and 31st March 2013

Figure 2. Kaplan-Meier curves for 2-year graft survival of deceased-donor kidney transplants

A. Peripheral vascular disease (PVD). B. Body mass index (BMI)

Figure 3. Kaplan-Meier curves for 2-year patient survival after deceased-donor kidney transplantation

A. Heart failure (HF). B. Cerebrovascular disease (CVD). C. Chronic liver disease (CLD)

ACCEPTED

Tables

Table 1. Characteristics of the study cohort

	DDKT recipients n=1288	LDKT recipients n=812	p-value
Recipient variables			
Recipient age, years (median, IQR)	54 (44 - 63)	46 (34 - 56)	<0.0001
Recipient age group, years (n, %)			<0.0001
18 – 34	132 (10.3)	229 (28.2)	
35 – 49	369 (28.7)	263 (32.4)	
50 – 64	543 (42.2)	252 (31.0)	
65 – 75	244 (18.9)	68 (8.4)	
Recipient gender (n, %)			0.267
Male	824 (64.0)	500 (61.6)	
Female	464 (36.0)	312 (38.4)	
Recipient ethnicity (n, %)			0.0002
White	1023 (79.7)	707 (87.1)	
Asian	140 (10.9)	62 (7.6)	
Black	96 (7.5)	35 (4.3)	
Other	25 (2.0)	8 (1.0)	
Primary renal disease (n, %)			<0.0001
Polycystic kidney disease	219 (17.0)	112 (13.9)	
Diabetic nephropathy	134 (10.4)	48 (5.9)	
Glomerulonephritis	320 (24.9)	232 (28.7)	
Pyelonephritis	138 (10.7)	128 (15.8)	
Hypertensive nephropathy	89 (6.9)	37 (4.6)	
Renal vascular disease	29 (2.3)	9 (1.1)	
Other	163 (12.7)	85 (10.5)	
Uncertain	194 (15.1)	157 (19.4)	
Time on dialysis (n, %)			<0.0001
Preemptive	137 (10.6)	279 (34.4)	

0 - 1 year	160 (12.4)	198 (24.4)	
1 - 3 years	366 (28.4)	185 (22.8)	
3 - 5 years	295 (22.9)	78 (9.6)	
> 5 years	330 (25.6)	72 (8.9)	
Previous transplant (n, %)	165 (12.9)	117 (14.5)	0.297
Highly sensitised, cRF \geq 85% (n, %)	126 (9.8)	95 (11.7)	0.163
Smoking status (n, %)			0.702
Nonsmoker	137 (11.7)	78 (10.7)	
Ex-smoker	325 (27.7)	212 (29.2)	
Smoker	710 (60.6)	437 (60.1)	

Donor variables

Donor age, years (median, IQR)	54 (43 - 64)	48 (39 - 57)	<0.0001
Donor age group, years (n, %)			<0.0001
<18	31 (2.4)	0	
18 - 34	160 (12.4)	143 (17.6)	
35 - 49	303 (23.5)	298 (36.7)	
50 - 64	512 (39.8)	308 (37.9)	
65 - 75	234 (18.2)	61 (7.5)	
>75	48 (3.7)	2 (0.3)	
Donor gender (n, %)			0.001
Male	696 (54.0)	379 (46.7)	
Female	592 (46.0)	432 (53.3)	
Donor ethnicity (n, %)			<0.0001
White	1208 (95.2)	720 (88.7)	
Asian	21 (1.7)	52 (6.4)	
Black	23 (1.8)	28 (3.5)	
Other	17 (1.3)	12 (1.5)	
Donor BMI, kg/m ² (n, %)			<0.0001
Underweight (<18.5)	0	0	
Normal (18.5 - 24.9)	463 (37.3)	254 (32.9)	
Overweight (25.0 - 29.9)	494 (39.7)	390 (50.5)	
Obese (\geq 30.0)	286 (23.0)	128 (16.6)	

Transplant variables

HLA MM level (n, %)			<0.0001
1	155 (12.0)	91 (11.2)	
2	355 (27.6)	105 (12.9)	
3	679 (52.7)	360 (44.3)	
4	99 (7.7)	256 (31.5)	
CIT, hours (median, IQR)	14.5 (11.4 - 17.3)	3.3 (2.4 - 4.1)	<0.0001
Delayed graft function (n, %)	378 (31.1)	30 (3.9)	<0.0001

DDKT; deceased-donor kidney transplant, LDKT; living-donor kidney transplant, IQR; interquartile range, cRF; calculated reaction frequency, BMI; body mass index, HLA MM; human leukocyte antigen mismatch, CIT; cold ischaemia time. Data are missing for some participants and excluded from percentage calculations. Number of missing data are shown in Table S2 (SDC, <http://links.lww.com/TP/B802>).

ACCEPTED

Table 2. Prevalence of recipient comorbidity

	DDKT recipients n=1288	LDKT recipients n=812	p-value
Diabetes (n, %)	205 (16.0)	83 (10.3)	0.0002
Ischaemic heart disease (n, %)	126 (9.8)	57 (7.0)	0.029
Heart failure (n, %)	40 (3.1)	13 (1.6)	0.033
Atrial fibrillation (n, %)	25 (1.9)	12 (1.5)	0.434
Cardiac valve replacement (n, %)	10 (0.8)	8 (1.0)	0.609
Pacemaker (n, %)	10 (0.8)	5 (0.6)	0.673
Cerebrovascular disease (n, %)	75 (5.8)	25 (3.1)	0.004
Peripheral vascular disease (n, %)	43 (3.3)	14 (1.7)	0.027
Abdominal aortic aneurysm (n, %)	4 (0.3)	2 (0.3)	0.790
Chronic respiratory disease (n, %)	108 (8.4)	59 (7.3)	0.359
Chronic liver disease (n, %)	25 (1.9)	14 (1.7)	0.722
Blood borne viruses (n, %)	38 (3.0)	13 (1.6)	0.051
Malignancy (n, %)	93 (7.2)	49 (6.1)	0.294
Mental illness (n, %)	75 (5.8)	41 (5.1)	0.453
BMI, kg/m ² (n, %)			0.121
Underweight (<18.5)	26 (2.1)	23 (3.0)	
Normal (18.5 - 24.9)	461 (37.5)	312 (40.8)	
Overweight (25.0 - 29.9)	462 (37.6)	282 (36.9)	
Obese (≥30.0)	281 (22.9)	147 (19.2)	
Number of comorbidities (n, %)			0.002
0	573 (46.7)	414 (54.4)	
1 - 2	579 (47.2)	316 (41.5)	
≥3	74 (6.0)	31 (4.1)	

DDKT; deceased-donor kidney transplant, LDKT; living-donor kidney

transplant, BMI; body mass index. Data are missing for some participants and excluded from percentage calculations. Number of missing data are shown in Table S2.

Table 3. Cox regression analysis for impact of comorbidity on 2-year survival outcomes of deceased donor kidney transplants

Variables	Transplant survival model		Graft survival model		Patient survival model	
	HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
Recipient comorbidity						
Heart failure	2.39 (1.30, 4.37)	0.005	-	-	3.77 (1.79, 7.95)	0.0005
Cerebrovascular disease	2.33 (1.40, 3.88)	0.001	-	-	3.45 (1.72, 6.92)	0.0005
Chronic liver disease	-	-	-	-	4.36 (1.29, 14.71)	0.018
Peripheral vascular disease	-	-	3.04 (1.37, 6.74)	0.006	-	-
BMI, kg/m²						
Underweight (<18.5)	-	-	0.86 (0.11, 6.49)	0.885	-	-
Normal (18.5 - 24.9)	-	-	1 (reference)	-	-	-
Overweight (25.0 - 29.9)	-	-	1.48 (0.84, 2.61)	0.180	-	-
Obese (≥30.0)	-	-	2.27 (1.27, 4.06)	0.006	-	-
Other variables						
Time on dialysis (years)						
< 3	1 (reference)	-	1 (reference)	-	1 (reference)	-
≥ 3	2.08 (1.41, 3.08)	0.0002	1.84 (1.11, 3.04)	0.018	2.47 (1.36, 4.50)	0.003
Recipient age (per 10 years)	1.10 (0.92, 1.30)	0.290	0.84 (0.68, 1.05)	0.128	1.67 (1.23, 2.25)	0.0009
Recipient ethnicity						
White	1 (reference)	-	1 (reference)	-	-	-
Asian	0.67 (0.35, 1.29)	0.228	0.76 (0.35, 1.69)	0.504	-	-
Black	1.23 (0.68, 2.21)	0.495	1.52 (0.77, 3.02)	0.228	-	-

Other	0.37 (0.05, 2.63)	0.317	0.62 (0.08, 4.53)	0.636	-	-
Highly sensitised (cRF \geq 85%)	1.47 (0.87, 2.47)	0.153	2.22 (1.18, 4.19)	0.014	-	-
Donor age (per 10 years)	1.14 (0.99, 1.31)	0.066	1.23 (1.02, 1.48)	0.028	1.11 (0.89, 1.39)	0.349
HLA MM level						
1	1 (reference)		1 (reference)		1 (reference)	
2	1.18 (0.62, 2.27)	0.612	2.94 (1.08, 7.98)	0.035	0.40 (0.16, 1.01)	0.052
3	1.05 (0.57, 1.94)	0.866	2.25 (0.85, 5.93)	0.103	0.46 (0.21, 1.01)	0.051
4	1.25 (0.53, 2.93)	0.612	2.78 (0.81, 9.59)	0.106	0.66 (0.22, 2.01)	0.467
Cold ischaemia time (per hour)	1.04 (1.01, 1.08)	0.028	1.01 (0.97, 1.06)	0.568	1.04 (0.99, 1.10)	0.118

HR; hazard ratio, CI; confidence interval, BMI; body mass index, cRF; calculated reaction frequency, HLA MM; human leukocyte antigen mismatch.

Table 4. Kaplan-Meier estimates for 2-year graft survival of deceased-donor kidney transplants

Comorbidity	Survival (95% CI)	p-value
Peripheral vascular disease		0.006
No	93.6 (92.0, 94.8)	
Yes	83.5 (68.5, 91.8)	
BMI, kg/m ²		0.012
Normal (18.5 - 24.9)	95.2 (92.7, 96.8)	
Obese (≥ 30.0)	90.1 (85.9, 93.1)	

p-value is for log-rank test.

ACCEPTED

Table 5. Cause of graft failure among DDKT cohort

Cause of graft failure	All patients	Obese patients	PVD patients
Acute rejection	26 (34.2%)	11 (44.0%)	1 (14.3%)
Vascular thrombosis	6 (7.9%)	0 (0%)	1 (14.3%)
Technical operative issues	9 (11.8%)	3 (12.0%)	3 (42.9%)
Nonviable kidney	9 (11.8%)	3 (12.0%)	1 (14.3%)
Infection	1 (1.3%)	0 (0%)	0 (0%)
Recurrent primary renal disease	4 (5.3%)	0 (0%)	0 (0%)
Other	21 (27.6%)	8 (32.0%)	1 (14.3%)

ACCEPTED

Table 6. Kaplan-Meier estimates for 2-year patient survival after deceased-donor kidney transplantation

Comorbidity	Survival (95% CI)	p-value
Heart failure		<0.0001
No	96.0 (94.8, 97.0)	
Yes	75.8 (58.5, 86.7)	
Cerebrovascular disease		<0.0001
No	96.2 (94.9, 97.1)	
Yes	82.7 (71.5, 89.8)	
Chronic liver disease		0.003
No	95.7 (94.3, 96.7)	
Yes	83.6 (62.0, 93.5)	

p-value is for log-rank test.

ACCEPTED

Table 7. Cox regression analysis for impact of comorbidity on 2-year transplant survival of living-donor kidney transplants

Variables	HR (95% CI)	p-value
Recipient comorbidity		
Heart failure	3.83 (1.15, 12.81)	0.029
Diabetes	2.23 (1.03, 4.81)	0.042
Other variables		
Time on dialysis (years)		
< 3	1 (reference)	
≥ 3	2.16 (1.13, 4.11)	0.019
Recipient age (per 10 years)	1.01 (0.80, 1.28)	0.926
Donor age (per 10 years)	1.03 (0.81, 1.31)	0.828
HLA MM level		
1	1 (reference)	
2	0.76 (0.23, 2.51)	0.657
3	0.74 (0.29, 1.86)	0.520
4	0.67 (0.25, 1.82)	0.428

HR; hazard ratio, CI; confidence interval, HLA MM; human leukocyte antigen mismatch.

Figure 1




Figure 2A





Figure 2b



ACCEPTED

Figure 3A






Figure 3B



ACCEPTED

Figure 3C



ACCEPTED