

1 **Navigating future uncertainty in marine protected area governance: Lessons from the**  
2 **Scottish MPA network**

3 Charlotte Rachael Hopkins<sup>a,\*</sup>, David Mark Bailey<sup>a</sup>, Tavis Potts<sup>b</sup>

4 <sup>a</sup> Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,  
5 Glasgow, G12 8QQ, UK

6 <sup>b</sup> Department of Geography and Environment, University of Aberdeen, AB24 3UF, UK

7 **Abstract**

8 As international pressure for marine protection has increased, Scotland has increased  
9 spatial protection through the development of a Marine Protected Area (MPA) network.  
10 Few MPA networks to date have included specific considerations of climate change in the  
11 design, monitoring or management of the network. The Scottish MPA network followed a  
12 feature-led approach to identify a series of MPAs across the Scottish marine area and  
13 incorporated the diverse views of many different stakeholders. This feature led approach  
14 has led to wide ranging opinions and understandings regarding the success of the MPA  
15 network. Translating ideas of success into a policy approach whilst also considering how  
16 climate change may affect these ideas of success is a complex challenge. This paper presents  
17 the results of a Delphi process that aimed to facilitate clear communication between  
18 academics, policy makers and stakeholders in order to identify specific climate change  
19 considerations applicable to the Scottish MPA network. This study engaged a group of  
20 academic and non-academic stakeholders to discuss potential options that could be  
21 translated into an operational process for management of the MPA network. The results of  
22 Delphi process discussion are presented with the output of a management matrix tool,  
23 which could aid in future decisions for MPA management under scenarios of climate  
24 change.

25 Key Words: climate change, Delphi technique, MPA management, marine protected area  
26 network, Scotland

## 27 **1. Introduction**

28 Marine ecosystems are facing a diverse range of threats, including climate change,  
29 prompting international efforts to safeguard marine biodiversity through the use of spatial  
30 management measures (Allison et al., 1998; Lubchenco et al., 2003; Chuenpagdee et al.,  
31 2013). Marine Protected Areas (MPAs) have been implemented as a conservation tool  
32 throughout the world, but their usefulness and effectiveness is strongly challenged by  
33 climate change (Harley et al., 2006; Andrello et al., 2015). Whilst MPAs cannot explicitly  
34 protect against climate change related disturbances (e.g. ocean acidification), MPAs can  
35 assist in sustaining biodiversity and ecosystem processes at regional and local scales (Levy  
36 and Ban, 2013). The reduction of other anthropogenic threats (e.g. overfishing) can  
37 minimise the synergistic impact of other stressors which may exacerbate detrimental  
38 changes to ecosystem health (Harley and Rogers-Bennett, 2004; Harley et al., 2006; Levy  
39 and Ban, 2013). The reduction of additional stressors could also contribute to increased  
40 ecosystem resilience in the face of climatic stress (see Bernhardt and Leslie, 2013).  
41 However, few MPA programmes have directly considered climate change in the design,  
42 management or monitoring of an MPA network (Hopkins et al., 2016a). Considering  
43 elements of design, management and monitoring that could enable an MPA network to  
44 perform effectively under scenarios of climate change, could also improve networks more  
45 generally.

46 Under international obligations, EU, UK and national targets (e.g. CBD, OSPAR), Scotland has  
47 developed an MPA network intended to protect marine biodiversity and contribute to the  
48 vision of a clean, healthy and productive marine environment (Scottish Government,  
49 2011a). The implementation of the Scottish MPA network has been a complex process  
50 requiring the consideration of stakeholder values and perceptions, scientific evidence and  
51 political factors (Hopkins et al., 2016b). There is a need to facilitate clear communication  
52 between academics, policy makers and stakeholders to progress MPA policy delivery and  
53 ensure decisions are jointly formed and therefore acceptable to multiple parties (Pollnac et  
54 al., 2010). The Scottish Nature Conservation MPA network consists of 30 MPAs designated  
55 in 2014: 17 MPAs under the Marine (Scotland) Act 2010 in Scottish territorial waters and 13  
56 MPAs under the Marine and Coastal Access Act 2009. Scottish Natural Heritage (SNH) and

57 the Joint Nature Conservation Committee (JNCC) submitted formal advice to parliament  
58 following a series of stakeholder workshops.

59 The Scottish MPA network (including other types of protected area designation) covers  
60 approximately 20% of the Scottish sea area. The Scottish MPA network is intended to  
61 contribute to an OSPAR ecologically coherent network and is part of the Scottish  
62 Government's three pillar approach to conservation, which includes spatial protection,  
63 wider seas measures and species-specific protection and management measures (Scottish  
64 Government, 2011a). Together, the three-pillar approach is intended to contribute to the  
65 achievement of Good Environmental Status (GES) under the Marine Strategy Framework  
66 Directive (MSFD). Therefore, it is important to assess the contribution that the MPA network  
67 makes towards protecting marine biodiversity and the delivery of GES. Furthermore, with  
68 increasing pressure from climate change on marine biodiversity, an effective MPA network  
69 will be crucial in providing climate change resilience. We define resilience here as the ability  
70 of an ecosystem to experience disturbance without substantial biological change (Holling,  
71 1973), a change that could result in an alternative state and loss of ecosystem function  
72 (Côté and Darling, 2010).

73 The Scottish MPA network was developed using a feature-based approach to site selection,  
74 whereby MPA sites were selected based on the “locations of habitats or species which are  
75 important, rare, threatened and/or representative of the range of features in the UK marine  
76 area” (Scottish Government, 2011b) termed Priority Marine Features (PMFs) (see Howson et  
77 al., 2012). It will be important to assess whether such a feature led approach is effective for  
78 selecting MPA sites that will remain resilient under climate change scenarios. Each Scottish  
79 MPA also has a Conservation Objective of either “conserve” or “recover” tying MPA  
80 management measures to the feature for which each site was designated. These objectives  
81 are vague and therefore difficult to measure under climate change scenarios where it may  
82 become unfeasible to achieve such an objective (Cliquet et al., 2009).

83 The aim of this study was to facilitate the identification of high level management options  
84 for Scottish MPA network in the context of potential climate change scenarios prior to the  
85 development of site specific management options. There are few examples of high level

86 MPA decision making, for example, under what circumstances should a new MPA be  
87 designated, or an MPA that is no longer effective or successful, de-designated. This study  
88 aimed to explore these options in the context of climate change, answering the following  
89 research questions:

90 Are there differences in the perceptions of MPA success between different stakeholder  
91 groups?

92 How can we effectively protect marine ecosystems under climate change scenarios?

93 What are feasible options for including climate change specific management and monitoring  
94 strategies?

95

## 96 **2. Materials and methods**

97 A Delphi method was devised in this study to elicit perceptions and options for climate  
98 change management scenarios. The Delphi method is becoming more frequently applied to  
99 conservation and biodiversity management issues due to their complex nature, involving a  
100 range of stakeholders and trade-offs (Hess and King, 2002; O'Neill et al., 2008; Gobbi et al.,  
101 2012). The Delphi method is a flexible methodology suitable for complex policy problems,  
102 particularly where there is significant uncertainty, lack of historical precedent and especially  
103 in situations where information is limited or conflicting (Mukherjee et al., 2015). Questions  
104 are posed and responses to those questions exchanged usually anonymously with other  
105 participations via a process facilitator and is an effective way for a group to deal with a  
106 complex issue either reaching consensus or identifying convergence of opinion (Linstone  
107 and Turoff, 2002; Hsu and Sandford, 2007). The benefit of the reflective deliberation of the  
108 Delphi method may also be the development of more creative solutions by groups of people  
109 (Reed, 2008). The Delphi method employed here did not seek consensus, seeking instead an  
110 improvement in understanding and clarification of the issue, therefore sharing similarities  
111 with Policy Delphi. As Rowe and Wright (2011) suggest, the most interesting and important  
112 issues often emerge where consensus is not evident.

113 MPA processes involve a complex range of stakeholders from various economic, social and  
114 environmental interest groups. As such, the panel was carefully selected to apply their  
115 knowledge and experience to the study issue and to reflect the diversity of stakeholders  
116 involved in the MPA process. Following Glass et al. (2013) a stakeholder map was created to  
117 identify a matrix of organisations and stakeholder interest groups related to the Scottish  
118 MPA process. Potential participants were selected if they met one or more of the following  
119 criteria: active role in the Scottish MPA process, relevant experience in other UK MPA  
120 processes, member of a representative body, and academically relevant research to MPAs  
121 and/or marine climate change. The size of the panel is not a critical feature of the Delphi  
122 method as participants are purposefully rather than randomly selected and reliable results  
123 can be obtained by choosing participants using strict inclusion criteria (Akins et al., 2005).

#### 124 2.1. Progression through rounds

125 The Delphi study began in January 2014 and consisted of two emailed questionnaires and a  
126 final focus group round that concluded the participant input process in September 2014.  
127 The focus group provided the participants with an opportunity for face to face interaction,  
128 encouraging motivation to remain engaged in the process. The participants had an adequate  
129 history of communication through the Scottish MPA process stakeholder workshops.  
130 Additionally, the use of the focus group further complemented the Delphi technique by  
131 emphasising the synergy of a group for producing ideas over and above individual  
132 contributions (Krueger and Casey, 2009). Results presented in this paper reflect final  
133 outcomes from the Delphi method, following the three rounds (Fig. 1.). Round One and Two  
134 identified potential management options and discussed the feasibility of these options.  
135 Recognising the feature-based approach to designation of the Scottish MPAs, the  
136 participants of the focus group were presented with a series of feature-based scenarios  
137 whereby the abundance or presence of a feature changed, to explore which possible  
138 management options were available and under which circumstances these were acceptable  
139 and feasible. The scenarios focused on the high level management options suggested by  
140 participants in previous rounds, rather than specific management relating to activities (e.g.  
141 types of gear restriction).




Figure 1. Overview of the Delphi process to identify management options under climate change scenarios for the Scottish MPA network. (Adapted from Lemieux and Scott (2011)).

143

## 144 2.2. Composition of the panel

145 Upon acceptance respondents from similar organisations nominated one person to speak  
146 on behalf of the interest group and this person became the point of contact (Participants 1,  
147 2 and 10). Reasons given for the collective input included the already heavy investment of  
148 relevant organisations involved in the on-going MPA designation process and reshuffling of  
149 employees within the relevant organisations to different policy areas. Six participants  
150 completed the Round One questionnaire and four participants responded to the Round Two  
151 questionnaire (Participants 1 and 8 did not complete). Whilst, this resulted in a low panel  
152 number for Round Two and a loss of two perspectives (policy maker and  
153 practitioner/professional), the information provided by the remaining four panellists was  
154 detailed and illustrated in-depth thinking concerning the feedback (from Round One) and  
155 resultant questions. Additionally, there was some overlap in the remaining participants with  
156 the non-respondents in terms of experience and background (i.e. a practitioner/professional  
157 and policy maker responded to Round Two). To counter-act the lower response rate of  
158 Round Two further action was taken: i) renewed efforts were made to contact the  
159 participants to encourage them to respond to the questionnaire and subsequent round; ii)  
160 additional potential participants from the stakeholder map having experience and  
161 knowledge in the research topic were invited to participate in the Delphi focus group.  
162 Subsequently, Participant 8 confirmed their acceptance of the invitation to attend the focus  
163 group with an additional four participants. The focus group was attended by ten participants  
164 (seven of whom had provided input into the preceding questionnaires (Table 1).

165

166 Table 1. Summary of participant characteristics and identification method.

Sub-Focus Group	Participant Number <sup>1</sup>	Organisation	(Group)	Identification Method
-	1*	Marine Scotland	Policy Makers and decision makers	Stakeholder Workshop Referral; reputation
-	2**	Scottish Environment Link	Representative Body; NGO	Stakeholder Workshop; reputation
1	3	Royal Society for the Protection of Birds (RSPB)	Representative Body; NGO	Stakeholder Workshop; reputation
1	4	Scottish Fishermen's Federation (SFF)	Representative Body	Stakeholder Workshop; referral
1	5†	Visit Scotland		Referral
1	6	Marine Conservation Society (MCS)	Representative Body; NGO	Stakeholder Workshop; reputation
1	7†	Sniffer (Registered charity)	Practitioner and Professional	Referral
2	8***	British Sub Aqua Club (BSAC), Academic	Practitioner and Professional	Referral; reputation
2	9	RSPB	Representative Body; NGO	Stakeholder Workshop; reputation
2	10	Scottish Natural Heritage (SNH)	Policy Makers and decision makers	Grey literature; Referral; reputation
2	11†	Academic	Practitioner and Professional	Referral; academic publications
2	12	Academic	Practitioner and Professional	Referral; academic publications

167 \*Participant completed Round 1 questionnaire but did not attend focus group

168 \*\* Participant completed questionnaires as collective (individual NGO members (RSPB and MCS)  
169 attended focus group)

170 \*\*\* Participant completed Round 1 questionnaire and attended focus group

171 †Participant attended focus group only

172 <sup>1</sup>The numbers used to list participants in the above table correspond to those used subsequently in  
173 this paper

174



## 175 2.3. Data collection and analysis

176 The questionnaire responses were imported into QSR International NVivo software (QSR  
177 International Pty Ltd, 2010) facilitating organisation, coding and retrieval of the data  
178 (Bazeley and Jackson, 2013). Analysis of questionnaire data followed a thematic content  
179 analysis to identify salient issues and key elements of the dataset (Green and Thorogood,  
180 2014). Data analysis broadly followed the steps suggested by Braun and Clarke (2006). Each  
181 questionnaire was firstly read through in detail with the addition of analytic notes and initial  
182 ideas regarding emerging themes. The data was then coded, grouping similar data segments  
183 (e.g. a particular sentence) together under each emergent code. Similar codes were  
184 combined under key themes that illustrated the perceptions of the participants for each  
185 question. All focus group sessions were audio-recorded and field notes were written by the  
186 researcher during and after the focus group. Additional field notes collected by the two  
187 facilitators, and flip charts produced by the participants were reviewed in the analysis  
188 process. The sessions were fully transcribed using NVivo software. Inductive open coding  
189 was used to generate codes and categories in the analysis providing a rich, in-depth and  
190 grounded account of the data (Corbin and Strauss, 2015). The results were interpreted by  
191 relating the categories to the research questions and theoretical ideas underpinning the  
192 research.

## 193 **3. Results**

### 194 3.1. Management success in the context of climate change

195 There were conflicting opinions as to whether the conservation objectives set for the MPA  
196 sites (conserve or recover for designated features), were ambitious enough in a climate  
197 change context. Opposing views were: MPAs should address wider ecological processes,  
198 improving the biodiversity of the designated site but also having wider benefits for the  
199 marine environment; and MPAs were designated for specific purposes (to conserve or  
200 recover specific species and habitats), therefore too high expectations were placed on what  
201 the network could successfully achieve.

202 “If the conservation objectives of an individual MPA are achieved then it could be argued  
203 that the MPA has been successful but you would maybe want to achieve more in terms of  
204 helping to increase resilience in the marine environment to climate change and other  
205 pressures.” **Participant 1.**

206 The difference between success of a single MPA site and the success of the network was  
207 highlighted, raising the question of how success of the network may be achieved if there are  
208 different objectives at a site and network level. Participants felt further work was needed to  
209 define ecological coherence and even a working definition of what is considered an MPA  
210 network in the context of the Scottish MPA sites.

211 “It is also not clear to what extent the network will be “ecologically coherent” given that it  
212 doesn't seem to have been designed with that in mind, but rather to protect a series of key  
213 (but at times isolated) features and species.” **Participant 11.**

214 There was concern that the network had not been designed to consider connectivity and  
215 therefore that success in terms of realising wider ecosystem health may not be  
216 accomplished. Participants recognised that enhancing ecosystem health would be important  
217 given the additional stress that climate change would likely have on the marine  
218 environment and that the network should not just keep the “status quo” by protecting  
219 residual populations. The concept of “status quo” was linked to ideas of dynamism in the  
220 marine environment, recognising that features may change in the face of climate change,  
221 i.e. it would not be possible to protect MPAs from sea temperature changes, as these wider  
222 processes would not recognise the site boundaries. Disagreement was evident; one  
223 participant was concerned with the approach recommended to protect areas for wider  
224 ecological processes.

225 This view reflects the feature based approach for the network yet appears to contradict  
226 with the original Scottish vision for the MPA network. The most widely mentioned factor for  
227 success was the ability of the management (as a result of the legislation underpinning the  
228 designations recognising climate change) to be adaptable. Participants were divided as to

229 whether planned management and monitoring (at the point of survey) would account for  
230 climate change.

231 “The planned management of MPAs in the Scottish MPA network is being driven by the  
232 sensitivity of the proposed protected features to pressures arising from activities known to  
233 be taking place within the sites. Climate change scenarios really aren't informing  
234 management at this stage.” **Participant 1.**

235 Overall, there was a dichotomy in participant opinion for a successful network: the  
236 protection of specific features and habitats of conservation interest versus wider  
237 improvement of the marine environment as a result of the protection and whether these  
238 are mutually achievable.

239

### 240 3.2. Management scenarios

241 The preceding questionnaires identified management options and discussed the feasibility  
242 of these options. These were reframed by the researchers into a matrix of high-level  
243 management actions in combination with possible climate change scenarios. For example, a  
244 feature is no longer present within the MPA, which possible management option is  
245 suitable/acceptable under this scenario. This approach was based on the discussions  
246 regarding feasible management options, and recognised the feature-based approach to  
247 designation of the Scottish MPAs. The participants were presented with a series of feature-  
248 based scenarios whereby the abundance or presence of the feature changed and each  
249 scenario was discussed by participants with the aim of deciding which possible management  
250 actions were available and under which circumstances these were acceptable and feasible.  
251 The matrix focused on the high level options suggested by participants in previous rounds,  
252 rather than specific management relating to activities (e.g. types of gear restriction). Sites  
253 with multiple designated features present were not considered, however, participants were  
254 given the option of considering wider biodiversity and whether this would affect their  
255 choice of management action.

256 The management scenarios matrix (Table 2) summarises the possible management options  
257 (from participant discussion) at a site and network level under five different scenarios of  
258 change for the MPA feature at the level of an individual MPA: i) the feature is no longer  
259 present ii) feature is decreasing iii) feature is stable/demonstrating no overall trend iv)  
260 feature is improving and v) the feature is recovered.<sup>1</sup> In terms of the matrix, the above  
261 change scenarios are in absolute terms (i.e. not compared to trends in other times and  
262 places). The scenarios are also further sub-categorised for site integrity (i.e. wider  
263 biodiversity of the site in addition to the status of the feature for which the site is  
264 designated) and how the MPA feature is performing at a network level i.e. whether it is  
265 stable/declining/increasing across the network. For all scenarios, participants suggested a  
266 “balanced review” would be required, and evidence to support decisions before deciding  
267 upon any action, taking into account the whole network at appropriate timescales, but did  
268 not elaborate on what would constitute a balanced review or what evidence would be  
269 needed. Participants suggested that a network review would be useful for a “recalibration”,  
270 identifying if any gaps in feature protection were present, or if broader network scale  
271 factors (i.e. climate change) were a cause of change. However, it was recognised that  
272 identifying causal factors was often incredibly difficult, highlighting the need for a strong  
273 monitoring programme. Therefore, some participants maintained a “precautionary”  
274 approach to management (i.e. stricter management measures); “precautionary” was also  
275 applied in reference to changing management, (i.e. ensuring a strong evidence base before  
276 changing current management measures).

277 Participants felt that a review of management measures would therefore be needed to  
278 answer whether the current management had fully removed the pressure. There was also  
279 recognition from participants that the dynamic nature of the marine environment would  
280 need to be reflected in adaptive management.

281 Control areas were mentioned in reference to understanding changes and linked to  
282 resilience. The option of a new MPA (or moving an MPA) was linked to recovering net loss of

---

<sup>1</sup> Researchers used the term “recovered” in reference to the draft definitions of MPA conservation objectives of either “conserve” or “recover” (Scottish Government, 2012). At the time of the research there was no quantitative definition or target of “recover” for the individual features.

283 a species where conditions were more favourable, or where suitable climatic conditions still  
284 prevailed. A more controversial option (from the participants) was MPA expansion, although  
285 mentioned in previous rounds, it was suggested that to expand the area a big change in  
286 policy would be needed as the boundaries of a site are tightly drawn around the feature of  
287 interest and legislatively implemented.

288 Problems with a feature based approach in a climate change context were identified by the  
289 researchers from the participant discussion; a summary of participant discussion and  
290 researcher comments around these problems is provided in Table 3.

291

292

293 Table 2. Summary Matrix of Management Options: Condition of MPA features under different scenarios of change

MPA feature Scenario at a site level <sup>1</sup>	Site Integrity <sup>2</sup>	MPA feature at a network level	Possible Management Actions (from participant discussion)	Decision Making Process (from participant discussion)
No Longer Present	Low quality	Still present	1. New MPA/Move MPA (Look to establish another MPA for the feature) Designate a new alternative area which may succeed, e.g. within new climatic window of feature.	<ul style="list-style-type: none"> <li>- Question whether the current management actions are/were appropriate</li> <li>- Is there an alternative feature within the MPA?</li> <li>- Would maintaining this MPA fill a gap in network wide protection?</li> </ul>
	Low/high quality	Still present	2. Reduce pressures in other MPAs. Look at other sites across the network where the PMF is still present within its climate window and reduce other stressors.	
	Low quality	Still present/no longer present across the network	3. De-designate the MPA <sup>3</sup> Option to give up on an area that has failed.	
	High quality site for biodiversity/other features	Still present/no longer present across the network	4. "Rebadge" the MPA (Look to designate the current site for another feature).	
Feature Decreasing	Low/high quality	Stable/Declining	1. Reduce pressures on PMF (further restriction to full ban on damaging activities).	<ul style="list-style-type: none"> <li>- Identify the causes of a decline</li> <li>- Look to recover net loss of the feature across the network</li> </ul>
	Low/High quality	Stable/Declining	2. Expand the area of the MPA	
	Low quality	Declining across the network	3. New MPA/Move MPA (Look to establish another MPA for the feature)	
Feature Stable	Low/High quality	Stable	1. Maintain current management measures	- Continue monitoring
Feature Improving	High quality site for feature	Stable across network/Feature common across network	1. Maintain current management measures	<ul style="list-style-type: none"> <li>- Review pressures across the network</li> <li>- Is there clear evidence of improvement? E.g. greater extent, higher biodiversity, better age structure</li> </ul>
	High quality for feature	Declining across the network	2. Expand the area of the MPA	
Feature Recovered	High quality for feature	Feature common across network	2. Review management of feature in other sites where it was not present previously	- Need for substantial evidence to reduce or change management

	High quality site for feature, biodiversity and other features	Feature common across network	3. Reduce or change management e.g. is there an option for sustainable use	<ul style="list-style-type: none"> <li>- Is there clear evidence that it was the management of an activity that led to that improvement?</li> <li>- Is there clear evidence of improvement? E.g. greater extent, higher biodiversity, better age structure</li> </ul>
--	--	-------------------------------	--	---

294 <sup>1</sup>Change scenarios are in absolute terms (i.e. not compared to trends in other times and places).

295 <sup>2</sup>Site Integrity: Quality of the site for wider biodiversity in addition to the status of the feature for which the site is designated. This was summarised as a qualitative  
296 statement of either “low quality” or “high quality”. Site integrity was mentioned by participants in reference to site condition monitoring for other nature conservation  
297 sites (i.e. SPAs and SACs) and therefore could be of future relevance to the MPA sites, whilst not referenced in MPA objectives.

298 <sup>3</sup>De-designate MPA: There is a provision to de-designate an MPA under the Marine Act (Scotland) 2010.

299  
300  
301  
302

Table 3. Summary of participant discussion around problems of a feature based approach in a climate change context

Researcher identified Problem from participant discussion	Participant Comments	Climate change scenario	Researcher Comments
Success judged on a single feature	Success of MPA will be dependent on state of that feature irrespective of wider biological health	Feature declines or is absent from site results in site viewed as failure irrespective of potential wider site improvement.	<i>Conceptually linked to valuation of marine biodiversity. Conflict between feature level objectives, wider pressures and an ecosystem or network level view of success</i>
MPA Management around a particular feature	Feature based management does not account for buffer zones or an ecosystem approach. Damaging activity is not precluded from the entire site, MPA is fragmented into various zones of management. Consequential protection of MPA designation is neglected.	An ecosystem approach required for climate change resilience at a network level is not considered. Wider biodiversity resilience to climate change impacts is not considered. Recovery (range expansion) of species and habitats is unlikely if management is tied to presence.	<i>Conflict between feature level objectives, wider pressures and an ecosystem or network level view of success</i>
"Rebadging" an MPA	A feature, for which the MPA is designated, is lost from the site. Potential for the site to be rebadged/repurposed for another feature.	If a feature is lost and you did not repurpose the MPA, you could lose consequential protection or any improvement in ecosystem health that resulted as a reduction in pressures. Secondly, there may be circumstances where data has improved and led to the identification of other Priority Marine Features (PMFs) or vulnerable species that could benefit from protection. Keep the site for monitoring purposes- resources dedicated	<i>Important that sites be retained for the right reasons which would require a network level review and stakeholder-determined reasons. There was a suggestion that it may be appropriate to look for a new area, although de-designation was seen as a last resort (species may not completely disappear or may have an opportunity to re-establish), but an option that should remain in the "management toolbox". Strong industry concerns in rebadging an MPA due to perceived lack of justification. A logical response from the MPA designation process would be to de-designate an MPA if it has not achieved its management objective (i.e conserve</i>



			<p><i>feature). By retaining the MPA for other reasons than the specific feature designation could be seen as “moving goalposts” by changing the rationale behind designation.</i></p> <p><i>However, there could be a trend towards loss of protection if failing MPAs are removed without seeking to understand why they are failing and seeking to rectify.</i></p> <p><i>Linked to the appropriate allocation of resources</i></p>
Features are not self-recruiting	Sites are not designed using connectivity principles.	Network is not designed as an ecologically coherent one and therefore does not consider potential climate change impacts	<i>Perceived limited consideration of connectivity across the network. Echoes concerns from MPA process stakeholder workshops</i>
Ecosystem health	A species cannot exist in isolation of its ecosystem. Lack of consideration for wider ecosystem health.	Network is not designed as an ecologically coherent one which takes into account wider ecosystem health and therefore does not consider potential climate change impacts	<i>Linked to the lack of connectivity principles across the site.</i>
Precautionary approach	Proposed management* is not optimal (or precautionary) and areas will be under protected. Considering wider ecosystem function and buffer zones of management and concern for whether the selection of features looked at richly biodiverse sites,	To ensure climate change resilience, effective management would be required.	<i>Effective management was considered by some participants as areas of strict protection surrounded by buffer zones</i>
Climate change not considered	Would more MPAs with features that are sensitive to climate change would have been established if climate change had been considered at the beginning of the process. Key features not considered in terms of their vulnerability to climate change	Under scenarios of loss, concerns were raised that if the success or quality of the site is to be judged solely on the status of the feature, and a site were designated for a climate sensitive species (e.g. maerl) which if declined or was lost from the site, the whole site would effectively be redundant. Therefore, it may be possible that a number of sites are potentially vulnerable to the feature being lost; the approach does not account for how assemblages of species in MPA sites may change under climate change scenarios.	<i>Some participants were reluctant to have the MPAs broadened, stating that they should be justified.</i>

		Suggested that sites identified for a specific habitat or biotope are unlikely to lose the whole interest under scenarios of decline. One solution proposed was to widen the designation of the site to incorporate more habitats and features	
--	--	--	--

303 \*Proposed Management: At the time of study management measures for the MPA sites were not in place

304 **4. Discussion**

305 Views of MPA success are likely to change under climate change scenarios (Hopkins et al., 2016a);  
306 this increases the complexity of applying legal definitions of success, which may become redundant  
307 under such scenarios. This study demonstrates the large fragmentation of opinion in what  
308 constitutes success even in the absence of considering climate change. As the discussion progressed  
309 from questionnaires to the focus group, the agreement of success in abstract principles broke down  
310 in the face of operational realities. A fundamental split was evident between participants  
311 sympathetic to the provision for sustainable use within the MPA network, and those participants  
312 stating that the MPA network should be primarily for conservation, enhancement of the wider  
313 marine environment and should contribute to climate change resilience. The different perceptions of  
314 MPA success influenced the subsequent discussions of management scenarios; whether participants  
315 felt the MPA network should strive for the minimum protection of species and habitats (features)  
316 versus MPAs enhancing the wider marine environment.

317 In the context of the Scottish feature-led MPA process, the approach to management resembles a  
318 discriminating approach using a feature sensitivity tool (FEAST),<sup>2</sup> which analyses the sensitivity of a  
319 designated feature to different types of human activity. Management measures based on this  
320 sensitivity may not be required across the entirety of the site if the feature is not present across the  
321 whole of the site. However, elsewhere there has been a move away from a species-by-species  
322 management towards broader ecosystem level strategies (Jentoft et al., 2007). By focusing  
323 management measures on one feature or species, impacts on other species (which may be of high  
324 ecological importance) are effectively ignored. Better protection of MPA features could be achieved  
325 by not only managing the direct impacts (i.e. habitat destruction) but also by considering the wider  
326 factors that influence their health (e.g. water quality, prey availability and trophic links). A review of  
327 scientific knowledge and international perceptions that informed the development of this study  
328 (Hopkins et al., 2016a) suggest management and protection should account for wider ecosystem  
329 links and concepts of resilience in the face of a large amount of uncertainty from climate change.

330 Participants noted that for MPAs to be successful under future scenarios of climate change,  
331 flexibility and adaptation were needed. However, although adaptive management is needed for  
332 climate change resilient MPAs (Davies et al., 2016; Hopkins et al., 2016a), there are few examples in  
333 practice. The importance of monitoring to inform adaptive management was noted whilst discussing

---

<sup>2</sup><http://www.marine.scotland.gov.uk/FEAST/>.

334 the scenarios to clearly evaluate the effect of protection and to discern the impacts of climate  
335 change. Proposed options for adaptively managing MPAs including: flexible boundaries, buffer zones  
336 of management, and temporary MPAs that track ecosystem processes or features were deemed far  
337 from a practical reality for MPAs at present. The iterative nature of the Delphi method highlighted  
338 the difference between proposing options and subsequently using these in a practical scenario. For  
339 example, changing MPA boundaries was proposed as an option in the questionnaire rounds, yet  
340 when confronted with implementing this option for a range expansion (for example), participants  
341 were reluctant to use boundary changes. Changing MPA boundaries was regarded by the  
342 environmental sector as too fluid a measure to provide effective long term protection, whilst the  
343 fishing sector were concerned that it would lead to long term financial uncertainty. Therefore, whilst  
344 most actors within the MPA process advocate adaptive management, it remains difficult to define  
345 how this will work in a practical sense.

346 The success of adaptive management is highly dependent on strong monitoring programmes that  
347 are consistent and well-funded (Mee et al., 2008) and the policy context. MPAs are likely to be  
348 implemented in the absence of high quality baseline information (Sale et al., 2005) and with a large  
349 uncertainty regarding how climate change will affect MPAs. Therefore, as more knowledge becomes  
350 available through targeted research and monitoring, adaptive management is a necessary  
351 mechanism for incorporating new information and refining management with regards to marine  
352 protection (Mee et al., 2008; McDonald and Styles, 2014). Participants highlighted their concerns  
353 that the monitoring task for the MPA network was overwhelming, both in terms of the scale of the  
354 information needed to be able to confidently state that the network was achieving its aims, and in  
355 terms of the amount of resources needed to monitor both at a site and network level. Whilst the  
356 political framework is in place for the Scottish network to be adapted in light of new knowledge via  
357 the network review process there is also the requirement of political will in order to implement  
358 suitable responses (Mee et al., 2008) and robust mechanisms that ensure action is taken in light of  
359 new information, rather than a continuation of monitoring.

360 Participants were concerned that the Scottish MPA network had not been designed to protect  
361 ecosystem function and wider biodiversity. MPA networks designed for protecting biodiversity are  
362 likely to be important in preserving ecological functioning and therefore contributing to ecosystem  
363 resilience (Steneck et al., 2002). A network consisting of strictly protected areas with no intense  
364 anthropogenic stressors (e.g. fishing) and that incorporate consideration of ecosystem function are  
365 likely to be the most resilient to climate change (Harley et al., 2006; Brock et al., 2012; Micheli et al.,  
366 2012). The feature based approach used in Scotland is therefore concerning because without a

367 coherent, connected MPA network, it is unlikely to be resilient to the impacts of climate change  
368 (Olds et al., 2012; Magris et al., 2014; Andrello et al., 2015). The approach taken by other countries  
369 (e.g. Australia) has been to incorporate multi-use at a network-scale but with a core of strictly  
370 protected no-take areas. Single MPAs that are small and not strictly protected, could be considered a  
371 false economy as larger well protected MPAs may be less costly in terms of reduced fisheries  
372 revenue by increasing the likelihood of spillover, stock recovery and a reduction in the variation of  
373 stock levels. However, fishers may not perceive the risk buffering capacity of larger MPAs sufficient  
374 to offset the value of foregone harvesting (Carter, 2003). Larger well protected MPAs may be less  
375 prone to sudden and unpredictable change (Edgar et al., 2014) and are likely easier to manage,  
376 requiring less adaptive management strategies and less detailed long term monitoring. However, at  
377 a network scale, there is potential for a portfolio of MPA design, with a range of protection from  
378 strict protection/no-take to multiple use. There is a useful opportunity for investigating varying  
379 levels of protection across the network, in the recently designated Fair Isle Demonstration and  
380 Research MPA as it is specifically targeted toward researching sustainable marine management  
381 approaches (FIMETI, 2015).

382 The restoration of marine habitats as outlined in the Marine Strategy Framework Directive (MSFD)  
383 and OSPAR guidelines, and a possible site level objective for an MPA feature in the Scottish MPA  
384 process recognises the need to increase resilience in degraded ecosystems. Whilst there are strong  
385 political foundations for restoration, these do not address the scientific (and socio-political)  
386 difficulties (Hopkins et al., 2016b). The use of feature presence is less ambiguous politically when  
387 compared to identifying and measuring overall ecosystem health. There are also technical  
388 uncertainties over whether a habitat will recover, how long it will take and non-linear recovery  
389 trajectories (Mee et al., 2008). Alternative stable states of an ecosystem may exist which make  
390 restoration attempts (to restore the ecosystem to the previous desirable state) unfeasible,  
391 impractical or too expensive (Hughes et al., 2005; Selkoe et al., 2015). The concept of shifting  
392 baselines (see Pauly (1995)) needs to be considered with regards to the desirable state of the  
393 ecosystem that the MPA should achieve. Suggestions from participants that qualitative discussions  
394 may need to occur to decide what past ecosystems looked like, echoed recommendations by  
395 Campbell et al. (2009) that marine restoration will need to explicitly recognise value laden  
396 judgements inherent in the decision context (Mee et al., 2008; Campbell et al., 2009). These value-  
397 laden judgements also extend into judgements of what future ecosystems will look like under  
398 climate change (as suggested in Hopkins et al., 2016a; b); reference states in this context are  
399 particularly contentious in marine systems (Mee et al., 2008).

400

401 Ecosystem Based Management (EBM) may provide a solution by integrating conservation with  
402 spatial ecology and ecosystem functioning. EBM focuses on the protection of multiple species,  
403 ecosystem processes and societal values, taking into account the wider effects of human use on the  
404 environment (Mee et al., 2008; Campbell et al., 2009; Olds et al., 2012). However, the data  
405 requirements for this and the current political landscape may mean that EBM approaches are  
406 unlikely to be implemented in the short term. The use of EBM as a solution was also not resolved in  
407 this study and remained part of the split in perceptions of whether the wider environment should be  
408 considered within the MPA designations. If EBM approaches are unfeasible at present, and feature-  
409 led approaches are inappropriate for climate change, management decisions need to be taken in  
410 light of data from reference sites and baseline for changes without the confounding influence of  
411 controllable (at least to some degree) or restrictable human stressors (e.g. fishing, dredging,  
412 development etc.). Without reference sites, “expert judgement” and human perceptions of change  
413 are used to make management decisions (Mee et al., 2008). As perceptions of quality can shift over  
414 each generation (Pauly, 1995) with each generation having its own reference state for what is high  
415 or “good” quality, these perceptions of quality may decrease as generally society becomes used to a  
416 lower level (Mee et al., 2008). Subjective management decisions are unlikely to be accurate and  
417 reference states of quality imply judgements of what is “good” or “bad” about the natural  
418 environment (Mee et al., 2008). The development of the MPA network is therefore recommended as  
419 a practical solution, but should include the implementation of strictly protected reference sites to  
420 allow more objective assessments of ecosystem health to be made (Mee et al., 2008) and  
421 importantly to increase resilience for climate change impacted species and habitats across the wider  
422 network.

## 423 **5. Conclusions**

424 The use of the Delphi method in this study enabled the researchers to include both stakeholders and  
425 decision makers to explore climate change adaptation options tailored to the Scottish MPA network.  
426 Continued dialogue between stakeholders, decision makers and scientists will be necessary to  
427 monitor, review and adaptively manage the MPA network in the context of climate change. The  
428 management framework presented here is intended to support the decision making process,  
429 recognising that some of the adaptation options may not be feasible or appropriate in a future  
430 context, and any decision should be made in response to new information and with consultation.

431 Over the course of the iterative process, a fundamental split between the perceptions of different  
432 stakeholder groups became evident. Those stakeholders, sympathetic to the provision of sustainable  
433 use (i.e the fishing sector representative) were supportive of the feature approach to conservation  
434 which underpins MPA designation in Scotland. Conversely, other stakeholders felt conservation  
435 through MPAs should contribute to wider ecosystem health requiring consideration of ecosystem  
436 links in the application of management. The process indicated that this difference in perception may  
437 be intractable between the two groups even within a carefully designed MPA process. The Scottish  
438 MPA process designated MPAs with an evidence base (feature presence and impact sensitivity) yet  
439 also specifies aiming to enhance ecosystem health and contribute to an ecologically coherent  
440 network but without a mechanism for Ecosystem Based Management (EBM) or a clear strategic  
441 ecosystem level vision. Proposed feasible options for including climate change specific management  
442 and monitoring strategies as a result of this study include the use of experimental reference areas  
443 (e.g. Fair Isle MPA). These areas could be used to monitor the impact of climate change on MPA  
444 species and habitats and the effect of varying levels of protection across the network on climate  
445 change resilience. Marine reserves are at this point considered politically unfeasible with some  
446 stakeholders, and the use of EBM as a solution appears unresolved.

447 From a scientific perspective strictly protected marine reserves are thought to be more resilient to  
448 climate change and reference areas will be critical to understand climate change impacts and effects  
449 supported by monitoring over medium to long-term timescales. Developing scenarios for MPAs  
450 under climate change is a useful exercise in developing potential management options and aiding  
451 decision making. For the Scottish MPA network, a key recommendation would be to develop  
452 research regarding how the MPA network at various scales will be affected by climate change, and  
453 use the outputs from this study to guide decisions regarding MPA management.

#### 454 **Acknowledgements**

455 Thank you to all the participants in this study and to Dr Sophie Elliot and Dr Joanne Clark for  
456 assistance in facilitation. Thank you to the reviewers of this manuscript for their insights. This  
457 research was supported and funded by ClimateXChange (referenceno:A10431853). ClimateXChange  
458 is a collaborative initiative between Scottish research and higher education institutes and is funded  
459 by the Scottish Government. The participant workshop was supported by a MASTS Small Grant  
460 SG215.

#### 461 **References**

- 462 Akins, R.B., Tolson, H., Cole, B.R., 2005. Stability of response characteristics of a Delphi panel :  
463 application of bootstrap data expansion. *BMC Med. Res. Meth.* 5.
- 464 Allison, G.W., Lubchenco, J., Carr, M.H., 1998. Marine reserves are necessary but not sufficient for  
465 marine conservation. *Ecol. Appl.* 8, 79–92.
- 466 Andrello, M., Mouillot, D., Somot, S., Thuiller, W., Manel, S., 2015. Additive effects of climate change  
467 on connectivity between marine protected areas and larval supply to fished areas. *Divers. Distrib.* 21,  
468 139–150.
- 469 Bazeley, P., Jackson, K., 2013. *Qualitative Data Analysis with NVivo*, second ed. Sage Publications,  
470 London. California. New Delhi. Singapore.
- 471 Bernhardt, J.R., Leslie, H.M., 2013. Resilience to climate change in coastal marine ecosystems. *Annu.*  
472 *Rev. Mar. Sci.* 5, 371–392.
- 473 Braun, V., Clarke, V., 2006. Using thematic analysis in psychology. *Qual. Res. Psychol.* 3, 77–101.
- 474 Brock, R.J., Kenchington, E., Martinez-Arroyo, A. (Eds.), 2012. *Scientific Guidelines for Designing*  
475 *Resilient Marine Protected Area Networks in a Changing Climate*. Commission for Environmental  
476 Cooperation, Montreal. Canada.
- 477 Campbell, L.M., Gray, N.J., Hazen, E.L., Shackeroff, J.M., 2009. Beyond baselines: rethinking priorities  
478 for ocean conservation. *Ecol. Soc.* 14 [www document]. <http://www.ecologyandsociety>.
- 479 Carter, D.W., 2003. *Protected Areas in Marine Resource Management: Another Look at the*  
480 *Economics and Research Issues*.
- 481 Chuenpagdee, R., Pascual-Fernández, J.J., Szeliánszky, E., Luis Alegret, J., Fraga, J., Jentoft, S., 2013.  
482 *Marine protected areas: Re-thinking their inception*. *Mar. Pol.* 39, 234–240.
- 483 Cliquet, A., Backes, C., Harris, J., Howsam, P., 2009. Adaptation to climate change – legal challenges  
484 for protected areas. *Utrecht Law Rev.* 5, 158–175.
- 485 Corbin, J.M., Strauss, A., 2015. *Basics of Qualitative Research. Techniques and Procedures for*  
486 *Developing Grounded Theory*, fourth ed. Sage Publications, UK.



487 Côté, I.M., Darling, E.S., 2010. Rethinking ecosystem resilience in the face of climate change. *PLoS*  
488 *Biol.* 8, e1000438.

489 Davies, H.N., Beckley, L.E., Kobryn, H.T., Lombard, A.T., 2016. Features into the incremental  
490 refinement of an existing marine park. *PLoS One* 11, 1–21.

491 Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S.,  
492 Becerro, M.A., Bernard, A.T.F., Berkhout, J., Buxton, C.D., Campbell, S.J., Cooper, A.T., Davey, M.,  
493 Edgar, S.C., Försterra, G., Galván, D.E., Irigoyen, A.J., Kushner, D.J., Moura, R., Parnell, P.E., Shears,  
494 N.T., Soler, G., Strain, E.M.A., Thomson, R.J., 2014. Global conservation outcomes depend on marine  
495 protected areas with five key features. *Nature* 506, 216–220.

496 FIMETI, 2015. Fair Isle Proposed Demonstration and Research MPA (Final Document). Available from:  
497 <http://www.gov.scot/Resource/0049/00494122.pdf>.

498 Glass, J.H., Scott, A.J., Price, M.F., 2013. The power of the process: Co-producing a sustainability  
499 assessment toolkit for upland estate management in Scotland. *Land Use Pol.* 30, 254–265.

500 Gobbi, M., Riservato, E., Bragalanti, N., Lencioni, V., 2012. An expert-based approach to invertebrate  
501 conservation: identification of priority areas in central-eastern Alps. *J. Nat. Conserv.* 20, 274–279.

502 Green, J., Thorogood, N., 2014. *Qualitative Methods for Health Research*. Sage Publications, UK.

503 Harley, C.D.G., Hughes, A.R., Hulgren, K.M., Miner, B.G., Sorte, C.J.B., Thornber, C.S., Rodriguez, L.F.,  
504 Tomanek, L., Williams, S.L., 2006. The impacts of climate change in coastal marine systems. *Ecol.*  
505 *Lett.* 9, 228–241.

506 Harley, C.D.G., Rogers-Bennett, L., 2004. The potential synergistic effects of climate change and  
507 fishing pressure on exploited invertebrates on rocky intertidal shores. *Calif. Coop. Ocean. Fish.*  
508 *Investig. Rep.* 45, 98–110.

509 Hess, G.R., King, T.J., 2002. Planning open spaces for wildlife. *Landsc. Urban Plann.* 58, 25–40.

510 Holling, C.S., 1973. Resilience and stability of ecological systems. *Annu. Rev. Ecol. Evol. Syst.* 4, 1–23.

- 511 Hopkins, C.R., Bailey, D.M., Potts, T., 2016a. Perceptions of practitioners: managing marine  
512 protected areas for climate change resilience. *Ocean Coast Manag.* 128, 18–28.
- 513 Hopkins, C.R., Bailey, D.M., Potts, T., 2016b. Scotland's Marine Protected Area network: reviewing  
514 progress towards achieving commitments for marine conservation. *Mar.Pol.* 71, 44–53.
- 515 Howson, C., Steel, L., Carruthers, M., Gillham, K., 2012. Identification of Priority Marine Features in  
516 Scottish Territorial Waters. Scottish Natural Heritage Commissioned Report No. 388.
- 517 Hsu, C., Sandford, B., 2007. The Delphi technique: making sense of consensus. Practical assessment.  
518 *Res. Eval.* 12, 1–8.
- 519 Hughes, T.P., Bellwood, D.R., Folke, C., Steneck, R.S., Wilson, J., 2005. New paradigms for supporting  
520 the resilience of marine ecosystems. *Trends Ecol. Evol.* 20, 380–386.
- 521 Jentoft, S., van Son, T.C., Bjorkan, M., 2007. Marine protected areas: a governance system analysis.  
522 *Hum. Ecol.* 35, 611–622.
- 523 Krueger, R.A., Casey, M.A., 2009. *Focus Groups: a Practical Guide for Applied Research.*
- 524 Lemieux, C.J., Scott, D.J., 2011. Changing climate, challenging choices: identifying and evaluating  
525 climate change adaptation options for protected areas management in Ontario, Canada. *Environ.*  
526 *Manag.* 48, 675–690.
- 527 Levy, J., Ban, N., 2013. A method for incorporating climate change modelling into marine  
528 conservation planning: an Indo-west Pacific example. *Mar. Pol.* 38, 16–24.
- 529 Linstone, H.A., Turoff, M., 2002. *The Delphi Method: Techniques and Applications.* Page Futures  
530 *Research Methodology.*
- 531 Lubchenco, J., Palumbi, S., Gaines, S.D., Andelman, S., 2003. Plugging a hole in the ocean: the  
532 emerging science of marine reserves 1. *Ecol. Appl.* 13, 3–7.
- 533 Magris, R.A., Pressey, R.L., Weeks, R., Ban, N.C., 2014. Integrating connectivity and climate change  
534 into marine conservation planning. *Biol. Conserv.* 170, 207–221.

535 McDonald, J., Styles, M.C., 2014. Legal strategies for adaptive management under climate change. J.  
536 Environ. Law 26, 25–53.

537 Mee, L.D., Jefferson, R.L., Laffoley, D.D.'A., Elliott, M., 2008. How good is good? Human values and  
538 Europe's proposed Marine Strategy Directive. Mar. Pollut. Bull. 56, 187–204.

539 Micheli, F., Saenz-Arroyo, A., Greenley, A., 2012. Evidence that marine reserves enhance resilience to  
540 climatic impacts. PLoS One 7, e40832.

541 Mukherjee, N., Hugel, J., Sutherland, W.J., McNeill, J., Van Opstal, M., Dahdouh-Guebas, F., Koedam,  
542 N., 2015. The Delphi technique in ecology and biological conservation: applications and guidelines.  
543 Met. Ecol. Evol. <http://dx.doi.org/10.1111/2041-210X>. 12387.

544 O'Neill, S.J., Osborn, T.J., Hulme, M., Lorenzoni, I., Watkinson, A.R., 2008. Using expert knowledge to  
545 assess uncertainties in future polar bear populations under climate change. J. Appl. Ecol. 45, 1649–  
546 1659.

547 Olds, A.D., Pitt, K.A., Maxwell, P.S., Connolly, R.M., 2012. Synergistic effects of reserves and  
548 connectivity on ecological resilience. J. Appl. Ecol. 49, 1195–1203.

549 Pauly, D., 1995. Anecdotes and the shifting baseline syndrome of fisheries. Trends Ecol. Evol. 10,  
550 430.

551 Pollnac, R., Christie, P., Cinner, J.E., Dalton, T., Daw, T.M., Forrester, G.E., Graham, N.A.J.,  
552 McClanahan, T.R., 2010. Marine reserves as linked social-ecological systems. Proc. Natl. Acad. Sci.  
553 U.S.A. 107, 18262–18265.

554 QSR International Pty Ltd, 2010. NVivo Qualitative Data Analysis Software, tenth ed.

555 Reed, M.S., 2008. Stakeholder participation for environmental management: a literature review.  
556 Biol. Conserv. 141, 2417–2431.

557 Rowe, G., Wright, G., 2011. The Delphi technique: past, present, and future prospects - introduction  
558 to the special issue. Technol. Forecast. Soc. Change 78, 1487–1490.

559 Sale, P.F., Cowen, R.K., Danilowicz, B.S., Jones, G.P., Kritzer, J.P., Lindeman, K.C., Planes, S., Polunin,  
560 N.V.C., Russ, G.R., Sadovy, Y.J., Steneck, R.S., 2005. Critical science gaps impede use of no-take  
561 fishery reserves. *Trends Ecol. Evol.* 20, 74–80.

562 Scottish Government, 2011a. A Strategy for Marine Nature Conservation in Scotland's Seas. Available  
563 from:  
564 <http://www.scotland.gov.uk/Topics/marine/marineenvironment/Conservationstrategy/marineconst>  
565 [rategy](http://www.scotland.gov.uk/Topics/marine/marineenvironment/Conservationstrategy/marineconst).

566 Scottish Government, 2011b. Marine Protected Areas in Scotland's Seas: Guidelines on the Selection  
567 of MPAs and the Development of the MPA Network. Available from:  
568 [http://www.scotland.gov.uk/Topics/marine/marine environment/mpanetwork/mpaguidelines](http://www.scotland.gov.uk/Topics/marine/marine%20environment/mpanetwork/mpaguidelines).

569 Scottish Government, 2012. Report to the Scottish parliament on progress to identify a Scottish  
570 network of marine protected areas. Available from:  
571 <http://www.scotland.gov.uk/Topics/marine/marine->  
572 [environment/mpanetwork/MPAParliamentReport](http://www.scotland.gov.uk/Topics/marine/marine-).

573 Selkoe, K.A., Blenckner, T., Caldwell, M.R., Crowder, L.B., Erickson, A.L., Essington, T.L., Estes, J.A.,  
574 Fujita, R., Halpern, B.S., Hunsicker, M.E., Mach, M.E., Martone, R.G., Mease, L.A., Salomon, A.K.,  
575 Samhuri, J.F., Scarborough, C., Stier, A.C., White, C., Zedler, J., 2015. Principles for managing marine  
576 ecosystems prone to tipping points. *Ecosys. Health Sustain.* 1, 17.

577 Steneck, R.S., Graham, M.H., Bourque, B.J., Corbett, D., Erlandson, J.M., Estes, J.A., Tegner, M.J.,  
578 2002. Kelp forest ecosystems: biodiversity, stability, resilience and future. *Environ. Conserv.* 29, 436–  
579 459.