



ARTICLE

Received 31 Jul 2012 | Accepted 10 Apr 2013 | Published 21 May 2013

DOI: [10.1038/ncomms2859](https://doi.org/10.1038/ncomms2859)

OPEN

Aurora-A controls pre-replicative complex assembly and DNA replication by stabilizing geminin in mitosis

Takaaki Tsunematsu¹, Yoshihiro Takihara², Naozumi Ishimaru³, Michele Pagano^{4,5}, Takashi Takata¹ & Yasusei Kudo^{1,3}

Geminin, an essential factor for DNA replication, directly binds to the licensing factor Cdt1 and inhibits pre-replicative complex formation to prevent re-replication. In G1, geminin levels are controlled by the anaphase-promoting complex/cyclosome (APC/C) ubiquitin ligase complex, which targets geminin for proteasomal degradation to allow pre-replicative complex formation. Conversely, from S to G2, geminin is stabilized due to APC/C ubiquitin ligase complex inhibition, ensuring the inhibition of pre-replicative complex formation. However, mitotic regulation of geminin has hitherto not been described. Here we show that Aurora-A phosphorylates geminin on Thr25 during M phase, and this event induces geminin stabilization by preventing its APC/C ubiquitin ligase complex-mediated degradation during mitosis. In turn, stabilized geminin inhibits SCF^{Skp2}-mediated degradation of Cdt1 to ensure pre-replicative complex formation in the ensuing S phase. The Aurora-A-geminin-Cdt1 axis therefore represents a critical regulator of proper DNA replication.

¹Department of Oral and Maxillofacial Pathobiology, Graduate School of Biomedical and Health Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan. ²Department of Stem Cell Biology, Research Institute for Radiation Biology and Medicine, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan. ³Department of Oral Molecular Pathology, Institute of Health Biosciences, University of Tokushima Graduate School, 3-18-15 Kuramoto, Tokushima 770-8504, Japan. ⁴Department of Pathology, New York University Cancer Institute, New York University School of Medicine, 522 First Avenue, Smilow Research Building 1107, New York, New York 10016, USA. ⁵Howard Hughes Medical Institute, New York, New York 10016, USA. Correspondence and requests for materials should be addressed to Y.K. (email: yasusei@tokushima-u.ac.jp).

by APC/C (Fig. 1b). To identify the phosphorylation site and further clarify the implication in geminin protein stability, we generated three phospho-mimicking mutants, geminin^{T25D}, geminin^{S32D} and geminin^{S60D}, in which serine/threonine was replaced with aspartic acid. Cells overexpressing either of these mutants were synchronized in M (noc 0h) and G1 (noc 4h). These mutants were highly expressed in M, but only geminin^{T25D} was abundant in G1 (Fig. 1c). Interestingly, Thr25 is a component of the D-box in geminin (Fig. 1b). We also generated a phospho-defective mutant (T25A) by replacing threonine with alanine. We examined the stability of these mutant molecules after the release from nocodazole (Noc) arrest. Geminin^{T25A} was already unstable during M, whereas geminin^{T25D} remained stable during G1 (Fig. 2a). We further confirmed this phenomenon through double

thymidine block release (Supplementary Fig. S2a). In G1 extracts, *in vitro*-translated geminin^{T25A} degraded faster than wild-type geminin, whereas geminin^{T25D} was stable (Fig. 2b). MG132 treatment increased wild-type geminin and geminin^{T25A}, but not geminin^{T25D}, protein levels (Supplementary Fig. S2b). These findings suggest that geminin is phosphorylated on Thr25 during mitosis and is degraded via APC/C^{Cdh1} in G1 phase after dephosphorylation. To demonstrate this hypothesis, we examined the effect of Thr25 phosphorylation on geminin degradation by APC/C^{Cdh1}. Cdh1 depletion increased wild-type geminin and geminin^{T25A}, but not geminin^{T25D}, protein levels (Supplementary Fig. S2c). As Emi1 depletion is known to give rise to ectopic APC/C^{Cdh1} activation⁵, we attempted to deplete Emi1, resulting in the reduction of wild-type geminin and geminin^{T25A}, but not




Figure 2 | Phosphorylation of geminin on Thr25 during mitosis prevents its proteasomal degradation by both APC/C^{Cdh1} and APC/C^{Cdc20}. (a) HeLa cells were transfected with either wild-type (WT) geminin, geminin^{T25A} or geminin^{T25D}, and synchronized in M-G1 by Noc. Cells were collected at 0, 2 and 4 h after mitotic shake-off. FLAG-tagged geminin, phospho-histone H3 (pHH3) and Cul1 expressions were examined by immunoblotting. (b) The half-life of IVT-myc-tagged geminin proteins (WT, T25A and T25D) were examined by the *in vitro* degradation assay. Cell extracts were obtained from synchronized HeLa cells in G1 (5 h after mitotic shake-off). Cell lysates with IVT-myc-tagged geminin protein were incubated for the indicated times. IVT-myc-tagged geminin protein expression was examined by immunoblotting with an anti-myc antibody. (c) An *in vitro* ubiquitylation assay was performed using ³⁵S-labelled IVT-geminin WT, geminin^{T25A} or geminin^{T25D}, and immunopurified APC/C^{Cdh1} from HeLa cells. ³⁵S-labelled IVT-geminin protein was incubated for the indicated times at 30 °C. The abundance of ³⁵S-labelled IVT-geminin proteins was detected by autoradiography. (d) FLAG-tagged geminin WT, geminin^{T25A} or geminin^{T25D}, with or without HA-tagged Cdc20 or Cdh1, were transfected into 293T cells. After 24 h, cells were treated with 25 μM MG132 for 7 h. Cell extracts were immunoprecipitated with an anti-HA antibody, and blotted with an anti-FLAG antibody and an anti-HA antibody. IP, immunoprecipitation; MW, molecular weight in kDa; WCE, whole cell extracts. (e) The half-life of IVT-myc-tagged geminin proteins (WT, T25A and T25D) was examined by the *in vitro* degradation assay. Cell extracts were obtained from synchronized HeLa cells in M (at mitotic shake-off). Cell lysates were incubated with IVT-myc-tagged geminin protein for the indicated times. IVT-myc-tagged geminin protein expression was examined by immunoblotting with an anti-myc antibody. (f) An *in vitro* ubiquitylation assay was performed using ³⁵S-labelled IVT-geminin WT, geminin^{T25A} or geminin^{T25D}, and immunopurified APC/C^{Cdc20} from HeLa cells. ³⁵S-labelled IVT-geminin proteins were incubated for the indicated times at 30 °C. The abundance of these proteins was detected by autoradiography. MW, molecular weight in kDa.

geminin^{T25D}, protein levels (Supplementary Fig. S2d). Moreover, the *in vitro* ubiquitylation of geminin^{T25D} via APC/C^{Cdh1} was not detectable, whereas that of wild-type geminin and geminin^{T25A} was observed (Fig. 2c). Interestingly, immunoprecipitation analysis revealed that HA-tagged Cdh1 interacted with wild-type geminin and geminin^{T25A}, but not with geminin^{T25D} (Fig. 2d), indicating that the inability of geminin^{T25D} to interact with APC/C^{Cdh1} may explain its resistance to APC/C^{Cdh1}-dependent proteolysis.

Geminin^{T25A} protein levels were lower than wild-type geminin protein levels at prometaphase (Fig. 2a), whereas *in vitro*-translated wild-type geminin, which was not phosphorylated on Thr25, and geminin^{T25A} were degraded in mitotic extracts (Fig. 2e), suggesting that phosphorylated geminin is protected from degradation via APC/C^{Cdc20}. As expected, the *in vitro* ubiquitylation of geminin^{T25D} via APC/C^{Cdc20} was not detectable (Fig. 2f). Moreover, HA-Cdc20 interacted with wild-type geminin and geminin^{T25A}, but not with geminin^{T25D} (Fig. 2d). Thus, phosphorylation of geminin on Thr25 also protects it from APC/C^{Cdc20}-mediated proteolysis in mitosis.

Next, to elucidate the physiological role of geminin phosphorylation on Thr25, we generated an antibody against geminin phosphorylated on Thr25. The λ-PPase treatment almost completely eliminated the recognition of geminin in mitotic extracts by this antibody (Supplementary Fig. S3a). Moreover, this antibody recognized the wild-type geminin, but not geminin^{T25A} (Supplementary Fig. S3b). We then used this antibody to probe for geminin phosphorylated on Thr25 at M and G1 phase. Thr25 phosphorylation of endogenous geminin was detectable exclusively in mitosis (Supplementary Fig. S3c).

Thr25 of geminin is phosphorylated by Aurora-A. To further deepen the understanding of geminin phosphorylation, we identified which kinase phosphorylates geminin on Thr25. We found that amino acids 23–26 (RRTL) within the D-box (which include Thr25) matched the consensus sequences (R-X-S/T-L/V) for phosphorylation by Aurora-A, a known mitotic kinase (Fig. 3a). The serine/threonine kinase Aurora-A is essential for mitotic entry, centrosome duplication, spindle formation, chromosome segregation and cytokinesis¹⁵. Geminin was co-localized with Aurora-A through mitotic progression (Fig. 3b and Supplementary Fig. S4a), but Aurora-A and phosphorylated geminin disappeared together in the G1 (Supplementary Fig. S4b). Interestingly, APC/C^{Cdh1} targets Aurora-A for the destruction at mitotic exit¹⁶.

We then investigated the direct interaction between geminin and Aurora-A. FLAG-tagged geminin co-immunoprecipitated with Xpress-tagged Aurora-A (Fig. 3c). Moreover, the endogenous interaction between Aurora-A and geminin was observed in mitosis, but not in interphase (Fig. 3d). Thus, Aurora-A exhibited mitosis-specific interaction with geminin. To determine the ability of Aurora-A to phosphorylate geminin, we performed an *in vitro* kinase assay. Aurora-A strongly phosphorylated purified wild-type glutathione S-transferase (GST)-geminin, but not GST-geminin^{T25A} (Fig. 3e). Moreover, a kinase-dead mutant (K162R) of Aurora-A displayed significantly reduced ability to phosphorylate geminin at Thr25, which further highlights the specificity of Aurora-A-mediated geminin phosphorylation (Fig. 3f). In fact, geminin phosphorylated on Thr25 co-localized with Aurora-A in mitotic cells (Supplementary Fig. S4b). Moreover, the depletion of Aurora-A impaired geminin phosphorylation (Fig. 3g). As shown in Fig. 2e, wild-type geminin and geminin^{T25A} were degraded in mitotic extracts, whereas wild-type geminin but not geminin^{T25A} was stabilized in mitotic extracts after *in vitro*

phosphorylation by Aurora-A (Fig. 3h). Moreover, a shorter half-life of geminin was observed in Noc-arrested cells after Aurora-A depletion (Supplementary Fig. S5). Cdc20 depletion stabilized geminin in mitotic Aurora-A-depleted cells (Supplementary Fig. S5), indicating that Aurora-A-mediated geminin phosphorylation prevents geminin degradation by APC/C^{Cdc20} at prometaphase.

Phosphorylated geminin stabilizes Cdt1 in mitosis. We then investigated the physiological role of geminin phosphorylation during mitosis. A previous report indicated that the inhibition of geminin synthesis during mitosis leads to impaired pre-RC formation and DNA replication in the ensuing cell cycle, and that geminin binds and protects Cdt1 from proteasome-dependent degradation to ensure the accumulation of Cdt1 in G₂ and M (ref. 3). Similarly, Cdt1 was reported to be rapidly downregulated after geminin was silenced in *Drosophila* cells¹⁷. As shown by previous studies^{3,18}, we confirmed that Cdt1 was hyperphosphorylated in mitosis and did not bind to chromatin (Supplementary Fig. S6). Moreover, dephosphorylation of Cdt1 coincided with chromatin association (Supplementary Fig. S6). A previous report revealed that the inhibition of CDK1 in mitosis induced an increase in the chromatin-associated Cdt1 fraction to the expense of the soluble Cdt1, and that geminin bound with Cdt1 interfered with this binding to the chromatin³. These findings suggest that Cdt1 recruitment to chromatin may be regulated by dephosphorylation in late M. Interestingly, geminin downregulation and Cdt1 recruitment to chromatin occur simultaneously (Supplementary Fig. S6). To understand the role of geminin in Cdt1 recruitment to chromatin, we examined whether mitotic depletion of geminin affected Cdt1 recruitment to chromatin (Fig. 4a). Consistent with that published before³, mitotic depletion of geminin decreased the levels of both soluble and chromatin-associated Cdt1 and impaired MCM6 loading (Fig. 4b). Interestingly, mitotic depletion of Aurora-A induced a similar decrease in the level of both soluble and chromatin-associated Cdt1 (Fig. 4c). These findings suggest that geminin phosphorylated by Aurora-A stabilizes soluble Cdt1, which, in turn, is recruited to the chromatin. In contrast to what was observed when geminin was depleted during interphase¹⁹, mitotic depletion of geminin or Aurora-A did not induce the DNA damage response, as shown by the lack of Chk2 phosphorylation (Supplementary Fig. S7). Moreover, the protein level and kinase activity of Aurora-A were not affected by the depletion of mitotic geminin (data not shown), and the phosphorylation of geminin on Thr25 did not affect its interaction with Cdt1 (Supplementary Fig. S8a).

Next, we examined whether introducing ectopic geminin^{T25D} rescued the downregulation of soluble Cdt1 protein at M phase in geminin- or Aurora-A-depleted cells. As we used siRNA targeting the 3'-untranslated region region to deplete geminin, exogenous geminin expression was not affected by this siRNA (Supplementary Fig. S8b). Interestingly, ectopic geminin^{T25D} in geminin- or Aurora-A-depleted cells induced a stabilization of soluble Cdt1 in M (Fig. 4b,c). As expected, the loading of Cdt1 and MCM in G1 was suppressed by ectopic geminin^{T25D} (Fig. 4b,c), indicating that phosphorylated geminin, which cannot be degraded via APC/C, prevents Cdt1 from loading onto chromatin during M. Indeed, ectopic Cdt1 rescued the impairment of MCM loading in geminin- or Aurora-A-depleted cells (Fig. 5a,b). Moreover, the depletion of geminin or Aurora-A in mitosis suppressed 5-bromodeoxyuridine (BrdU) incorporation in the ensuing S phase (Supplementary Fig. S9a,b and Fig. 5c,d), even though the kinetics of mitotic exit was not affected, as shown by the phosphorylation of histone H3 on Ser10




Figure 3 | Phosphorylation of geminin on Thr25 by Aurora-A kinase during mitosis. (a) Comparison of the amino acid sequences of human geminin from 22 to 27 with those of phosphorylation sites of other known Aurora-A substrates. (b) Subcellular co-localization of geminin and Aurora-A during mitosis. Expression of geminin (green) and Aurora-A (red) was visualized in HeLa cells by immunofluorescence analysis. Scale bar, 5 μ m. (c) FLAG-tagged geminin wild-type (WT) or empty vector was transfected into 293T cells with or without Xpress-tagged Aurora-A. Cell lysates were immunoprecipitated with an anti-Xpress antibody. IP, immunoprecipitation; MW, molecular weight in kDa; WCE, whole cell extracts. (d) Noc-arrested HeLa cells were used as arrested mitotic cells (at mitotic shake-off) and adherent cells after mitotic shake-off. Cell lysates were immunoprecipitated with an anti-geminin polyclonal antibody. Noc, arrested mitotic cells (at mitotic shake-off); Ad, adherent cells after mitotic shake-off; pHH, phospho-histone H3 (pHH3). (e) *In vitro* Aurora-A kinase assay using GST-tagged recombinant geminin protein (WT or T25A) and γ ³²-ATP. The phosphorylation was detected by autoradiography. (f) *In vivo* Aurora-A kinase assay using 293T cells transfected with Xpress-tagged Aurora-A WT or the kinase-dead mutant (K/R). After 24 h, cells were collected and lysed. Phosphorylation status was examined by immunoblotting with an anti-phosphorylated geminin Thr25-specific antibody (P-geminin^{Thr25}). (g) Levels of geminin phosphorylation on Thr25 in Aurora-A-depleted cells. 293T cells were co-transfected with FLAG-tagged geminin WT and control or Aurora-A siRNA. Expression of phospho-geminin (P-geminin^{Thr25}), FLAG-tagged geminin, Aurora-A and Cul1 was examined by immunoblotting. (h) The half-life of IVT-myc-tagged geminin proteins (WT and T25A) after Aurora-A-mediated phosphorylation was examined by an *in vitro* degradation assay. Cell extracts were obtained from synchronized HeLa cells in M phase (at mitotic shake-off). IVT-myc-tagged geminin protein (WT or T25A) was incubated with recombinant Aurora-A protein in kinase buffer at 30 °C for 1 h. Then, IVT-myc-tagged geminin protein was incubated with cell lysates for the indicated times. Expression of IVT-myc-tagged geminin protein was examined by immunoblotting with an anti-myc antibody. MW, molecular weight in kDa.




Figure 5 | Depletion of geminin or Aurora-A during mitosis impaired DNA replication. (a) Geminin depletion during mitosis impaired licensing at mitotic exit. U2OS cells were treated with geminin siRNA during release from the second thymidine block and collected at 0, 1 and 2 h after release from the Noc block as described in Fig. 4a. After release from the first thymidine block, cells were transfected with HA-tagged Cdt1. The soluble and chromatin fractions were separated and immunoblotted for the indicated proteins. MW, molecular weight in kDa; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; pHH, phospho-histone H3 (pHH3). (b) Aurora-A depletion during mitosis impaired licensing at mitotic exit. U2OS cells were treated with Aurora-A siRNA during release from the second thymidine block and collected at 0, 1 and 2 h after release from the Noc block as described in Fig. 4a. After release from the first thymidine block, cells were transfected with HA-tagged Cdt1. The soluble and chromatin fractions were separated and immunoblotted for the indicated proteins. MW, molecular weight in kDa. (c) U2OS cells plated on glass coverslips were synchronized as described in Fig. 4a and labelled with BrdU for 30 min at 10.5 h after release from the Noc block. Cells were then fixed and stained for BrdU (green) by indirect immunofluorescence analysis. Figure shows representative images obtained by indirect immunofluorescence analysis of the cells. The cell nuclei were stained with DAPI (4',6-diamidino-2-phenylindole; blue). Scale bar, 200 μ m. (d) Graph shows the average number of BrdU-positive cells in geminin or Aurora-A mitotic-depleted cells with or without HA-tagged Cdt1. Three independent counts ($n = 150$ cells each count) were used to calculate mean \pm s.d.




Figure 6 | Downregulation of Cdt1 by geminin or Aurora-A mitotic depletion is caused by SCF^{Skp2}. (a) U2OS cells were treated with geminin or Aurora-A siRNA during release from the second thymidine block and collected immediately after release from the Noc block as described in Fig. 4a. Cells were treated with 25 μ M MG132 for 5 h before mitotic shake-off. The soluble fraction was separated and immunoblotted for the indicated proteins. MW, molecular weight in kDa. (b) U2OS cells were treated with geminin or Aurora-A siRNA during release from the second thymidine block and collected immediately after release from the Noc block as described in Fig. 4a. Cells were treated with 1 μ M MLN4924 for 2 h before mitotic shake-off. The soluble fraction was separated and immunoblotted for the indicated proteins. (c) U2OS cells were treated with geminin siRNA, with or without Skp2 or Cdt2 siRNA, during release from the second thymidine block and collected immediately after release from the Noc block as described in Fig. 4a. The soluble fraction was separated and immunoblotted for the indicated proteins. MW, molecular weight in kDa.




Figure 7 | Geminin or Aurora-A depletion during mitosis induces SCF^{Skp2}-dependent proteolysis of soluble Cdt1 protein. (a) Geminin depletion during mitosis impaired licensing at mitotic exit. U2OS cells were treated with geminin and/or Skp2 siRNA during release from the second thymidine block and collected at 0 and 2 h after release from the Noc block as described in Fig. 4a. The soluble and chromatin fractions were separated and immunoblotted for the indicated proteins. MW, molecular weight in kDa; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; pHH, phospho-histone H3 (pHH3). (b) Aurora-A depletion during mitosis impaired licensing at mitotic exit. U2OS cells were treated with Aurora-A and/or Skp2 siRNA during release from the second thymidine block and collected at 0 and 2 h after release from the Noc block. Soluble and chromatin fractions were separated and immunoblotted for the indicated proteins. (c) Cells were treated with the geminin siRNA during release from second thymidine block and collected immediately after mitotic shake-off. After release from the first thymidine block, cells were transfected with 0.5, 1 or 2 μ g of FLAG-tagged geminin^{T25D}. Cdt1, FLAG-tagged geminin, Skp2 and Cul1 expression was examined by immunoblotting. (d) An *in vitro* ubiquitylation assay was performed using wild-type (WT) or T29A mutant IVT-HA-tagged Cdt1 and immunopurified SCF^{Skp2} from 293T cells. WT IVT-HA-tagged Cdt1 protein was incubated with or without IVT-myc-tagged geminin^{T25D} protein for the indicated times. As a negative control, IVT-HA-tagged Cdt1^{T29A} protein known as a non-degradable mutant for SCF^{Skp2}-mediated degradation was also used. The abundance of IVT-HA-tagged Cdt1 was detected by SDS-PAGE with a HA antibody. (e) Schematic model for pre-RC formation regulated by the Aurora-A-geminin-Cdt1 axis. Aurora-A protects geminin from APC/C^{Cdc20}- and APC/C^{Cdh1}-mediated proteolysis through phosphorylation. Phosphorylated geminin may stabilize soluble Cdt1 protein by inhibiting SCF^{Skp2}-mediated proteolysis. Stabilized soluble Cdt1 may load onto chromatin after dephosphorylation and then form the pre-RC. MW, molecular weight in kDa.

Geminin restricts nuclear DNA replication to one round per cell cycle, which ensures genome stability and prevents cells from becoming aneuploid¹. Re-replication may occur when the pathways controlling pre-RC formation are impaired. Indeed, geminin depletion in mammalian cells is sufficient to cause re-replication^{19,31}, indicating that geminin has a critical role in replication control. The pre-RC assembly reaction, known as licensing, involves the loading of the Mcm2-7 complex in an ATP-dependent reaction that requires ORC and two essential factors, Cdc6 and Cdt1 (ref. 1). It is suggested that the binding of geminin to Cdt1 protects the latter from ubiquitylation and subsequent proteolysis³. In this study, we found that introduction of *in vitro*-translated geminin^{T25D} protein suppressed SCF^{Skp2}-dependent Cdt1 ubiquitylation (Fig. 7d), suggesting that geminin protects soluble Cdt1 from SCF^{Skp2}-dependent proteolysis by interfering with the binding between Cdt1 and Skp2. However, as

Cdt2 depletion slightly increased soluble Cdt1 protein levels (Fig. 6c), the possibility of Cdt1 degradation by CRL4^{Cdt2} in mitotic geminin-depleted cells cannot be denied. Further detailed studies on the mechanism of inhibition of Cdt1 ubiquitylation may be required.

In *Xenopus* eggs, geminin is suggested to inhibit licensing from S phase to metaphase and seems to be inactivated in interphase through activation of the licensing process^{8,32–34}. However, a significant proportion of molecules seems to escape degradation at the metaphase–anaphase transition in *Xenopus* egg extracts³⁵, and the Cdt1-inhibiting activity of these remaining molecules must be suppressed through mechanisms other than by proteolysis. Indeed, exogenous geminin associates with and inhibits Cdt1 in the interphase egg extracts^{4,8,35}. Although geminin rapidly disappears at the exit of mitosis and accumulates during S phase in cultured cells^{2,4}, the remaining

pool of geminin in the egg extracts may be regulated by Aurora-A-mediated phosphorylation through escape from APC/C-mediated degradation. It is interesting to examine the precise molecular mechanism of this phenomenon in *Xenopus* eggs.

In S phase, chromatin-associated geminin presents even in the absence of Cdt1, possibly as a precaution to prevent Cdt1 from gaining access to chromatin if it accidentally accumulates during S phase³⁶. In this situation, geminin is phosphorylated at Ser45 and Ser49 in HeLa cells³⁷, although the physiological relevance of this modification remains to be clarified. Indeed, we identified geminin phosphorylation on Thr25 by Aurora-A kinase during mitosis. In Fig. 1c, the band of geminin^{T25D} was still shifted at G1 similar to that of wild-type geminin and other geminin mutants (geminin^{S32D} and geminin^{S60D}), suggesting that geminin is phosphorylated at multiple sites during mitosis. These findings indicate that phosphorylation may be important for the functional regulation of geminin in mammalian cells. Furthermore, we found that abundant soluble geminin and Cdt1 protein expression was observed at prometaphase, and that abundant chromatin-associated Cdt1 expression was observed during the G1 phase. Although it is still unclear how the molecular interaction between Cdt1 and geminin can be regulated during M to G1 in mammalian cells, recent studies revealed that (i) the geminin-Cdt1 complex can exist in two distinct forms, a licensing-permissive heterotrimer and a licensing-inhibitory heterohexamer, to regulate a molecular switch between licensing-competent and licensing-defective states in *Xenopus* and mammalian cells³⁸ and (ii) the two forms of the Cdt1-geminin complex act as active or inactive complexes for licensing in *Xenopus* egg extracts³⁹. In addition, geminin-mediated nuclear Cdt1 foci formation supports the presence of local interactions between geminin-Cdt1 complexes on chromatin in *Xenopus* egg extracts⁴⁰. Ode *et al.*⁴⁰ suggested that the higher-order interactions of Cdt1-geminin complexes are cooperatively stabilized on chromatin. Overall, these findings suggest that licensing for proper DNA replication may be strictly regulated by the geminin protein level during cell cycle progression. The amount of geminin, which is determined by Emi1- and Aurora-A-mediated protection from APC/C-mediated proteolysis, may thus be essential for proper DNA replication.

Methods

Reagents and antibodies. Reagents were from the following suppliers: MG132 (Z-Leu-Leu-Leu-CHO) (Peptide Institute Inc., Osaka, Japan); CHX (Sigma); λ -PPase (New England Bio Labs); MLN4924 (Millennium Pharmaceuticals). The geminin phospho-Thr25-specific antibody was generated by immunizing rabbits with the synthetic peptide CVPRR*TLKMIQ, corresponding to amino acids 21–30 of the human geminin sequence, with a phospho-threonine at position 25 (*T). The antibody was purified from serum by two rounds of affinity chromatography, on a phospho-Thr25 peptide column and on a non-phosphopeptide column. The anti-geminin polyclonal antibody was generated using GST fusion recombinant molecules⁴¹, and the anti-Cdt1 monoclonal antibody was a gift from Dr Kristian Helin (University of Copenhagen). Commercial antibodies were from the following suppliers: anti-p27, anti-Aurora-A and anti-Aurora-B antibodies (Transduction Laboratories); anti-HA (Y-11), anti-cyclin A, anti-cyclin B, anti-MCM6, anti-Cdc25A, anti-Cdc6 and anti-GAPDH antibodies (Santa-Cruz Biotechnology); anti-phospho-Histone H3 (Ser10) antibody (Upstate); anti-Cul1, anti-Emi1, anti-Skp2 and anti-PLK1 antibodies (Zymed); anti-Xpress antibody (Invitrogen); anti-Cdh1 antibody (MBL); anti-geminin (2H7) antibody (Abnova); anti-histone H2A antibody (Abcam); anti-FLAG (M2) and anti- β -actin antibodies (Sigma); anti-phospho-tyrosine antibody (Merck Millipore); anti-Cdt1 and anti-Cdt2 antibodies (Bethyl).

Plasmids. pEF-FLAG-tagged geminin and pcDNA-HA-tagged Cdt1 were kind gifts from Dr Anindya Dutta (University of Virginia). To produce GST-tagged recombinant proteins, geminin complementary DNA was subcloned into pGEX 4T-1 vectors (GE Healthcare). For *in vitro* translation, geminin cDNA was also subcloned into pcDNA3.1 His/myc-tagged vectors (Invitrogen). The pcDNA3.1 His/Xpress-tagged Aurora-A was previously described¹⁶, pCMV HA-tagged Cdt1 and HA-tagged Cdh1 expression vectors were gifts from Dr Kristian Helin

(University of Copenhagen). The pEF, pGEX and pcDNA3.1 vectors encoding the FLAG-tagged geminin substitution mutants (T25A and T25D), the pcDNA3.1 vector encoding His/Xpress-tagged Aurora-A kinase-dead mutant (K162R) and the pcDNA3 vector encoding HA-tagged Cdt1 substitution mutant (T29A) were generated using a KOD plus mutagenesis kit (Toyobo).

Cell culture and transient transfection. HeLa, U2OS and 293T cells were routinely maintained in DMEM (Nissui Pharmaceutical Co. Ltd, Tokyo, Japan) supplemented with 10% heat-inactivated fetal bovine serum (Gibco) and 100 U ml⁻¹ penicillin-streptomycin (Gibco) at 37 °C and 5% CO₂ in air. For experiments, cells were grown to subconfluence in this medium. Transient transfections were performed using FuGENE HD (Roche) following the manufacturer's instructions. For synchronization, Noc (Sigma) and thymidine (Sigma) were used. HeLa cells were synchronized at prometaphase using 50 ng ml⁻¹ Noc treatment followed by mitotic shake-off. Subsequently, synchronized cells were released from the mitotic arrest by washing out Noc. Adherent cells remaining after mitotic shake-off were also used. Adherent cells dominantly included the cells arrested at G₂, as confirmed by FACS analysis (data not shown). U2OS were synchronized at the G1/S border by a double thymidine block. In brief, cells were incubated for 20 h in 2 mM thymidine (Sigma), released for 8 h and then incubated for another 20 h in 2 mM thymidine.

Immunoprecipitation and immunoblot analysis. Cells were transfected with vectors by using FuGENE HD (Roche). Cell lysis and immunoprecipitation were performed as described¹³. Twenty micrograms of protein were subjected to 10% PAGE followed by electroblotting onto a nitrocellulose membrane. To detect the immunocomplex, an ECL Western blotting detection system (Amersham) was used. The immunoprecipitates were subjected to immunoblot analysis. To detect phospho-Thr25 geminin, we performed immunoprecipitation with a phospho-specific antibody against Thr25 of geminin, followed by immunoblotting analysis with a monoclonal antibody against endogenous geminin or an anti-FLAG antibody against FLAG-tagged exogenous geminin.

In vitro ubiquitylation assay. Ubiquitylation assays were performed as previously described¹⁶. [³⁵S]methionine-labelled human geminin protein was prepared by coupled transcription-translation reactions in wheat germ extract (Promega). GST-tagged Cdh1 (Abnova) or cold *in vitro*-translated human Cdc20 protein was also used. The extracts from HeLa cells were immunoprecipitated with an anti-Cdc27 antibody (Sigma). Immunoprecipitates were incubated with Cdh1 or Cdc20 for 1 h at 37 °C and washed four times in QA buffer (10 mM Tris-HCl pH 7.5, 100 mM KCl, 1 mM MgCl₂, 0.1 mM CaCl₂, 1 mM dithiothreitol (DTT)). The purified APC/C^{Cdh1} or APC/C^{Cdc20} complexes were incubated in reaction mixtures containing the following components in a volume of 7 μ l: 40 mM Tris-HCl (pH 7.6), 1 mg ml⁻¹ carboxymethyl BSA, 1 mM DTT, 5 mM MgCl₂, 10 mM phosphocreatine, 50 μ g ml⁻¹ creatine phosphokinase, 0.5 mM ATP, 50 μ M ubiquitin, 1 μ M ubiquitin aldehyde, 1 pmol of E1, 5 pmol of E2-C, 1 μ M okadaic acid and 1–2 pmol of [³⁵S]methionine-labelled human geminin protein. Following incubation at 30 °C for 2 h, samples were loaded for electrophoresis on an 8% polyacrylamide-SDS gel.

In vitro ubiquitylation assays for Cdt1 were performed in a volume of 30 μ l containing 0.1 μ M E1 (Boston Biochem), 10 ng ml⁻¹ Ubch3, 10 ng ml⁻¹ Ubch5c, 1 μ M ubiquitin aldehyde, 2.5 μ g ml⁻¹ ubiquitin (Sigma) and SCF^{Skp2} complex in a ubiquitylation buffer (50 mM Tris (pH 7.6), 2 mM ATP, 5 mM MgCl₂, 0.6 mM DTT, 0.1 μ M okadaic acid). *In vitro*-translated HA-tagged wild-type Cdt1 or Cdt1^{T29A} was used as the substrate. The reactions were incubated at 30 °C for the indicated times and analysed by SDS-PAGE with an anti-HA antibody. The SCF^{Skp2} complex was obtained from the lysates of 293T cells after transfection with FLAG-tagged Skp2 together with other SCF components. After 24 h of transfection, cells were incubated with MG132 for 3 h before lysis. Anti-FLAG antibody was used to immunoprecipitate the SCF^{Skp2} complex. The immunoprecipitates were washed four times in lysis buffer and two times in ubiquitylation reaction buffer (10 mM Tris-HCl pH 7.5, 100 mM NaCl, 5 mM MgCl₂ and 1 mM DTT). Finally, beads were then used for *in vitro* ubiquitylation assays.

Chromatin fractionation. For chromatin fractionation, we separated the soluble and insoluble fractions as previously described⁴². Briefly, cells were lysed in CSK buffer and centrifuged for 4 min at 3,200 r.p.m. After centrifugation, the supernatant contained the soluble fraction. The pellet was resuspended in 2 \times Laemmli buffer after washing with CSK buffer and then sonicated. This lysate contained the chromatin-enriched fraction.

RNA interference. The following siRNA oligonucleotides were obtained from Qiagen: geminin, 5'-CCUUAUGCAUUAAGUACA-3'; Aurora-A, 5'-CACCUUCGGCAUCCUAAUA-3'; and Cdh1, 5'-UCUGGUGGACUGGUCGUCC-3'. Emi1 siRNA was obtained from Dharmacon as the following sequence: 5'-GAUUGUGAUCUCUUAUUA-3'. Cdt2 siRNA was obtained from Bonac (Kurume, Japan) as the following sequence: 5'-CAAUGGACACCAGAACUCUACUUU-3'. The negative control (siTrio-negative control siRNA) was obtained from Cosmo

Bio Inc. (Tokyo, Japan). Transfections were performed using 20 nM siRNA and Oligofectamine RNAi MAX (Invitrogen), according to the manufacturer's instructions.

In vitro degradation assay. For M or G1 extracts, HeLa cells were synchronized in prometaphase by Noc treatment and collected at 0 or 5 h after mitotic shake-off. The collected cells were lysed in hypotonic buffer (25 mM HEPES pH 7.5, 1.5 mM MgCl₂, 5 mM KCl, 1 mM DTT, 15 mM creatine phosphate, 2 mM ATP) by ten freeze-and-thaw cycles and centrifuged for 5 min at 5,000 r.p.m. and subsequently for 60 min at 13,200 r.p.m. As a substrate, cold IVT-myc-geminin (wild-type, T25A and T25D) was prepared by coupled transcription-translation reactions in wheat germ extracts (Promega). The substrate was incubated in 20 µl of cell extract and a degradation cocktail containing the following components in a volume of 20 µl: 1.5 mg ml⁻¹ ubiquitin, 7.5 mM creatine phosphate, 1 mM ATP, 1 mM MgCl₂ and 25 µg ml⁻¹ CHX. Following incubation at 30 °C for 2, 4 and 6 h, samples were loaded for electrophoresis on a 12% polyacrylamide-SDS gel.

Immunofluorescence. Cells were grown on glass coverslips, fixed in 4% paraformaldehyde for 10 min and permeabilized with PBS containing 0.1% Triton X-100 for 20 min. After three rinses with PBS, the coverslips were incubated with the primary antibodies. Anti-geminin, anti-geminin phospho-Thr25 and anti-Aurora-A antibodies were used. Alexa Fluor 488-conjugated anti-mouse or rabbit IgG and Alexa Fluor 594-conjugated anti-mouse or rabbit IgG were used as secondary antibodies (Invitrogen). DNA was visualized by DAPI (4',6-diamidino-2-phenylindole) staining. Immunostaining of the cell preparations was recorded by using an epifluorescence Zeiss Axioplan 2 microscope (Zeiss, Inc., Thornwood, NY) attached to a charge-coupled-device camera.

In vitro kinase assay. GST-tagged wild-type and geminin^{T25A} mutant geminin were subcloned into pGEX, expressed in *E. coli* and purified. Aurora-A kinase was obtained from Millipore. Purified GST-tagged wild-type geminin or geminin^{T25A} was incubated with Aurora-A in a kinase reaction buffer (25 mM Tris-HCl pH 7.5, 5 mM β-glycerophosphate, 2 mM DTT, 0.1 mM Na₃VO₄, 10 mM MgCl₂ and 0.1 mM ATP) supplemented with 5 µCi of [³²P] ATP at 30 °C for 60 min. The reactions were stopped with Laemmli buffer for 5 min at 95 °C. After the reaction mixtures were resolved by SDS-PAGE, the gels were stained with Coomassie brilliant blue, and were dried and subjected to autoradiography.

BrdU staining. Cells were treated with 33 µM BrdU for 30 min at 10.5 h after double thymidine block release (Fig. 4a). Cells were then fixed with 70% ethanol containing 10 mM glycine for 10 min, stained with mouse anti-BrdU primary antibody (Roche) and fluorescein-conjugated anti-mouse IgG secondary antibody (Roche), and analysed by indirect immunofluorescence. Nuclei were simultaneously counterstained with DAPI. The BrdU-positive nuclei were counted and plotted. Three independent counts (*n* = 150 cells each count) were used to calculate mean ± s.d.

References

- Dutta, A. & Bell, S. P. Initiation of DNA replication in eukaryotic cells. *Annu. Rev. Cell Dev. Biol.* **13**, 293–332 (2002).
- Wohlschlegel, J. A. *et al.* Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Science* **290**, 2309–2312 (2000).
- Ballabeni, A. *et al.* Human geminin promotes pre-RC formation and DNA replication by stabilizing CDT1 in mitosis. *EMBO J.* **23**, 3122–3132 (2004).
- McGarry, T. J. & Kirschner, M. W. Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell* **93**, 1043–1053 (1998).
- Di Fiore, B. & Pines, J. Emi1 is needed to couple DNA replication with mitosis but does not regulate activation of the mitotic APC/C. *J. Cell Biol.* **177**, 425–437 (2007).
- Machida, Y. J. & Dutta, A. The APC/C inhibitor, Emi1, is essential for prevention of rereplication. *Genes Dev.* **21**, 184–194 (2007).
- Peters, J. M. The anaphase promoting complex/cyclosome: a machine designed to destroy. *Nat. Rev. Mol. Cell Biol.* **7**, 644–656 (2006).
- Tada, S., Li, A., Maiorano, D., Mechali, M. & Blow, J. J. Repression of origin assembly in metaphase depends on inhibition of RLF-B/Cdt1 by geminin. *Nat. Cell Biol.* **3**, 107–113 (2001).
- Listovsky, T. *et al.* Mammalian Cdh1/Fzr mediates its own degradation. *EMBO J.* **23**, 1619–1626 (2004).
- Rape, M. & Kirschner, M. W. Autonomous regulation of the anaphase-promoting complex couples mitosis to S-phase entry. *Nature* **432**, 588–595 (2004).
- Pefani, D. E. *et al.* Idas, a novel phylogenetically conserved geminin-related protein, binds to geminin and is required for cell cycle progression. *J. Biol. Chem.* **286**, 23234–23246 (2011).
- Hsu, J. Y., Reimann, J. D., Sørensen, C. S., Lukas, J. & Jackson, P. K. E2F-dependent accumulation of hEmi1 regulates S phase entry by inhibiting APC^{Cdh1}. *Nat. Cell Biol.* **4**, 358–366 (2002).
- Guardavaccaro, D. *et al.* Control of meiotic and mitotic progression by the F box protein β-Trcp1 in vivo. *Dev. Cell* **4**, 799–812 (2003).
- Margottin-Goguet, F. *et al.* Prophase destruction of Emi1 by the SCF^{βTrCP/Slimb} ubiquitin ligase activates the anaphase promoting complex to allow progression beyond prometaphase. *Dev. Cell* **4**, 813–826 (2003).
- Marumoto, T., Zhang, D. & Saya, H. Aurora-A—a Guardian of poles. *Nat. Rev. Cancer* **5**, 42–50 (2005).
- Kitajima, S. *et al.* Constitutive phosphorylation of aurora-a on ser51 induces its stabilization and consequent overexpression in cancer. *PLoS ONE* **2**, e944 (2007).
- Mihaylov, I. S. *et al.* Control of DNA replication and chromosome ploidy by geminin and cyclin A. *Mol. Cell Biol.* **22**, 1868–1880 (2002).
- Nishitani, H., Taraviras, S., Lygerou, Z. & Nishimoto, T. The human licensing factor for DNA replication Cdt1 accumulates in G1 and is destabilized after initiation of S-phase. *J. Biol. Chem.* **276**, 44905–44911 (2001).
- Zhu, W., Chen, Y. & Dutta, A. Rereplication by depletion of geminin is seen regardless of p53 status and activates a G2/M checkpoint. *Mol. Cell Biol.* **24**, 7140–7150 (2004).
- Nishitani, H. *et al.* Two E3 ubiquitin ligases, SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis. *EMBO J.* **25**, 1126–1136 (2006).
- Hall, J. R. *et al.* Cdt1 and Cdc6 are destabilized by rereplication-induced DNA damage. *J. Biol. Chem.* **283**, 25356–25363 (2008).
- Sugimoto, N. *et al.* Cdt1 phosphorylation by cyclin A-dependent kinases negatively regulates its function without affecting geminin binding. *J. Biol. Chem.* **279**, 19691–19697 (2004).
- Mailand, N. & Diffley, J. F. CDKs promote DNA replication origin licensing in human cells by protecting Cdc6 from APC/C-dependent proteolysis. *Cell* **122**, 915–926 (2005).
- Lin, H. K. *et al.* Phosphorylation-dependent regulation of cytosolic localization and oncogenic function of Skp2 by Akt/PKB. *Nat. Cell Biol.* **11**, 420–432 (2009).
- Rodier, G., Coulombe, P., Tanguay, P. L., Boutonnet, C. & Meloche, S. Phosphorylation of Skp2 regulated by CDK2 and Cdc14B protects it from degradation by APC(Cdh1) in G1 phase. *EMBO J.* **27**, 679–691 (2008).
- Choi, E. *et al.* BubR1 acetylation at prometaphase is required for modulating APC/C activity and timing of mitosis. *EMBO J.* **28**, 2077–2089 (2009).
- Rape, M., Reddy, S. K. & Kirschner, M. W. The processivity of multiubiquitination by the APC determines the order of substrate degradation. *Cell* **124**, 89–103 (2006).
- Wolthuis, R. *et al.* Cdc20 and Cks direct the spindle checkpoint-independent destruction of cyclin A. *Mol. Cell* **30**, 290–302 (2008).
- Montanari, M. *et al.* Increased expression of geminin stimulates the growth of mammary epithelial cells and is a frequent event in human tumors. *J. Cell. Physiol.* **202**, 215–222 (2005).
- Bravou, V., Nishitani, H., Song, S. Y., Taraviras, S. & Varakis, J. Expression of the licensing factors, Cdt1 and Geminin, in human colon cancer. *Int. J. Oncol.* **27**, 1511–1518 (2005).
- Melixetian, M. *et al.* Loss of Geminin induces rereplication in the presence of functional p53. *J. Cell Biol.* **165**, 473–482 (2004).
- Blow, J. J. Preventing re-replication of DNA in a single cell cycle: evidence for a replication licensing factor. *J. Cell Biol.* **122**, 993–1002 (1993).
- Hodgson, B., Li, A., Tada, S. & Blow, J. J. Geminin becomes activated as an inhibitor of Cdt1/RLF-B following nuclear import. *Curr. Biol.* **12**, 678–683 (2002).
- Mahbubani, H. M., Chong, J. P., Chevalier, S., Thommes, P. & Blow, J. J. Cell cycle regulation of the replication licensing system: involvement of a Cdk-dependent inhibitor. *J. Cell Biol.* **136**, 125–135 (1997).
- Li, A. & Blow, J. J. Non-proteolytic inactivation of geminin requires CDK-dependent ubiquitination. *Nat. Cell Biol.* **6**, 260–267 (2004).
- Kulartz, M. & Knippers, R. The replicative regulator protein geminin on chromatin in the HeLa cell cycle. *J. Biol. Chem.* **279**, 41686–41694 (2004).
- Kulartz, M. *et al.* Expression and phosphorylation of the replication regulator protein geminin. *Biochem. Biophys. Res. Commun.* **305**, 412–420 (2003).
- De Marco, V. *et al.* Quaternary structure of the human Cdt1-Geminin complex regulates DNA replication licensing. *Proc. Natl Acad. Sci. USA* **106**, 19807–19812 (2009).
- Lutzmann, M., Maiorano, D. & Mechali, M. A Cdt1-geminin complex licenses chromatin for DNA replication and prevents rereplication during S phase in *Xenopus*. *EMBO J.* **25**, 5764–5774 (2006).
- Ode, K. L., Fujimoto, K., Kubota, Y. & Takisawa, H. Inter-origin cooperativity of geminin action establishes an all-or-none switch for replication origin licensing. *Genes Cells* **16**, 380–396 (2011).
- Ohtsubo, M. *et al.* Polycomb-group complex 1 acts as an E3 ubiquitin ligase for Geminin to sustain hematopoietic stem cell activity. *Proc. Natl Acad. Sci. USA* **105**, 10396–10401 (2010).

42. Bassermann, F. *et al.* The Cdc14B-Cdh1-Plk1 axis controls the G2 DNA damage response checkpoint. *Cell* **134**, 256–267 (2008).

Acknowledgements

We thank Drs Anindya Dutta (University of Virginia) and Hidehiko Kawai (Hiroshima University) for helpful discussions and Dr Rieko Arakaki (Tokushima University) for technical advice. We thank Dr Kristian Helin (University of Copenhagen) and Dr Anindya Dutta (University of Virginia) for providing materials. This work was supported by the Ministry of Education, Science and Culture of Japan grants-in-aid to Y.K. and T.Ta., and by a grant-in-aid for JSPS Fellows to T.Ts.

Author contributions

T.Ts. performed most of the experiments. Y.K. and M.P. designed the experiments. Y.K., M.P. and Y.T. wrote the manuscript. All authors discussed the experiments and provided suggestions for the manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Tsunematsu, T. *et al.* Aurora-A controls pre-replicative complex assembly and DNA replication by stabilizing geminin in mitosis. *Nat. Commun.* **4**:1885 doi: 10.1038/ncomms2859 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>