

MAPT p.V363I mutation

A rare cause of corticobasal degeneration

Sarah Ahmed, BS, Monica Diez Fairen, MS, Marya S. Sabir, BS, Pau Pastor, MD, PhD, Jinhui Ding, PhD, Lourdes Ispierto, MD, Ankur Butala, MD, Christopher M. Morris, PhD, Claudia Schulte, PhD, Thomas Gasser, MD, Edwin Jabbari, MD, Olga Pletnikova, MD, Huw R. Morris, MD, PhD, Juan Troncoso, MD, Ellen Gelpi, MD, PhD, Alexander Pantelyat, MD, and Sonja W. Scholz, MD, PhD

Correspondence
Dr. Scholz
sonja.scholz@nih.gov

Neurol Genet 2019;5:e347. doi:10.1212/NXG.0000000000000347

Abstract

Objective

Patients with corticobasal syndrome (CBS) present with heterogeneous clinical features, including asymmetric parkinsonism, dyspraxia, aphasia, and cognitive impairment; to better understand the genetic etiology of this rare disease, we undertook a genetic analysis of microtubule-associated protein tau (*MAPT*).

Methods

We performed a genetic evaluation of *MAPT* mutations in 826 neurologically healthy controls and 173 cases with CBS using the Illumina NeuroChip genotyping array.

Results

We identified 2 patients with CBS heterozygous for a rare mutation in *MAPT* (p.V363I) that is located in the highly conserved microtubule-binding domain. One patient was pathologically confirmed and demonstrated extensive 4-repeat-tau-positive thread pathology, achromatic neurons, and astrocytic plaques consistent with corticobasal degeneration (CBD).

Conclusions

We report 2 CBS cases carrying the rare p.V363I *MAPT* mutation, one of which was pathologically confirmed as CBD. Our findings support the notion that this rare coding change is pathogenic.

From the Neurodegenerative Disease Research Unit (S.A., M.S.S., S.W.S.), National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD; Movement Disorders and Memory Unit (M.D.F., P.P.), Department of Neurology, University Hospital Mutua de Terrassa, and Fundació per la Recerca Biomèdica i Social Mútua Terrassa, Barcelona, Spain; Laboratory of Neurogenetics (J.D.), National Institutes on Aging, National Institutes of Health, Bethesda, MD; Neurology Service (L.I.), Hospital Universitari Germans Trias i Pujol, Badalona, Spain; Department of Neurology (A.B., A.P., S.W.S.), Johns Hopkins University Medical Center, Baltimore, MD; Newcastle Institute for Ageing (C.M.M.), Campus for Ageing and Vitality, Newcastle University, Newcastle upon Tyne, United Kingdom; Department of Neurodegenerative Diseases (C.S., T.G.), Center of Neurology, Hertie-Institute for Clinical Brain Research, University of Tuebingen, and German Center for Neurodegenerative Diseases, Germany; Department of Molecular and Clinical Neuroscience (E.J., H.R.M.), Institute of Neurology, University College London, United Kingdom; Department of Pathology (Neuropathology) (O.P., J.T.), Johns Hopkins University Medical Center, Baltimore, MD; Department of Clinical Neurosciences (H.R.M.), Royal Free Campus UCL, Institute of Neurology, London, United Kingdom; Neurological Tissue Bank (E.G.), University of Barcelona-Hospital Clinic, IDIBAPS, Barcelona, Spain; and Institute of Neurology (E.G.), Medical University of Vienna, Austria.

Go to Neurology.org/NG for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the NIH.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

CBD = corticobasal degeneration; **CBS** = corticobasal syndrome; **FTD** = frontotemporal dementia; **MAPT** = microtubule-associated protein tau; **PGRN** = progranulin; **PSP** = progressive supranuclear palsy.

Corticobasal syndrome (CBS) is a rare neurologic disease that presents with heterogeneous motor symptoms and cognitive impairment.¹ A high misdiagnosis rate due to clinical heterogeneity limits efforts to extend disease-modifying therapy trials to this patient population. Improving the diagnostic accuracy of complex neurodegenerative syndromes is an important, yet unmet need in the research community.

Although understanding of the genetic underpinnings of CBS is limited, rare mutations in the microtubule-associated protein tau (*MAPT*) gene are implicated as a cause of CBS and related tauopathy spectrum disorders.^{2–6} One of these *MAPT* mutations is the variant p.V363I (rs63750869; c.1087G>A; NM_005910.5), located in the *MAPT* microtubule-binding domain. Previously described in a small number of patients with clinical tauopathy phenotypes (table 1), the mutation is present at a very low frequency in population databases and is hypothesized to be a disease-causing mutation with decreased penetrance rather than a rare polymorphism.^{6–10} The rare nature of the mutation makes it difficult to demonstrate disease segregation, and in silico prediction algorithms of this mutation are inconclusive (table e-1, links.lww.com/NXG/A162).

We describe 2 CBS cases who were found to carry the rare p.V363I *MAPT* mutation. In addition, we summarize the clinicopathologic features of previously reported cases with a coding mutation at the *MAPT* p.V363 residue. One of our CBS cases had postmortem confirmation, which found abundant four-repeat tau accumulations consistent with corticobasal degeneration (CBD). As a pathologically confirmed case with this rare missense mutation, this case provides supporting evidence for the pathogenic nature of the p.V363I *MAPT* mutation.

Methods

Study population

Case 1 is a 73-year-old, right-handed, white woman who presented to the NIH Clinical Center in Bethesda, MD, for participation in genetic research. She was diagnosed with probable CBS based on the consensus criteria.¹¹ A commercial genetic panel (Invitae, San Francisco, CA) that included screening of the genes *CHCHD10*, *DCTN*, *FUS*, *GRN*, *TARDBP*, *VCP*, *UBQLN2*, *TBK1*, *PSEN1*, *PSEN2*, *APP*, and *MAPT* had previously identified that she was a carrier of the *MAPT* p.V363I variant. Case 2 was identified by querying a research database for the presence of the *MAPT* p.V363I variant. This database contains genotype information on European-ancestry individuals, including 826 neurologically healthy controls and 961 patients with frontotemporal

dementia (FTD) spectrum disorders (n = 772 cases with progressive supranuclear palsy [PSP], n = 173 patients with CBS/CBD, n = 41 patients with FTD; sample characteristics are summarized in table e-2 [links.lww.com/NXG/A162]; source of samples and number of samples per disease are described in table e-3). Case 2 was clinically diagnosed with hemiparkinsonism, primary progressive aphasia, and probable CBS.

Standard protocol approvals, registrations, and patient consents

The study was approved by the respective institutional review boards. Written informed consent for research participation was obtained from all participants.

Genetic analysis and validation

For each participant, DNA was extracted from blood or brain tissue using standard methods and followed by genotyping on the NeuroChip platform (Illumina, San Diego, CA). This affordable genotyping array contains a tagging single nucleotide polymorphism backbone combined with high-yield custom content that allows for rapid screening of ~180,000 mutations and risk variants previously implicated in neurologic diseases, including the *MAPT* p.V363I variant. The detailed contents of this versatile genotyping platform have been described elsewhere.¹² The *MAPT* p.V363I mutation was only present in 2 patients (henceforth referred to as case 1 and case 2), and we validated the mutation via direct Sanger sequencing using the following primers: forward 5'-GTGGCCAGGTGGAAGTAAAA, reverse 5'-ACATC-CAGCCAGTCAACACA. To rule out other possible pathogenic mutations in these 2 patients, we assessed the NeuroChip data for damaging progranulin (*PGRN*) gene mutations. We also performed repeat-primed PCR screening of the *C9orf72* repeat using methods described elsewhere.¹³ *APOE* genotypes were determined by extracting rs7412 and rs429358 as previously described.¹² *MAPT* haplotype status was determined by imputation of the polymorphism rs1052553 ($R^2 = 0.99494$), with the "A" allele determining the H1 haplotype and the "G" allele segregating with the H2 haplotype.¹⁴

Bioinformatic analysis


To better understand the effects of the p.V363I variant, a systematic literature review was conducted and summarized in table 1. In silico predictive tools (SIFT, PolyPhen2, FATHMM-XF, M-CAP, MutationTaster, CADD, ClinVar, and ClinPred) were applied to classify the *MAPT* p.V363I mutation.^{15–21} Sequence conservation analyses were performed in T-Coffee.²² A previously described, cryo-electron microscopy structure of the tau protofibril was used for 3-dimensional protein modeling (figure 1).²³ Allele frequency differences between CBS cases and neurologically healthy

Table 1 Clinicopathologic features of patients with a mutation at the highly conserved p.V363 residue of *MAPT*

No.	Clinical features						Genetics				
	Clinical diagnosis	Sex	AAO	AAD	FH	Neuroimaging finding(s)	Mutation	Haplotype	Pathology	Country	Reference
1	CBS	F	70	NA	+	MRI: bilateral parietal atrophy	p.V363I	H1/H1	NA	United States	This report
2	CBS and PPA	F	Late 50s	62	-	NA	p.V363I	H1/H1	CBD	Spain	This report
3	PPA (nonfluent variant)	F	69	NA	+	SPECT: bilateral Sylvian hypoperfusion	p.V363I	H1/H1	NA	Spain	Munoz et al. ⁹
4	FTD (behavioral variant)	F	53	61	+	MRI: bilateral frontotemporal atrophy	p.V363I	H1/H1	NA	Italy	Anfossi et al. ⁸
5	PPA (semantic variant)	F	46	NA	-	MRI: asymmetric temporopolar atrophy	p.V363I	NA	NA	Italy	Bessi et al. ⁶
6	FTD and PPA (nonfluent variant)	F	55	NA	NA	SPECT: bilateral Sylvian hypoperfusion	p.V363I	NA	NA	Italy	Rossi et al. ⁷
7	PCA	F	54	NA	NA	NA	p.V363I	NA	NA	Italy	Rossi et al. ⁷
8	FTD, PPA (nonfluent variant), and CBS	F	55	NA	NA	MRI: mild left frontal atrophy SPECT: left frontotemporal predominant hypoperfusion FDG-PET: left parietal hypometabolism	p.V363I	H1/H1	NA	Italy	Rossi et al. ¹⁰
9	PCA	F	51	N/A	+	MRI: slight, asymmetric atrophy of posterior temporoparietal and occipital lobes; white matter abnormalities FDG-PET: bilateral posterior temporo-occipital and right posterior frontoparietal hypometabolism	p.V363I	H1/H1	N/A	Italy	Rossi et al. ¹⁰
10	PSP	M	53	NA	+	MRI: midbrain atrophy DAT scan: bilateral dopaminergic denervation	p.V363A	H1/H1	NA	Italy	Rossi et al. ¹⁰

Abbreviations: AAD = age at death; AAO = age at onset; CBS = corticobasal syndrome; CBD = corticobasal degeneration; DAT scan = dopamine transporter scan; FDG = fluorodeoxyglucose PET; FH = family history; +/- = present/absent; FTD = frontotemporal dementia; *MAPT* = microtubule-associated protein tau; NA = not available or not applicable; PCA = posterior cortical atrophy; PPA = primary progressive aphasia; PSP = progressive supranuclear palsy; SPECT = single-photon emission computed tomography.

Figure 1 This schematic representation illustrates the location of the *MAPT* p.V363I mutation



controls were determined using a Fisher exact test with a significance threshold of 0.05.

Neuropathology

The brain of case 2 was pathologically evaluated at the Neurological Tissue Bank of the IDIBPAS Biobank in Barcelona, Spain, after obtaining written informed consent from the patient's relatives for use of tissue for diagnostic and research purposes. Hematoxylin and eosin staining was performed after standard formalin fixation and paraffin block sectioning of multiple cortical and subcortical brain areas. Immunohistochemistry was performed using phospho-tau (Ser202 and Thr205) monoclonal antibodies (AT8; 1:2000; Thermo Scientific, Rockford, IL) and anti-4R-tau (RD4) antibodies. In addition, selected areas were stained for β A4-amyloid (6F/3D 1:400; Dako, Glostrup, Denmark), α -synuclein (KM51 2:200; Novocastra, Newcastle upon Tyne, UK), and TDP43 protein (2E2-E3 1:500; Abnova, Taipei, Taiwan) for identification of concomitant pathologies.

Data availability

Deidentified data are available upon request from qualified investigators.

Results

Genetic characteristics

In a cohort of 173 CBS cases, we identified 2 patients who were heterozygous for the rare p.V363I (c.1087G>A: NM_005910.5) mutation located in the highly conserved microtubule-binding domain of *MAPT* (Fisher exact test comparing CBS cases with neurologically healthy controls $p = 0.0299$). Both patients were homozygous for the H1 *MAPT* haplotype and carried no pathogenic mutations in *PGRN* or *C9orf72*. The patients' *APOE* genotypes were $\epsilon 3/\epsilon 3$. The *MAPT* p.V363I mutation was absent in $\sim 1,800$ additional samples, including 826 neurologically healthy controls and 984 cases with diverse frontotemporal degeneration spectrum disorders. Bioinformatic predictions demonstrated that SIFT, PolyPhen2, MutationTaster, and ClinPred categorized the variant as tolerated and benign, whereas ClinVar, FATHMM-XF, M-CAP, and CADD predictions suggested a likely pathogenic mutation (table e-1, links.lww.com/NXG/A162).

Clinicopathologic features


Case 1 is a 73-year-old, right-handed, white woman with a medical history of hypertension, coronary artery disease, an

old segmental left parietal stroke at age 54 years that resolved without residual neurologic deficits, and major depressive disorder. She was diagnosed with CBS at age 70 years after developing progressive right-sided impairment of her dexterity, slowed gait, and imbalance resulting in backward falls. A levodopa trial up to a maximum dose of 450 mg daily yielded no benefits. Over the course of 3 years, she gradually developed dysarthria, severe gait dysfunction rendering her wheelchair-bound, asymmetric parkinsonism, hand dystonia, apraxia, impaired word retrieval, and executive dysfunction. Her neurologic examination demonstrated bradyphrenia with a tendency to perseverate. She had severe ideomotor apraxia that was more prominent in her dominant hand. She was neglecting her right-sided space. Her speech was moderately dysarthric. Cranial nerve examination showed slowed, hypometric saccades (vertical more than horizontal), severe axial and right-sided rigidity with only mild rigidity on the left, bradykinesia, and dystonia with high-frequency/low-amplitude tremor in her right hand. She had agraphesthesia and astereognosis in her right hand. Reflexes were brisk

throughout. Primitive reflexes, including grasp and palmo-mental reflexes, were present. She was unable to stand without assistance and would spontaneously fall without support. MRI of the brain demonstrated bilateral parietal atrophy with proportional, ex vacuo dilatation of the lateral ventricles. Her family history was notable for parkinsonism in her father (age at onset ~65 years). No DNA was available from her father to test for segregation. The patient is alive after a 3-year disease duration.

Case 2 was a white woman who presented in her late 50s with primary progressive aphasia, left-sided parkinsonism, and CBS. The disease progressed to complete anarthria and severe dysphagia. She died at age 62 years. Clinical data on this case were limited. She had no known family history of dementia. The patient's neuropathologic findings were notable for widespread, 4-repeat-tau-positive inclusions in cortical and subcortical regions, including neurons and glial cells, consistent with CBD (figure 2). Frequent achromatic neurons were detected in frontal, parietal, and cingular cortices. These were

Figure 2 These images showcase the pertinent neuropathologic findings of case 2



Hematoxylin and eosin staining shows superficial spongiosis in the postcentral region (A), a large achromatic or ballooned cell (highlighted by asterisk in B), and prominent nigral degeneration with severe neuronal loss and abundant extracellular neuromelanin pigment (C). (D–I) Abnormal pTau (AT8) and 4-repeat-tau-positive protein deposition on immunohistochemistry. Notable abnormal histopathologic findings included astrocytic plaques (D), frequent pretangles with some focal cytoplasmic condensations (E), tangles, pretangles, and abundant threads in the substantia nigra (F), very abundant threads (arrow heads) and coiled bodies (arrow) in the white matter (G and H), and abundant threads and pretangles in the striatum (I), overall consistent with the neuropathologic findings observed in corticobasal degeneration. Magnification scale bars are indicated in the bottom right corner of each panel.

associated with focal superficial spongiosis, diffuse neuronal loss, astrogliosis, and microglial activation in cortical areas, including the motor cortex, and diffuse gliosis of the underlying white matter. Prominent neuronal loss was noted in the globus pallidus and in the substantia nigra. Immunohistochemistry revealed astrocytic plaques, abundant pretangles, ballooned neurons, and neuropil threads in the cortex, abundant threads and pretangles in her basal ganglia, and prominent white matter pathology with widespread threads and coiled bodies involving also the brainstem. Remarkably, there was also prominent involvement of the hippocampus, including the granule cells of the dentate gyrus, without grain pathology. Co-comitant pathologies included a moderate amount of diffuse β A4-amyloid deposits and few cored plaques in cortical areas, as well as few neuronal and glial cytoplasmic TDP43 protein inclusions in the globus pallidus, without frontal, temporal, or hippocampal involvement. No α -synuclein aggregates were identified.

Discussion

We describe 2 CBS cases carrying the rare p.V363I *MAPT* mutation located in the conserved microtubule-binding domain. One of the 2 cases was pathologically confirmed, demonstrating widespread, 4-repeat-tau-positive neuronal and glial pathology consistent with CBD. This report describes the pathology present in a p.V363I *MAPT* mutation carrier providing further support for the notion that this variant is likely disease causing. To date, this coding mutation has been described in 7 neurodegenerative disease cases with heterogeneous presentations, including FTD, primary progressive aphasia, and posterior cortical atrophy (table 1).^{6,8–10} Another mutation at the same residue (p.V363A) has been described in a single case with clinically diagnosed PSP.¹⁰ Of interest, all cases were female, had ϵ 3/ ϵ 3 *APOE* genotypes, and were homozygous for the *MAPT* H1 haplotype. The average age at onset was 57 years, ranging from 46 to 70 years. The 2 CBS cases presented here extend the disease onset. This wide age spectrum is consistent with the pattern seen in tauopathies associated with *MAPT* mutations and could indicate a decreased, age-related penetrance.²⁴

Among the cases, the initial disease manifestations were quite varied, including gait disturbances, memory deficits, and personality changes. This heterogeneity is not unusual for patients with *MAPT* mutations.²⁵ The p.V363I mutation is present in 3 of 62,784 people in the NHLBI TopMed Bravo database (bravo.sph.umich.edu/freeze5/hg38/; allele frequency: 0.0000239; date accessed: October 14, 2018) and in 2 of 60,702 individuals in ExAC (allele frequency: 0.0000167, data accessed: October 14, 2018).²⁶ The very rare presence within population databases might be explained by incomplete penetrance and late disease onset. In addition, limited in vitro analyses in 1 case demonstrated that this mutation leads to an increased propensity for microtubule polymerization and the formation of tau protein oligomers.¹⁰

Because of the lack of familial genetic data, we were not able to test for disease segregation, and this has not yet been reported by other investigators.

We present a pathologically confirmed patient with a p.V363I *MAPT* mutation. The neuropathologic findings of this case were consistent with CBD. An additional p.V363I carrier was identified with a CBS phenotype. This mutation was absent in neurologically healthy controls. Considering previous reports on mutation carriers with information about sequence conservation, functional studies, and pathologic confirmation, we nominate the *MAPT* p.V363I change as a likely disease-causing mutation. Identifying additional cases with this mutation will be important to understand the natural history and penetrance of this familial disease.

Acknowledgment

The authors thank all the subjects who donated their time and biological samples to be a part of this study. They thank the NIH NeuroBrainBank and the IDIBAPS Brain Bank, Barcelona, Spain, for contributing brain tissue samples. This study used tissue samples and data provided by the Michigan Brain Bank, the Michigan Alzheimer's Disease Center (SP30 AG053760), and the Protein Folding Disorders Program. This study used samples from the NINDS Repository at Coriell (catalog.coriell.org) and clinical data. The authors are grateful to the Banner Sun Health Research Institute Brain and Body Donation Program of Sun City, Arizona, for the provision of human brain tissue. The Brain and Body Donation Program is supported by the NINDS (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the Arizona Department of Health Services (contract 211002, Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-0901, and 1001 to the Arizona Parkinson's Disease Consortium), and the Michael J. Fox Foundation for Parkinson's Research. The authors thank the Kathleen Price Bryan Brain Bank at Duke University Medical Center, the Harvard Brain Bank, and the Georgetown University Brain Bank for the provision of tissue and DNA samples. Tissue for this study was provided by the Newcastle Brain Tissue Resource, which is funded in part by a grant from the UK Medical Research Council (G0400074), by NIHR Newcastle Biomedical Research Centre and Unit awarded to the Newcastle upon Tyne NHS Foundation Trust and Newcastle University, and as part of the Brains for Dementia Research Programme jointly funded by Alzheimer's Research UK and Alzheimer's Society. The authors thank members of the North American Brain Expression Consortium for providing DNA samples on neurologically healthy controls. Tissue samples for genotyping were provided by the Johns Hopkins Morris K. Udall Center of Excellence for Parkinson's Disease Research (NIH P50 NS38377) and the Johns Hopkins Alzheimer Disease Research Center (NIH P50 AG05146).

Study funding

This work was supported in part by the Intramural Research Programs of the NINDS and the National Institute on Aging (NIA) (project numbers Z01-AG000949 and 1ZIA NS003154). This study was partially supported by a CurePSP research grant (Cure PSP grant no.: 515-14; 2013-2015) to PP. EG received support from a grant from the Marató de TV3 (grant no. 20141610).

Disclosure

Disclosures available: Neurology.org/NG.

Publication history

Received by *Neurology: Genetics* March 10, 2019. Accepted in final form May 15, 2019.

Appendix Authors

Name	Location	Role	Contributions
Sarah Ahmed, BS	National Institutes of Health	Author	Drafting of the manuscript; genetic assessment and analysis; and critical review
Monica Diez Fairen, MS	University Hospital Mutua de Terrassa, and Fundació per la Recerca Biomèdica i Social Mútua Terrassa	Author	Genetic assessment and critical review
Marya S. Sabir, BS	National Institutes of Health	Author	Genetic assessment and critical review
Pau Pastor, MD, PhD	University Hospital Mutua de Terrassa, and Fundació per la Recerca Biomèdica i Social Mútua Terrassa	Author	Clinical/pathologic characterization and critical review
Jinhui Ding, PhD	National Institutes of Health	Author	Genetic assessment and critical review
Lourdes Ispierto, MD	Hospital Universitari Germans Trias	Author	Clinical/pathologic characterization and critical review
Ankur Butala, MD	Johns Hopkins University Medical Center	Author	Clinical/pathologic characterization and critical review
Christopher M. Morris, PhD	Newcastle University	Author	Clinical/pathologic characterization and critical review
Claudia Schulte, PhD	University of Tuebingen	Author	Clinical/pathologic characterization and critical review
Thomas Gasser, MD	University of Tuebingen	Author	Clinical/pathologic characterization and critical review
Edwin Jabbari, MD	University College London	Author	Clinical/pathologic characterization and critical review
Olga Pletnikova, MD	Johns Hopkins University Medical Center	Author	Clinical/pathologic characterization and critical review

Appendix (continued)

Name	Location	Role	Contributions
Huw R. Morris, MD, PhD	University College London, and Royal Free Campus	Author	Clinical/pathologic characterization and critical review
Juan C. Troncoso, MD	Johns Hopkins University Medical Center	Author	Clinical/pathologic characterization and critical review
Ellen Gelpi, MD, PhD	University of Barcelona-Hospital Clinic	Author	Neuropathologic assessment and critical review
Alexander Pantelyat, MD	Johns Hopkins University Medical Center	Author	Conceptualization and design; clinical/pathologic characterization; and critical review
Sonja W. Scholz, MD, PhD	National Institutes of Health and Johns Hopkins University Medical Center	Author	Drafting of the manuscript; conceptualization and design; clinical/pathologic characterization; and critical review

References

1. Kouri N, Murray ME, Hassan A, et al. Neuropathological features of corticobasal degeneration presenting as corticobasal syndrome or Richardson syndrome. *Brain* 2011;134:3264–3275.
2. Poorkaj P, Bird TD, Wijsman E, et al. Tau is a candidate gene for chromosome 17 frontotemporal dementia. *Ann Neurol* 1998;43:815–825.
3. Hutton M, Lendon CL, Rizzu P, et al. Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 1998;393:702–705.
4. Spillantini MG, Murrell JR, Goedert M, Farlow MR, Klug A, Ghetti B. Mutation in the tau gene in familial multiple system tauopathy with presenile dementia. *Proc Natl Acad Sci U S A* 1998;95:7737–7741.
5. Pickering-Brown SM, Rollinson S, Du Plessis D, et al. Frequency and clinical characteristics of progranulin mutation carriers in the Manchester frontotemporal lobar degeneration cohort: comparison with patients with MAPT and no known mutations. *Brain* 2008;131:721–731.
6. Bessi V, Bagnoli S, Nacmias B, Tedde A, Sorbi S, Bracco L. Semantic dementia associated with mutation V363I in the tau gene. *J Neurol Sci* 2010;296:112–114.
7. Rossi G, Conconi D, Panzeri E, et al. Mutations in MAPT gene cause chromosome instability and introduce copy number variations widely in the genome. *J Alzheimers Dis* 2013;33:969–982.
8. Anfossi M, Bernardi L, Gallo M, et al. MAPT V363I variation in a sporadic case of frontotemporal dementia: variable penetrant mutation or rare polymorphism?. *Alzheimer Dis Assoc* 2011;25:96–99.
9. Munoz DG, Ros R, Fatas M, Bermejo F, de Yébenes JG. Progressive nonfluent aphasia associated with a new mutation V363I in tau gene. *Am J Alzheimers Dis Other Demen* 2007;22:294–299.
10. Rossi G, Bastone A, Piccoli E, et al. Different mutations at V363 MAPT codon are associated with atypical clinical phenotypes and show unusual structural and functional features. *Neurobiol Aging* 2014;35:408–417.
11. Armstrong MJ, Litvan I, Lang AE, et al. Criteria for the diagnosis of corticobasal degeneration. *Neurology* 2013;80:496–503.
12. Blauwendraat C, Faghri F, Pihlstrom L, et al. NeuroChip, an updated version of the NeuroX genotyping platform to rapidly screen for variants associated with neurological diseases. *Neurobiol Aging* 2017;57:247 e9–247 e13.
13. Renton AE, Majounie E, Waite A, et al. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257–268.
14. Poorkaj P, Grossman M, Steinbart E, et al. Frequency of tau gene mutations in familial and sporadic cases of non-Alzheimer dementia. *Arch Neurol* 2001;58:383–387.
15. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res* 2003;31:3812–3814.
16. Jagadeesh KA, Wenger AM, Berger MJ, et al. M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. *Nat Genet* 2016;48:1581–1586.
17. Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging missense mutations. *Nat Methods* 2010;7:248–249.

18. Landrum MJ, Lee JM, Benson M, et al. ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic Acids Res* 2016;44:D862–D868.
19. Rogers MF, Shihab HA, Mort M, Cooper DN, Gaunt TR, Campbell C. FATHMM-XF: accurate prediction of pathogenic point mutations via extended features. *Bioinformatics* 2018;34:511–513.
20. Kircher M, Witten DM, Jain P, O’Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet* 2014;46:310–315.
21. Alirezaie N, Kernohan KD, Hartley T, Majewski J, Hocking TD. ClinPred: prediction tool to identify disease-relevant nonsynonymous single-nucleotide variants. *Am J Hum Genet* 2018;103:474–483.
22. Rausch T, Emde AK, Weese D, Doring A, Notredame C, Reinert K. Segment-based multiple sequence alignment. *Bioinformatics* 2008;24:i187–192.
23. Fitzpatrick AWP, Falcon B, He S, et al. Cryo-EM structures of tau filaments from Alzheimer’s disease. *Nature* 2017;547:185–190.
24. Irwin DJ. Tauopathies as clinicopathological entities. *Parkinsonism Relat Disord* 2016;22(suppl 1):S29–S33.
25. Ghetti B, Oblak AL, Boeve BF, Johnson KA, Dickerson BC, Goedert M. Invited review: frontotemporal dementia caused by microtubule-associated protein tau gene (MAPT) mutations: a chameleon for neuropathology and neuroimaging. *Neuro-pathol Appl Neurobiol* 2015;41:24–46.
26. Lek M, Karczewski KJ, Minikel EV, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–291.

Neurology[®] Genetics

***MAPT* p.V363I mutation: A rare cause of corticobasal degeneration**

Sarah Ahmed, Monica Diez Fairen, Marya S. Sabir, et al.

Neurol Genet 2019;5;

DOI 10.1212/NXG.0000000000000347

This information is current as of June 25, 2019

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/5/4/e347.full.html
References	This article cites 26 articles, 1 of which you can access for free at: http://ng.neurology.org/content/5/4/e347.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics Corticobasal degeneration http://ng.neurology.org/cgi/collection/corticobasal_degeneration Parkinson's disease/Parkinsonism http://ng.neurology.org/cgi/collection/parkinsons_disease_parkinsonism
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

