

Article

Looking, seeing and believing in autism: Eye movements reveal how subtle cognitive processing differences impact in the social domain

Benson, Valerie, Castelhana, MS, Howard, PL, Latif, N and Rayner, K

Available at <http://clock.uclan.ac.uk/28048/>

Benson, Valerie ORCID: 0000-0002-0351-4563, Castelhana, MS, Howard, PL, Latif, N and Rayner, K (2016) Looking, seeing and believing in autism: Eye movements reveal how subtle cognitive processing differences impact in the social domain. Autism research : official journal of the International Society for Autism Research, 9 (8). pp. 879-887. ISSN 1939-3792

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1002/aur.1580>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



**Looking, Seeing and Believing in Autism: Eye Movements
reveal how subtle cognitive processing differences impact in
the social domain.**

Journal:	<i>Autism Research</i>
Manuscript ID	AUR-15-0133.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Benson, Valerie; University of Southampton, School of Psychology Castelhano, Monica; Queens University, Psychology Howard, Phillipa; University of Southampton, Psychology Latif, Nida; Queens University, Psychology Rayner, Keith; University of California, San Diego, Psychology
Keywords:	Developmental Psychology, Neuropsychology, Autism, Eye movements, cognitive processing differences

SCHOLARONE™
Manuscripts

1
2
3
4
5 Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle
6
7
8 cognitive processing differences impact in the social domain.
9

10 Valerie Benson ^{1*}, Monica S. Castelhana ², Philippa L. Howard ¹, Nida Latif ² & Keith Rayner ³
11
12
13

14
15
16 ¹ AU Psychology, Shackleton Building, Highfield Campus, University of Southampton, UK.
17

18 ² Queen's University, Kingston, ON, Canada
19

20 ³ University of California, San Diego, USA
21
22

23
24 *Correspondence to: vb1@soton.ac.uk
25

26 Telephone: +44 (0) 2380594596
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Lay Abstract

Adults with High Functioning Autism (ASD) viewed scenes with people in them, whilst having their eye movements recorded. The task was to indicate, using a button press, whether the pictures were normal, or in some way weird or odd. Oddities in the pictures were categorized as violations of either perceptual or social norms. Compared to a Typically Developed (TD) control group, the ASD participants were equally able to categorise the scenes as odd or normal, but they took longer to respond. The eye movement patterns showed that the ASD group made more fixations and revisits to the target areas in the odd scenes compared to the TD group. Additionally, when the ASD group first fixated the target areas in the scenes, they failed to initially detect the social oddities. These two findings have clear implications for processing difficulties in ASD for the social domain, where it is important to detect social cues on-line, and where there is little opportunity to go back and recheck possible cues in fast dynamic interactions.

Scientific Abstract

1
2
3
4
5
6
7
8 Autistic Spectrum Disorder (ASD) and Typically Developed (TD) adult participants had to
9 decide whether scenes were 'odd', 'unusual' or 'weird'. All scenes contained people, with
10 'oddness' operationally defined as either perceptual or social. There were no group differences in
11 accuracy, but the ASD group were slower overall to respond manually. Eye movements revealed
12 that the ASD group made more fixations overall and more re-fixations into the target regions of
13 the pictures, for both conditions. Importantly, as indexed by the first fixation duration the ASD
14 group failed to 'pick up' immediately on what was 'odd' for the social violations. The propensity
15 to 'go back' and re-fixate targets, coupled with a failure to initially detect social violations, has
16 obvious significance for ASD in fast dynamic social communication.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 Key phrases: Autism Spectrum Disorder, Eye movements, On-line cognitive processing, Social
33 and Perceptual Oddities.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle cognitive
4
5
6 processing differences impact in the social domain.
7

8
9 Autism spectrum disorders (ASD) are lifelong neuro-developmental conditions
10 characterized by repetitive behaviours and impairments in interaction and communication
11 (American Psychiatric Association, 2013). Incidence rates are rising across the globe with the
12 prevalence rate in the UK approximately 10 per 1000 (1%) individuals (Brugha et al., 2011) and
13 the UK National Audit office recently called for more research into adults with ASD (Clark et
14 al., 2009).
15
16
17
18
19
20
21
22

23 High functioning adults with ASD are similar to typically developed (TD) adults in terms
24 of performance on pencil and paper and IQ tests, however in the social domain evidence suggests
25 an inability to process social information in a typical fashion. Eye movement studies report
26 unusual gaze behaviour for faces appearing on their own (Dalton et al., 2005), in social scenes
27 (Riby & Hancock, 2009) or in dynamic movie clips (Klin & Jones, 2006; Speer et al., 2007).
28 Research consistently reports a lack of spontaneous gaze fixation towards the eyes (Jones et al.,
29 2008; Klin et al., 2002a; Pelphrey et al., 2002). However, where one study reports impairments,
30 say for face processing (Klin et al., 2002b), another study does not (Fletcher-Watson et al.,
31 2009). Moreover, lack of attentional modulation for social stimuli (Bird et al., 2006) may be
32 overridden, as when cued, attention may be allocated to faces in a more 'typical' manner (Bar-
33 Haim et al., 2006).
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 A recent scene perception study (Louwerse et al., 2014) reported that abnormal
49 autonomic responses in ASD are unlikely to underpin social processing impairments. It is
50 interesting then to ask whether impairments in the cognitive domain in ASD (Happé & Frith,
51 2006; Minshew et al., 1997; Russell, 1997) could account for impairments in the social domain?
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Current cognitive theories of ASD include, among others, Weak Central Coherence (WCC) theory (Frith & Happé, 1994; Happé & Frith, 2006) which proposes that a reduced sensitivity to global context in favour of focussing on details, results in integration difficulties in ASD. Executive function theory (Russell, 1997), proposes that ASD are impaired at switching their focus of attention, and may engage in perseverative behaviour. Alternatively, the Complex Information Processing theory (Minshew & Goldstein, 1998) proposes that individuals with ASD have impairments in processing complex information with intact processing for tasks that require simple (rule based) processing. Finally, the Extreme Male Brain (EMB) theory (Baron-Cohen, 2002) posits that individuals with ASD have above average levels of systemizing, that contribute to the non-social characteristics of the disorder, coupled with below average levels of empathizing, which results in atypical social functioning. These contemporary theories of ASD can, for different reasons, offer some account of ineffective social functioning in ASD. However, to date very few investigations have examined processing differences as these occur on-line in ASD.

Our studies investigating cognitive processing in adults with ASD (Au-Yeung et al., 2011; Au-Yeung et al., 2013; Benson et al., 2012; Benson et al., 2009; Benson & Fletcher-Watson, 2011) have found subtle on-line processing differences, as revealed by eye movements, for a range of tasks that are more in line with Minshew and Goldsteins (1998) theory, than with any other. For example, there are more similarities than differences in eye movements between TD and ASD groups for simple rule based tasks in the attentional (Kuhn et al., 2010) and the social domains (Benson & Fletcher-Watson, 2011). Whereas for more complex abstract reasoning tasks where there is ambiguity with relation to target items (Au-Yeung et al., 2013), or where value judgements rely on top down knowledge (Benson et al., 2012), there are early

1
2
3 processing differences indicating that initial orienting to, and immediate detection of weird target
4 items in scenes is absent in ASD. Moreover, the time to respond and to locate the weird target
5 items was modulated by whether or not the pictures had people in them (Benson et al., 2012).
6
7
8 These differences have potential for impact on everyday activities, since the time course for
9 detection of socially relevant information is crucial for successful everyday interactions.
10
11
12
13
14

15 The aim of the present study was to examine whether increased times to find and initially
16 detect an unusual target in a scene would be modulated by whether the violation that rendered
17 the target unusual or odd could be categorised as social or perceptual. We employed a similar
18 paradigm to that used in a previously reported study (Rayner et al., 2009) where single scenes,
19 all with people in them, were presented in isolation and the task was to indicate whether each
20 picture was normal or in some way weird unusual or odd.
21
22
23
24
25
26
27
28
29
30
31

32 Insert Figure 1. about here
33
34
35

36 Social oddities in the pictures were operationally defined as violations of social norms
37 e.g. someone in the picture throwing a baby in the air. Access to top down knowledge is required
38 in that condition to inform a subjective value judgement as to whether a social violation is
39 present in the picture. No value judgement as to whether a violation infringes some implicit or
40 explicit social rule is needed for the perceptual oddities in this study, which are operationally
41 defined as physical violations e.g. someone in the picture has a limb positioned back to front. See
42 supplementary information for Materials. Figure 1 shows an example of the stimuli.
43
44
45
46
47
48
49
50
51

52 Method

53 Participants

54
55
56
57
58
59
60

1
2
3 ASD participants ($n = 24$ age 18 – 52 years, 20 males), with a formal diagnosis of either
4
5 Autistic Disorder ($n = 3$) or Asperger's Syndrome were recruited from local charitable
6
7 organisations. TD participants ($n = 24$, age 20– 52 years, 20 males) were volunteers from the
8
9 local community. All participants were paid and had normal or corrected to normal vision. Each
10
11 ASD participant completed module 4 of the Autism Diagnostic Observation Scale 2 (Lord et al.,
12
13 2012) and five (3 males and 2 females with a diagnosis of Asperger's Syndrome) failed to meet
14
15 the Autism Spectrum criteria. To ensure maximum homogeneity in our study those five
16
17 participants were excluded from the analyses, resulting in a final sample of 19 ASD participants.
18
19 Both groups completed the Autism-spectrum Quotient (AQ; Baron-Cohen et al., 2001) and the
20
21 ASD participants reported a higher number of autistic traits $t(41) = 8.04, p < .001$ (TD $M =$
22
23 15.42, $SD = 7.55$; ASD $M = 35.32, SD = 8.45$). The two groups did not differ in age $t(41) =$
24
25 0.31, $p = .759$ (TD $M = 30.29, SD = 10.20$; ASD $M = 31.26, SD = 10.29$), performance IQ $t(41) =$
26
27 = 0.21, $p = .832$ (TD $M = 111.79, SD = 12.01$; ASD $M = 110.95, SD = 13.84$), verbal IQ $t(41) =$
28
29 1.21, $p = .234$ (TD $M = 116.96, SD = 10.65$, ASD $M = 112.26, SD = 14.84$), or full scale IQ t
30
31 (46) = 0.87, $p = .392$ (TD $M = 116.04, SD = 9.98$, ASD $M = 112.95, SD = 13.48$), as measured
32
33 by two performance (Block Design and Matrix Reasoning) and verbal components (Vocabulary
34
35 and Similarity) of the Weschler Abbreviated Scale of Intelligence (The Psychological
36
37 Corporation, 1999).
38
39
40
41
42
43
44
45
46
47

48 **Eye movement recording**

49

50 Eye movements were sampled at 1000Hz using an Eyelink tracker with spatial accuracy
51
52 of less than 1/4th degree (S.R. Research Ltd, Canada). The start and end of saccades were
53
54 detected automatically using a default velocity criterion of 30degrees per second using the
55
56
57
58
59
60

1
2
3 Eyelink software. See Stampe (1993) for an in depth technical description of how the software
4 detects saccades and fixations. A chin rest and forehead support, were used to stabilize head
5 position. Viewing was binocular but data were analysed for the right eye only. Individual
6 participants were calibrated using a nine point matrix that covered the dimensions of the screen
7 (1024 x 768 pixels), with each point fixated sequentially, followed by a validation procedure to
8 ensure that fixation was within 0.5 degrees of each calibration point. Calibration was checked
9 using a validation procedure prior to each trial, whereby participants fixated five dots, presented
10 at five different locations that covered the dimensions of the screen and recalibration was
11 performed if the eyes were off centre for any of these dots.
12
13
14
15
16
17
18
19
20
21
22
23

24 25 26 27 **Materials**

28
29 In total, 96 full-colour photographs were used as stimuli, all of which contained people.
30
31 In both the perceptual and social oddity conditions there were 48 pictures (24 weird and 24
32 normal). Perceptual oddities in the pictures violated a physical norm e.g. someone in the picture
33 has a limb positioned back to front, or a vehicle was in an unexpected position e.g midway
34 through the upper wall of a building. Social oddities in the pictures violated a social norm e.g.
35 someone in the picture throwing a baby in the air, or someone with a cabbage on a lead. A small
36 number of the pictures were from a previous study (Benson et al., 2012) and the remainder were
37 collected from the internet and included either a perceptual or social oddity. Adobe Photoshop
38 was used to prepare a normal version of each scene in which the oddity was replaced with either
39 the background elements or with another suitable object. The scenes typically had a focal object
40 or a central group of objects. Figure 1 shows an example of a normal and a socially odd scene,
41 and below those examples, a normal and a perceptually odd scene. The target regions for these
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 scenes are outlined in black – but these black outlines were not visible during the presentation of
4
5 the scenes in the experiment.
6
7

8 A norming study was conducted to verify the validity of the images used to examine the
9
10 processing of social and perceptual violations. Twenty undergraduates from Queen's university
11 participated (M age: 19.8years, SD : 1.71, 10 Females) with five participants in each of the four
12 conditions (counterbalanced conditions: weird/normal and right/left button response). Stimuli
13 were displayed on a 21inch CRT monitor with a refresh rate of 100Hz and the experiment was
14 run using E-Prime software. Participants were presented with a fixation cross for 500ms
15 followed by the presentation of the image for 20 seconds or until a keyboard "Y" (yes) or "N"
16 (N) was made. Participants had to indicate as quickly as possible whether the presented image
17 was "odd", "weird" or "unusual" in some way. The proportion of identified images indicated that
18 on average, the social normal ($M = 0.86$, $SD = 0.15$) and social weird ($M = 0.84$, $SD = 0.17$),
19 perceptual normal ($M = 0.82$, $SD = 0.17$) and perceptual weird ($M = 0.77$, $SD = 0.19$) images
20 were identified correctly.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Design

40 For the behavioural measures of Accuracy and RT a mixed design with Picture Type
41 (weird versus normal) and Oddity Type (social versus perceptual) as within participants
42 variables, and Group (ASD versus TD) as a between participants variable was employed for the
43 experiment. On condition that accuracy was equivalent for both normal and weird pictures, for
44 the eye movement measures the main focus of interest was to compare social versus perceptual
45 oddities in both groups. Therefore a mixed design with Oddity Type (social versus perceptual) as
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 a within participants variable, and Group (ASD versus TD) as a between participants variable
4
5 was employed in the eye movement analysis.
6
7
8
9

10 **Procedure**

11
12 All participants were shown a hard copy of an example picture from each condition prior
13 to completing the task, to ensure comprehension of the task and the ability to respond
14 appropriately. The specific instruction was ‘for each of the following pictures, please indicate
15 whether you think the picture is odd, unusual or strange in some way’.
16
17
18
19
20
21

22 For the task, 48 pictures were presented randomly to either the left or the right of the
23 centre of a 19inch colour monitor, at a distance of 65cm for 20s or until a button press response
24 was made. Prior to the onset of a picture, participants fixated a cross that was presented on the
25 opposite side of the screen to where the picture was set to appear, to ensure all participants
26 started at the same point in each trial. Once participants fixated this cross, the experimenter
27 initiated each trial. Picture type (normal vs. weird) and, the position of the picture (left or right
28 on the display) were counterbalanced across items. The buttons used to respond either yes or no
29 (left or right trigger button) was counterbalanced across participants.
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 **Data preparation and analyses**

45 For each ‘odd’ picture we created an interest region around the target item and the
46 corresponding region of the same size in the counterpart ‘normal’ picture. Regions of interest
47 were created using the approximate average regions from those identified and manually selected
48 by participants who took part in the norming experiment.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Eyelink software was used to detect saccades and fixations using a default velocity criterion and
4
5 in-house software was used to calculate the distribution of fixations falling into the target region.
6
7
8 Five items were removed from analysis because for these scenes, participant accuracy scores
9
10 were significantly below chance¹. In addition, each participant was asked whether they had seen
11
12 any of the scenes previously, and if so these scenes were also removed. This resulted in a loss of
13
14 2.3% of trials. Fixations below 90ms and above 1200ms were classed as outliers and removed;
15
16
17 resulting in a loss of 5.13% of the fixation data.
18
19

20 21 22 Results

23
24 For the principal analyses we compared manual reaction times and accuracy (see Table
25
26 1), as well as a number of eye movement measures, such as, the time taken to begin fixating in
27
28 the target region (ELT), the duration of the first fixation when the eyes landed in the target
29
30 region (FFD) and the mean time (MFD) spent fixating in the target area (see Table. 2). Global
31
32 eye movement measures for the target region, such as mean fixation duration and mean number
33
34 of fixations over all trials were also analysed, Finally, baseline measures (not just target related),
35
36 were calculated across the duration of the trials, to check for overall sampling differences
37
38 between the ASD and TD participants (see Table 2). See supplementary information for methods
39
40 and materials.
41
42
43
44
45
46
47

48 Behavioural Data

49
50 **Accuracy.** There was no effect of oddity type (perceptual vs. social) $F(1, 41) = 1.44, p =$
51
52 $.237, \eta_p^2 = .03$, picture type (normal or weird) $F(1, 41) = 0.94, p = .339, \eta_p^2 = .02$, or group, $F(1,$
53
54 $41) = .06, p = .808, \eta_p^2 < .01$ on accuracy rates, and no interactions ($F_s < 2.39, p's > .130$). At
55
56
57
58
59
60

1
2
3 least in terms of accuracy, performance was the same for both groups. For means and standard
4
5
6 deviations for behavioural data see Table 1.

7
8 **Response Time (RT).** A main effect of oddity type $F(1, 41) = 8.16, p = .007, \eta_p^2 = .17$,
9
10 showed participants took longer to respond when presented with a perceptual oddity in
11
12 comparison to a social oddity. Longer response times for perceptual oddities could reflect that
13
14 these ‘impossible’, physical violations were so unusual that they resulted in more time spent
15
16 looking at them prior to responding. See below for confirmation of that in the eye movement
17
18 data. There was also a main effect of picture type $F(1, 41) = 24.03, p < .001, \eta_p^2 = .37$, with
19
20 participants taking longer to respond when the picture was normal in comparison to when it was
21
22 weird, in line with longer reaction times for target absent trials in standard visual search tasks
23
24 (Wolfe, 1998). A significant oddity type x picture type interaction $F(1, 41) = 4.22, p = .046, \eta_p^2$
25
26 $= .09$, demonstrated that the difference between RTs for perceptual and social pictures was
27
28 reliable for odd (TD; $t(23) = 2.62, p = .015$, ASD; $t(18) = 2.98, p = .008$) but not for normal
29
30 pictures (TD; $t(23) = 0.94, p = .355$, ASD; $t(18) = 0.08, p = .935$). A main effect of group $F(1,$
31
32 $41) = 4.07, p = .050, \eta_p^2 = .09$, showed the ASD participants had longer reaction times, in line
33
34 with a previous study (Benson et al., 2012). There were no significant group interactions ($F_s <$
35
36 1) demonstrating both groups to be slower for normal versus odd pictures, and for perceptual vs
37
38 social oddities. See Table 1 for the means. The eye movement data below reveals subtle
39
40 processing differences that contribute to this overall slowed manual responding in the ASD
41
42 group.
43
44
45
46
47
48
49
50
51
52

53 Eye Movement Data

54
55
56
57
58
59
60

1
2
3 Having shown that all participants could distinguish between ‘odd’ and ‘normal’ pictures,
4 normal picture trials were excluded from the eye movement analyses resulting in a 2 (group:
5 ASD versus TD) X 2 (odddity type: social versus perceptual) design to investigate on-line
6 processing differences between the two groups for the two types of oddities.
7

8
9
10
11
12 Baseline eye movement measures, calculated across the full length of trials indicated no
13 group difference in mean fixation duration $F(1, 41) = 1.14, p = .293, \eta_p^2 = .03$, but the ASD
14 group made more fixations overall $F(1, 41) = 9.44, p = .004, \eta_p^2 = .19$, corresponding to the
15 increased RT’s reported above. Table 2 presents a summary of all the eye movement measures.
16
17
18
19
20
21

22
23
24 **Elapsed time to the target region (ELT).** A marginal effect of oddity type $F(1, 41) = 3.99, p =$
25 $.053, \eta_p^2 = .09$, on the time taken from the onset of each trial display to the start of the first
26 fixation on the target region, revealed both groups took less time to fixate the target region when
27 the oddity was perceptual in comparison to social in nature. There was no effect of group $F(1,$
28 $41) = 0.59, p = .446, \eta_p^2 = .01$, and no interaction $F(1, 41) = 0.72, p = .401, \eta_p^2 = .02$.
29
30
31
32
33
34
35
36
37
38

39 **First Fixation Duration (FFD) on target region.** There was no effect of oddity type $F(1, 41) =$
40 $1.84, p = .182, \eta_p^2 = .04$ or group $F(1, 41) = 0.15, p = .699, \eta_p^2 < .01$ but a significant interaction
41 $F(1, 41) = 6.72, p = .013, \eta_p^2 = .14$ (See Figure 2) revealed that FFDs were equivalent for
42 perceptual and social oddities for TD participants $t(23) = 0.86, p = .399, d = .16$, but for ASD
43 participants FFDs for social oddities were significantly shorter, in comparison to perceptual
44 oddities, $t(18) = 2.99, p = .008, d = .65$, see Figure 2. Given that the FFD is a measure of early
45 processing (Liversedge & Findlay, 2000; Rayner, 1998; Rayner, 2009). This finding indicates
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that the ASD group did not detect the social oddities as quickly as the perceptual oddities on
4
5 initial fixation, whereas TD participants detected both types of oddity equally fast.
6
7
8
9

10
11 Insert Figure 2. about here
12
13
14

15 **First gaze duration (FGD) in the target region.** The FGD is the elapsed time from when the
16 target region is first fixated until the eye moves on to a new location. FGD includes multiple
17 fixations in the target region reflects an overt strategy to re-examine an area. A main effect of
18 oddity type, $F(1, 41) = 12.10, p = .001, \eta_p^2 = .23$ demonstrated that both groups spent more time
19 inspecting perceptual oddities, again highlighting the salience of the perceptual oddities in
20 comparison to social oddities in this study. However, there was no group effect $F(1, 41) = 2.43,$
21 $p = .127, \eta_p^2 = .06,$ or interaction $F(1, 41) = .08, p = .785, \eta_p^2 < .01.$
22
23
24
25
26
27
28
29
30
31
32
33

34 **Mean Fixation Count in the target region.** If the ASD are slower to respond manually, but do
35 not take longer to begin inspecting the target region, then extra time to respond might reflect
36 more fixations in the target region. A main effect of oddity type $F(1, 41) = 64.27, p < .001, \eta_p^2 =$
37 $.61,$ demonstrated more fixations in the target region when the oddity was perceptual, rather than
38 social. A main effect of group $F(1, 41) = 6.99, p = .012, \eta_p^2 = .15$ showed that the ASD group
39 made more fixations than the TD group in the target regions, for both types of oddities. However
40 no interaction was observed $F(1, 41) = 0.77, p = .386, \eta_p^2 = .02$ indicating that the ASD group
41 made more fixations in the target regions for both types of oddities.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Mean Fixation Duration in the target region.** Longer mean fixation durations indicate that
4 more processing has been done during the task. A main effect of oddity type $F(1, 41) = 8.22, p =$
5 $.007, \eta_p^2 = .17$ showed significantly longer fixations for social, compared to perceptual oddity
6 target regions. This suggests that processing social oddities was more effortful, and since there
7 was no main effect of group $F(1, 41) = 0.34, p = .562, \eta_p^2 < .01$, and no interaction $F(1, 41) =$
8 $1.12, p = .297, \eta_p^2 = .03$, both groups on average fixated the social oddity target region for longer
9 than the perceptual target region. Thus, while the perceptual oddities attracted more fixations
10 overall, individual fixations were longer for the social target region.
11
12
13
14
15
16
17
18
19
20
21

22 This interesting juxtaposition, although unrelated to the main aim of this study, suggests
23 that difficulty associated with processing each oddity type manifests in different eye movement
24 patterns: one reflecting an overt strategy (fixation count) to re-inspect the target area, and one
25 reflecting more in depth on-line cognitive processing (fixation duration) on the target areas.
26
27
28
29
30
31
32
33

34 **Mean Number of Entries to the target region.** Since there were no group differences in FGD,
35 and no increased MFD, both reported above, the delayed RT in ASD might reflect repeated visits
36 to inspect the target region?
37
38
39
40

41 A main effect of oddity type was found $F(1, 41) = 9.83, p = .003, \eta_p^2 = .19$, with the
42 target region entered more often for perceptual oddities in comparison to social oddities. In
43 addition, a main effect of group was found $F(1, 41) = 5.72, p = .021, \eta_p^2 = .12$, with ASD
44 participants entering the target regions significantly more than the TD group, before making a
45 response. There was no interaction $F(1, 41) = 0.13, p = .717, \eta_p^2 < .01$.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Mean Total Time in the target region.** A higher mean total time means that more attention to,
4 and processing of, information in that region was carried out overall. This measure should
5 complement the measures reported above – namely, that if there are more fixations overall in
6 ASD then the Total Time spent fixating in the target region should be longer in ASD.
7
8
9

10
11
12 A main effect of oddity type $F(1, 41) = 34.89, p < .001, \eta_p^2 = .46$, showed perceptual
13 oddities received more overall processing time, in line with the previous measures for the
14 perceptual oddities. In addition, a main effect of group $F(1, 41) = 6.41, p = .015, \eta_p^2 = .14$
15 revealed that the ASD participants spent a longer period of time inspecting the target regions for
16 both perceptual and social oddities. There was no interaction $F(1, 41) = 0.51, p = .481, \eta_p^2 = .01$.
17
18
19

20
21 As above, this finding reflects a greater number of entries to the target region overall by
22 the ASD group prior to making their decision as to whether the picture under inspection was
23 weird, odd or unusual, irrespective of the nature of the violation (perceptual or social).
24
25
26

27 **Discussion**

28
29 A principal aim of the study was to investigate whether increased times in ASD to find
30 and initially detect an unusual target in a scene containing people would be modulated by
31 whether the violation could be categorised as social or perceptual.
32
33

34
35 Accuracy rates were equivalent and high for both groups, but the ASD group took longer
36 to respond manually. It was hypothesised that the eye movement data would reveal the cause of
37 this delayed response in ASD, since eye movements reflect how information is processed on-line
38 (Liversedge & Findlay, 2000; Rayner, 1998; 2009). The data showed that our ASD group did not
39 take longer to find the target regions, but that they scanned the scenes more prior to responding
40 manually, as shown by an increased number of fixations throughout each scene presentation.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Furthermore, these increased fixations were not random, but were made to the target regions of
4
5 both social and perceptually odd pictures. Thus, increased fixations reflect a propensity to re-
6
7 inspect the target regions prior to making a response in ASD, regardless as to the type of oddity
8
9 in the scenes. It should be noted here that the oddity detection task in the current study provides
10
11 the ASD participants with structure that is not typically available in real-life. As such, the task is
12
13 not implicit and the knowledge that oddities might be present could also be a factor that drives
14
15 the repeated scanning in ASD.
16
17
18
19

20 The eye movement patterns also showed that our ASD group failed to immediately detect
21
22 social oddities, thus looking at but not seeing them. An obvious question from this finding is
23
24 whether the failure to quickly spot subtle social oddities in ASD reflects an inability to react or
25
26 respond appropriately to social cues in everyday communication? This one impairment in itself
27
28 could potentially account for the significant problems experienced in dynamic social interactions
29
30 in this high functioning population. For example, a failure to immediately spot social cues would
31
32 result in a failure to react or respond to such cues thus resulting in an inability to both keep track
33
34 of what is going on in social interactions, and a failure to engage appropriately or effectively
35
36 with other people.
37
38
39
40

41 If social oddities are more complex to detect in ASD, then the Complex Information
42
43 Processing theory (Minschew & Goldstein, 1998) might explain the failure to detect the social
44
45 oddity on first fixation. This lack of immediate detection could reflect the less obvious and more
46
47 'inferential' nature of the social oddities, which required more complex reasoning, at least in the
48
49 ASD group, for detection. It is important to note that social rules are often implicit, sometimes
50
51 ambiguous, and cannot always be understood by following a script or flow chart as in the case
52
53 for logical rule based tasks. At a neurobiological level, it has been postulated that that
54
55
56
57
58
59
60

1
2
3 impairments in cognitive processing in ASD reflect slow information transmission rates for high
4
5 level cognitive tasks, resulting from underconnectivity between frontal-posterior cortical regions
6
7 (Just et al., 2013; Just et al., 2012; Minshew & Williams, 2007). Since underconnectivity reduces
8
9 the rate of inter-region information transfer, it could be that in social communication tasks that
10
11 rely on the fast detection of implicit, often ambiguous cues, a failure to be able to transmit
12
13 information across different cortical areas quickly, might mean that these cues are missed,
14
15 rendering inappropriate responses in ASD.
16
17

18
19
20 The findings of repeated scanning of the target areas, coupled with an inability to
21
22 immediately recognise social violations upon initial fixation, reflect how eye movements can
23
24 reveal subtle processing differences in ASD. In everyday social communication in the real world
25
26 there is no opportunity to 'go back' to check if one has missed an important cue. Social
27
28 processing in such circumstances must be carried out on-line, and our eye movement data clearly
29
30 show that ASD are unable to do this quickly. The strategy of re-checking their interpretation, and
31
32 thus not believing what they have seen, before responding, may be related to cortical
33
34 underconnectivity in ASD. In everyday terms, these findings have obvious consequences for fast
35
36 dynamic social interactions, and offer an account as to why high functioning adults with ASD
37
38 experience difficulty in the social domain.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

This work was supported by a grant awarded to the second author from the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the Ontario Ministry of Research and Innovation, and Queen’s University Senate Advisory Research Committee, and an Experimental Psychology Society study visit grant awarded to the third author.

For Peer Review

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Au-Yeung, S. K., Benson, V., Castelhana, M. S., Rayner, K. (2011). Eye movement sequences during simple versus complex information processing of scenes in autism spectrum disorder. *Autism Research and treatment*, 2011.
- Au-Yeung, S. K., Kaakinen, J. K., Benson, V. (2013). Cognitive perspective taking during scene perception in autism spectrum disorder: evidence from eye movements. *Autism Research*, 7, 84-93.
- Bar-Haim, Y., Shulman, C., Lamy, D., & Reuveni, A. (2006). Attention to eyes and mouth in high-functioning children with autism, *Journal of Autism and Developmental Disorders*, 36, 131-137.
- Baron-Cohen, S. (2002). The extreme male brain theory of autism. *Trends in Cognitive Science*, 6, 248-254.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5-17.
- Benson, V., & Fletcher-Watson, S. (2012). Eye Movements in Autism. In: *Liversedge, S. P., Gilchrist, I. D., Everling, S., editors. The Oxford Handbook of Eye Movements*, Oxford UK: Oxford University Press, chap 39.
- Benson, V., Castelhana, M. S., Au-Yeung, S. K., & Rayner K. (2012). Eye movements reveal no immediate "WOW" ("which one's weird") effect in autism spectrum disorder, *Quarterly*

1
2
3 Journal of Experimental Psychology, 65, 1139-1150.

4
5 Benson, V., Piper, J., & Fletcher-Watson, S. (2009). Atypical saccadic sampling in autistic
6 spectrum disorder, *Neuropsychologia*, 47, 1178-1182, (2009).

7
8 Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate neural
9 responses to social stimuli in autism spectrum disorders. *Neuroimaging*, 15, 1614-1624.

10
11 Brugha, T. S., McManus, S., Bankart, J., Scott, F., Purdon S., et al. (2011). Epidemiology of
12 autism spectrum disorders in adults in the community in England. *Archives of General*
13 *Psychiatry*, 68, 459-65.

14
15 Clark, F., Scharaschkin, A., Xu, D. (2009). Supporting autism through adulthood, National Audit
16 Office.

17
18 Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith,
19 H. H., Alexander, A. L., & Davidson, R. J. (2005). Gaze fixation and the neural circuitry
20 of face processing in autism. *Nature Neuroscience*, 8, 519-526.

21
22 Fletcher-Watson, S., Leekam, S. R., Benson, V., Frank, M. C., & Findlay, J. M. (2009). Eye-
23 movements reveal attention to social information in autism spectrum disorder.
24
25 *Neuropsychologia*, 47, 248-257.

26
27 Fletcher-Watson., Leekam, S.R., Benson, V., Frank, M.C., & Findlay, J.M. (2009). Eye-
28 movements reveal attention to social information in autism spectrum disorder.
29
30 *Neuropsychologia*, 47, 248-257.

31
32 Frith, U., & Happé, F. (1994). Autism; Beyond the theory of mind. *Cognition*, 50, 115-132.

33
34 Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in
35 autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 36, 5-25.

- 1
2
3 Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of
4
5 approaching adults predicts level of social disability in 2-year-old toddlers with autism
6
7 spectrum disorder. *Archives of General Psychiatry*, 65, 946-954.
8
9
- 10 Just, M. A., Keller, T. A., & Kana, R. K. (2013). A theory of autism based on frontal-posterior
11
12 underconnectivity. In M. A. Just & K. A. Pelphrey (Eds.), *Development and brain*
13
14 *systems in autism* (pp. 35-63). New York: Psychology Press.
15
16
- 17 Just, M. A., Keller, T. A., Malave, V. L., Kana, R. K., & Varma, S. (2012). Autism as a neural
18
19 systems disorder: A theory of frontal-posterior underconnectivity. *Neuroscience and*
20
21 *Biobehavioral Reviews*, 36, 1292-1313.
22
23
- 24 Klin, A., & Jones, W. (2006). Attributing social and physical meaning to ambiguous visual
25
26 displays in individuals with higher functioning autism spectrum disorders. *Brain &*
27
28 *Cognition*, 61, 40-53.
29
30
- 31 Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen D. J. (2002a). Defining and
32
33 quantifying the social phenotype in autism, *American Journal of Psychiatry*, 159, 895-
34
35 908.
36
37
- 38 Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen, D. J. (2002b). Visual fixation
39
40 patterns during viewing of naturalistic social situations as predictors of social competence
41
42 in individuals with autism. *Archives of General Psychiatry*, 59, 809-816.
43
44
- 45 Kuhn, G. Benson, V., Fletcher-Watson, S., Kovshoff, H., McCormick, C., & Kirkby, J. (2010)
46
47 Eye movements affirm: Automatic overt gaze and arrow cueing for typical adults and
48
49 adults with autism spectrum disorder. *Experimental Brain Research*, 201, 155-165.
50
51
- 52 Liversedge, S. P., & Findlay, J. M. (2000). Saccadic eye movements and cognition. *Trends in*
53
54 *Cognitive Sciences*, 4, 6-14.
55
56
57
58
59
60

- 1
2
3 Lord C., Rutter M., DiLavore P. C., Risi S., Gotham K., Bishop S. (2012). Autism diagnostic
4 observation schedule, second edition. Torrance, CA: Western Psychological Services.
5
6
7
8 Louwrese, A., Tulen, J. H. M., van der Geest, J. N., van der Ende, J., Verhulst, F. C., & Greaves-
9
10 Lord, K. (2014). Autonomic Responses to Social and Nonsocial Pictures in Adolescents
11 With Autism Spectrum Disorder. *Autism Research*, 7, 17-27. doi:10.1002/aur.1327
12
13
14
15 Minshew, N. J., & Goldstein, G. (1998). Autism as a disorder of complex information
16 processing. *Mental Retardation and Developmental Disabilities Research Reviews*, 4,
17 129-136.
18
19
20
21
22 Minshew, N. J., & Williams, D. L., (2007). The new neurobiology of autism: cortex,
23 connectivity, and neuronal organization. *Archives of Neurology*, 64, 945-950.
24
25
26
27 Minshew, N. J., Goldstein, G., & Siegel, D. J. (1997). Neuropsychologic functioning in autism:
28 Profile of complex information processing disorder. *Journal of the International*
29 *Neuropsychological Society*, 3, 303-316.
30
31
32
33
34 Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002).
35 Visual scanning of faces in autism, *Journal of Autism and Developmental Disorders*, 32,
36 249-261.
37
38
39
40
41 Rayner, K. (2009). Eye movements and attention in reading, scene perception, and visual search.
42 *Quarterly Journal of Experimental Psychology*, 62, 1457-1506.
43
44
45
46 Rayner, K., Castelhana, M. S., & Yang J. (2009). Eye Movements when Looking at
47 Unusual/Weird Scenes: Are there Cultural Differences? *Journal of Experimental*
48 *Psychology: Language, Memory & Cognition*, 35, 254-259.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Rayner, K., Warren, T., Juhasz, B. J., & Liversedge, S. P. (2004). The effect of plausibility on
4
5 eye movements in reading. *Journal of Experimental Psychology-Learning Memory and*
6
7
8 *Cognition*, 30(6), 1290-1301.

9
10 Riby, D. M., & Hancock, P. J. B. (2009). Do faces capture the attention of individuals with
11
12 Williams syndrome or Autism? Evidence from tracking eye movements, *Journal of*
13
14 *Autism and Developmental Disorders*, 39, 421-431. doi:10.1007/s10803-008-0641-z

15
16
17 Russell, J. (1997). *Autism as an executive disorder*. Oxford, UK: Oxford University Press.

18
19
20 Speer, L. L., Cook, A. E., McMahon, W. M., & Clark, E. (2007). Face processing in children
21
22 with autism: effects of stimulus contents and type, *Autism*, 11, 263-277.

23
24
25 Stampe, D. M. (1993). Heuristic filtering and reliable calibration methods for video-based pupil-
26
27 tracking systems. *Behavior Research Methods, Instruments, & Computers*. 25, 137-
28
29 142.

30
31
32 The Psychological Corporation, Wechsler Abbreviated Scale of Intelligence (WASI) Manual,
33
34 (The Psychological Corporation, San Antonio, TX, 1999).

35
36
37 Wolfe, J. M. (1998) What can one million trials tell us about visual search? *Psychological*
38
39 *Science*, 9, 33-39.

Footnotes

1
2
3
4
5
6 1. Accuracy errors for the excluded scenes were either a result of a failure to detect a subtle
7
8 oddity, or were a result of participants thinking the control version of the scenes were odd.
9
10 Removing these items from analysis did not change the pattern of effects, but reduced variation
11
12 in reaction time and eye movement data.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1.

Accuracy and reaction time means (standard deviations).

	Perceptual				Social			
	Normal		Weird		Normal		Weird	
	TD	ASD	TD	ASD	TD	ASD	TD	ASD
Accuracy	0.81 (0.20)	0.86 (0.10)	0.84 (0.20)	0.77 (0.17)	0.80 (0.17)	0.83 (0.12)	0.80 (0.21)	0.76 (0.24)
Reaction Time (ms)	5289 (2372)	6024 (2560)	3697 (1703)	5184 (2228)	5117 (2443)	6053 (2884)	3056 (1270)	4633 (2131)

Table 2.

Eye movement measure means in ms (standard deviations).

		Perceptual Oddity		Social Oddity	
		TD	ASD	TD	ASD
Mean Fixation Duration (overall)		250 (31)	257 (29)	241 (26)	252 (33)
Mean Fixation Count (overall)		11 (4)	15 (6)	9 (4)	13 (7)
Elapsed time to target region	ELT	551 (235)	557 (144)	600 (235)	679 (284)
Fixation count before target region	FCT	2.32 (0.72)	2.49 (0.67)	2.62 (0.76)	2.68 (0.73)
First fixation duration in target region	FFD	224 (52)	237 (42)	232 (45)	210 (41)
First gaze duration in target region	FGD	808 (250)	924 (437)	630 (163)	715 (208)
Mean fixation duration in target region	MFD	262 (43)	274 (34)	281 (38)	282 (45)
Mean fixation count in target region	MFC	5.78 (2.28)	7.72 (2.84)	3.97 (1.81)	5.47 (2.14)
Mean number of entries into target region	MNE	1.93 (0.64)	2.32 (0.73)	1.62 (0.52)	2.07 (0.66)
Mean total time in target region	TT	1522 (729)	2065 (809)	1040 (527)	1450 (664)

Figures captions

Figure 1. Example of social normal (A), social weird (B), perceptual normal (C) and perceptual weird (D) stimuli. The black square represents the target region. Note the black square is not visible during the experiment.

Figure 2. Interaction for first fixation duration (ms) between group and oddity type. Error bars denote \pm 1 SE.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RUNNING HEAD: LOOKING, SEEING AND BELIEVING IN AUTISM.

1

Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle
cognitive processing differences impact in the social domain.

Valerie Benson ^{1*}, Monica S. Castelhana ², Philippa L. Howard ¹, Nida Latif ² & Keith Rayner ³

¹ AU Psychology, Shackleton Building, Highfield Campus, University of Southampton, UK.

² Queen’s University, Kingston, ON, Canada

³ University of California, San Diego, USA

*Correspondence to: vb1@soton.ac.uk

Telephone: +44 (0) 2380594596

Lay Abstract

Adults with High Functioning Autism (ASD) viewed scenes with people in them, whilst having their eye movements recorded. The task was to indicate, using a button press, whether the pictures were normal, or in some way weird or odd. Oddities in the pictures were categorized as violations of either perceptual or social norms. Compared to a Typically Developed (TD) control group, the ASD participants were equally able to categorise the scenes as odd or normal, but they took longer to respond. The eye movement patterns showed that the ASD group made more fixations and revisits to the target areas in the odd scenes compared to the TD group. Additionally, when the ASD group first fixated the target areas in the scenes, they failed to initially detect the social oddities. These two findings have clear implications for processing difficulties in ASD for the social domain, where it is important to detect social cues on-line, and where there is little opportunity to go back and recheck possible cues in fast dynamic interactions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

3

Scientific Abstract

Autistic Spectrum Disorder (ASD) and Typically Developed (TD) adult participants had to decide whether scenes were 'odd', 'unusual' or 'weird'. All scenes contained people, with 'oddness' operationally defined as either perceptual or social. There were no group differences in accuracy, but the ASD group were slower overall to respond manually. Eye movements revealed that the ASD group made more fixations overall and more re-fixations into the target regions of the pictures, for both conditions. Importantly, as indexed by the first fixation duration the ASD group failed to 'pick up' immediately on what was 'odd' for the social violations. The propensity to 'go back' and re-fixate targets, coupled with a failure to initially detect social violations, has obvious significance for ASD in fast dynamic social communication.

Key phrases: Autism Spectrum Disorder, Eye movements, On-line cognitive processing, Social and Perceptual Oddities.

Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle cognitive processing differences impact in the social domain.

Autism spectrum disorders (ASD) are lifelong neuro-developmental conditions characterized by repetitive behaviours and impairments in interaction and communication (American Psychiatric Association, 2013). Incidence rates are rising across the globe with the prevalence rate in the UK approximately 10 per 1000 (1%) individuals (Brugha et al., 2011) and the UK National Audit office recently called for more research into adults with ASD (Clarke et al., 2009).

High functioning adults with ASD are similar to typically developed (TD) adults in terms of performance on pencil and paper and IQ tests, however in the social domain evidence suggests an inability to process social information in a typical fashion. Eye movement studies report unusual gaze behaviour for faces appearing on their own (Dalton et al., 2005), in social scenes (Riby & Hancock, 2009) or in dynamic movie clips (Klin & Jones, 2006; Speer et al., 2007). Research consistently reports a lack of spontaneous gaze fixation towards the eyes (Jones et al., 2008; Klin et al., 2002a; Pelphrey et al., 2002). However, where one study reports impairments, say for face processing (Klin et al., 2002b), another study does not (Fletcher-Watson et al., 2009). Moreover, lack of attentional modulation for social stimuli (Bird et al., 2006) may be overridden, as when cued, attention may be allocated to faces in a more 'typical' manner (Bar-Haim et al., 2006).

A recent scene perception study (Louwse et al., 2014) reported that abnormal autonomic responses in ASD are unlikely to underpin social processing impairments. It is interesting then to ask whether impairments in the cognitive domain in ASD (Happé & Frith, 2006; Minshew et al., 1997; Russell, 1997) could account for impairments in the social domain?

Current cognitive theories of ASD include among others, Weak Central Coherence (WCC) theory (Frith & Happé, 1994; Happé & Frith, 2006) which proposes that a reduced sensitivity to global context in favour of focussing on details, results in integration difficulties in ASD. Executive function theory (Russell, 1997), proposes that ASD are impaired at switching their focus of attention, and may engage in perseverative behaviour. Alternatively, the Complex Information Processing theory (Minshew & Goldstein, 1998) proposes that individuals with ASD have impairments in processing complex information with intact processing for tasks that require simple (rule based) processing. Finally, the Extreme Male Brain (EMB) theory (Baron-Cohen, 2002) posits that individuals with ASD have above average levels of systemizing, that contribute to the non-social characteristics of the disorder, coupled with below average levels of empathizing, which results in atypical social functioning. These contemporary theories of ASD can, for different reasons, offer some account offer ineffective social functioning in ASD. ~~However, but sparse research~~ to date has very few investigations ~~have examined~~ time course processing differences as these occur on-line in ASD, ~~and how these might impact in the social domain.~~

Our studies investigating cognitive processing in adults with ASD (Au-Yeung et al., 2011; Au-Yeung et al., 2013; Benson et al., 2012; Benson et al., 2009; Benson & Fletcher-Watson, 2011) have found subtle on-line processing differences, as revealed by eye movements, for a range of tasks that are more in line with Minshew and Goldsteins (1998) theory, than with any other. For example, there are more similarities than differences in eye movements between TD and ASD groups for simple rule based tasks in the attentional (Kuhn et al., 2010) and the social domains (Benson & Fletcher-Watson, 2011). Whereas for more complex abstract reasoning tasks where there is ambiguity with relation to target items (Au-Yeung et al., 2013), or

where value judgements rely on top down knowledge (Benson et al., 2012), there are early processing differences indicating that initial orienting to, and immediate detection of weird target items in scenes is absent in ASD. Moreover, the time to respond and to locate the weird target items was modulated by whether or not the pictures had people in them (Benson et al., 2012).

These differences have potential ~~forte~~ impact ~~on~~ everyday activities, since the time course for detection of socially relevant information is crucial for successful everyday interactions.

The aim of the present study was to examine whether increased times to find and initially detect an unusual target in a scene would be modulated by whether the violation that rendered the target unusual or odd could be categorised as social or perceptual. We employed a similar paradigm to that used in a previously reported study (Rayner et al., 2009) where single scenes, all with people in them, were presented in isolation and the task was to indicate whether each picture was normal or in some way weird unusual or odd.

Insert Figure 1. about here

Social oddities in the pictures were operationally defined as violations of social norms e.g. someone in the picture throwing a baby in the air. Access to top down knowledge is required in that condition to inform a subjective value judgement as to whether a social violation is present in the picture. No value judgement as to whether a violation infringes some implicit or explicit social rule is needed for the perceptual oddities in this study, which are operationally defined as physical violations e.g. someone in the picture has a limb positioned back to front. See supplementary information for Materials ~~and Methods~~. Figure 1 shows an example of the stimuli.

Method

Participants

ASD participants ($n = 24$ age 18 – 52 years, 20 males), with a formal diagnosis of either Autistic Disorder ($n = 3$) or Asperger's Syndrome were recruited from local charitable organisations. TD participants ($n = 24$, age 20– 52 years, 20 males) were volunteers from the local community. All participants were paid and had normal or corrected to normal vision. Each ASD participant completed module 4 of the Autism Diagnostic Observation Scale 2 (Lord et al., 2012) and five (3 males and 2 females with a diagnosis of Asperger's Syndrome) failed to meet the Autism Spectrum criteria. To ensure maximum homogeneity in our study those five participants were excluded from the analyses, resulting in a final sample of 19 ASD participants. Both groups completed the Autism-spectrum Quotient (AQ; Baron-Cohen et al., 2001) and the ASD participants reported a higher number of autistic traits $t(41) = 8.04, p < .001$ (TD $M = 15.42, SD = 7.55$; ASD $M = 35.32, SD = 8.45$). The two groups did not differ in age $t(41) = 0.31, p = .759$ (TD $M = 30.29, SD = 10.20$; ASD $M = 31.26, SD = 10.29$), performance IQ $t(41) = 0.21, p = .832$ (TD $M = 111.79, SD = 12.01$; ASD $M = 110.95, SD = 13.84$), verbal IQ $t(41) = 1.21, p = .234$ (TD $M = 116.96, SD = 10.65$, ASD $M = 112.26, SD = 14.84$), or full scale IQ $t(46) = 0.87, p = .392$ (TD $M = 116.04, SD = 9.98$, ASD $M = 112.95, SD = 13.48$), as measured by two performance (Block Design and Matrix Reasoning) and verbal components (Vocabulary and Similarity) of the Weschler Abbreviated Scale of Intelligence (The Psychological Corporation, 1999).

Eye movement recording

Eye movements were sampled at 1000Hz using an Eyelink tracker with spatial accuracy of less than $1/4^{\text{th}}$ degree (S.R. Research Ltd, Canada). The start and end of saccades were

Formatted: Font: (Default) Times New Roman, 12 pt

detected automatically using a default velocity criterion of 30degrees per second using the Eyelink software. See Stampe (1993) for an in depth technical description of how the software detects saccades and fixations. A chin rest and forehead support, were used to stabilize head position. Viewing was binocular but data were analysed for the right eye only. Individual participants were calibrated using a nine point matrix that covered the dimensions of the screen (1024 x 768 pixels), with each point fixated sequentially, followed by a validation procedure to ensure that fixation was within 0.5 degrees of each calibration point. Calibration was checked using a validation procedure prior to each trial, whereby participants fixated five dots, presented at five different locations that covered the dimensions of the screen and recalibration was performed if the eyes were off centre for any of these dots.

Materials

In total, 96 full-colour photographs were used as stimuli, all of which contained people. In both the perceptual and social oddity conditions there were 48 pictures (24 weird and 24 normal). Perceptual oddities in the pictures violated a physical norm e.g. someone in the picture has a limb positioned back to front, or a vehicle was in an unexpected position e.g midway through the upper wall of a building. Social oddities in the pictures violated a social norm e.g. someone in the picture throwing a baby in the air, or someone with a cabbage on a lead. A small number of the pictures were from a previous study (Benson et al., 2012) and the remainder were collected from the internet and included either a perceptual or social oddity. Adobe Photoshop was used to prepare a normal version of each scene in which the oddity was replaced with either the background elements or with another suitable object. The scenes typically had a focal object or a central group of objects. Figure 1 shows an example of a normal and a socially odd scene,

and below those examples, a normal and a perceptually odd scene. The target regions for these scenes are outlined in black – but these black outlines were not visible during the presentation of the scenes in the experiment.

A norming study was conducted to verify the validity of the images used to examine the processing of social and perceptual violations. Twenty undergraduates from Queen's university participated (M age: 19.8years, SD : 1.71, 10 Females) with five participants in each of the four conditions (counterbalanced conditions: weird/normal and right/left button response). Stimuli were displayed on a 21 inch CRT monitor with a refresh rate of 100Hz and the experiment was run using E-Prime software. Participants were presented with a fixation cross for 500ms followed by the presentation of the image for 20 seconds or until a keyboard "Y" (yes) or "N" (N) was made. Participants had to indicate as quickly as possible whether the presented image was "odd", "weird" or "unusual" in some way. Results-The proportion of identified images indicated that on average, the social normal ($M = 0.86$, $SD = 0.15$) and social weird ($M = 0.84$, $SD = 0.17$), perceptual normal ($M = 0.82$, $SD = 0.17$) and perceptual weird ($M = 0.77$, $SD = 0.19$) images were identified correctly.

Design

For the behavioural measures of Accuracy and RT a mixed design with Picture Type (weird versus normal) and Oddity Type (social versus perceptual) as within participants variables, and Group (ASD versus TD) as a between participants variable was employed for the experiment. On condition that accuracy was equivalent for both normal and weird pictures, for the eye movement measures the main focus of interest was to compare social versus perceptual oddities in both groups. Therefore a mixed design with Oddity Type (social versus perceptual) as

a within participants variable, and Group (ASD versus TD) as a between participants variable was employed in the eye movement analysis.

Procedure

All participants were shown a hard copy of an example picture from each condition prior to completing the task, to ensure comprehension of the task and the ability to respond appropriately. The specific instruction was 'for each of the following pictures, please indicate whether you think the picture is odd, unusual or strange in some way'.

For the task, 48 pictures were presented randomly to either the left or the right of the centre of a 19inch colour monitor, at a distance of 65cm for 20s or until a button press response was made. Prior to the onset of a picture, participants fixated a cross that was presented on the opposite side of the screen to where the picture was set to appear, to ensure all participants started at the same point in each trial. Once participants fixated this cross, the experimenter initiated each trial. Picture type (normal vs. weird) and, the position of the picture (left or right on the display) were counterbalanced across items. The buttons used to respond either yes or no (left or right trigger button) was counterbalanced across participants.

Data preparation and analyses

For each 'odd' picture we created an interest region around the target item and the corresponding region of the same size in the counterpart 'normal' picture. Regions of interest were created using the approximate average regions from those identified and manually selected by participants who took part in the norming experiment.

Formatted: Line spacing: Double

Formatted: Font: (Default) Times New Roman, 12 pt

Eyelink software was used to detect saccades and fixations using a default velocity criterion and in-house software was used to calculate the distribution of fixations falling into the target region. Five items were removed from analysis because for these scenes, participant accuracy scores were significantly below chance¹. In addition, each participant was asked whether they had seen any of the scenes previously, and if so these scenes were also removed. This resulted in a loss of 2.3% of trials. Fixations below 90ms and above 1200ms were classed as outliers and removed; resulting in a loss of 5.13% of the fixation data.

Results

For the principal analyses we compared manual reaction times and accuracy (see Table 1), as well as a number of eye movement measures, such as, the time taken to begin fixating in the target region (ELT), the duration of the first fixation when the eyes landed in the target region (FFD) and the mean time (MFD) spent fixating in the target area (see Table. 2). Global eye movement measures for the target region, such as mean fixation duration and mean number of fixations over all trials were also analysed, Finally, baseline measures (not just target related), were calculated across the duration of the trials, to check for overall sampling differences between the ASD and TD participants (see Table 2). See supplementary information for methods and materials.

Behavioural Data

Accuracy. There was no effect of oddity type (perceptual vs. social) $F(1, 41) = 1.44, p = .237, \eta_p^2 = .03$, picture type (normal or weird) $F(1, 41) = 0.94, p = .339, \eta_p^2 = .02$, or group, $F(1, 41) = .06, p = .808, \eta_p^2 < .01$ on accuracy rates, and no interactions ($F_s < 2.39, p's > .130$). At

Formatted: Font: Italic

least in terms of accuracy, performance was the same for both groups. For means and standard deviations for behavioural data see Table 1.

Response Time (RT). A main effect of oddity type $F(1, 41) = 8.16, p = .007, \eta_p^2 = .17$, showed participants took longer to respond when presented with a perceptual oddity in comparison to a social oddity. Longer response times for perceptual oddities could reflect that these ‘impossible’, physical violations were so unusual that they resulted in more time spent looking at them prior to responding. See below for confirmation of that in the eye movement data. There was also a main effect of picture type $F(1, 41) = 24.03, p < .001, \eta_p^2 = .37$, with participants taking longer to respond when the picture was normal in comparison to when it was weird, in line with longer reaction times for target absent trials in standard visual search tasks (Wolfe, 1998). A significant oddity type x picture type interaction $F(1, 41) = 4.22, p = .046, \eta_p^2 = .09$, demonstrated that the difference between RTs for perceptual and social pictures was reliable for odd (TD; $t(23) = 2.62, p = .015$, ASD; $t(18) = 2.98, p = .008$) but not for normal pictures (TD; $t(23) = 0.94, p = .355$, ASD; $t(18) = 0.08, p = .935$). A main effect of group $F(1, 41) = 4.07, p = .050, \eta_p^2 = .09$, showed the ASD participants had longer reaction times, in line with a previous study (Benson et al., 2012, 25). There were no significant group interactions ($F_s < 1$) demonstrating both groups to be slower for normal versus odd pictures, and for perceptual vs social oddities. See Table 1 for the means. The eye movement data below reveals subtle processing differences that contribute to this overall slowed manual responding in the ASD group.

Formatted: Font: Not Italic

Eye Movement Data

Having shown that all participants could distinguish between 'odd' and 'normal' pictures, normal picture trials were excluded from the eye movement analyses resulting in a 2 (group: ASD versus TD) X 2 (oddy type: social versus perceptual) design to investigate on-line processing differences between the two groups for the two types of oddities.

Baseline eye movement measures, calculated across the full length of trials indicated no group difference in mean fixation duration $F(1, 41) = 1.14, p = .293, \eta_p^2 = .03$, but the ASD group made more fixations overall $F(1, 41) = 9.44, p = .004, \eta_p^2 = .19$, corresponding to the increased RT's reported above. Table 2 presents a summary of all the eye movement measures.

Elapsed time to the target region (ELT). A marginal effect of oddity type $F(1, 41) = 3.99, p = .053, \eta_p^2 = .09$, on the time taken from the onset of each trial display to the start of the first fixation on the target region, revealed both groups took less time to fixate the target region when the oddity was perceptual in comparison to social in nature. ~~The impossible nature of the perceptual violation may have increased the salience of the target region but there~~ there was no effect of group $F(1, 41) = 0.59, p = .446, \eta_p^2 = .01$, and no interaction $F(1, 41) = 0.72, p = .401, \eta_p^2 = .02$.

First Fixation Duration (FFD) on target region. There was no effect of oddity type $F(1, 41) = 1.84, p = .182, \eta_p^2 = .04$ or group $F(1, 41) = 0.15, p = .699, \eta_p^2 < .01$ but a significant interaction $F(1, 41) = 6.72, p = .013, \eta_p^2 = .14$ (See Figure 2) revealed that FFDs were equivalent for perceptual and social oddities for TD participants $t(23) = 0.86, p = .399, d = .16$, but for ASD participants FFDs for social oddities were significantly shorter, in comparison to perceptual oddities, $t(18) = 2.99, p = .008, d = .65$, see Figure 2. Given that the FFD is a measure of early

processing (Liversedge & Findlay, 2000; Rayner, 1998; Rayner, 2009), this finding indicates that the ASD group did not detect the social oddities as quickly as the perceptual oddities on initial fixation, whereas TD participants detected both types of oddity equally fast.

Insert Figure 2. about here

First gaze duration (FGD) in the target region. The FGD is the elapsed time from when the target region is first fixated until the eye moves on to a new location. FGD includes multiple fixations in the target region reflects an overt strategy to re-examine an area. A main effect of oddity type, $F(1, 41) = 12.10, p = .001, \eta_p^2 = .23$ demonstrated that both groups spent more time inspecting perceptual oddities, again highlighting the salience of the perceptual oddities in comparison to social oddities in this study. However, there was no group effect $F(1, 41) = 2.43, p = .127, \eta_p^2 = .06$, or interaction $F(1, 41) = .08, p = .785, \eta_p^2 < .01$.

Mean Fixation Count in the target region. If the ASD are slower to respond manually, but do not take longer to begin inspecting the target region, then extra time to respond might reflect more fixations in the target region. A main effect of oddity type $F(1, 41) = 64.27, p < .001, \eta_p^2 = .61$, demonstrated more fixations in the target region when the oddity was perceptual, rather than social. A main effect of group $F(1, 41) = 6.99, p = .012, \eta_p^2 = .15$ showed that the ASD participants made more fixations than the TD group in the target regions, for both types of oddities overall. However no interaction was observed $F(1, 41) = 0.77, p = .386, \eta_p^2 = .02$ indicating that the ASD group made more fixations in the target regions for both types of oddities.

Formatted: Highlight

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Highlight

Mean Fixation Duration in the target region. Longer mean fixation durations indicate that more processing has been done during the task. A main effect of oddity type $F(1, 41) = 8.22, p = .007, \eta_p^2 = .17$ showed significantly longer fixations for social, compared to perceptual oddity target regions. This suggests that processing social oddities was more effortful, and since there was no main effect of group $F(1, 41) = 0.34, p = .562, \eta_p^2 < .01$, and no interaction $F(1, 41) = 1.12, p = .297, \eta_p^2 = .03$, both groups on average fixated the social oddity target region for longer than the perceptual target region. Thus, while the perceptual oddities attracted more fixations overall, individual fixations were longer for the social target region.

This interesting juxtaposition, although unrelated to the main aim of this study, suggests that difficulty associated with processing each oddity type manifests in different eye movement patterns: one reflecting an overt strategy (fixation count) to re-inspect the target area, and one reflecting more in depth on-line cognitive processing (fixation duration) on the target areas.

Mean Number of Entries to the target region. Since there were no group differences in FGD, and no increased MFD, both reported above, the delayed RT in ASD might reflect repeated visits to inspect the target region?

A main effect of oddity type was found $F(1, 41) = 9.83, p = .003, \eta_p^2 = .19$, with the target region entered more often for perceptual oddities in comparison to social oddities. In addition, a main effect of group was found $F(1, 41) = 5.72, p = .021, \eta_p^2 = .12$, with ASD participants entering the target regions significantly more than the TD group, before making a response. There was no interaction $F(1, 41) = 0.13, p = .717, \eta_p^2 < .01$.

Mean Total Time in the target region. A higher mean total time means that more attention to, and processing of, information in that region was carried out overall. This measure should complement the measures reported above – namely, that if there are more fixations overall in ASD then the Total Time spent fixating in the target region should be longer in ASD.

A main effect of oddity type $F(1, 41) = 34.89, p < .001, \eta_p^2 = .46$, showed perceptual oddities received more overall processing time, in line with the previous measures for the perceptual oddities. In addition, a main effect of group $F(1, 41) = 6.41, p = .015, \eta_p^2 = .14$ revealed that the ASD participants spent a longer period of time inspecting the target regions for both perceptual and social oddities. There was no interaction $F(1, 41) = 0.51, p = .481, \eta_p^2 = .01$.

As above, this finding reflects a greater number of entries to the target region overall by the ASD group prior to making their decision as to whether the picture under inspection was weird, odd or unusual, irrespective of the nature of the violation (perceptual or social).

Discussion

A principal aim of the study was to investigate whether increased times in ASD to find and initially detect an unusual target in a scene containing people would be modulated by whether the violation could be categorised as social or perceptual.

Accuracy rates were equivalent and high for both groups, but the ASD group took longer to respond manually. It was hypothesised that the eye movement data would reveal the cause of this delayed response in ASD, since eye movements reflect how information is processed on-line (Liversedge & Findlay, 2000; Rayner, 1998; 2009). The data showed that our ASD group did not take longer to find the target regions, but that they scanned the scenes more prior to responding manually, as shown by an increased number of fixations throughout each scene presentation.

Furthermore, these increased fixations were not random, but were made to the target regions of both social and perceptually odd pictures. Thus, increased fixations reflect a propensity to re-inspect the target regions prior to making a response in ASD, regardless as to the type of oddity in the scenes. It should be noted here that the oddity detection task in the current study provides the ASD participants with structure that is not typically available in real-life. As such, the task is not implicit and the knowledge that oddities might be present could also be a factor that drives the repeated scanning in ASD.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

The eye movement patterns also showed that our ASD group failed to immediately detect social oddities, thus looking at but not seeing them. An obvious question from this finding is whether the failure to quickly spot subtle social oddities in ASD reflects an inability to react or respond appropriately to social cues in everyday communication? This one impairment in itself could potentially account for the significant problems experienced in dynamic social interactions in this high functioning population. For example, a failure to immediately spot social cues would result in a failure to react or respond to such cues thus resulting in an inability to both keep track of what is going on in social interactions, and a failure to engage appropriately or effectively with other people.

~~In relation to current theories that underpin ASD our findings do not fit with WCC theory (Frith & Happé, 1994; Happé & Frith, 2006), since clearly context can be taken into account in ASD during task completion. Moreover, the slowness to respond in ASD is not reflected in longer times to find the target, and the increased fixations in ASD were seen for both perceptual and social oddities in the scenes. It may be that repeated visits to the target area indicate integration problems in ASD, but even if that were the case, such problems are not restricted to social processing per se. Neither do they support the executive function theory (Russell, 1997), as~~

~~ASD participants showed no indication of increased mean fixation durations, thus no evidence of perseveration or any inability to switch focus throughout the task. Our findings could however be interpreted to support the EMB (Baron-Cohen, 2002), with the detection of social oddities that required empathy being delayed in ASD, however note that higher re-inspection of perceptual oddities that could be detected on the basis of systematic rules, was also found for ASD participants.~~

If social oddities are more complex to detect in ASD, then the Complex Information Processing theory (Minsheu & Goldstein, 1998) might explain the failure to detect the social oddity on first fixation. This lack of immediate detection could reflect the less obvious and more 'inferential' nature of the social oddities, which required more complex reasoning, at least in the ASD group, for detection. It is important to note that social rules are often implicit, sometimes ambiguous, and cannot always be understood by following a script or flow chart as in the case for logical rule based tasks. At a neurobiological level, it has been postulated that that impairments in cognitive processing in ASD reflect slow information transmission rates for high level cognitive tasks, resulting from underconnectivity between frontal-posterior cortical regions (Just et al., 2013; Just et al., 2012; Minsheu & Williams, 2007). Since underconnectivity reduces the rate of inter-region information transfer, it could be that in social communication tasks that rely on the fast detection of implicit, often ambiguous cues, a failure to be able to transmit information across different cortical areas quickly, might mean that these cues are missed, rendering inappropriate responses in ASD.

The findings of repeated scanning of the target areas, coupled with an inability to immediately recognise social violations upon initial fixation, reflect how eye movements can reveal subtle processing differences in ASD. In everyday social communication in the real world

Formatted: Not Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Not Highlight

Formatted: Not Highlight

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

there is no opportunity to ‘go back’ to check if one has missed an important cue. Social processing in such circumstances must be carried out on-line, and our eye movement data clearly show that ASD are unable to do this quickly. The strategy of re-checking their interpretation, and thus not believing what they have seen, before responding, may be related to cortical underconnectivity in ASD. In everyday terms, these findings have obvious consequences for fast dynamic social interactions, and offer an account as to why high functioning adults with ASD experience difficulty in the social domain.

For Peer Review

Acknowledgments

This work was supported by a grant awarded to the second author from the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the Ontario Ministry of Research and Innovation, and Queen’s University Senate Advisory Research Committee, and an Experimental Psychology Society study visit grant awarded to the third author.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Au-Yeung, S. K., Benson, V., Castelhana, M. S., Rayner, K. (2011). Eye movement sequences during simple versus complex information processing of scenes in autism spectrum disorder. *Autism Research and treatment*, 2011.
- Au-Yeung, S. K., Kaakinen, J. K., Benson, V. (2013). Cognitive perspective taking during scene perception in autism spectrum disorder: evidence from eye movements. *Autism Research*, 7, 84-93.
- Bar-Haim, Y., Shulman, C., Lamy, D., & Reuveni, A. (2006). Attention to eyes and mouth in high-functioning children with autism, *Journal of Autism and Developmental Disorders*, 36, 131-137.
- Baron-Cohen, S. (2002). The extreme male brain theory of autism. *Trends in Cognitive Science*, 6, 248-254.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5-17.
- Benson, V., & Fletcher-Watson, S. (2012). Eye Movements in Autism. In: Liversedge, S. P., Gilchrist, I. D., Everling, S., editors. *The Oxford Handbook of Eye Movements*, Oxford UK: Oxford University Press, chap 39.
- Benson, V., Castelhana, M. S., Au-Yeung, S. K., & Rayner K. (2012). Eye movements reveal no immediate "WOW" ("which one's weird") effect in autism spectrum disorder, *Quarterly*

Journal of Experimental Psychology, 65, 1139-1150.

Benson, V., Piper, J., & Fletcher-Watson, S. (2009). Atypical saccadic sampling in autistic spectrum disorder, *Neuropsychologia*, 47, 1178-1182, (2009).

Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate neural responses to social stimuli in autism spectrum disorders. *Neuroimaging*, 15, 1614-1624.

Brugha, T. S., McManus, S., Bankart, J., Scott, F., Purdon S., et al. (2011). Epidemiology of autism spectrum disorders in adults in the community in England. *Archives of General Psychiatry*, 68, 459-65.

Clark, F., Scharaschkin, A., Xu, D. (2009). Supporting autism through adulthood, National Audit Office.

Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., Alexander, A. L., & Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nature Neuroscience*, 8, 519-526.

Fletcher-Watson, S., Leekam, S. R., Benson, V., Frank, M. C., & Findlay, J. M. (2009). Eye-movements reveal attention to social information in autism spectrum disorder. *Neuropsychologia*, 47, 248-257.

Fletcher-Watson, S., Leekam, S.R., Benson, V., Frank, M.C., & Findlay, J.M. (2009). Eye-movements reveal attention to social information in autism spectrum disorder. *Neuropsychologia*, 47, 248-257.

Frith, U., & Happé, F. (1994). Autism; Beyond the theory of mind. *Cognition*, 50, 115-132.

Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 36, 5-25.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

23

Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder. *Archives of General Psychiatry*, 65, 946-954.

Just, M. A., Keller, T. A., & Kana, R. K. (2013). A theory of autism based on frontal-posterior underconnectivity. In M. A. Just & K. A. Pelphrey (Eds.), *Development and brain systems in autism* (pp. 35-63). New York: Psychology Press.

Just, M. A., Keller, T. A., Malave, V. L., Kana, R. K., & Varma, S. (2012). Autism as a neural systems disorder: A theory of frontal-posterior underconnectivity. *Neuroscience and Biobehavioral Reviews*, 36, 1292-1313.

Klin, A., & Jones, W. (2006). Attributing social and physical meaning to ambiguous visual displays in individuals with higher functioning autism spectrum disorders. *Brain & Cognition*, 61, 40-53.

Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen D. J. (2002a). Defining and quantifying the social phenotype in autism, *American Journal of Psychiatry*, 159, 895-908.

Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen, D. J. (2002b). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59, 809-816.

Kuhn, G., Benson, V., Fletcher-Watson, S., Kovshoff, H., McCormick, C., & Kirkby, J. (2010). Eye movements affirm: Automatic overt gaze and arrow cueing for typical adults and adults with autism spectrum disorder. *Experimental Brain Research*, 201, 155-165.

Liversedge, S. P., & Findlay, J. M. (2000). Saccadic eye movements and cognition. *Trends in Cognitive Sciences*, 4, 6-14.

- 1
2
3
4
5
6
7
8
9 Lord C., Rutter M., DiLavore P. C., Risi S., Gotham K., Bishop S. (2012). Autism diagnostic
10 observation schedule, second edition. Torrance, CA: Western Psychological Services.
11
12 Louwerse, A., Tulen, J. H. M., van der Geest, J. N., van der Ende, J., Verhulst, F. C., & Greaves-
13 Lord, K. (2014). Autonomic Responses to Social and Nonsocial Pictures in Adolescents
14 With Autism Spectrum Disorder. *Autism Research*, 7, 17-27. doi:10.1002/aur.1327
15
16 Minshew, N. J., & Goldstein, G. (1998). Autism as a disorder of complex information
17 processing. *Mental Retardation and Developmental Disabilities Research Reviews*, 4,
18 129-136.
19
20 Minshew, N. J., & Williams, D. L., (2007). The new neurobiology of autism: cortex,
21 connectivity, and neuronal organization. *Archives of Neurology*, 64, 945-950.
22
23 Minshew, N. J., Goldstein, G., & Siegel, D. J. (1997). Neuropsychologic functioning in autism:
24 Profile of complex information processing disorder. *Journal of the International*
25 *Neuropsychological Society*, 3, 303-316.
26
27 Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002).
28 Visual scanning of faces in autism, *Journal of Autism and Developmental Disorders*, 32,
29 249-261.
30
31 Rayner, K. (2009). Eye movements and attention in reading, scene perception, and visual search.
32 *Quarterly Journal of Experimental Psychology*, 62, 1457-1506.
33
34 Rayner, K., Castelano, M. S., & Yang J. (2009). Eye Movements when Looking at
35 Unusual/Weird Scenes: Are there Cultural Differences? *Journal of Experimental*
36 *Psychology: Language, Memory & Cognition*, 35, 254-259.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rayner, K., Warren, T., Juhasz, B. J., & Liversedge, S. P. (2004). The effect of plausibility on eye movements in reading. *Journal of Experimental Psychology-Learning Memory and Cognition*, 30(6), 1290-1301.

Riby, D. M., & Hancock, P. J. B. (2009). Do faces capture the attention of individuals with Williams syndrome or Autism? Evidence from tracking eye movements, *Journal of Autism and Developmental Disorders*, 39, 421-431. doi:10.1007/s10803-008-0641-z

Russell, J. (1997). *Autism as an executive disorder*. Oxford, UK: Oxford University Press.

Speer, L. L., Cook, A. E., McMahon, W. M., & Clark, E. (2007). Face processing in children with autism: effects of stimulus contents and type, *Autism*, 11, 263-277.

Stampe, D. M. (1993). Heuristic filtering and reliable calibration methods for video-based pupil-tracking systems. *Behavior Research Methods, Instruments, & Computers*, 25, 137-142.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Indent: Left: 0", Hanging: 0.5", Line spacing: Double

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

The Psychological Corporation, Wechsler Abbreviated Scale of Intelligence (WASI) Manual, (The Psychological Corporation, San Antonio, TX, 1999).

Wolfe, J. M. (1998) What can one million trials tell us about visual search? *Psychological Science*, 9, 33-39.

Footnotes

1. Accuracy errors for the excluded scenes were either a result of a failure to detect a subtle oddity, or were a result of participants thinking the control version of the scenes were odd. Removing these items from analysis did not change the pattern of effects, but reduced variation in reaction time and eye movement data.

For Peer Review

Tables

Table 1.

Accuracy and reaction time means (standard deviations).

	Perceptual				Social			
	Normal		Weird		Normal		Weird	
	TD	ASD	TD	ASD	TD	ASD	TD	ASD
Accuracy	0.81 (0.20)	0.86 (0.10)	0.84 (0.20)	0.77 (0.17)	0.80 (0.17)	0.83 (0.12)	0.80 (0.21)	0.76 (0.24)
Reaction Time (ms)	5289 (2372)	6024 (2560)	3697 (1703)	5184 (2228)	5117 (2443)	6053 (2884)	3056 (1270)	4633 (2131)

Peer Review

Table 2.

Eye movement measure means in ms (standard deviations).

		Perceptual Oddity		Social Oddity	
		TD	ASD	TD	ASD
Mean Fixation Duration (overall)		250 (31)	257 (29)	241 (26)	252 (33)
Mean Fixation Count (overall)		11 (4)	15 (6)	9 (4)	13 (7)
Elapsed time to target region	ELT	551 (235)	557 (144)	600 (235)	679 (284)
Fixation count before target region	FCT	2.32 (0.72)	2.49 (0.67)	2.62 (0.76)	2.68 (0.73)
First fixation duration in target region	FFD	224 (52)	237 (42)	232 (45)	210 (41)
First gaze duration in target region	FGD	808 (250)	924 (437)	630 (163)	715 (208)
Mean fixation duration in target region	MFD	262 (43)	274 (34)	281 (38)	282 (45)
Mean fixation count in target region	MFC	5.78 (2.28)	7.72 (2.84)	3.97 (1.81)	5.47 (2.14)
Mean number of entries into target region	MNE	1.93 (0.64)	2.32 (0.73)	1.62 (0.52)	2.07 (0.66)
Mean total time in target region	TT	1522 (729)	2065 (809)	1040 (527)	1450 (664)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RUNNING HEAD: LOOKING, SEEING AND BELIEVING IN AUTISM.

29

Figures captions

Figure 1. Example of social normal (A), social weird (B), perceptual normal (C) and perceptual weird (D) stimuli. The black square represents the target region. Note the black square is not visible during the experiment.

Figure 2. Interaction for first fixation duration (ms) between group and oddity type. Error bars denote ± 1 SE.

For Peer Review



Figure 1. Example of social normal (A), social weird (B), perceptual normal (C) and perceptual weird (D) stimuli. The black square represents the target region. Note the black square is not visible during the experiment.
149x129mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Interaction for first fixation duration (ms) between group and oddity type. Error bars denote +/- 1 SE.
228x158mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RUNNING HEAD: LOOKING, SEEING AND BELIEVING IN AUTISM

1

Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle cognitive processing differences impact in the social domain.

Valerie Benson ¹*, Monica S. Castelhana ², Philippa L. Howard ¹, Nida Latif ² & Keith Rayner ³

¹ AU Psychology, Shackleton Building, Highfield Campus, University of Southampton, UK.

² Queen's University, Kingston, ON, Canada

³ University of California, San Diego, USA

*Correspondence to: vb1@soton.ac.uk

Telephone: +44 (0) 2380594596

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

2

Lay Abstract

Adults with High Functioning Autism (ASD) viewed scenes with people in them, whilst having their eye movements recorded. The task was to indicate, using a button press, whether the pictures were normal, or in some way weird or odd. Oddities in the pictures were categorized as violations of either perceptual or social norms. Compared to a Typically Developed (TD) control group, the ASD participants were equally able to categorise the scenes as odd or normal, but they took longer to respond. The eye movement patterns showed that the ASD group made more fixations and revisits to the target areas in the odd scenes compared to the TD group. Additionally, when the ASD group first fixated the target areas in the scenes, they failed to initially detect the social oddities. These two findings have clear implications for processing difficulties in ASD for the social domain, where it is important to detect social cues on-line, and where there is little opportunity to go back and recheck possible cues in fast dynamic interactions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

3

Scientific Abstract

Autistic Spectrum Disorder (ASD) and Typically Developed (TD) adult participants had to decide whether scenes were 'odd', 'unusual' or 'weird'. All scenes contained people, with 'oddness' operationally defined as either perceptual or social. There were no group differences in accuracy, but the ASD group were slower overall to respond manually. Eye movements revealed that the ASD group made more fixations overall and more re-fixations into the target regions of the pictures, for both conditions. Importantly, as indexed by the first fixation duration the ASD group failed to 'pick up' immediately on what was 'odd' for the social violations. The propensity to 'go back' and re-fixate targets, coupled with a failure to initially detect social violations, has obvious significance for ASD in fast dynamic social communication.

Key phrases: Autism Spectrum Disorder, Eye movements, On-line cognitive processing, Social and Perceptual Oddities.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

4

Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle cognitive processing differences impact in the social domain.

Autism spectrum disorders (ASD) are lifelong neuro-developmental conditions characterized by repetitive behaviours and impairments in interaction and communication (American Psychiatric Association, 2013). Incidence rates are rising across the globe with the prevalence rate in the UK approximately 10 per 1000 (1%) individuals (Brugha et al., 2011) and the UK National Audit office recently called for more research into adults with ASD (Clark et al., 2009).

High functioning adults with ASD are similar to typically developed (TD) adults in terms of performance on pencil and paper and IQ tests, however in the social domain evidence suggests an inability to process social information in a typical fashion. Eye movement studies report unusual gaze behaviour for faces appearing on their own (Dalton et al., 2005), in social scenes (Riby & Hancock, 2009) or in dynamic movie clips (Klin & Jones, 2006; Speer et al., 2007). Research consistently reports a lack of spontaneous gaze fixation towards the eyes (Jones et al., 2008; Klin et al., 2002a; Pelphrey et al., 2002). However, where one study reports impairments, say for face processing (Klin et al., 2002b), another study does not (Fletcher-Watson et al., 2009). Moreover, lack of attentional modulation for social stimuli (Bird et al., 2006) may be overridden, as when cued, attention may be allocated to faces in a more 'typical' manner (Bar-Haim et al., 2006).

A recent scene perception study (Louwerse et al., 2014) reported that abnormal autonomic responses in ASD are unlikely to underpin social processing impairments. It is interesting then to ask whether impairments in the cognitive domain in ASD (Happé & Frith, 2006; Minshew et al., 1997; Russell, 1997) could account for impairments in the social domain?

LOOKING, SEEING AND BELIEVING IN AUTISM

5

Current cognitive theories of ASD include, among others, Weak Central Coherence (WCC) theory (Frith & Happé, 1994; Happé & Frith, 2006) which proposes that a reduced sensitivity to global context in favour of focussing on details, results in integration difficulties in ASD. Executive function theory (Russell, 1997), proposes that ASD are impaired at switching their focus of attention, and may engage in perseverative behaviour. Alternatively, the Complex Information Processing theory (Minshew & Goldstein, 1998) proposes that individuals with ASD have impairments in processing complex information with intact processing for tasks that require simple (rule based) processing. Finally, the Extreme Male Brain (EMB) theory (Baron-Cohen, 2002) posits that individuals with ASD have above average levels of systemizing, that contribute to the non-social characteristics of the disorder, coupled with below average levels of empathizing, which results in atypical social functioning. These contemporary theories of ASD can, for different reasons, offer some account of ineffective social functioning in ASD. However, but sparse research to date has very few investigations have examined online-time course processing differences as these occur on-line in ASD, and how these might impact in the social domain.

Our studies investigating cognitive processing in adults with ASD (Au-Yeung et al., 2011; Au-Yeung et al., 2013; Benson et al., 2012; Benson et al., 2009; Benson & Fletcher-Watson, 2011) have found subtle on-line processing differences, as revealed by eye movements, for a range of tasks that are more in line with Minshew and Goldstein's (1998) theory, than with any other. For example, there are more similarities than differences in eye movements between TD and ASD groups for simple rule based tasks in the attentional (Kuhn et al., 2010) and the social domains (Benson & Fletcher-Watson, 2011). Whereas for more complex abstract reasoning tasks where there is ambiguity with relation to target items (Au-Yeung et al., 2013), or

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

6

where value judgements rely on top down knowledge (Benson et al., 2012), there are early processing differences indicating that initial orienting to, and immediate detection of weird target items in scenes is absent in ASD. Moreover, the time to respond and to locate the weird target items was modulated by whether or not the pictures had people in them (Benson et al., 2012).

These differences have potential ~~for to~~ impact ~~on to~~ everyday activities, since the time course for detection of socially relevant information is crucial for successful everyday interactions.

The aim of the present study was to examine whether increased times to find and initially detect an unusual target in a scene would be modulated by whether the violation that rendered the target unusual or odd could be categorised as social or perceptual. We employed a similar paradigm to that used in a previously reported study (Rayner et al., 2009) where single scenes, all with people in them, were presented in isolation and the task was to indicate whether each picture was normal or in some way weird unusual or odd.

Insert Figure 1. about here

Social oddities in the pictures were operationally defined as violations of social norms e.g. someone in the picture throwing a baby in the air. Access to top down knowledge is required in that condition to inform a subjective value judgement as to whether a social violation is present in the picture. No value judgement as to whether a violation infringes some implicit or explicit social rule is needed for the perceptual oddities in this study, which are operationally defined as physical violations e.g. someone in the picture has a limb positioned back to front. See supplementary information for ~~Materials and Methods~~. Figure 1 shows an example of the stimuli.

Method

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

7

Participants

ASD participants (n = 24 age 18–52 years, 20 males), with a formal diagnosis of either Autistic Disorder (n = 3) or Asperger’s Syndrome were recruited from local charitable organisations. TD participants (n = 24, age 20–52 years, 20 males) were volunteers from the local community. All participants were paid and had normal or corrected to normal vision. Each ASD participant completed module 4 of the Autism Diagnostic Observation Scale 2 (Lord et al., 2012) and five (3 males and 2 females with a diagnosis of Asperger’s Syndrome) failed to meet the Autism Spectrum criteria. To ensure maximum homogeneity in our study those five participants were excluded from the analyses, resulting in a final sample of 19 ASD participants. Both groups completed the Autism-spectrum Quotient (AQ; Baron-Cohen et al., 2001) and the ASD participants reported a higher number of autistic traits $t(41) = 8.04, p < .001$ (TD $M = 15.42, SD = 7.55$; ASD $M = 35.32, SD = 8.45$). The two groups did not differ in age $t(41) = 0.31, p = .759$ (TD $M = 30.29, SD = 10.20$; ASD $M = 31.26, SD = 10.29$), performance IQ $t(41) = 0.21, p = .832$ (TD $M = 111.79, SD = 12.01$; ASD $M = 110.95, SD = 13.84$), verbal IQ $t(41) = 1.21, p = .234$ (TD $M = 116.96, SD = 10.65$, ASD $M = 112.26, SD = 14.84$), or full scale IQ $t(46) = 0.87, p = .392$ (TD $M = 116.04, SD = 9.98$, ASD $M = 112.95, SD = 13.48$), as measured by two performance (Block Design and Matrix Reasoning) and verbal components (Vocabulary and Similarities) of the Weschler Abbreviated Scale of Intelligence (The Psychological Corporation, 1999).

Eye movement recording

Eye movements were sampled at 1000Hz using an Eyelink tracker with spatial accuracy of less than 1/4th degree (S.R. Research Ltd, Canada). The start and end of saccades were

Formatted: Font: (Default) Times New Roman, 12 pt

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

8

[detected automatically using a default velocity criterion of 30degrees per second using the EyeLink software. See Stampe \(1993\) for an in depth technical description of how the software detects saccades and fixations.](#) A chin rest and forehead support, were used to stabilize head position. Viewing was binocular but data were analysed for the right eye only. Individual participants were calibrated using a nine point matrix that covered the dimensions of the screen (1024 x 768 pixels), with each point fixated sequentially, followed by a validation procedure to ensure that fixation was within 0.5 degrees of each calibration point. Calibration was checked using a validation procedure prior to each trial, whereby participants fixated five dots, presented at five different locations that covered the dimensions of the screen and recalibration was performed if the eyes were off centre for any of these dots.

Materials

In total, 96 full-colour photographs were used as stimuli, all of which contained people. In both the perceptual and social oddity conditions there were 48 pictures (24 weird and 24 normal). Perceptual oddities in the pictures violated a physical norm e.g. someone in the picture has a limb positioned back to front, or a vehicle was in an unexpected position e.g midway through the upper wall of a building. Social oddities in the pictures violated a social norm e.g. someone in the picture throwing a baby in the air, or someone with a cabbage on a lead. A small number of the pictures were from a previous study (Benson et al., 2012) and the remainder were collected from the internet and included either a perceptual or social oddity. Adobe Photoshop was used to prepare a normal version of each scene in which the oddity was replaced with either the background elements or with another suitable object. The scenes typically had a focal object or a central group of objects. Figure 1 shows an example of a normal and a socially odd scene,

and below those examples, a normal and a perceptually odd scene. The target regions for these scenes are outlined in black – but these black outlines were not visible during the presentation of the scenes in the experiment.

A norming study was conducted to verify the validity of the images used to examine the processing of social and perceptual violations. Twenty undergraduates from Queen's university participated (M age: 19.8 years, SD : 1.71, 10 Females) with five participants in each of the four conditions (counterbalanced conditions: weird/normal and right/left button response). Stimuli were displayed on a 21 inch CRT monitor with a refresh rate of 100Hz and the experiment was run using E-Prime software. Participants were presented with a fixation cross for 500ms followed by the presentation of the image for 20 seconds or until a keyboard "Y" (yes) or "N" (N) was made. Participants had to indicate as quickly as possible whether the presented image was "odd", "weird" or "unusual" in some way. ~~Results-~~ The proportion of identified images indicated that on average, the social normal ($M = 0.86$, $SD = 0.15$) and social weird ($M = 0.84$, $SD = 0.17$), perceptual normal ($M = 0.82$, $SD = 0.17$) and perceptual weird ($M = 0.77$, $SD = 0.19$) images were identified correctly.

Design

For the behavioural measures of Accuracy and RT a mixed design with Picture Type (weird versus normal) and Oddity Type (social versus perceptual) as within participants variables, and Group (ASD versus TD) as a between participants variable was employed for the experiment. On condition that accuracy was equivalent for both normal and weird pictures, for the eye movement measures the main focus of interest was to compare social versus perceptual oddities in both groups. Therefore a mixed design with Oddity Type (social versus perceptual) as

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

10

a within participants variable, and Group (ASD versus TD) as a between participants variable was employed in the eye movement analysis.

Procedure

All participants were shown a hard copy of an example picture from each condition prior to completing the task, to ensure comprehension of the task and the ability to respond appropriately. The specific instruction was 'for each of the following pictures, please indicate whether you think the picture is odd, unusual or strange in some way'.

For the task, 48 pictures were presented randomly to either the left or the right of the centre of a 19inch colour monitor, at a distance of 65cm for 20s or until a button press response was made. Prior to the onset of a picture, participants fixated a cross that was presented on the opposite side of the screen to where the picture was set to appear, to ensure all participants started at the same point in each trial. Once participants fixated this cross, the experimenter initiated each trial. Picture type (normal vs. weird) and, the position of the picture (left or right on the display) were counterbalanced across items. The buttons used to respond either yes or no (left or right trigger button) was counterbalanced across participants.

Data preparation and analyses

For each 'odd' picture we created an interest region around the target item and the corresponding region of the same size in the counterpart 'normal' picture. Regions of interest were created using the approximate average regions from those identified and manually selected by participants who took part in the norming experiment.

Formatted: Line spacing: Double

Formatted: Font: (Default) Times New Roman, 12 pt

LOOKING, SEEING AND BELIEVING IN AUTISM

11

Eyelink software was used to detect saccades and fixations using a default velocity criterion and in-house software was used to calculate the distribution of fixations falling into the target region. Five items were removed from analysis because for these scenes, participant accuracy scores were significantly below chance¹. In addition, each participant was asked whether they had seen any of the scenes previously, and if so these scenes were also removed. This resulted in a loss of 2.3% of trials. Fixations below 90ms and above 1200ms were classed as outliers and removed; resulting in a loss of 5.13% of the fixation data.

Results

For the principal analyses we compared manual reaction times and accuracy (see Table 1), as well as a number of eye movement measures, such as, the time taken to begin fixating in the target region (ELT), the duration of the first fixation when the eyes landed in the target region (FFD) and the mean time (MFD) spent fixating in the target area (see Table 2). Global eye movement measures for the target region, such as mean fixation duration and mean number of fixations over all trials were also analysed. Finally, baseline measures (not just target related), were calculated across the duration of the trials, to check for overall sampling differences between the ASD and TD participants (see Table 2). See supplementary information for methods and materials.

Behavioural Data

Accuracy. There was no effect of oddity type (perceptual vs. social) $F(1, 41) = 1.44, p = .237, \eta_p^2 = .03$, picture type (normal or weird) $F(1, 41) = 0.94, p = .339, \eta_p^2 = .02$, or group, $F(1, 41) = .06, p = .808, \eta_p^2 < .01$ on accuracy rates, and no interactions ($F_s < 2.39, p_s > .130$). At

Formatted: Font: Italic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

least in terms of accuracy, performance was the same for both groups. For means and standard deviations for behavioural data see Table 1.

Response Time (RT). A main effect of oddity type $F(1, 41) = 8.16, p = .007, \eta_p^2 = .17$, showed participants took longer to respond when presented with a perceptual oddity in comparison to a social oddity. Longer response times for perceptual oddities could reflect that these 'impossible', physical violations were so unusual that they resulted in more time spent looking at them prior to responding. See below for confirmation of that in the eye movement data. There was also a main effect of picture type $F(1, 41) = 24.03, p < .001, \eta_p^2 = .37$, with participants taking longer to respond when the picture was normal in comparison to when it was weird, in line with longer reaction times for target absent trials in standard visual search tasks (Wolfe, 1998) A significant oddity type x picture type interaction $F(1, 41) = 4.22, p = .046, \eta_p^2 = .09$, demonstrated that the difference between RTs for perceptual and social pictures was reliable for odd (TD; $t(23) 2.62, p = .015$, ASD; $t(18) = 2.98, p = .008$) but not for normal pictures (TD; $t(23) 0.94, p = .355$, ASD; $t(18) = 0.08, p = .935$). A main effect of group $F(1, 41) = 4.07, p = .050, \eta_p^2 = .09$, showed the ASD participants had longer reaction times, in line with a previous study (Benson et al., 2012, 25). There were no significant group interactions ($F_s < 1$) demonstrating both groups to be slower for normal versus odd pictures, and for perceptual vs social oddities. See Table 1 for the means. The eye movement data below reveals subtle processing differences that contribute to this overall slowed manual responding in the ASD group.

Eye Movement Data

Formatted: Font: Not Italic

Having shown that all participants could distinguish between 'odd' and 'normal' pictures, normal picture trials were excluded from the eye movement analyses resulting in a 2 (group: ASD versus TD) X 2 (oddy type: social versus perceptual) design to investigate on-line processing differences between the two groups for the two types of oddities.

Baseline eye movement measures, calculated across the full length of trials indicated no group difference in mean fixation duration $F(1, 41) = 1.14, p = .293, \eta_p^2 = .03$, but the ASD group made more fixations overall $F(1, 41) = 9.44, p = .004, \eta_p^2 = .19$, corresponding to the increased RT's reported above. Table 2 presents a summary of all the eye movement measures.

Elapsed time to the target region (ELT). A marginal effect of oddity type $F(1, 41) = 3.99, p = .053, \eta_p^2 = .09$, on the time taken from the onset of each trial display to the start of the first fixation on the target region, revealed both groups took less time to fixate the target region when the oddity was perceptual in comparison to social in nature. ~~The impossible nature of the perceptual violation may have increased the salience of the target region but~~ there was no effect of group $F(1, 41) = 0.59, p = .446, \eta_p^2 = .01$, and no interaction $F(1, 41) = 0.72, p = .401, \eta_p^2 = .02$.

First Fixation Duration (FFD) on target region. There was no effect of oddity type $F(1, 41) = 1.84, p = .182, \eta_p^2 = .04$ or group $F(1, 41) = 0.15, p = .699, \eta_p^2 < .01$ but a significant interaction $F(1, 41) = 6.72, p = .013, \eta_p^2 = .14$ (See Figure 2) revealed that FFDs were equivalent for perceptual and social oddities for TD participants $t(23) = 0.86, p = .399, d = .16$, but for ASD participants FFDs for social oddities were significantly shorter, in comparison to perceptual oddities, $t(18) = 2.99, p = .008, d = .65$, see Figure 2. Given that the FFD is a measure of early

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

processing (Aversedge & Findlay, 2000; Rayner, 1998; Rayner, 2009) this finding indicates that the ASD group did not detect the social oddities as quickly as the perceptual oddities on initial fixation, whereas TD participants detected both types of oddity equally fast.

Insert Figure 2. about here

First gaze duration (FGD) in the target region. The FGD is the elapsed time from when the target region is first fixated until the eye moves on to a new location. FGD includes multiple fixations in the target region reflects an overt strategy to re-examine an area. A main effect of oddity type, $F(1, 41) = 12.10, p = .001, \eta_p^2 = .23$ demonstrated that both groups spent more time inspecting perceptual oddities, again highlighting the salience of the perceptual oddities in comparison to social oddities in this study. However, there was no group effect $F(1, 41) = 2.43, p = .127, \eta_p^2 = .06$, or interaction $F(1, 41) = .08, p = .785, \eta_p^2 < .01$.

Mean Fixation Count in the target region. If the ASD are slower to respond manually, but do not take longer to begin inspecting the target region, then extra time to respond might reflect more fixations in the target region. A main effect of oddity type $F(1, 41) = 64.27, p < .001, \eta_p^2 = .61$, demonstrated more fixations in the target region when the oddity was perceptual, rather than social. A main effect of group $F(1, 41) = 6.99, p = .012, \eta_p^2 = .15$ showed that the ASD participants made more fixations than the TD group in the target regions for both types of oddities overall. However no interaction was observed $F(1, 41) = 0.77, p = .386, \eta_p^2 = .02$ indicating that the ASD group made more fixations in the target regions for both types of oddities.

Formatted: Highlight
Formatted: Font: (Default) Times New Roman, 12 pt
Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic
Formatted: Font: Not Italic
Formatted: Font: (Default) Times New Roman, 12 pt
Formatted: Highlight

Mean Fixation Duration in the target region. Longer mean fixation durations indicate that more processing has been done during the task. A main effect of oddity type $F(1, 41) = 8.22, p = .007, \eta_p^2 = .17$ showed significantly longer fixations for social, compared to perceptual oddity target regions. This suggests that processing social oddities was more effortful, and since there was no main effect of group $F(1, 41) = 0.34, p = .562, \eta_p^2 < .01$, and no interaction $F(1, 41) = 1.12, p = .297, \eta_p^2 = .03$, both groups on average fixated the social oddity target region for longer than the perceptual target region. Thus, while the perceptual oddities attracted more fixations overall, individual fixations were longer for the social target region.

This interesting juxtaposition, although unrelated to the main aim of this study, suggests that difficulty associated with processing each oddity type manifests in different eye movement patterns: one reflecting an overt strategy (fixation count) to re-inspect the target area, and one reflecting more in depth on-line cognitive processing (fixation duration) on the target areas.

Mean Number of Entries to the target region. Since there were no group differences in FGD, and no increased MFD, both reported above, the delayed RT in ASD might reflect repeated visits to inspect the target region?

A main effect of oddity type was found $F(1, 41) = 9.83, p = .003, \eta_p^2 = .19$, with the target region entered more often for perceptual oddities in comparison to social oddities. In addition, a main effect of group was found $F(1, 41) = 5.72, p = .021, \eta_p^2 = .12$, with ASD participants entering the target regions significantly more than the TD group, before making a response. There was no interaction $F(1, 41) = 0.13, p = .717, \eta_p^2 < .01$.

Mean Total Time in the target region. A higher mean total time means that more attention to, and processing of, information in that region was carried out overall. This measure should complement the measures reported above – namely, that if there are more fixations overall in ASD then the Total Time spent fixating in the target region should be longer in ASD.

A main effect of oddity type $F(1, 41) = 34.89, p < .001, \eta_p^2 = .46$, showed perceptual oddities received more overall processing time, in line with the previous measures for the perceptual oddities. In addition, a main effect of group $F(1, 41) = 6.41, p = .015, \eta_p^2 = .14$ revealed that the ASD participants spent a longer period of time inspecting the target regions for both perceptual and social oddities. There was no interaction $F(1, 41) = 0.51, p = .481, \eta_p^2 = .01$.

As above, this finding reflects a greater number of entries to the target region overall by the ASD group prior to making their decision as to whether the picture under inspection was weird, odd or unusual, [irrespective of the nature of the violation \(perceptual or social\)](#).

Discussion

A principal aim of the study was to investigate whether increased times in ASD to find and initially detect an unusual target in a scene containing people would be modulated by whether the violation could be categorised as social or perceptual.

Accuracy rates were equivalent and high for both groups, but the ASD group took longer to respond manually. It was hypothesised that the eye movement data would reveal the cause of this delayed response in ASD, since eye movements reflect how information is processed on-line (Liversedge & Findlay, 2000; Rayner, 1998; 2009). The data showed that our ASD group did not take longer to find the target regions, but that they scanned the scenes more prior to responding manually, as shown by an increased number of fixations throughout each scene presentation.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Furthermore, these increased fixations were not random, but were made to the target regions of both social and perceptually odd pictures. Thus, increased fixations reflect a propensity to re-inspect the target regions prior to making a response in ASD, regardless as to the type of oddity

in the scenes. ~~It should be noted here that the oddity detection task in the current study provides the ASD participants with structure that is not typically available in real-life. As such, the task is not implicit and the knowledge that oddities might be present could also be a factor that drives the repeated scanning in ASD.~~

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

The eye movement patterns also showed that our ASD group failed to immediately detect social oddities, thus looking at but not seeing them. An obvious question from this finding is whether the failure to quickly spot subtle social oddities in ASD reflects an inability to react or respond appropriately to social cues in everyday communication? This one impairment in itself could potentially account for the significant problems experienced in dynamic social interactions in this high functioning population. For example, a failure to immediately spot social cues would result in a failure to react or respond to such cues thus resulting in an inability to both keep track of what is going on in social interactions, and a failure to engage appropriately or effectively with other people.

~~In relation to current theories that underpin ASD our findings do not fit with WCC theory (Frith & Happé, 1994; Happé & Frith, 2006), since clearly context can be taken into account in ASD during task completion. Moreover, the slowness to respond in ASD is not reflected in longer times to find the target, and the increased fixations in ASD were seen for both perceptual and social oddities in the scenes. It may be that repeated visits to the target area indicate integration problems in ASD, but even if that were the case, such problems are not restricted to social processing per se. Neither do they support the executive function theory (Russell, 1997), as~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

18

ASD participants showed no indication of increased mean fixation durations, thus no evidence of perseveration or any inability to switch focus throughout the task. Our findings could however be interpreted to support the EMB (Baron-Cohen, 2002), with the detection of social-oddities that required empathy being delayed in ASD, however note that higher re-inspection of perceptual oddities that could be detected on the basis of systematic rules, was also found for ASD participants.

If social oddities are more complex to detect in ASD, then the Complex Information Processing theory (Minshew & Goldstein, 1998) might explain the failure to detect the social oddity on first fixation. This lack of immediate detection could reflect the less obvious and more 'inferential' nature of the social oddities, which required more complex reasoning, at least in the ASD group, for detection. It is important to note that social rules are often implicit, sometimes ambiguous, and cannot always be understood by following a script or flow chart as in the case for logical rule based tasks. At a neurobiological level, it has been postulated that that impairments in cognitive processing in ASD reflect slow information transmission rates for high level cognitive tasks, resulting from underconnectivity between frontal-posterior cortical regions (Just et al., 2013; Just et al., 2012; Minshew & Williams, 2007). Since underconnectivity reduces the rate of inter-region information transfer, it could be that in social communication tasks that rely on the fast detection of implicit, often ambiguous cues, a failure to be able to transmit information across different cortical areas quickly, might mean that these cues are missed, rendering inappropriate responses in ASD.

The findings of repeated scanning of the target areas, coupled with an inability to immediately recognise social violations upon initial fixation, reflect how eye movements can reveal subtle processing differences in ASD. In everyday social communication in the real world


Formatted: Not Highlight
Formatted: Font: (Default) Times New Roman, 12 pt, Not Highlight
Formatted: Not Highlight

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

19

there is no opportunity to 'go back' to check if one has missed an important cue. Social processing in such circumstances must be carried out on-line, and our eye movement data clearly show that ASD are unable to do this quickly. The strategy of re-checking their interpretation, and thus not believing what they have seen, before responding, may be related to cortical underconnectivity in ASD. In everyday terms, these findings have obvious consequences for fast dynamic social interactions, and offer an account as to why high functioning adults with ASD experience difficulty in the social domain.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

20

Acknowledgments

This work was supported by a grant awarded to the second author from the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the Ontario Ministry of Research and Innovation, and Queen's University Senate Advisory Research Committee, and an Experimental Psychology Society study visit grant awarded to the third author.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

21

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Au-Yeung, S. K., Benson, V., Castelhana, M. S., Rayner, K. (2011). Eye movement sequences during simple versus complex information processing of scenes in autism spectrum disorder. *Autism Research and treatment*, 2011.
- Au-Yeung, S. K., Kaakinen, J. K., Benson, V. (2013). Cognitive perspective taking during scene perception in autism spectrum disorder: evidence from eye movements. *Autism Research*, 7, 84-93.
- Bar-Haim, Y., Shulman, C., Lamy, D., & Reuveni, A. (2006). Attention to eyes and mouth in high-functioning children with autism. *Journal of Autism and Developmental Disorders*, 36, 131-137.
- Baron-Cohen, S. (2002). The extreme male brain theory of autism. *Trends in Cognitive Science*, 6, 248-254.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5-17.
- Benson, V., & Fletcher-Watson, S. (2012). Eye Movements in Autism. In: Liversedge, S. P., Gilchrist, I. D., Everling, S., editors. *The Oxford Handbook of Eye Movements*, Oxford UK: Oxford University Press, chap 39.
- Benson, V., Castelhana, M. S., Au-Yeung, S. K., & Rayner K. (2012). Eye movements reveal no immediate "WOW" ("which one's weird") effect in autism spectrum disorder. *Quarterly*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

22

- Journal of Experimental Psychology, 65, 1139-1150.
- Benson, V., Piper, J., & Fletcher-Watson, S. (2009). Atypical saccadic sampling in autistic spectrum disorder, *Neuropsychologia*, 47, 1178-1182, (2009).
- Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate neural responses to social stimuli in autism spectrum disorders. *Neuroimaging*, 15, 1614-1624.
- Brugha, T. S., McManus, S., Bankart, J., Scott, F., Purdon S., et al. (2011). Epidemiology of autism spectrum disorders in adults in the community in England. *Archives of General Psychiatry*, 68, 459-65.
- Clark, F., Scharaschkin, A., Xu, D. (2009). Supporting autism through adulthood, National Audit Office.
- Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., Alexander, A. L., & Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nature Neuroscience*, 8, 519-526.
- Fletcher-Watson, S., Leekam, S. R., Benson, V., Frank, M. C., & Findlay, J. M. (2009). Eye-movements reveal attention to social information in autism spectrum disorder. *Neuropsychologia*, 47, 248-257.
- Fletcher-Watson, S., Leekam, S. R., Benson, V., Frank, M. C., & Findlay, J. M. (2009). Eye-movements reveal attention to social information in autism spectrum disorder. *Neuropsychologia*, 47, 248-257.
- Frith, U., & Happé, F. (1994). Autism; Beyond the theory of mind. *Cognition*, 50, 115-132.
- Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 36, 5-25.

LOOKING, SEEING AND BELIEVING IN AUTISM

23

- Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder. *Archives of General Psychiatry*, 65, 946-954.
- Just, M. A., Keller, T. A., & Kana, R. K. (2013). A theory of autism based on frontal-posterior underconnectivity. In M. A. Just & K. A. Pelphrey (Eds.), *Development and brain systems in autism* (pp. 35-63). New York: Psychology Press.
- Just, M. A., Keller, T. A., Malave, V. L., Kana, R. K., & Varma, S. (2012). Autism as a neural systems disorder: A theory of frontal-posterior underconnectivity. *Neuroscience and Biobehavioral Reviews*, 36, 1292-1313.
- Klin, A., & Jones, W. (2006). Attributing social and physical meaning to ambiguous visual displays in individuals with higher functioning autism spectrum disorders. *Brain & Cognition*, 61, 40-53.
- Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen D. J. (2002a). Defining and quantifying the social phenotype in autism. *American Journal of Psychiatry*, 159, 895-908.
- Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen, D. J. (2002b). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59, 809-816.
- Kuhn, G., Benson, V., Fletcher-Watson, S., Kovshoff, H., McCormick, C., & Kirkby, J. (2010). Eye movements affirm: Automatic overt gaze and arrow cueing for typical adults and adults with autism spectrum disorder. *Experimental Brain Research*, 201, 155-165.
- Liversedge, S. P., & Findlay, J. M. (2000). Saccadic eye movements and cognition. *Trends in Cognitive Sciences*, 4, 6-14.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

24

Lord C., Rutter M., DiLavore P. C., Risi S., Gotham K., Bishop S. (2012). Autism diagnostic observation schedule, second edition. Torrance, CA: Western Psychological Services.

Louwerse, A., Tulen, J. H. M., van der Geest, J. N., van der Ende, J., Verhulst, F. C., & Greaves-Lord, K. (2014). Autonomic Responses to Social and Nonsocial Pictures in Adolescents With Autism Spectrum Disorder. *Autism Research*, 7, 17-27. doi:10.1002/aur.1327

Minshew, N. J., & Goldstein, G. (1998). Autism as a disorder of complex information processing. *Mental Retardation and Developmental Disabilities Research Reviews*, 4, 129-136.

Minshew, N. J., & Williams, D. L., (2007). The new neurobiology of autism: cortex, connectivity, and neuronal organization. *Archives of Neurology*, 64, 945-950.

Minshew, N. J., Goldstein, G., & Siegel, D. J. (1997). Neuropsychologic functioning in autism: Profile of complex information processing disorder. *Journal of the International Neuropsychological Society*, 3, 303-316.

Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual scanning of faces in autism. *Journal of Autism and Developmental Disorders*, 32, 249-261.

Rayner, K. (2009). Eye movements and attention in reading, scene perception, and visual search. *Quarterly Journal of Experimental Psychology*, 62, 1457-1506.

Rayner, K., Castelano, M. S., & Yang J. (2009). Eye Movements when Looking at Unusual/Weird Scenes: Are there Cultural Differences? *Journal of Experimental Psychology: Language, Memory & Cognition*, 35, 254-259.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

25

Rayner, K., Warren, T., Juhasz, B. J., & Liversedge, S. P. (2004). The effect of plausibility on eye movements in reading. *Journal of Experimental Psychology-Learning Memory and Cognition*, 30(6), 1290-1301.

Riby, D. M., & Hancock, P. J. B. (2009). Do faces capture the attention of individuals with Williams syndrome or Autism? Evidence from tracking eye movements. *Journal of Autism and Developmental Disorders*, 39, 421-431. doi:10.1007/s10803-008-0641-z

Russell, J. (1997). *Autism as an executive disorder*. Oxford, UK: Oxford University Press.

Speer, L. L., Cook, A. E., McMahon, W. M., & Clark, E. (2007). Face processing in children with autism: effects of stimulus contents and type. *Autism*, 11, 263-277.

Stampa, D. M. (1993). Heuristic filtering and reliable calibration methods for video-based pupil-tracking systems. *Behavior Research Methods, Instruments, & Computers*, 25, 137-142.

The Psychological Corporation, Wechsler Abbreviated Scale of Intelligence (WASI) Manual, (The Psychological Corporation, San Antonio, TX, 1999).

Wolfe, J. M. (1998) What can one million trials tell us about visual search? *Psychological Science*, 9, 33-39.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic
Formatted: Indent: Left: 0", Hanging: 0.5", Line spacing: Double
Formatted: Font: (Default) Times New Roman, 12 pt
Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

26

Footnotes

1. Accuracy errors for the excluded scenes were either a failure to detect a subtle oddity, or were a result of participants thinking the control version of the scenes were odd. Removing these items from analysis did not change the pattern of effects, but reduced variation in reaction time and eye movement data.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LOOKING, SEEING AND BELIEVING IN AUTISM

27

Tables

Table 1.
Accuracy and reaction time means (standard deviations).

	Perceptual				Social			
	Normal		Weird		Normal		Weird	
	TD	ASD	TD	ASD	TD	ASD	TD	ASD
Accuracy	0.81 (0.20)	0.86 (0.10)	0.84 (0.20)	0.77 (0.17)	0.80 (0.17)	0.83 (0.12)	0.80 (0.21)	0.76 (0.24)
Reaction Time (ms)	5289 (2372)	6024 (2560)	3697 (1703)	5184 (2228)	5117 (2443)	6053 (2884)	3056 (1270)	4633 (2131)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

RUNNING HEAD: LOOKING, SEEING AND BELIEVING IN AUTISM.

28

Table 2.
Eye movement measure means in ms (standard deviations).

	Perceptual Oddity		Social Oddity	
	TD	ASD	TD	ASD
Mean Fixation Duration (overall)	250 (31)	257 (29)	241 (26)	252 (33)
Mean Fixation Count (overall)	11 (4)	15 (6)	9 (4)	13 (7)
Elapsed time to target region	ELT 551 (235)	557 (144)	600 (235)	679 (284)
Fixation count before target region	FCT 2.32 (0.72)	2.49 (0.67)	2.62 (0.76)	2.68 (0.73)
First fixation duration in target region	FFD 224 (52)	237 (42)	232 (45)	210 (41)
First gaze duration in target region	FGD 808 (250)	924 (437)	630 (163)	715 (208)
Mean fixation duration in target region	MFD 262 (43)	274 (34)	281 (38)	282 (45)
Mean fixation count in target region	MFC 5.78 (2.28)	7.72 (2.84)	3.97 (1.81)	5.47 (2.14)
Mean number of entries into target region	MNE 1.93 (0.64)	2.32 (0.73)	1.62 (0.52)	2.07 (0.66)
Mean total time in target region	TT 1522 (729)	2065 (809)	1040 (527)	1450 (664)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RUNNING HEAD: LOOKING, SEEING AND BELIEVING IN AUTISM

29

Figures captions

Figure 1. Example of social normal (A), social weird (B), perceptual normal (C) and perceptual weird (D) stimuli. The black square represents the target region. Note the black square is not visible during the experiment.

Figure 2. Interaction for first fixation duration (ms) between group and oddity type. Error bars denote ± 1 SE.

1
2
3
4
5 Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle
6
7
8 cognitive processing differences impact in the social domain.
9

10 Valerie Benson ^{1*}, Monica S. Castelhana ², Philippa L. Howard ¹, Nida Latif ² & Keith Rayner ³
11
12

13
14
15
16 ¹ AU Psychology, Shackleton Building, Highfield Campus, University of Southampton, UK.
17

18 ² Queen's University, Kingston, ON, Canada
19

20 ³ University of California, San Diego, USA
21
22

23
24 *Correspondence to: vb1@soton.ac.uk
25

26 Telephone: +44 (0) 2380594596
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Lay Abstract

Adults with High Functioning Autism (ASD) viewed scenes with people in them, whilst having their eye movements recorded. The task was to indicate, using a button press, whether the pictures were normal, or in some way weird or odd. Oddities in the pictures were categorized as violations of either perceptual or social norms. Compared to a Typically Developed (TD) control group, the ASD participants were equally able to categorise the scenes as odd or normal, but they took longer to respond. The eye movement patterns showed that the ASD group made more fixations and revisits to the target areas in the odd scenes compared to the TD group. Additionally, when the ASD group first fixated the target areas in the scenes, they failed to initially detect the social oddities. These two findings have clear implications for processing difficulties in ASD for the social domain, where it is important to detect social cues on-line, and where there is little opportunity to go back and recheck possible cues in fast dynamic interactions.

Scientific Abstract

1
2
3
4
5
6
7
8 Autistic Spectrum Disorder (ASD) and Typically Developed (TD) adult participants had to
9 decide whether scenes were 'odd', 'unusual' or 'weird'. All scenes contained people, with
10 'oddness' operationally defined as either perceptual or social. There were no group differences in
11 accuracy, but the ASD group were slower overall to respond manually. Eye movements revealed
12 that the ASD group made more fixations overall and more re-fixations into the target regions of
13 the pictures, for both conditions. Importantly, as indexed by the first fixation duration the ASD
14 group failed to 'pick up' immediately on what was 'odd' for the social violations. The propensity
15 to 'go back' and re-fixate targets, coupled with a failure to initially detect social violations, has
16 obvious significance for ASD in fast dynamic social communication.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 Key phrases: Autism Spectrum Disorder, Eye movements, On-line cognitive processing, Social
33 and Perceptual Oddities.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Looking, Seeing and Believing in Autism: Eye Movements reveal how subtle cognitive
4
5 processing differences impact in the social domain.
6
7

8
9 Autism spectrum disorders (ASD) are lifelong neuro-developmental conditions
10 characterized by repetitive behaviours and impairments in interaction and communication
11 (American Psychiatric Association, 2013). Incidence rates are rising across the globe with the
12 prevalence rate in the UK approximately 10 per 1000 (1%) individuals (Brugha et al., 2011) and
13 the UK National Audit office recently called for more research into adults with ASD (Clark et
14 al., 2009).
15
16
17
18
19
20
21
22

23 High functioning adults with ASD are similar to typically developed (TD) adults in terms
24 of performance on pencil and paper and IQ tests, however in the social domain evidence suggests
25 an inability to process social information in a typical fashion. Eye movement studies report
26 unusual gaze behaviour for faces appearing on their own (Dalton et al., 2005), in social scenes
27 (Riby & Hancock, 2009) or in dynamic movie clips (Klin & Jones, 2006; Speer et al., 2007).
28 Research consistently reports a lack of spontaneous gaze fixation towards the eyes (Jones et al.,
29 2008; Klin et al., 2002a; Pelphrey et al., 2002). However, where one study reports impairments,
30 say for face processing (Klin et al., 2002b), another study does not (Fletcher-Watson et al.,
31 2009). Moreover, lack of attentional modulation for social stimuli (Bird et al., 2006) may be
32 overridden, as when cued, attention may be allocated to faces in a more 'typical' manner (Bar-
33 Haim et al., 2006).
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 A recent scene perception study (Louwerse et al., 2014) reported that abnormal
49 autonomic responses in ASD are unlikely to underpin social processing impairments. It is
50 interesting then to ask whether impairments in the cognitive domain in ASD (Happé & Frith,
51 2006; Minshew et al., 1997; Russell, 1997) could account for impairments in the social domain?
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
Current cognitive theories of ASD include, among others, Weak Central Coherence (WCC) theory (Frith & Happé, 1994; Happé & Frith, 2006) which proposes that a reduced sensitivity to global context in favour of focussing on details, results in integration difficulties in ASD. Executive function theory (Russell, 1997), proposes that ASD are impaired at switching their focus of attention, and may engage in perseverative behaviour. Alternatively, the Complex Information Processing theory (Minshew & Goldstein, 1998) proposes that individuals with ASD have impairments in processing complex information with intact processing for tasks that require simple (rule based) processing. Finally, the Extreme Male Brain (EMB) theory (Baron-Cohen, 2002) posits that individuals with ASD have above average levels of systemizing, that contribute to the non-social characteristics of the disorder, coupled with below average levels of empathizing, which results in atypical social functioning. These contemporary theories of ASD can, for different reasons, offer some account of ineffective social functioning in ASD. However, to date very few investigations have examined online processing differences in ASD.

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Our studies investigating cognitive processing in adults with ASD (Au-Yeung et al., 2011; Au-Yeung et al., 2013; Benson et al., 2012; Benson et al., 2009; Benson & Fletcher-Watson, 2011) have found subtle on-line processing differences, as revealed by eye movements, for a range of tasks that are more in line with Minshew and Goldstein's (1998) theory, than with any other. For example, there are more similarities than differences in eye movements between TD and ASD groups for simple rule based tasks in the attentional (Kuhn et al., 2010) and the social domains (Benson & Fletcher-Watson, 2011). Whereas for more complex abstract reasoning tasks where there is ambiguity with relation to target items (Au-Yeung et al., 2013), or where value judgements rely on top down knowledge (Benson et al., 2012), there are early processing differences indicating that initial orienting to, and immediate detection of weird target

1
2
3 items in scenes is absent in ASD. Moreover, the time to respond and to locate the weird target
4
5 items was modulated by whether or not the pictures had people in them (Benson et al., 2012).
6
7
8 These differences have potential for impact on everyday activities, since the time course for
9
10 detection of socially relevant information is crucial for successful everyday interactions.
11

12
13 The aim of the present study was to examine whether increased times to find and initially
14
15 detect an unusual target in a scene would be modulated by whether the violation that rendered
16
17 the target unusual or odd could be categorised as social or perceptual. We employed a similar
18
19 paradigm to that used in a previously reported study (Rayner et al., 2009) where single scenes,
20
21 all with people in them, were presented in isolation and the task was to indicate whether each
22
23 picture was normal or in some way weird unusual or odd.
24
25
26
27
28

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Insert Figure 1. about here

33
34 Social oddities in the pictures were operationally defined as violations of social norms
35
36 e.g. someone in the picture throwing a baby in the air. Access to top down knowledge is required
37
38 in that condition to inform a subjective value judgement as to whether a social violation is
39
40 present in the picture. No value judgement as to whether a violation infringes some implicit or
41
42 explicit social rule is needed for the perceptual oddities in this study, which are operationally
43
44 defined as physical violations e.g. someone in the picture has a limb positioned back to front. See
45
46 supplementary information for Materials. Figure 1 shows an example of the stimuli.
47
48
49

50 51 52 53 54 55 56 57 58 59 60 **Method**

Participants

55 ASD participants (n = 24 age 18 – 52 years, 20 males), with a formal diagnosis of either
56
57 Autistic Disorder (n = 3) or Asperger's Syndrome were recruited from local charitable
58
59
60

1
2
3 organisations. TD participants ($n = 24$, age 20– 52 years, 20 males) were volunteers from the
4
5 local community. All participants were paid and had normal or corrected to normal vision. Each
6
7 ASD participant completed module 4 of the Autism Diagnostic Observation Scale 2 (Lord et al.,
8
9 2012) and five (3 males and 2 females with a diagnosis of Asperger’s Syndrome) failed to meet
10
11 the Autism Spectrum criteria. To ensure maximum homogeneity in our study those five
12
13 participants were excluded from the analyses, resulting in a final sample of 19 ASD participants.
14
15 Both groups completed the Autism-spectrum Quotient (AQ; Baron-Cohen et al., 2001) and the
16
17 ASD participants reported a higher number of autistic traits $t(41) = 8.04, p < .001$ (TD $M =$
18
19 $15.42, SD = 7.55$; ASD $M = 35.32, SD = 8.45$). The two groups did not differ in age $t(41) =$
20
21 $0.31, p = .759$ (TD $M = 30.29, SD = 10.20$; ASD $M = 31.26, SD = 10.29$), performance IQ $t(41) =$
22
23 $0.21, p = .832$ (TD $M = 111.79, SD = 12.01$; ASD $M = 110.95, SD = 13.84$), verbal IQ $t(41) =$
24
25 $1.21, p = .234$ (TD $M = 116.96, SD = 10.65$, ASD $M = 112.26, SD = 14.84$), or full scale IQ t
26
27 $(46) = 0.87, p = .392$ (TD $M = 116.04, SD = 9.98$, ASD $M = 112.95, SD = 13.48$), as measured
28
29 by two performance (Block Design and Matrix Reasoning) and verbal components (Vocabulary
30
31 and Similarities) of the Weschler Abbrevated Scale of Intelligence (The Psychological
32
33 Corporation, 1999).

44 Eye movement recording

45
46 Eye movements were sampled at 1000Hz using an Eyelink tracker with spatial accuracy
47
48 of less than $1/4^{\text{th}}$ degree (S.R. Research Ltd, Canada). The start and end of saccades were
49
50 detected automatically using a default velocity criterion of 30degrees per second using the
51
52 Eyelink software. See Stampe (1993) for an in depth technical description of how the software
53
54 detects saccades and fixations. A chin rest and forehead support, were used to stabilize head
55
56
57
58
59
60

1
2
3 position. Viewing was binocular but data were analysed for the right eye only. Individual
4
5 participants were calibrated using a nine point matrix that covered the dimensions of the screen
6
7 (1024 x 768 pixels), with each point fixated sequentially, followed by a validation procedure to
8
9 ensure that fixation was within 0.5 degrees of each calibration point. Calibration was checked
10
11 using a validation procedure prior to each trial, whereby participants fixated five dots, presented
12
13 at five different locations that covered the dimensions of the screen and recalibration was
14
15 performed if the eyes were off centre for any of these dots.
16
17
18
19

20 21 22 **Materials**

23
24 In total, 96 full-colour photographs were used as stimuli, all of which contained people.
25
26 In both the perceptual and social oddity conditions there were 48 pictures (24 weird and 24
27
28 normal). Perceptual oddities in the pictures violated a physical norm e.g. someone in the picture
29
30 has a limb positioned back to front, or a vehicle was in an unexpected position e.g. midway
31
32 through the upper wall of a building. Social oddities in the pictures violated a social norm e.g.
33
34 someone in the picture throwing a baby in the air, or someone with a cabbage on a lead. A small
35
36 number of the pictures were from a previous study (Benson et al., 2012) and the remainder were
37
38 collected from the internet and included either a perceptual or social oddity. Adobe Photoshop
39
40 was used to prepare a normal version of each scene in which the oddity was replaced with either
41
42 the background elements or with another suitable object. The scenes typically had a focal object
43
44 or a central group of objects. Figure 1 shows an example of a normal and a socially odd scene,
45
46 and below those examples, a normal and a perceptually odd scene. The target regions for these
47
48 scenes are outlined in black – but these black outlines were not visible during the presentation of
49
50 the scenes in the experiment.
51
52
53
54
55
56
57
58
59
60

1
2
3 A norming study was conducted to verify the validity of the images used to examine the
4 processing of social and perceptual violations. Twenty undergraduates from Queen's university
5 participated (M age: 19.8years, SD : 1.71, 10 Females) with five participants in each of the four
6 conditions (counterbalanced conditions: weird/normal and right/left button response). Stimuli
7 were displayed on a 21inch CRT monitor with a refresh rate of 100Hz and the experiment was
8 run using E-Prime software. Participants were presented with a fixation cross for 500ms
9 followed by the presentation of the image for 20 seconds or until a keyboard "Y" (yes) or "N"
10 (N) was made. Participants had to indicate as quickly as possible whether the presented image
11 was "odd", "weird" or "unusual" in some way. The proportion of identified images indicated that
12 on average, the social normal ($M = 0.86$, $SD = 0.15$) and social weird ($M = 0.84$, $SD = 0.17$),
13 perceptual normal ($M = 0.82$, $SD = 0.17$) and perceptual weird ($M = 0.77$, $SD = 0.19$) images
14 were identified correctly.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 Design

35
36 For the behavioural measures of Accuracy and RT a mixed design with Picture Type
37 (weird versus normal) and Oddity Type (social versus perceptual) as within participants
38 variables, and Group (ASD versus TD) as a between participants variable was employed for the
39 experiment. On condition that accuracy was equivalent for both normal and weird pictures, for
40 the eye movement measures the main focus of interest was to compare social versus perceptual
41 oddities in both groups. Therefore a mixed design with Oddity Type (social versus perceptual) as
42 a within participants variable, and Group (ASD versus TD) as a between participants variable
43 was employed in the eye movement analysis.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Procedure

All participants were shown a hard copy of an example picture from each condition prior to completing the task, to ensure comprehension of the task and the ability to respond appropriately. The specific instruction was ‘for each of the following pictures, please indicate whether you think the picture is odd, unusual or strange in some way’.

For the task, 48 pictures were presented randomly to either the left or the right of the centre of a 19inch colour monitor, at a distance of 65cm for 20s or until a button press response was made. Prior to the onset of a picture, participants fixated a cross that was presented on the opposite side of the screen to where the picture was set to appear, to ensure all participants started at the same point in each trial. Once participants fixated this cross, the experimenter initiated each trial. Picture type (normal vs. weird) and, the position of the picture (left or right on the display) were counterbalanced across items. The buttons used to respond either yes or no (left or right trigger button) was counterbalanced across participants.

Data preparation and analyses

For each ‘odd’ picture we created an interest region around the target item and the corresponding region of the same size in the counterpart ‘normal’ picture. Regions of interest were created using the approximate average regions from those identified and manually selected by participants who took part in the norming experiment. Eyelink software was used to detect saccades and fixations using a default velocity criterion and in-house software was used to calculate the distribution of fixations falling into the target region. Five items were removed from analysis because for these scenes, participant accuracy scores were significantly below chance¹. In addition, each participant was asked whether they had seen

1
2
3 any of the scenes previously, and if so these scenes were also removed. This resulted in a loss of
4
5 2.3% of trials. Fixations below 90ms and above 1200ms were classed as outliers and removed;
6
7
8 resulting in a loss of 5.13% of the fixation data.
9

10 11 12 13 Results

14
15 For the principal analyses we compared manual reaction times and accuracy (see Table
16
17 1), as well as a number of eye movement measures, such as, the time taken to begin fixating in
18
19 the target region (ELT), the duration of the first fixation when the eyes landed in the target
20
21 region (FFD) and the mean time (MFD) spent fixating in the target area (see Table. 2). Global
22
23 eye movement measures for the target region, such as mean fixation duration and mean number
24
25 of fixations over all trials were also analysed, Finally, baseline measures (not just target related),
26
27 were calculated across the duration of the trials, to check for overall sampling differences
28
29 between the ASD and TD participants (see Table 2). See supplementary information for methods
30
31 and materials.
32
33
34
35
36
37
38

39 Behavioural Data

40
41 **Accuracy.** There was no effect of oddity type (perceptual vs. social) $F(1, 41) = 1.44, p =$
42
43 $.237, \eta_p^2 = .03$, picture type (normal or weird) $F(1, 41) = 0.94, p = .339, \eta_p^2 = .02$, or group, $F(1,$
44
45 $41) = .06, p = .808, \eta_p^2 < .01$ on accuracy rates, and no interactions ($F_s < 2.39, p's > .130$). At
46
47 least in terms of accuracy, performance was the same for both groups. For means and standard
48
49 deviations for behavioural data see Table 1.
50
51

52
53 **Response Time (RT).** A main effect of oddity type $F(1, 41) = 8.16, p = .007, \eta_p^2 = .17$,
54
55 showed participants took longer to respond when presented with a perceptual oddity in
56
57
58
59
60

1
2
3 comparison to a social oddity. Longer response times for perceptual oddities could reflect that
4 these ‘impossible’, physical violations were so unusual that they resulted in more time spent
5
6 these ‘impossible’, physical violations were so unusual that they resulted in more time spent
7
8 looking at them prior to responding. See below for confirmation of that in the eye movement
9
10 data. There was also a main effect of picture type $F(1, 41) = 24.03, p < .001, \eta_p^2 = .37$, with
11
12 participants taking longer to respond when the picture was normal in comparison to when it was
13
14 weird, in line with longer reaction times for target absent trials in standard visual search tasks
15
16 (Wolfe, 1998). A significant oddity type x picture type interaction $F(1, 41) = 4.22, p = .046, \eta_p^2$
17
18 $= .09$, demonstrated that the difference between RTs for perceptual and social pictures was
19
20 reliable for odd (TD; $t(23) = 2.62, p = .015$, ASD; $t(18) = 2.98, p = .008$) but not for normal
21
22 pictures (TD; $t(23) = 0.94, p = .355$, ASD; $t(18) = 0.08, p = .935$). A main effect of group $F(1,$
23
24 $41) = 4.07, p = .050, \eta_p^2 = .09$, showed the ASD participants had longer reaction times, in line
25
26 with a previous study (Benson et al., 2012). There were no significant group interactions ($F_s <$
27
28 1) demonstrating both groups to be slower for normal versus odd pictures, and for perceptual vs
29
30 social oddities. See Table 1 for the means. The eye movement data below reveals subtle
31
32 processing differences that contribute to this overall slowed manual responding in the ASD
33
34 group.

Eye Movement Data

35
36
37
38
39
40
41
42
43
44
45
46 Having shown that all participants could distinguish between ‘odd’ and ‘normal’ pictures,
47
48 normal picture trials were excluded from the eye movement analyses resulting in a 2 (group:
49
50 ASD versus TD) X 2 (oddity type: social versus perceptual) design to investigate on-line
51
52 processing differences between the two groups for the two types of oddities.
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Baseline eye movement measures, calculated across the full length of trials indicated no group difference in mean fixation duration $F(1, 41) = 1.14, p = .293, \eta_p^2 = .03$, but the ASD group made more fixations overall $F(1, 41) = 9.44, p = .004, \eta_p^2 = .19$, corresponding to the increased RT's reported above. Table 2 presents a summary of all the eye movement measures.

Elapsed time to the target region (ELT). A marginal effect of oddity type $F(1, 41) = 3.99, p = .053, \eta_p^2 = .09$, on the time taken from the onset of each trial display to the start of the first fixation on the target region, revealed both groups took less time to fixate the target region when the oddity was perceptual in comparison to social in nature. There was no effect of group $F(1, 41) = 0.59, p = .446, \eta_p^2 = .01$, and no interaction $F(1, 41) = 0.72, p = .401, \eta_p^2 = .02$.

First Fixation Duration (FFD) on target region. There was no effect of oddity type $F(1, 41) = 1.84, p = .182, \eta_p^2 = .04$ or group $F(1, 41) = 0.15, p = .699, \eta_p^2 < .01$ but a significant interaction $F(1, 41) = 6.72, p = .013, \eta_p^2 = .14$ (See Figure 2) revealed that FFDs were equivalent for perceptual and social oddities for TD participants $t(23) = 0.86, p = .399, d = .16$, but for ASD participants FFDs for social oddities were significantly shorter, in comparison to perceptual oddities, $t(18) = 2.99, p = .008, d = .65$, see Figure 2. Given that the FFD is a measure of early processing (Liversedge & Findlay, 2000; Rayner, 1998; Rayner, 2009). this finding indicates that the ASD group did not detect the social oddities as quickly as the perceptual oddities on initial fixation, whereas TD participants detected both types of oddity equally fast.

Insert Figure 2. about here

1
2
3 **First gaze duration (FGD) in the target region.** The FGD is the elapsed time from when the
4 target region is first fixated until the eye moves on to a new location. FGD includes multiple
5 fixations in the target region reflects an overt strategy to re-examine an area. A main effect of
6 oddity type, $F(1, 41) = 12.10, p = .001, \eta_p^2 = .23$ demonstrated that both groups spent more time
7 inspecting perceptual oddities, again highlighting the salience of the perceptual oddities in
8 comparison to social oddities in this study. However, there was no group effect $F(1, 41) = 2.43,$
9 $p = .127, \eta_p^2 = .06,$ or interaction $F(1, 41) = .08, p = .785, \eta_p^2 < .01.$

10
11
12
13
14
15
16
17
18
19
20
21
22 **Mean Fixation Count in the target region.** If the ASD are slower to respond manually, but do
23 not take longer to begin inspecting the target region, then extra time to respond might reflect
24 more fixations in the target region. A main effect of oddity type $F(1, 41) = 64.27, p < .001, \eta_p^2 =$
25 $.61,$ demonstrated more fixations in the target region when the oddity was perceptual, rather than
26 social. A main effect of group $F(1, 41) = 6.99, p = .012, \eta_p^2 = .15$ showed that the ASD group
27 made more fixations than the TD group in the target regions, for both types of oddities. However
28 no interaction was observed $F(1, 41) = 0.77, p = .386, \eta_p^2 = .02$ indicating that the ASD group
29 made more fixations in the target regions for both types of oddities.

30
31
32
33
34
35
36
37
38
39
40
41
42
43 **Mean Fixation Duration in the target region.** Longer mean fixation durations indicate that
44 more processing has been done during the task. A main effect of oddity type $F(1, 41) = 8.22, p =$
45 $.007, \eta_p^2 = .17$ showed significantly longer fixations for social, compared to perceptual oddity
46 target regions. This suggests that processing social oddities was more effortful, and since there
47 was no main effect of group $F(1, 41) = 0.34, p = .562, \eta_p^2 < .01,$ and no interaction $F(1, 41) =$
48 $1.12, p = .297, \eta_p^2 = .03,$ both groups on average fixated the social oddity target region for longer
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 than the perceptual target region. Thus, while the perceptual oddities attracted more fixations
4
5 overall, individual fixations were longer for the social target region.
6
7

8 This interesting juxtaposition, although unrelated to the main aim of this study, suggests
9
10 that difficulty associated with processing each oddity type manifests in different eye movement
11
12 patterns: one reflecting an overt strategy (fixation count) to re-inspect the target area, and one
13
14 reflecting more in depth on-line cognitive processing (fixation duration) on the target areas.
15
16
17

18
19
20 **Mean Number of Entries to the target region.** Since there were no group differences in FGD,
21
22 and no increased MFD, both reported above, the delayed RT in ASD might reflect repeated visits
23
24 to inspect the target region?
25
26

27 A main effect of oddity type was found $F(1, 41) = 9.83, p = .003, \eta_p^2 = .19$, with the
28
29 target region entered more often for perceptual oddities in comparison to social oddities. In
30
31 addition, a main effect of group was found $F(1, 41) = 5.72, p = .021, \eta_p^2 = .12$, with ASD
32
33 participants entering the target regions significantly more than the TD group, before making a
34
35 response. There was no interaction $F(1, 41) = 0.13, p = .717, \eta_p^2 < .01$.
36
37
38
39
40

41 **Mean Total Time in the target region.** A higher mean total time means that more attention to,
42
43 and processing of, information in that region was carried out overall. This measure should
44
45 complement the measures reported above – namely, that if there are more fixations overall in
46
47 ASD then the Total Time spent fixating in the target region should be longer in ASD.
48
49

50 A main effect of oddity type $F(1, 41) = 34.89, p < .001, \eta_p^2 = .46$, showed perceptual
51
52 oddities received more overall processing time, in line with the previous measures for the
53
54 perceptual oddities. In addition, a main effect of group $F(1, 41) = 6.41, p = .015, \eta_p^2 = .14$
55
56
57
58
59
60

1
2
3 revealed that the ASD participants spent a longer period of time inspecting the target regions for
4
5 both perceptual and social oddities. There was no interaction $F(1, 41) = 0.51, p = .481, \eta_p^2 = .01$.
6
7

8 As above, this finding reflects a greater number of entries to the target region overall by
9
10 the ASD group prior to making their decision as to whether the picture under inspection was
11
12 weird, odd or unusual, irrespective of the nature of the violation (perceptual or social).
13
14
15
16
17

18 Discussion

19
20 A principal aim of the study was to investigate whether increased times in ASD to find
21
22 and initially detect an unusual target in a scene containing people would be modulated by
23
24 whether the violation could be categorised as social or perceptual.
25
26

27 Accuracy rates were equivalent and high for both groups, but the ASD group took longer
28
29 to respond manually. It was hypothesised that the eye movement data would reveal the cause of
30
31 this delayed response in ASD, since eye movements reflect how information is processed on-line
32
33 (Liversedge & Findlay, 2000; Rayner, 1998; 2009). The data showed that our ASD group did not
34
35 take longer to find the target regions, but that they scanned the scenes more prior to responding
36
37 manually, as shown by an increased number of fixations throughout each scene presentation.
38
39 Furthermore, these increased fixations were not random, but were made to the target regions of
40
41 both social and perceptually odd pictures. Thus, increased fixations reflect a propensity to re-
42
43 inspect the target regions prior to making a response in ASD, regardless as to the type of oddity
44
45 in the scenes. It should be noted here that the oddity detection task in the current study provides
46
47 the ASD participants with structure that is not typically available in real-life. As such, the task is
48
49 not implicit and the knowledge that oddities might be present could also be a factor that drives
50
51 the repeated scanning in ASD.
52
53
54
55
56
57
58
59
60

1
2
3 The eye movement patterns also showed that our ASD group failed to immediately detect
4 social oddities, thus looking at but not seeing them. An obvious question from this finding is
5 whether the failure to quickly spot subtle social oddities in ASD reflects an inability to react or
6 respond appropriately to social cues in everyday communication? This one impairment in itself
7 could potentially account for the significant problems experienced in dynamic social interactions
8 in this high functioning population. For example, a failure to immediately spot social cues would
9 result in a failure to react or respond to such cues thus resulting in an inability to both keep track
10 of what is going on in social interactions, and a failure to engage appropriately or effectively
11 with other people.
12
13
14
15
16
17
18
19
20
21
22
23

24 If social oddities are more complex to detect in ASD, then the Complex Information
25 Processing theory (Minshew & Goldstein, 1998) might explain the failure to detect the social
26 oddity on first fixation. This lack of immediate detection could reflect the less obvious and more
27 'inferential' nature of the social oddities, which required more complex reasoning, at least in the
28 ASD group, for detection. It is important to note that social rules are often implicit, sometimes
29 ambiguous, and cannot always be understood by following a script or flow chart as in the case
30 for logical rule based tasks. At a neurobiological level, it has been postulated that that
31 impairments in cognitive processing in ASD reflect slow information transmission rates for high
32 level cognitive tasks, resulting from underconnectivity between frontal-posterior cortical regions
33 (Just et al., 2013; Just et al., 2012; Minshew & Williams, 2007). Since underconnectivity reduces
34 the rate of inter-region information transfer, it could be that in social communication tasks that
35 rely on the fast detection of implicit, often ambiguous cues, a failure to be able to transmit
36 information across different cortical areas quickly, might mean that these cues are missed,
37 rendering inappropriate responses in ASD.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The findings of repeated scanning of the target areas, coupled with an inability to
4 immediately recognise social violations upon initial fixation, reflect how eye movements can
5 reveal subtle processing differences in ASD. In everyday social communication in the real world
6 there is no opportunity to 'go back' to check if one has missed an important cue. Social
7 processing in such circumstances must be carried out on-line, and our eye movement data clearly
8 show that ASD are unable to do this quickly. The strategy of re-checking their interpretation, and
9 thus not believing what they have seen, before responding, may be related to cortical
10 underconnectivity in ASD. In everyday terms, these findings have obvious consequences for fast
11 dynamic social interactions, and offer an account as to why high functioning adults with ASD
12 experience difficulty in the social domain.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

This work was supported by a grant awarded to the second author from the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the Ontario Ministry of Research and Innovation, and Queen’s University Senate Advisory Research Committee, and an Experimental Psychology Society study visit grant awarded to the third author.

For Peer Review

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Au-Yeung, S. K., Benson, V., Castelhana, M. S., Rayner, K. (2011). Eye movement sequences during simple versus complex information processing of scenes in autism spectrum disorder. *Autism Research and treatment*, 2011.
- Au-Yeung, S. K., Kaakinen, J. K., Benson, V. (2013). Cognitive perspective taking during scene perception in autism spectrum disorder: evidence from eye movements. *Autism Research*, 7, 84-93.
- Bar-Haim, Y., Shulman, C., Lamy, D., & Reuveni, A. (2006). Attention to eyes and mouth in high-functioning children with autism, *Journal of Autism and Developmental Disorders*, 36, 131-137.
- Baron-Cohen, S. (2002). The extreme male brain theory of autism. *Trends in Cognitive Science*, 6, 248-254.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5-17.
- Benson, V., & Fletcher-Watson, S. (2012). Eye Movements in Autism. In: *Liversedge, S. P., Gilchrist, I. D., Everling, S., editors. The Oxford Handbook of Eye Movements*, Oxford UK: Oxford University Press, chap 39.
- Benson, V., Castelhana, M. S., Au-Yeung, S. K., & Rayner K. (2012). Eye movements reveal no immediate "WOW" ("which one's weird") effect in autism spectrum disorder, *Quarterly*

1
2
3 Journal of Experimental Psychology, 65, 1139-1150.

4
5 Benson, V., Piper, J., & Fletcher-Watson, S. (2009). Atypical saccadic sampling in autistic
6 spectrum disorder, *Neuropsychologia*, 47, 1178-1182, (2009).

7
8 Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate neural
9 responses to social stimuli in autism spectrum disorders. *Neuroimaging*, 15, 1614-1624.

10
11 Brugha, T. S., McManus, S., Bankart, J., Scott, F., Purdon S., et al. (2011). Epidemiology of
12 autism spectrum disorders in adults in the community in England. *Archives of General*
13 *Psychiatry*, 68, 459-65.

14
15 Clark, F., Scharaschkin, A., Xu, D. (2009). Supporting autism through adulthood, National Audit
16 Office.

17
18 Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith,
19 H. H., Alexander, A. L., & Davidson, R. J. (2005). Gaze fixation and the neural circuitry
20 of face processing in autism. *Nature Neuroscience*, 8, 519-526.

21
22 Fletcher-Watson, S., Leekam, S. R., Benson, V., Frank, M. C., & Findlay, J. M. (2009). Eye-
23 movements reveal attention to social information in autism spectrum disorder.
24
25 *Neuropsychologia*, 47, 248-257.

26
27 Fletcher-Watson., Leekam, S.R., Benson, V., Frank, M.C., & Findlay, J.M. (2009). Eye-
28 movements reveal attention to social information in autism spectrum disorder.
29
30 *Neuropsychologia*, 47, 248-257.

31
32 Frith, U., & Happé, F. (1994). Autism; Beyond the theory of mind. *Cognition*, 50, 115-132.

33
34 Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in
35 autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 36, 5-25.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of
4
5 approaching adults predicts level of social disability in 2-year-old toddlers with autism
6
7 spectrum disorder. *Archives of General Psychiatry*, 65, 946-954.
8
9
- 10 Just, M. A., Keller, T. A., & Kana, R. K. (2013). A theory of autism based on frontal-posterior
11
12 underconnectivity. In M. A. Just & K. A. Pelphrey (Eds.), *Development and brain*
13
14 *systems in autism* (pp. 35-63). New York: Psychology Press.
15
16
- 17 Just, M. A., Keller, T. A., Malave, V. L., Kana, R. K., & Varma, S. (2012). Autism as a neural
18
19 systems disorder: A theory of frontal-posterior underconnectivity. *Neuroscience and*
20
21 *Biobehavioral Reviews*, 36, 1292-1313.
22
23
- 24 Klin, A., & Jones, W. (2006). Attributing social and physical meaning to ambiguous visual
25
26 displays in individuals with higher functioning autism spectrum disorders. *Brain &*
27
28 *Cognition*, 61, 40-53.
29
30
- 31 Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen D. J. (2002a). Defining and
32
33 quantifying the social phenotype in autism, *American Journal of Psychiatry*, 159, 895-
34
35 908.
36
37
- 38 Klin, A., Jones, W., Schultz, R. T., Volkmar, F. R., & Cohen, D. J. (2002b). Visual fixation
39
40 patterns during viewing of naturalistic social situations as predictors of social competence
41
42 in individuals with autism. *Archives of General Psychiatry*, 59, 809-816.
43
44
- 45 Kuhn, G. Benson, V., Fletcher-Watson, S., Kovshoff, H., McCormick, C., & Kirkby, J. (2010)
46
47 Eye movements affirm: Automatic overt gaze and arrow cueing for typical adults and
48
49 adults with autism spectrum disorder. *Experimental Brain Research*, 201, 155-165.
50
51
- 52 Liversedge, S. P., & Findlay, J. M. (2000). Saccadic eye movements and cognition. *Trends in*
53
54 *Cognitive Sciences*, 4, 6-14.
55
56
57
58
59
60

1
2
3 Lord C., Rutter M., DiLavore P. C., Risi S., Gotham K., Bishop S. (2012). Autism diagnostic
4 observation schedule, second edition. Torrance, CA: Western Psychological Services.
5
6

7
8 Louwrese, A., Tulen, J. H. M., van der Geest, J. N., van der Ende, J., Verhulst, F. C., & Greaves-

9
10 Lord, K. (2014). Autonomic Responses to Social and Nonsocial Pictures in Adolescents
11 With Autism Spectrum Disorder. *Autism Research*, 7, 17-27. doi:10.1002/aur.1327
12
13

14
15 Minshew, N. J., & Goldstein, G. (1998). Autism as a disorder of complex information
16 processing. *Mental Retardation and Developmental Disabilities Research Reviews*, 4,
17 129-136.
18
19

20
21
22 Minshew, N. J., & Williams, D. L., (2007). The new neurobiology of autism: cortex,
23 connectivity, and neuronal organization. *Archives of Neurology*, 64, 945-950.
24
25

26
27 Minshew, N. J., Goldstein, G., & Siegel, D. J. (1997). Neuropsychologic functioning in autism:
28 Profile of complex information processing disorder. *Journal of the International*
29 *Neuropsychological Society*, 3, 303-316.
30
31
32

33
34 Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002).
35 Visual scanning of faces in autism, *Journal of Autism and Developmental Disorders*, 32,
36 249-261.
37
38
39

40
41 Rayner, K. (2009). Eye movements and attention in reading, scene perception, and visual search.
42 *Quarterly Journal of Experimental Psychology*, 62, 1457-1506.
43
44

45
46 Rayner, K., Castelhana, M. S., & Yang J. (2009). Eye Movements when Looking at
47 Unusual/Weird Scenes: Are there Cultural Differences? *Journal of Experimental*
48 *Psychology: Language, Memory & Cognition*, 35, 254-259.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Rayner, K., Warren, T., Juhasz, B. J., & Liversedge, S. P. (2004). The effect of plausibility on
4
5 eye movements in reading. *Journal of Experimental Psychology-Learning Memory and*
6
7
8 *Cognition*, 30(6), 1290-1301.

9
10 Riby, D. M., & Hancock, P. J. B. (2009). Do faces capture the attention of individuals with
11
12 Williams syndrome or Autism? Evidence from tracking eye movements, *Journal of*
13
14 *Autism and Developmental Disorders*, 39, 421-431. doi:10.1007/s10803-008-0641-z

15
16
17 Russell, J. (1997). *Autism as an executive disorder*. Oxford, UK: Oxford University Press.

18
19
20 Speer, L. L., Cook, A. E., McMahon, W. M., & Clark, E. (2007). Face processing in children
21
22 with autism: effects of stimulus contents and type, *Autism*, 11, 263-277.

23
24
25 Stampe, D. M. (1993). Heuristic filtering and reliable calibration methods for video-based pupil-
26
27 tracking systems. *Behavior Research Methods, Instruments, & Computers*. 25, 137-
28
29 142.

30
31
32 The Psychological Corporation, Wechsler Abbreviated Scale of Intelligence (WASI) Manual,
33
34 (The Psychological Corporation, San Antonio, TX, 1999).

35
36
37 Wolfe, J. M. (1998) What can one million trials tell us about visual search? *Psychological*
38
39 *Science*, 9, 33-39.

Footnotes

1
2
3
4
5
6 1. Accuracy errors for the excluded scenes were either a result of a failure to detect a subtle
7
8 oddity, or were a result of participants thinking the control version of the scenes were odd.
9
10 Removing these items from analysis did not change the pattern of effects, but reduced variation
11
12 in reaction time and eye movement data.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Tables

Table 1.

Accuracy and reaction time means (standard deviations).

	Perceptual				Social			
	Normal		Weird		Normal		Weird	
	TD	ASD	TD	ASD	TD	ASD	TD	ASD
Accuracy	0.81 (0.20)	0.86 (0.10)	0.84 (0.20)	0.77 (0.17)	0.80 (0.17)	0.83 (0.12)	0.80 (0.21)	0.76 (0.24)
Reaction Time (ms)	5289 (2372)	6024 (2560)	3697 (1703)	5184 (2228)	5117 (2443)	6053 (2884)	3056 (1270)	4633 (2131)

Table 2.

Eye movement measure means in ms (standard deviations).

		Perceptual Oddity		Social Oddity	
		TD	ASD	TD	ASD
Mean Fixation Duration (overall)		250 (31)	257 (29)	241 (26)	252 (33)
Mean Fixation Count (overall)		11 (4)	15 (6)	9 (4)	13 (7)
Elapsed time to target region	ELT	551 (235)	557 (144)	600 (235)	679 (284)
Fixation count before target region	FCT	2.32 (0.72)	2.49 (0.67)	2.62 (0.76)	2.68 (0.73)
First fixation duration in target region	FFD	224 (52)	237 (42)	232 (45)	210 (41)
First gaze duration in target region	FGD	808 (250)	924 (437)	630 (163)	715 (208)
Mean fixation duration in target region	MFD	262 (43)	274 (34)	281 (38)	282 (45)
Mean fixation count in target region	MFC	5.78 (2.28)	7.72 (2.84)	3.97 (1.81)	5.47 (2.14)
Mean number of entries into target region	MNE	1.93 (0.64)	2.32 (0.73)	1.62 (0.52)	2.07 (0.66)
Mean total time in target region	TT	1522 (729)	2065 (809)	1040 (527)	1450 (664)

Figures captions

Figure 1. Example of social normal (A), social weird (B), perceptual normal (C) and perceptual weird (D) stimuli. The black square represents the target region. Note the black square is not visible during the experiment.

Figure 2. Interaction for first fixation duration (ms) between group and oddity type. Error bars denote \pm 1 SE.

For Peer Review