

# Using dengue epidemics and local weather in Bali, Indonesia to predict imported dengue in Australia

Zhiwei Xu<sup>1,2</sup>, Hilary Bambrick<sup>1,2</sup>, Laith Yakob<sup>3</sup>, Gregor Devine<sup>4</sup>, Francesca D. Frentiu<sup>2,5</sup>, Rina Marina<sup>6</sup>, Pandji Wibawa Dhewantara<sup>7,8</sup>, Roy Nusa<sup>9</sup>, R. Tedjo Sasmono<sup>10</sup>, Wenbiao Hu<sup>1,2\*</sup>

## Author affiliations

<sup>1</sup> School of Public Health and Social Work, Queensland University of Technology, Brisbane 4059, Australia;

<sup>2</sup> Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane 4059, Australia;

<sup>3</sup> Department of Disease Control, London School of Hygiene and Tropical Medicine, London WC1H 9SH, UK;

<sup>4</sup> Mosquito Control Laboratory, QIMR Berghofer Medical Research Institute, Brisbane 4006, Australia;

<sup>5</sup> School of Biomedical Sciences, Queensland University of Technology, Brisbane 4059, Australia;

<sup>6</sup> Center of Public Health Effort Research and Development, National Institute of Health Research and Development, Jakarta 10560, Indonesia;

<sup>7</sup> UQ Spatial Epidemiology Laboratory, School of Veterinary Science, University of Queensland, Gatton 4343, Australia;

<sup>8</sup> Pangandaran Unit for Health Research and Development, National Institute of Health Research and Development, Ministry of Health of Indonesia, Pangandaran 46396, Indonesia;

<sup>9</sup> Indonesian Ministry of Health, Jakarta 12950, Indonesia;

<sup>10</sup> Eijkman Institute for Molecular Biology, Jakarta 10430, Indonesia.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**\*Correspondence to:**

Prof. Wenbiao Hu, School of Public Health and Social Work, Queensland University of  
Technology, Victoria Park Road, Kelvin Grove, Brisbane, Queensland 4059, Australia.

Telephone number: +61 7 3138 5724. Email address: [w2.hu@qut.edu.au](mailto:w2.hu@qut.edu.au)

**Conflict of interest:**

The authors report no conflicts of interest.

1  
2  
3  
4 **Abstract**  
5  
6  
7

8 **Background:** Although the association between dengue in Bali, Indonesia, and imported dengue  
9  
10 in Australia has been widely asserted, no study has quantified this association so far.  
11  
12

13 **Methods:** Monthly data on dengue and climatic factors over the past decade for Bali and Jakarta  
14  
15 as well as monthly data on imported dengue in Australia underwent a three-stage analysis. Stage  
16  
17 I: a quasi-Poisson regression with distributed lag non-linear model was used to assess the  
18  
19 associations of climatic factors with dengue in Bali. Stage II: a generalized additive model was  
20  
21 used to quantify the association of dengue in Bali with imported dengue in Australia with and  
22  
23 without including the number of travelers in log scale as an offset. Stage III: the associations of  
24  
25 mean temperature and rainfall (two climatic factors identified in stage I) in Bali with imported  
26  
27 dengue in Australia were examined using stage I approach.  
28  
29  
30  
31  
32

33 **Results:** The number of dengue cases in Bali increased with increasing mean temperature, and,  
34  
35 up to a certain level, it also increased with increasing rainfall but dropped off for high levels of  
36  
37 rainfall. Above a monthly incidence of 1.05 cases per 100,000, dengue in Bali was almost  
38  
39 linearly associated with imported dengue in Australia at a lag of one month. Mean temperature  
40  
41 (relative risk (RR) per 0.5 °C increase: 2.95, 95% confidence interval (CI): 1.87, 4.66) and  
42  
43 rainfall (RR per 7.5 mm increase: 3.42, 95% CI: 1.07, 10.92) in Bali were significantly  
44  
45 associated with imported dengue in Australia at a lag of four months.  
46  
47  
48  
49  
50

51 **Conclusions:** This study suggests that climatic factors (i.e., mean temperature and rainfall)  
52  
53 known to be conducive of dengue transmission in Bali can provide an early warning with 4-  
54  
55 month lead time for Australia in order to mitigate future outbreaks of local dengue in Australia.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

This study also provides a template and framework for future surveillance of travel-related infectious diseases globally.

**Keywords:** Australia; Bali; dengue; early warning

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## **Funding sources**

This work was supported by the National Health and Medical Research Council (APP  
1138622\_Hu).

# 1. Introduction

Dengue is the most common and geographically fastest-spreading mosquito borne disease in the world, and is now present in more than 126 countries (Furuya-Kanamori et al. 2016). It has been estimated that there are 390 million dengue infections per year globally and 96 million of them manifest apparently (Bhatt et al. 2013). A large proportion of global dengue burden is borne by people living in Southeast Asia and Western Pacific region (WHO Western Pacific Region 2014). In Australia, local dengue transmission is restricted to northern Queensland and is seeded by importations, and dengue infections in other Australian states and territories are imported cases (Huang et al. 2016).

Indonesia, a dengue endemic country, has been considered as an important hub for dengue transmission in Asia Pacific region (Vainio et al. 2010; Yuan and Nishiura 2018). Indonesia is the most frequent international travel destination of Australians (Australian Bureau of Statistics 2018), and prior studies have suggested that a large proportion of Australian dengue cases acquired in Indonesia are from the island of Bali (Masyeni et al. 2018), the most famous tourist spot destination in Indonesia (Knope et al. 2013). Further, a considerable number of residents in Western Australia work in Bali, with these “fly-in-fly-out” workers traveling between Bali and Perth frequently due to cheap four-hour flights (Butt 2017). In Western Australia, during 2010 and 2011, more than 80% of dengue cases acquired in Indonesia were acquired in Bali (Knope et al. 2013). Although the relationship between dengue in Bali and imported dengue in Australia has been widely asserted (Asma et al. 2019; Ernst et al. 2015), no study has yet quantified this relationship, and the risk posed to Australia by local factors affecting endemicity in Bali remains unclear.

1  
2  
3  
4 Climatic factors, including temperature, rainfall, relative humidity and wind speed, affect the  
5  
6 occurrence of dengue cases through their impacts on the replication rate and transmission of  
7  
8 dengue virus, mosquito ecology and human behaviors (Morin et al. 2013). The effect of climatic  
9  
10 factors on the dengue transmission is therefore usually delayed (i.e., lag effect), and the lag  
11  
12 period ranges from a few weeks to a few months (Xiang et al. 2017; Xu et al. 2017). This lag  
13  
14 period potentially enables prediction of dengue occurrence a few months in advance and  
15  
16 provides the opportunity to take proactive measures in a timely manner (Lowe et al. 2018).  
17  
18

19  
20  
21 If dengue in Bali can predict imported dengue in Australia, and climatic factors in Bali can  
22  
23 predict dengue in Bali, our hypothesis is that imported dengue in Australia can be predicted  
24  
25 using Balinese climatic data. The present study attempted to test this hypothesis by achieving  
26  
27 two research objectives: 1) to quantify the temporal relationship between dengue in Bali and  
28  
29 imported dengue in Australia; and 2) to explore the associations of climatic factors in Bali and  
30  
31 imported dengue in Australia. To test the uniqueness of Bali in the role of spreading dengue to  
32  
33 Australia, we also used data from Jakarta (the capital city of Indonesia) and examined the  
34  
35 relationship between dengue in Jakarta and imported dengue in Australia.  
36  
37  
38  
39  
40  
41  
42  
43  
44

## 45 **2. Materials and methods**

### 46 47 48 **2.1 Data collection**

49  
50  
51 Figure S1 (supplementary material) shows the geographical locations of Bali, Jakarta and  
52  
53 Australia. It also shows the locations of Bali districts. Monthly dengue data and yearly  
54  
55 population data in Bali and its districts from January 2007 to December 2017 as well as in  
56  
57 Jakarta from January 2001 to December 2017 were provided by the Ministry of Health,  
58  
59  
60  
61

1  
2  
3  
4 Indonesia. Dengue data in Karangasem (a Bali district) from August to December 2010 were  
5  
6 missing, and dengue data in all Bali districts in December 2010 were missing except for  
7  
8 Klungkung. Daily weather data, including mean temperature, maximum temperature, minimum  
9  
10 temperature, relative humidity, rainfall, and wind speed, from January 2007 to December 2017 in  
11  
12 Bali were obtained from Meteorological, Climatological and Geophysical Agency, Indonesia  
13  
14 (<http://dataonline.bmkg.go.id/home>). Daily weather data were merged into monthly climate data  
15  
16  
17 by averaging daily values.  
18  
19  
20  
21

22 Monthly dengue data (including both local dengue and imported dengue) in Australia and its  
23  
24 eight states and territories (Australian Capital Territory, New South Wales, Northern Territory,  
25  
26 Queensland, South Australia, Tasmania, Victoria, and Western Australia) from January 2001 to  
27  
28 May 2018 were obtained from the National Notifiable Diseases Surveillance System  
29  
30 (<http://www9.health.gov.au/cda/source/cda-index.cfm>). Queensland has both local and imported  
31  
32 dengue cases (Hu et al. 2012). Monthly local dengue data and imported dengue data in  
33  
34 Queensland from January 2007 to December 2013 were provided by Queensland Health, and we  
35  
36 used this data to examine the relationships of dengue in Bali with imported dengue and local  
37  
38 dengue in Queensland. This study aimed to quantify the temporal relationship between dengue in  
39  
40 Bali and imported dengue in Australia, and there is regular conflation between imported dengue  
41  
42 and local dengue in Queensland resulting in misclassification (Huang et al. 2013; McKerr et al.  
43  
44 2015). Thus, results for Queensland are presented as supplementary material in this study. In this  
45  
46 paper, the term “imported dengue in Australia” refers to “imported dengue in Australia excluding  
47  
48 Queensland”, unless stated otherwise. Data on monthly number of Australian residents returning  
49  
50 from Indonesia from January 2001 to May 2018 were obtained from the Australia Bureau of  
51  
52 Statistics (Australian Bureau of Statistics 2018).  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 Ethics approval was granted by University Human Research Ethics Committee of Queensland  
5  
6  
7 University of Technology prior to the data collection.  
8  
9

## 10 **2.2 Data analysis**

11  
12 A three-stage analysis was conducted. In the first stage, there were two steps. Step I: we  
13 examined the relationship between each climatic factor and occurrence of dengue cases in Bali  
14 (i.e., one-climatic-factor model) and selected those climatic factors which were significantly  
15 associated with the occurrence of dengue cases. We found significant associations between each  
16 of mean temperature, maximum temperature, minimum temperature, relative humidity, and  
17 rainfall with the occurrence of dengue cases in Bali (Figure S2 (supplementary material)). Step II:  
18 using these significant climatic factors identified in step I, we built up models including two  
19 climatic factors. Because temperature indicators (i.e., mean temperature, maximum temperature,  
20 and minimum temperature) are highly correlated with each other and relative humidity and  
21 rainfall are also highly correlated with each other, we chose one variable from temperature  
22 indicators and one variable from relative humidity and rainfall to build up the models to avoid  
23 multicollinearity issues. By comparing the values of quasi-AIC (Akaike information criterion) of  
24 all models in step II (Gasparrini and Armstrong 2013), we found that the model with mean  
25 temperature and rainfall performed the best (Table S1 (supplementary material)). The optimal  
26 lags for mean temperature and rainfall in the model (i.e., three months) were selected based on  
27 the greatest values of pre-whitened cross correlation (Figures S3A and S3B (supplementary  
28 material)) (Chuang et al. 2017). For both two steps of the first stage, we used Poisson regression  
29 allowing for over-dispersion with distributed lag non-linear model (Gasparrini et al. 2010) as  
30 prior studies have reported non-linear climate-dengue associations (Lowe et al. 2018; Wu et al.  
31 2018). We herewith use mean temperature as an example to elaborate the modeling details. A  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 cross-basis defined by natural cubic spline with two degrees of freedom (*dfs*) for the space of  
5  
6 mean temperature was used. A maximum lag of three months was adopted in the analysis based  
7  
8 on the findings of previous studies quantifying climate-dengue relationships (Lowe et al. 2018;  
9  
10 Wu et al. 2018; Xiang et al. 2017) and model fit results. Month and year were included as  
11  
12 dummy variables in the model to control for seasonality and long-term trend.  
13  
14

15  
16  
17 In the second stage, there were two steps. Step I: we assessed the correlation between dengue in  
18  
19 Bali and imported dengue in Australia at different lags to ascertain whether there may be a  
20  
21 quantitative association between dengue in Bali and imported dengue in Australia. Significant  
22  
23 correlations between dengue in Bali and imported dengue in Australia at all lags were observed,  
24  
25 with correlation coefficients at a lag of one month having the strongest correlation (Table S2  
26  
27 (supplementary material)). We've also calculated the pre-whitened cross correlation between  
28  
29 dengue in Bali and imported dengue in Australia and found the correlation coefficient value was  
30  
31 the greatest at a lag of one month (Figure S4 (supplementary material)). This finding echoes the  
32  
33 result from Asma et al. whereby dengue cases in Australia were typically presented to  
34  
35 emergency departments within 14 days of return from Bali (Asma et al. 2019). Step II: based on  
36  
37 the findings of step I in the second stage analysis, we used a generalized additive model to  
38  
39 quantify the relationship between dengue in Bali and dengue in Australia at a lag of one month.  
40  
41 In this model, month and year were included as dummy variables to control for seasonality and  
42  
43 long-term trend. The number of travelers in a log scale was included in the model as an offset.  
44  
45 To make sure that a lag of one month was the optimal lag in the model, we compared the values  
46  
47 of un-biased risk estimator (UBRE) of three models using three different lags (i.e., one month,  
48  
49 two months, and three months), and found that the UBRE values of the three models were  
50  
51 5.4875 (one month), 6.1577 (two months), and 6.5031 (three months), respectively. The lowest  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 value of UBRE was observed for the one-month-lag model, suggesting that the optimal lag was  
5  
6 one month.  
7

8  
9  
10 In the third stage, we used Poisson regression allowing for over-dispersion with a distributed lag  
11  
12 non-linear model to assess the associations between mean temperature and rainfall in Bali with  
13  
14 the occurrence of dengue cases in Australia. A lag of four months was used in the third stage  
15  
16 analysis, and the choice of this lag period was based on the analyses results of the first and  
17  
18 second stages. The logic behind this is: 1) Dengue in Bali drives the number of Australian  
19  
20 imported dengue cases. 2) Number of dengue cases in Bali can be predicted by climatic factors  
21  
22 in Bali several months in advance. 3) Therefore, import risk of dengue in Australia can be  
23  
24 predicted by climatic factors in Bali several months in advance.  
25  
26  
27  
28

29  
30 We replicated above mentioned second stage analysis in Jakarta to examine the relationship  
31  
32 between dengue in Jakarta and imported dengue in Australia and test the specificity of the  
33  
34 observed Bali association. All analyses in the present study were conducted in R statistical  
35  
36 environment (version 3.4.4) with “dlnm” package used to fit the distributed lag non-linear model  
37  
38 (Gasparini 2011) and “mgcv” package used to fit the generalized additive model.  
39  
40  
41  
42  
43  
44

### 45 **3. Results**

#### 46 ***3.1 Monthly distributions of dengue cases in Australia, Bali and Jakarta***

47  
48  
49  
50 Figures S5A, S5B, and S5C (supplementary material) present the monthly distributions of  
51  
52 imported dengue cases in Australia and dengue cases in Bali and Jakarta. The number of  
53  
54 imported dengue cases in Australia increased after 2010. Dengue in Bali had a distinct seasonal  
55  
56  
57  
58  
59  
60  
61  
62

1  
2  
3  
4 pattern, with the biggest dengue outbreak occurring in 2016. The biggest dengue outbreak in  
5  
6 Jakarta occurred in 2004.  
7  
8

### 9 10 ***3.2 Associations of climatic factors with dengue in Bali***

11  
12 Figure 1 shows the associations between mean temperature and rainfall with the occurrence of  
13 dengue cases in Bali, suggesting that the number of dengue cases increased with increased mean  
14  
15 temperature. Number of dengue cases also increased with increasing rainfall, but only up to 16.2  
16  
17 mm beyond which cases started dropping off.  
18  
19  
20  
21  
22

### 23 24 ***3.3 Association between dengue in Bali and dengue in Australia***

25  
26 Figure 2A illustrates a significant association between dengue in Bali and imported dengue in  
27  
28 Australia at a lag of one month. At the Australian state level, dengue in Bali was also  
29  
30 significantly associated with dengue in New South Wales, Northern Territory, Tasmania,  
31  
32 Victoria, and Western Australia at a lag of one month. Most of the associations were practically  
33  
34 linear. Figure S6 (supplementary material) shows that, from 2007 to 2013, there were significant  
35  
36 relationships of dengue in Bali with imported dengue in Australia (including Queensland  
37  
38 imported dengue) and local dengue in Queensland at a lag of one month.  
39  
40  
41  
42  
43

44 Dengue in Jakarta and imported dengue in Australia at a lag of one month is presented in Figure  
45  
46 2B, and demonstrates no clear association.  
47  
48  
49

### 50 51 ***3.4 Association between Australian residents returning from Indonesia and dengue in*** 52 53 ***Australia***

54  
55 Table S3 (supplementary material) shows the number of Australian residents returning from  
56  
57 Indonesia and the number of imported dengue cases in Australia has the strongest correlation  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 without a time lag and this was statistically significant (see Figure 3). Sensitivity analyses  
5  
6 assessing the association of number of Australian residents returning from Indonesia with  
7  
8 numbers of imported dengue cases in Australia at a lag of one month confirmed that the findings  
9  
10 in Figure 3 were robust (Figure S7 (supplementary material)).  
11  
12  
13

### 14 ***3.5 Association between dengue in Bali and dengue in Australia after including the number of*** 15 ***Australian residents returning from Indonesia in log scale as an offset in the model*** 16 17 18

19  
20 Figure 4 shows the association between dengue in Bali and imported dengue in Australia at a lag  
21  
22 of one month after including the number of Australian residents returning from Indonesia in log  
23  
24 scale as an offset in the model. Figure 5 disaggregates this to the Balinese districts level and  
25  
26 indicates that dengue in all Bali districts (except for Bangli and Denpasar) was significantly  
27  
28 associated with imported dengue in Australia. Table S4 (supplementary material) presents the  
29  
30 thresholds of monthly dengue incidence in Bali and its districts beyond which imported dengue  
31  
32 in Australia started to increase substantially, suggesting that when monthly dengue incidence in  
33  
34 Bali was above 1.05 per 100,000, dengue in Australia started to increase. The thresholds in Bali  
35  
36 districts including Badung, Klungkung, and Tabanan were stable.  
37  
38  
39  
40  
41  
42

### 43 ***3.6 Associations of climatic factors in Bali with dengue in Australia*** 44 45

46 Figure 6A and Figure 6B show the significant associations between mean temperature and  
47  
48 rainfall in Bali with occurrence of imported dengue cases in Australia, with the most pronounced  
49  
50 association found for the state of Victoria.  
51  
52  
53  
54  
55  
56

## 57 **4. Discussion** 58 59 60 61 62 63 64 65

1  
2  
3  
4 Indonesia, in particular Bali, has been recognized as a primary source for dengue importations  
5  
6 into Australia. Despite this, little has been done in terms of using case notification data from  
7  
8 Indonesia to inform public health measures to mitigate dengue importation into Australia. In  
9  
10 tweezing apart the associations between imported dengue, local Balinese endemicity and easily  
11  
12 accessible climatic variables, this study identifies a statistically robust early warning signal for  
13  
14 dengue imports into Australia with remarkably long lead-time. The mechanisms behind this lead-  
15  
16 time may include a few weeks between suitable climate and increased mosquito activity and  
17  
18 local dengue transmission in Bali, and the incubation time for infections in Australian travelers  
19  
20 as well as time to diagnosis in Australia.  
21  
22  
23  
24

25  
26 The life cycle of dengue viruses and the growth and survival of dengue mosquitos are  
27  
28 inextricably linked to climate (Li et al. 2019; Morin et al. 2013; Naish et al. 2014). In this study,  
29  
30 we observed that, in Bali, the number of dengue cases increased with the increase of mean  
31  
32 temperature, which is partially consistent with prior findings in China (Wu et al. 2018),  
33  
34 Singapore (Earnest et al. 2011), Sri Lanka (Goto et al. 2013), and Vietnam (Pham et al. 2011).  
35  
36 The number of dengue cases generally started dropping when mean temperature reached a high  
37  
38 level in prior studies but we did not observe this pattern in our study. We also found that the  
39  
40 relationship between rainfall and the number of dengue cases was bell-shaped, which is in line  
41  
42 with the findings in China (Yi et al. 2019) and the Philippines (Iguchi et al. 2018). Previous  
43  
44 studies examining the associations of climatic factors with dengue in Indonesia found contrasting  
45  
46 results in different regions. Tosepu et al. and Bangs et al. found that mean temperature was  
47  
48 significantly associated with the occurrence of dengue cases in Sulawesi Province and  
49  
50 Palembang (Indonesia), respectively (Bangs et al. 2006; Tosepu et al. 2018). However,  
51  
52 Kesetyaningsih et al. reported that rainfall and relative humidity, rather than temperature, were  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 associated with dengue in Yogyakarta, Indonesia (Kesetyaningsih et al. 2018). The reasons for  
5  
6 the discrepancy in dominant climatic driver(s) of dengue across different regions still remain  
7  
8 largely unclear (Morin et al. 2013). Moreover, the different analytical approaches used in these  
9  
10 studies and our study might also result in the different findings as prior studies in Indonesia  
11  
12 tended to assume a linear relationship between climatic factors and dengue. In recent years the  
13  
14 biological mechanisms underlying the effects of temperature and rainfall on dengue have been  
15  
16 increasingly explored. Specifically, increased temperature facilitates dengue virus replication  
17  
18 within the vectors and shortens the extrinsic incubation period of dengue viruses (Morin et al.  
19  
20 2013). Temperature also affects the development and survival of dengue mosquitos (Morin et al.  
21  
22 2013). Adequate rainfall provides enough water for potential breeding sites of the principal  
23  
24 vector mosquito *Aedes aegypti* (i.e, water containers), but very heavy rainfall may flush away  
25  
26 larvae and pupae from containers which restricts the development of mosquitoes (Sarfraz et al.  
27  
28 2012). This may explain our finding that the number of dengue cases increased when rainfall  
29  
30 increased but started dropping when rainfall exceeded 16.2 mm in Bali. The positive association  
31  
32 between mean temperature and number of dengue cases is of additional concern as global surface  
33  
34 temperature will continue to increase as climate change proceeds, and future temperature-related  
35  
36 dengue burden in Bali might consequently change and necessitate changes in vector control (Ebi  
37  
38 and Nealon 2016).

39  
40  
41 International travel related transmission of diseases has become a major global public health  
42  
43 issue (Fang et al. 2018; Findlater and I 2018). There were more than 1.2 billion international  
44  
45 tourists in 2016, and this figure is still growing (UNTWO 2017). Australia witnessed a  
46  
47 substantial increase in the number of residents travelling internationally over the past decade, and  
48  
49 the monthly number of Australian residents returning from Indonesia has increased substantially  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61

1  
2  
3  
4 from an average of 28,100 in the first five months of 2008 to an average of 98,980 in the first  
5  
6 five months of 2018 (Australian Bureau of Statistics 2018). International travel plays an  
7  
8 important role in the dispersal of dengue across different countries (e.g., Europe) (Choe et al.  
9  
10 2017; Semenza et al. 2014; Tian et al. 2017). An appreciable proportion of returning Australian  
11  
12 travelers hospitalized for dengue had unrecognized signs of severe disease (Tai et al. 2017). It  
13  
14 has been reported that travelers with dengue infection returning from Bali (Indonesia) to  
15  
16 Queensland (Australia) may present a risk of starting a local dengue outbreak (David et al. 2012).  
17  
18 In this study, we observed that, from 2007 to 2013, there was a significant relationship between  
19  
20 dengue in Bali and local dengue in Queensland at a lag of one month, and the magnitude of this  
21  
22 relationship was much greater than that found between dengue in Bali and imported dengue in  
23  
24 Australia (Figure S4 (supplementary material)). Identifying these relationships is of great  
25  
26 importance when considered in the context of providing more solid evidence for the efficacy in  
27  
28 ongoing and upscaling novel dengue vector control strategies employing *Wolbachia* (O'Neill  
29  
30 2018; O'Neill et al. 2018).

31  
32  
33  
34  
35  
36  
37  
38  
39 The first dengue case in Australia was reported in 1873 in Victoria (Medical Society of Victoria  
40  
41 1873), and local dengue cases were prevalent in Queensland, New South Wales, Northern  
42  
43 Territory, and Western Australia until the 1940s (Mackenzie et al. 1996). Dengue reappeared in  
44  
45 Queensland in 1981 and only Queensland has local dengue cases nowadays (Mackenzie et al.  
46  
47 1996). Nevertheless, existing literature reported that there is a potential of *Aedes aegypti*  
48  
49 incursion into northern parts of Western Australia and Northern Territory as climate change  
50  
51 continues (Russell et al. 2009). Another dengue vector mosquito, *Aedes albopictus*, may have  
52  
53 been introduced into Australia due to illegal fishing activity originating from Indonesia (Beebe et  
54  
55 al. 2013), implying that imported cases in Western Australia and Northern Territory may not just  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 pose extra burden to local healthcare system, but also may present a risk of dengue endemic in  
5 these regions if climatic conditions are favorable for dengue mosquitos in the future. The  
6  
7 quantitative associations of dengue in Bali with imported dengue in New South Wales, Northern  
8  
9 Territory, Tasmania, Victoria, and Western Australia that we observed in this study suggest that  
10  
11 routine surveys in Australian international airports including asking returned travelers about  
12  
13 countries they visited and their clinical manifestations as well as follow-ups of travelers with  
14  
15 warning signs of dengue (especially those who returned from Bali) can help identify dengue  
16  
17 patients in a timely manner and alert public health communities to take measures accordingly.  
18  
19 Intriguingly, we observed that the dengue incidence thresholds in Badung, Klungkung and  
20  
21 Tabanan were stable (Table S4), and this may be due to the fact that most tourists visit places in  
22  
23 these three districts because popular resorts, beaches, temples and natural parks are located in  
24  
25 these districts (e.g., Uluwatu Beach, Kuta Beach, Nusa Dua, Tanah Lot, Bedugul, and Nusa  
26  
27 Penida). Unexpectedly, we did not observe a significant relationship between dengue in  
28  
29 Denpasar and imported dengue in Australia (Figure 5), although Denpasar is an important  
30  
31 sightseeing destination for tourists.  
32  
33

34  
35  
36 This study quantified the relationship between dengue in Bali and dengue in Australia, and it  
37  
38 provides novel insight to inform the development of dengue early warning systems in Australia.  
39  
40 There are studies calling for actions to incorporate climate information into the development of  
41  
42 dengue early warning systems (Lee et al. 2017; Morin et al. 2018), and there are studies  
43  
44 elucidating the importance of using travel-related data in dengue early warning (Racloz et al.  
45  
46 2012). Our study suggested that the information on climatic conditions in and travelers returning  
47  
48 from one city may be used as important early warning signals for another city/region. It also calls  
49  
50 for future endeavors assessing how infectious disease outbreaks overseas affect infectious  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 disease importation into Australia. However, several limitations of this current study should be  
5  
6 acknowledged. First, because of data availability we used data on number of returning residents  
7  
8 in the whole of Australia rather than the number in each state/territory, which may not be  
9  
10 accurate as some states/territories may have relatively more or fewer travelers. Second, data on  
11  
12 number of residents returning specifically from Bali were not available, and thus we could only  
13  
14 use data on number of residents returning from the whole of Indonesia. Third, sensitivity of any  
15  
16 associations detected using our methods is compromised by under-reporting which is  
17  
18 characteristic of infections such as dengue that are highly asymptomatic or have non-specific  
19  
20 symptoms. Finally, local dengue outbreaks in northern Queensland were triggered annually by  
21  
22 the arrival of infected international travelers. In this study, we were only able to assess the  
23  
24 relationships of dengue in Bali with imported dengue and local dengue in the whole of  
25  
26 Queensland. Future research should try to improve the spatial resolution to allow for improved  
27  
28 targeting of surveillance and control. In this study, we were only able to analyze the dengue data  
29  
30 in Queensland from 2007 to 2013 due to data availability issue, and more up-to-date data on  
31  
32 imported dengue and local dengue in Queensland will be collected in the future.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

## 44 **5. Conclusion**

45  
46  
47  
48 Dengue in Bali was quantitatively associated with imported dengue in Australia at a lag of one  
49  
50 month. Incorporating information on dengue incidence in Bali one month ahead and information  
51  
52 on mean temperature and rainfall in Bali four months ahead into the development of dengue  
53  
54 early warning systems may assist the control of imported dengue and mitigate outbreaks of local  
55  
56 dengue in Australia. Our findings suggest that Australia's biosecurity program may need to look  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 beyond its own borders and contribute to dengue control overseas. This study also provides a  
5  
6 template and framework for future surveillance of travel-related infectious diseases globally.  
7  
8  
9

## 10 11 12 13 **Acknowledgements** 14

15  
16 We would like to thank Dr. Jian Cheng for his assistance in figure plotting.  
17  
18  
19  
20  
21

## 22 23 **References** 24

25  
26 Asma, S.; L, M.S.; Rachel, L.; Karin, L.; Bismi, J.; Christopher A.B.; Anton, Y.P. 2019.  
27

28         Spectrum of illness among returned Australian travellers from Bali, Indonesia: a 5-year  
29         retrospective observational study. Intern Med J. 49, 34-40.  
30  
31

32  
33 Australian Bureau of Statistics. 2018. 3401.0. - Overseas arrivals and departures, Australia, May  
34  
35 2018.  
36

37  
38         [http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/3401.0Explanatory%20Notes1Ma](http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/3401.0Explanatory%20Notes1May%202018?OpenDocument)  
39  
40         [y%202018?OpenDocument](http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/3401.0Explanatory%20Notes1May%202018?OpenDocument).  
41  
42

43  
44 Bangs, M.; Larasati, R.; Corwin, A.; Wuryadi, S. 2006. Climatic factors associated with  
45  
46 epidemic dengue in Palembang, Indonesia: implications of short-term meteorological  
47  
48 events on virus transmission. Southeast Asian J Trop Med Public Health. 37, 1103-1116.  
49

50  
51 Beebe, N.W.; Ambrose, L.; Hill, L.A.; Davis, J.B.; Hapgood, G.; Cooper, R.D.; Russell, R.C.;

52  
53 Ritchie, S.A.; Reimer, L.J.; Lobo, N.F.; Syafruddin, D.; van den Hurk, A.F. 2013.  
54

55  
56 Tracing the tiger: Population genetics provides valuable insights into the Aedes  
57  
58  
59  
60  
61

1  
2  
3  
4 (Stegomyia) albopictus invasion of the Australasian region. PLoS Negl Trop Dis. 7,  
5  
6 e2361.  
7

8  
9 Bhatt, S.; Gething, P.W.; Brady, O.J.; Messina, J.P.; Farlow, A.W.; Moyes, C.L.; Drake, J.M.;  
10  
11 Brownstein, J.S.; Hoen, A.G.; Sankoh, O.; Myers, M.F.; George, D.B.; Jaenisch, T.; Wint,  
12  
13 G.R.W.; Simmons, C.P.; Scott, T.W.; Farrar, J.J.; Hay, S.I. 2013. The global distribution  
14  
15 and burden of dengue. Nature. 496, 504-507.  
16  
17

18  
19 Butt, C. 2017. Dengue fever cases hit 20-year high in Australia. The Sydney Morning Herald  
20  
21 Sydney Fairfax Media.  
22

23  
24 Choe, Y.J.; Choe, S.A.; Cho, S.I. Importation of travel-related infectious diseases is increasing in  
25  
26 South Korea: An analysis of salmonellosis, shigellosis, malaria, and dengue surveillance  
27  
28 data. Travel Med Infect Dis 2017;19:22-27.  
29

30  
31 Chuang, T.W.; Chaves, L.F.; Chen, P.J. Effects of local and regional climatic fluctuations on  
32  
33 dengue outbreaks in southern Taiwan. PLOS ONE 2017;12:e0178698  
34

35  
36 David, W.; Judith, A.N.; Alyssa, T.P. 2012. Sources of dengue viruses imported into Queensland,  
37  
38 Australia, 2002–2010. Emerg Infect Dis. 18, 1850  
39

40  
41 Earnest, A.; Tan, S.B.; Wilder-Smith, A. Meteorological factors and El Niño Southern  
42  
43 Oscillation are independently associated with dengue infections. Epidemiol Infect  
44  
45 2011;140:1244-1251  
46

47  
48 Ebi, K.L.; Nealon, J. 2016. Dengue in a changing climate. Environ Res. 151, 115-123.  
49

50  
51 Ernst, T.; McCarthy, S.; Chidlow, G.; Luang-Suarkia, D.; Holmes, E.C.; Smith, D.W.; Imrie, A.  
52  
53 2015. Emergence of a new lineage of dengue virus type 2 identified in travelers entering  
54  
55 Western Australia from Indonesia, 2010-2012. PLoS Negl Trop Dis. 9, e0003442.  
56  
57  
58  
59  
60  
61

- 1  
2  
3  
4 Fang, L.Q.; Sun, Y.; Zhao, G.P.; Liu, L.J.; Jiang, Z.J.; Fan, Z.W.; Wang, J.X.; Ji, Y.; Ma, M.J.;  
5  
6 Teng, J.; Zhu, Y.; Yu, P.; Li, K.; Tian, Y.J.; Cao, W.C. 2018. Travel-related infections in  
7  
8 mainland China, 2014–16: an active surveillance study. *Lancet Public Health*. 3, e385-  
9  
10 e394.  
11  
12  
13  
14 Findlater, A.; Bogoch, I. 2018. Human mobility and the global spread of infectious diseases: a  
15  
16 focus on air travel. *Trends in Parasitology*. 34, 772-83.  
17  
18  
19 Furuya-Kanamori, L.; Liang, S.; Milinovich, G.; Soares Magalhaes, R.J.; Clements, A.C.A.; Hu,  
20  
21 W.; Brasil, P.; Frentiu, F.D.; Dunning, R.; Yakob, L. 2016. Co-distribution and co-  
22  
23 infection of chikungunya and dengue viruses. *BMC Infect Dis*. 16, 84.  
24  
25  
26 Gasparini, A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J Stat*  
27  
28 *Softw* 2011;43:1-20  
29  
30  
31 Gasparrini, A.; Armstrong, B. Reducing and meta-analysing estimates from distributed lag non-  
32  
33 linear models. *BMC Med Res Methodol* 2013;13:1  
34  
35  
36 Gasparini, A. 2011. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J*  
37  
38 *Stat Softw*. 43, 1-20.  
39  
40  
41 Goto, K.; Kumarendran, B.; Mettananda, S.; Gunasekara, D.; Fujii, Y.; Kaneko, S. Analysis of  
42  
43 effects of meteorological factors on dengue incidence in Sri Lanka using time series data.  
44  
45 *PLOS ONE* 2013;8:e63717.  
46  
47  
48 Hu, W.; Clements, A.; Williams, G.; Tong, S.; Mengersen, K. Spatial patterns and  
49  
50 socioecological drivers of dengue fever transmission in Queensland, Australia. *Environ*  
51  
52 *Health Perspect* 2012;120:260-266  
53  
54  
55 Huang, X.; Williams, G.; Clements, A.C.A.; Hu, W. Imported dengue cases, weather variation  
56  
57 and autochthonous dengue incidence in Cairns, Australia. *PLOS ONE* 2013;8:e81887  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Huang, X.; Yakob, L.; Devine, G.; Frentiu, F.D.; Fu, S.Y.; Hu, W. 2016. Dynamic  
5  
6 spatiotemporal trends of imported dengue fever in Australia. *Sci Rep.* 6, 30360.  
7  
8  
9 Iguchi, J.A.; Seposo, X.T.; Honda, Y. Meteorological factors affecting dengue incidence in  
10  
11 Davao, Philippines. *BMC Public Health* 2018;18:629  
12  
13  
14 Kesetyaningsih, T.; Andarini, S.; Sudarto; Pramoedyo, H. 2018. Determination of environmental  
15  
16 factors affecting dengue incidence in Selman District, Yogyakarta, Indonesia. *Afr J Infect*  
17  
18 *Dis.* 12, 13-25.  
19  
20  
21 Knope, K.; Committee, N.A.a.M.A.; Giele, C. 2013. Increasing notifications of dengue in  
22  
23 Australia related to overseas travel, 1991 to 2012. *Commun Dis Intell Q Rep.* 37, E55-  
24  
25 E59.  
26  
27  
28 Lee, J.S.; Carabali, M.; Lim, J.K.; Herrera, V.M.; Park, I.Y.; Villar, L.; Farlow, A. Early warning  
29  
30 signal for dengue outbreaks and identification of high risk areas for dengue fever in  
31  
32 Colombia using climate and non-climate datasets. *BMC Infect Dis* 2017;17:480-480  
33  
34  
35 Li, R.; Xu, L.; Bjørnstad, O.N.; Liu, K.; Song, T.; Chen, A.; Xu, B.; Liu, Q.; Stenseth, N.C.  
36  
37 Climate-driven variation in mosquito density predicts the spatiotemporal dynamics of  
38  
39 dengue. *Proc Natl Acad Sci U S A* 2019;116:3624-3629  
40  
41  
42  
43 Lowe, R.; Gasparrini, A.; Van Meerbeeck, C.J.; Lippi, C.A.; Mahon, R.; Trotman, A.R.; Rollock,  
44  
45 L.; Hinds, A.Q.J.; Ryan, S.J.; Stewart-Ibarra, A.M. Nonlinear and delayed impacts of  
46  
47 climate on dengue risk in Barbados: A modelling study. *PLOS Med* 2018;15:e1002613  
48  
49  
50 Mackenzie, J.; Brooy, J.; Hueston, L.; Cunningham, A. 1996. Dengue in Australia. *J Med*  
51  
52 *Microbiol.* 45, 159-161  
53  
54  
55 Masyeni, S.; Yohan, B.; Biomed, M.; Somia, I.K.A.; Myint, K.S.A.; Sasmono, R.T. 2018.  
56  
57 Dengue infection in international travellers visiting Bali, Indonesia. *J Travel Med.* 25, 1.  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 McKerr, C.; Lo, Y.C.; Edeghere, O.; Bracebridge, S. 2015. Evaluation of the national Notifiable  
5  
6 Diseases Surveillance System for dengue fever in Taiwan, 2010–2012. *PLoS Negl Trop*  
7  
8 *Dis.* 9, e0003639.  
9  
10  
11 Medical Society of Victoria. 1873. Local topics. *The Australian Medical Journal*:[https://digitised-](https://digitised-collections.unimelb.edu.au/bitstream/handle/11343/23137/267571_UDS2010779-2010209.pdf?sequence=2010775&isAllowed=y)  
12  
13 [collections.unimelb.edu.au/bitstream/handle/11343/23137/267571\\_UDS2010779-](https://digitised-collections.unimelb.edu.au/bitstream/handle/11343/23137/267571_UDS2010779-2010209.pdf?sequence=2010775&isAllowed=y)  
14  
15 [2010209.pdf?sequence=2010775&isAllowed=y](https://digitised-collections.unimelb.edu.au/bitstream/handle/11343/23137/267571_UDS2010779-2010209.pdf?sequence=2010775&isAllowed=y) (Accessed on 19th August 2018).  
16  
17  
18  
19 Morin, C.; Comrie, A.; Ernst, K. 2013. Climate and dengue transmission: evidence and  
20  
21 implications. *Environ Health Perspect.* 121, 1264-1272.  
22  
23  
24 Morin, C.W.; Semenza, J.C.; Trtanj, J.M.; Glass, G.E.; Boyer, C.; Ebi, K.L. Unexplored  
25  
26 opportunities: Use of climate- and weather-driven early warning systems to reduce the  
27  
28 burden of infectious diseases. *Curr Environ Health Rep* 2018;5:430-438  
29  
30  
31 Naish, S.; Dale, P.; Mackenzie, J.S.; McBride, J.; Mengersen, K.; Tong, S. Climate change and  
32  
33 dengue: a critical and systematic review of quantitative modelling approaches. *BMC*  
34  
35 *Infect Dis* 2014;14:167  
36  
37  
38 O'Neill, S. 2018. The use of Wolbachia by the World Mosquito Program to interrupt  
39  
40 transmission of *Aedes aegypti* transmitted viruses. *Adv Exp Med Biol.* 1062, 355-360  
41  
42  
43 O'Neill, S.; Ryan, P.; Turley, A.; Wilson, G.; Retzki, K.; Iturbe-Ormaetxe, I.; Dong, Y.; Kenny,  
44  
45 N.; Paton, C.; Ritchie, S.; Brown-Kenyon, J.; Stanford, D.; Wittmeier, N.; Anders, K.;  
46  
47 Simmons, C. 2018. Scaled deployment of Wolbachia to protect the community from  
48  
49 *Aedes* transmitted arboviruses. *Gates Open Res.* 2, 36.  
50  
51  
52  
53 Pham, H.V.; Doan, H.T.M.; Phan, T.T.T.; Minh, N.N.T. Ecological factors associated with  
54  
55 dengue fever in a Central Highlands province, Vietnam. *BMC Infect Dis* 2011;11:172-  
56  
57 172  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Racloz, V.; Ramsey, R.; Tong, S.; Hu, W. Surveillance of dengue fever virus: a review of  
5  
6 epidemiological models and early warning systems. *PLoS Negl Trop Dis* 2012;6:e1648-  
7  
8 e1648  
9  
10  
11 Russell, R.; Currie, B.; Lindsay, M.; Mackenzie, J.; Ritchie, S.; Whelan, P. 2009. Dengue and  
12  
13 climate change in Australia: predictions for the future should incorporate knowledge  
14  
15 from the past. *Med J Aust.* 190, 265-268.  
16  
17  
18 Sarfraz, M.S.; Tripathi, N.K.; Tipdecho, T.; Thongbu, T.; Kerdthong, P.; Souris, M. 2012.  
19  
20 Analyzing the spatio-temporal relationship between dengue vector larval density and  
21  
22 land-use using factor analysis and spatial ring mapping. *BMC Public Health.* 12, 853.  
23  
24  
25 Semenza, J.C.; Sudre, B.; Miniota, J.; Rossi, M.; Hu, W.; Kossowsky, D.; Suk, J.E.; Van Bortel,  
26  
27 W.; Khan, K. International dispersal of dengue through air travel: importation risk for  
28  
29 Europe. *PLoS Negl Trop Dis* 2014;8:e3278-e3278  
30  
31  
32  
33 Tai, A.Y.; McGuinness, S.L.; Robosa, R.; Turner, D.; Huang, G.K.L.; Leder, K.; Korman, T.M.;  
34  
35 Thevarajan, I.; Stewardson, A.J.; Padiglione, A.A.; Johnson, D.F. Management of dengue  
36  
37 in Australian travellers: a retrospective multicentre analysis. *Med J Aust* 2017;206:295-  
38  
39 300  
40  
41  
42  
43 Tian, H.; Sun, Z.; Faria, N.R.; Yang, J.; Cazelles, B.; Huang, S.; Xu, B.; Yang, Q.; Pybus, O.G.;  
44  
45 Xu, B. Increasing airline travel may facilitate co-circulation of multiple dengue virus  
46  
47 serotypes in Asia. *PLoS Negl Trop Dis* 2017;11:e0005694-e0005694  
48  
49  
50  
51 Tosepu, R.; Tantrakarnapa, K.; Nakhapakorn, K.; Worakhunpiset, S. 2018. Climate variability  
52  
53 and dengue hemorrhagic fever in Southeast Sulawesi Province, Indonesia. *Environ Sci*  
54  
55 *Pollut Res Int.* 25, 14944-14952.  
56  
57  
58 UNTWO. 2017. Compendium of Tourism Statistics, Data 2011 - 2015.  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 Vainio, K.; Noraas, S.; Holmberg, M.; Fremstad, H.; Wahlstrøm, M.; Ånestad, G.; Dudman, S.  
5  
6 2010. Fatal and mild primary dengue virus infections imported to Norway from Africa  
7  
8 and south-east Asia, 2008-2010. *Eurosurveillance*. 15, 19666.  
9

10  
11 WHO Western Pacific Region. 2014. Dengue fact sheet. Available at  
12  
13 [http://www.wpro.who.int/mediacentre/factsheets/fs\\_09032012\\_Dengue/en/](http://www.wpro.who.int/mediacentre/factsheets/fs_09032012_Dengue/en/) (accessed 25  
14  
15 July 2018).  
16  
17

18  
19 Wu, X.; Lang, L.; Ma, W.; Song, T.; Kang, M.; He, J.; Zhang, Y.; Lu, L.; Lin, H.; Ling, L. 2018.  
20  
21 Non-linear effects of mean temperature and relative humidity on dengue incidence in  
22  
23 Guangzhou, China. *Sci Total Environ*. 628-629, 766-771.  
24

25  
26 Xiang, J.; Hansen, A.; Liu, Q.; Liu, X.; Tong, M.X.; Sun, Y.; Cameron, S.; Hanson-Easey, S.;  
27  
28 Han, G.-S.; Williams, C.; Weinstein, P.; Bi, P. 2017. Association between dengue fever  
29  
30 incidence and meteorological factors in Guangzhou, China, 2005–2014. *Environ Res*. 153,  
31  
32 17-26.  
33  
34

35  
36 Xu, L.; Stige, L.C.; Chan, K.-S.; Zhou, J.; Yang, J.; Sang, S.; Wang, M.; Yang, Z.; Yan, Z.; Jiang,  
37  
38 T.; Lu, L.; Yue, Y.; Liu, X.; Lin, H.; Xu, J.; Liu, Q.; Stenseth, N.C. 2017. Climate  
39  
40 variation drives dengue dynamics. *Proc Natl Acad Sci U S A*. 114, 113-118.  
41  
42



43  
44 Yi, L.; Xu, X.; Ge, W.; Xue, H.; Li, J.; Li, D.; Wang, C.; Wu, H.; Liu, X.; Zheng, D.; Chen, Z.;  
45  
46 Liu, Q.; Bi, P.; Li, J. The impact of climate variability on infectious disease transmission  
47  
48 in China: Current knowledge and further directions. *Environ Res* 2019;173:255-261  
49

50  
51 Yuan, B.; Nishiura, H. Estimating the actual importation risk of dengue virus infection among  
52  
53 Japanese travelers. *PLOS ONE* 2018;13:e0198734  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure

[Click here to download high resolution image](#)


**Figure 1.** Associations of mean temperature and rainfall with the occurrence of dengue cases in Bali in the optimal model



Figure

[Click here to download high resolution image](#)


**Figure 2A.** Relationship between dengue in Bali and imported dengue in Australia (excluding Queensland) at lag one month




Figure

[Click here to download high resolution image](#)

**Figure 2B.** Relationship between dengue in Jakarta and imported dengue in Australia (excluding Queensland) at lag one month




**Figure 3.** Relationship between number of Australians returning from Indonesia and imported dengue in Australia (excluding Queensland)



Figure

[Click here to download high resolution image](#)

**Figure 4.** Relationship between dengue in Bali and imported dengue in Australia (excluding Queensland) at lag one month after including the number of travelers in log scale as an offset in the model






Figure

[Click here to download high resolution image](#)


**Figure 5.** Relationship between dengue in Bali districts and imported dengue in Australia (excluding Queensland) at lag one month after including the number of travelers in log scale as an offset in the model



Figure

[Click here to download high resolution image](#)

**Figure 6A.** Relationship between mean temperature in Bali and imported dengue in Australia (excluding Queensland)






# Figure

[Click here to download high resolution image](#)

**Figure 6B.** Relationship between rainfall in Bali and imported dengue in Australia (excluding Queensland)



**Supplementary Material**

[Click here to download Supplementary Material: Supporting information.docx](#)

**Supplementary Material**

[Click here to download Supplementary Material: Table S1.docx](#)

**Supplementary Material**

[Click here to download Supplementary Material: Table S2.docx](#)

**Supplementary Material**

[Click here to download Supplementary Material: Table S3.docx](#)

**Supplementary Material**

[Click here to download Supplementary Material: Table S4.docx](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S1.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S2.jpg](#)



**Supplementary Material**

[Click here to download Supplementary Material: Figure S3A.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S3B.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S4.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S5A.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S5B.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S5C.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S6.jpg](#)

**Supplementary Material**

[Click here to download Supplementary Material: Figure S7.jpg](#)