

Sains Malaysiana 48(3)(2019): 607–612
<http://dx.doi.org/10.17576/jsm-2019-4803-13>

Muscle Size and Strength Benefits of Functional Electrical Stimulation-Evoked Cycling Dosage in Spinal Cord Injury: A Narrative Review

(Faedah Dos Rangsangan-Elektrik Berfungsi Berbasikal terhadap Saiz dan Kekuatan Otot pada Kecederaan Saraf Tunjang: Suatu Kajian Naratif)

NURHAIDA ROSLEY, NUR AZAH HAMZAID, NAZIRAH HASNAN, GLEN M DAVIS & HAIDZIR MANAF*

ABSTRACT

Loss of sensory motor function is one of the main causes of physical and activity limitations among individual with spinal cord injury (SCI). SCI may lead to muscle paralysis, weakness and disused muscle atrophy. Evidences have shown electrical stimulation and strengthening exercise might improve lower limb muscle strength and size among individual with SCI. Functional electrical stimulation (FES) evoked cycling is one of the methods that can elicit leg muscle contractions in order to produce a cycling motion and promote the integrity of the involved muscles. Therefore, this review is to synthesize the scientific literature regarding the effects of multiple dosages of FES-evoked lower limb cycling on muscle properties. A systematic literature search from 1946 to 2016 was performed. From over 1,139 articles retrieved from the database, about 31 potentially relevant articles were retained for possible inclusion. However, only 10 articles out of 31 articles fulfilled the inclusion criteria. Although the available evidence is compelling, there is insufficient quantity and quality evidence to draw conclusions regarding the specific parameter of FES-CE that may optimally increase muscle strength, mass, and circumference. However, it can be safely concluded that an effective training session would spend for 45-60 min, 3 times a week for at least 4 weeks to see changes in muscle size and strength.

Keywords: Cycling exercise; functional electrical stimulation; muscle strength; spinal cord injury

ABSTRAK

Kehilangan fungsi deria motor adalah salah satu punca kepada pembatasan fizikal dan aktiviti dalam kalangan individu yang mengalami kecederaan saraf tunjang. Kecederaan saraf tunjang boleh mengakibatkan kelumpuhan otot, kelemahan dan ketidakgunaan otot atau atrofi. Bukti menunjukkan stimulasi elektrik dan senaman kekuatan otot boleh memperbaiki saiz dan kekuatan otot bahagian kaki dalam kalangan individu yang mengalami kecederaan saraf tunjang. Sistem berbasikal rangsangan FES adalah salah satu metod yang boleh merangsang kontraksi otot kaki untuk menghasilkan pergerakan berbasikal dan meningkatkan integriti otot yang terlibat. Oleh itu, kajian ini adalah bertujuan untuk mensintesis kepustakaan saintifik untuk mengkaji kesan pelbagai dos rangsangan FES untuk anggota bawah berbasikal pada sifat otot. Satu carian kepustakaan secara sistematik telah dijalankan dari 1946 hingga 2016. Daripada 1,139 artikel yang didapati daripada pangkalan data, 31 artikel yang berpotensi telah dikekalkan mengikut kebarangkalian inklusi. Walau bagaimanapun, hanya 10 artikel daripada 31 artikel memenuhi kriteria rangkuman. Walaupun bukti yang didapati sangat menarik perhatian, namun kuantiti dan kualitinya tidak mencukupi untuk dijadikan kesimpulan berkaitan dengan parameter khusus FES-CE yang mampu meningkatkan kekuatan, jisim dan ukur lilit otot secara optimum. Namun, ia selamat untuk disimpulkan bahawa keberkesanan suatu sesi latihan mengambil masa 45-60 minit, 3 kali seminggu untuk sekurang-kurangnya 4 minggu untuk melihat perubahan saiz dan kekuatan otot.

Kata kunci: Kecederaan saraf tunjang; kekuatan otot; rangsangan elektrik berfungsi; senaman berbasikal

INTRODUCTION

Spinal cord injury (SCI) is one of the leading causes of physical disability worldwide. In the USA, approximately 17,000 new cases of SCI were reported each year due to road traffic accident, falls, and sports (National Spinal Cord Injury Database 2015). It can be pronounced that the consequences after SCI is usually a profound reduction in the muscle mass, strength, and muscle volume (Dionyssiotis et al. 2015). Thus, as a result of this muscle integrity degeneration, it could lead to lower limb muscle weakness and muscle atrophy. These can affect their quality

of life in which long-term recovery and rehabilitation process is required (LiVecchi 2011; Westcott & Rosa 2010).

It has been widely established that exercise and training programs for individual with SCI could enhance their muscle power function and prevent muscle atrophy (American College of Sports Medicine 2010). Thus, a suitable exercise regime can help in achieving these goals and reduce the risk of secondary health problems accompanying with physical inactivity. One of the technique is by applying the functional electrical stimulation (FES). This device will assist exercise through the application of a low-level

electrical stimulation current on muscle motor control to evoke muscle contraction (Lynch et al. 2008; Peckham & Knutson 2005). There have been so many method with FES, however, FES evoked cycling exercise (FES-CE) has been widely used in clinical settings for aerobic and strength training applications (Estigoni et al. 2011; Fornusek et al. 2013; Kuhn et al. 2014; Thrasher et al. 2013). Technically, FES-CE is a computer-controlled electrical stimulation that elicits leg muscle contractions in order to produce a cycling motion and promote the integrity of the involved muscles (Ragnarsson 2008). Based on a study by Hamzaid and Davis (2009), the FES-CE could increase muscle size and volume, and prevent the alteration of muscle morphology in individuals with lower motor neuron lesions SCI (Hamzaid & Davis 2009). Therefore, the ultimate goal of this intervention is to provide therapeutic gains for persons with SCI (Estigoni et al. 2011; Fornusek & Davis 2004; Hasnan et al. 2013)

Despite numerous studies of FES-CE effect on muscle size and muscle strength among individual with SCI, no evidence concerning about its appropriate dosage of training. In this narrative review, inclusive review of obtainable literature was provided to verify the effects of FES-CE on muscle size and muscle strengths among individual with SCI. By providing the quality of evidence and dosage for this kind of training, it may help the medical practitioners to decide on its proper application. Thus, the aim of this review was to synthesize the scientific literature to explore the multiple dosages of FES-CE that is adequate to improve muscle properties and characteristics including muscle mass, volume, and strength for individuals with SCI.

METHODS: STUDY SELECTION

A systematic literature search was conducted by using electronic databases: ProQuest, Science Direct, MEDLINE between 1946 and October 2016; Ovid MEDLINE Daily Update; and Web of Science between 1980 and 2016. For a wider coverage, the literature searches also included manual searches of clinical rehabilitation specific journals including Clinical Rehabilitation, Spinal Cord, Physiotherapy, Physical therapy and multidisciplinary journals for examples, BMJ, Lancet, and JAMA. The search strategy for reviews on our topic was also carried out using the Cochrane Library and Cochrane Database for Systematic Reviews. The literature search was, however, limited to human research studies, based on any language if the translation of English abstract provides a clear description of the study and the outcome measures. All systematic literature findings were explored by the author based on certain inclusion criteria until it reaches 'saturation point.'

The inclusion criteria for this review was that the studies must involve the following: functional electrical stimulation (FES); or functional neuromuscular stimulation (FNS) evoked cycling; lower limbs cycling; outcome should be measured on the effects of the cycling

exercise on improvement or otherwise of muscle strength, mass, and volume; and in persons with spinal cord injury. In addition, studies that involved a combination of FES-evoked upper limb and lower limb exercise were included, if they provide lower limbs muscle strength as one of the outcome measures. Keywords used in this review were: FES, FNS, functional electrical stimulation, functional neuromuscular stimulation, electrical stimulation, functional stimulation, neuromuscular stimulation, neurostimulation; training, exercise, cycling; SCI, spinal cord survivors, paraplegic, paraplegia, quadriplegic, tetraplegia, tetraplegic, paralyzed, paralysis; subjects, participants, patients, human, persons; muscle strength, muscle atrophy, muscle mass, muscle volume, muscle circumference and muscle properties. The title of all the potentially eligible studies were scanned manually using Google Scholar search engine. On the first round of search, 1,139 experimental studies were discovered. Thus, the second round of search included randomized controlled trial (RCT), quasi-experimental research (randomized only or controlled only), case-controlled study, observation studies and crossover designs. As highlighted earlier only studies that recruited persons with SCI for the experiment and control group were retained. Studies that recruited able-bodied participants as the control were excluded. This was due to the knowledge that the inference or conclusion from such studies may not be sufficient to verify the proposed hypothesis (Hamzaid & Davis 2009). In the present review, the RCT and quasi-RCT retained were meant to provide a strong evidence whereas the other studies were used as supporting evidence for the hypothesis of the study.

RESULTS

The articles from RCT, quasi-RCT, nonrandomized or controlled studies and cross-sectional or crossover designs were retained for this review as they provide the scientific evidence regarding the FES-CE. The article search was conducted from February 2016 until October 2016. From over 1,139 articles retrieved from the database, about 31 potentially relevant articles were retained for possible inclusion. However, only 10 articles out of 31 articles fulfilled the inclusion criteria. All the 10 studies reported muscle size and strength benefits effect following FES-CE in individuals with SCI (Table 1). All studies had samples sizes ranging from 8-45 participants.

Muscle size has generally been measured as body lean mass by using bioelectrical impedance analysis and Lunar DPX x-ray to determine lesser fat or bigger muscle after FES-CE training (Baldi et al. 1998; Liu et al. 2007). Muscle circumference usually measured in the supine position using a tape measure (Fornusek et al. 2013; Kuhn et al. 2014; Liu et al. 2007). The measurements were usually reported before and after the FES-CE training based on the anatomical landmarks chosen by researchers. Significant improvement of muscle peak torque has been reported in individuals with SCI after FES-CE training

TABLE 1. Studies reported muscle size and strength benefits effect following FES-evoked lower limb cycling exercise in individuals with SCI

References	Sample	Duration	Intervention	Main result
Johnston et al. (2015)	N= 17 Complete SCI	6 months	FES cycling 1-hour × 3 times per week Low cadence vs. high cadence	Muscle volume Low cadence group increased by 19%, high cadence group increase by 10%
Sköld et al. (2002)	N= 15 Complete SCI	6 months	FES cycling 30 min × 3 times per week Freq= 30 Hz PW= 350 µsec Stim= up to 130 mA Pedal cadence= 35 rpm	Muscle volume Group trained with FES increased 1300 cm ³ ($p < 0.001$) than those in the control group
Kuhn et al. (2014)	N=30 Complete SCI vs. Incomplete SCI	4 weeks	FES-CE 20 min × 2 times per week Freq= 30 Hz PW= 250 µsec Stim= 10- 130 mA Pedal cadence= 15-55 rpm	Circumferences Greater improvement in incomplete group ($p < 0.05$) Manual muscle testing Those in incomplete group reported 43.5% increase ($p < 0.001$) in quadriceps femoris muscle and 25% ($p < 0.001$) in gluteus maximus muscle
Thrasher et al. (2013)	N= 11 Complete SCI vs. incomplete SCI	13 weeks	FES-LCE training 1 h × 3 times per week, 40 sessions Freq= 30 Hz PW= 250 µsec Stim= up to 140 mA Pedal cadence= 45 rpm	Peak torque Both groups had improved knee extension torque 3.8 to 16.9 Nm ($p = 0.006$). However, those in incomplete group shown greater improvement
Fornusek et al. (2013)	N=8 Complete and Incomplete SCI	6 weeks	FES-CE 30 min × 3 times per week 1 leg on low cadence 10 rpm and the other leg on high cadence 50 rpm Freq= 35 Hz PW= 250 µsec Stim= 40- 140 mA	Thigh girth Both groups reported greater improvement on low cadence Quadriceps peak torque Quadriceps peak torque has increased 87% in low cadence
Sadowsky et al. (2013)	N= 45 Complete and Incomplete SCI	29.1 month	FES-CE 50-60 min × 3 times per week Freq= 100 Hz PW= 500 µsec Stim= up to 140 mA Pedal cadence= 50 rpm	Muscle volume Those in FES group had significantly higher anterior (36%; 96.3 cc; $p \leq 0.001$) and posterior (30%; 63.9 cc; $p = 0.005$) thigh compartment muscles than those in the control group Muscle strength trained muscles of the FES group was significantly greater (quadriceps, $p = 0.006$; hamstrings, $p = 0.011$) than in controls
Duffell et al. (2008)	N= 11 Complete SCI	1 year	FES cycling 1 h × 5 times/week Freq= 50 Hz if rapid fatigue 10 Pedal cadence= 45-55 rpm	Quadriceps Maximal Torque Increased progressively and significantly throughout training ($p = 0.012$). It had increased by 399% and 673% after 3 and 12 months
Liu et al. (2007)	N=18 Complete and Incomplete SCI	8 weeks	FES-CE 30 min × 3 times per week Freq= 30 Hz PW= 300 µsec Stim= 10-130 mA Pedal cadence= 45 rpm	Body lean mass Mild increase after 8 weeks training ($p=0.03$) Limb girth Significant increase in both upper and lower limb girth after 4 and 8 weeks of training ($p < 0.05$) Peak torque Shown improvement only after 8 weeks training ($p < 0.05$)
Demchak et al. (2005)	N= 10 Complete SCI	13 weeks	FES-CE 30 min 3x per week Freq= 35 Hz Pedal cadence= 35 rpm	Cross-sectional area (muscle biopsy) Group trained with FES-CE increased CSAf 171% greater than the CSAf in the control group ($p = 0.05$)
Baldi et al. (1998)	N=26 Complete SCI	3 month + 6 month	FES-CE 30 min × 3 times per week, FES-IC 1 h × 5 times per week Freq= 60 Hz PW= 375 µsec Stim= up to 140 mA Pedal cadence= 35-50 rpm	Lean body mass FES-CE showed a significant hypertrophy after 6 months on both lower limb and gluteal lean body mass in comparison to FES alone

FES-CE = Functional Electrical Stimulation-Cycling Ergometer; FES-LCE= Functional electrical Stimulation Leg Cycling Ergometer; FES-IC= Functional Electrical Stimulation Isometric Contraction, N= Number of participants, Freq= Frequency, PW= Pulse Width, Stim= Stimulation

(Duffell et al. 2008; Fornusek et al. 2013; Liu et al. 2007; Sadowsky et al. 2013; Thrasher et al. 2006). However, individuals with incomplete SCI has been found to have a greater improvement than those with complete SCI (Fornusek et al. 2013; Sadowsky et al. 2013; Thrasher et al. 2013). Therefore, available evidence showed that the strength gains during FES-CE training were positively related to the spared muscle function after SCI trauma. This may suggest that SCI individuals with higher residual muscle strength might achieve better results from the cycling training.

DISCUSSION

This review yielded only 10 articles that emphasize the effect of FES-evoked cycling exercise on muscle size and muscle strength among individual with SCI. It has been proven that this type of exercise is highly relevant modality for individual with chronic SCI as it has a significant impact on their capability to perform activities of daily living. Additionally, it has been identified that the FES-CE does not only improve muscle size, it also promotes the quality of life of incomplete SCI individuals in particular.

EFFECTS OF FES-CE ON MUSCLE SIZE

Individuals who were trained with FES-CE had shown significant improvement in muscle mass and muscle girth after 4 weeks (Kuhn et al. 2014). However, it has been identified that the training should start within the first 3 months after a SCI to ensure the efficacy of FES-CE training (Baldi et al. 1998). Baldi et al. (1998) have shown that FES-CE could prevent the progression of muscle mass reduction after 6 months of training. Paralysis and episode of sedentary lifestyle after chronic SCI could lead to elevation of fat mass and reduction of lean body mass percentage (Giangregorio et al. 2005). Notably, individuals with acute SCI only have 60-65% lean mass than their able-bodied counterparts and their fat gain can be up to 100-113% of their able-bodied counterparts (Giangregorio et al. 2005). Therefore, this may lead to a serious long-term health complication that may compromise with their quality of life (Hicks et al. 2011).

Nevertheless, the significant changes in leg lean mass following FES-CE training for incomplete SCI has been reported to be greater than in complete SCI individuals even in a short-term (8 weeks) cycling training (Liu et al. 2007). Hypertrophy of stimulated muscle has also been observed after both short and long-duration FES-CE protocol (Fornusek et al. 2013; Kuhn et al. 2014). Additionally, although a non-significant general increase of body lean mass in completely paralyzed individuals has been reported, such increase has a positive effect in preventing muscle atrophy of stimulated muscles (Baldi et al. 1998). Based on the literature reviewed, we can safely infer that FES cycling with 3 days per week training for more than 4 weeks could prevent the loss of muscle mass (Fornusek et al. 2013; Kuhn et al. 2014).

EFFECTS OF FES-CE ON MUSCLE STRENGTH

Evidence suggested that in order to achieve greater dramatic effects of muscle strength, the stimulation parameters used in the FES-CE training must be able to increase fatigue-resistance of muscles (Fornusek et al. 2013; Johnston et al. 2015). Low-cadence (10 rpm) but not high-cadence (i.e. 50 rpm) cycling has been shown to promote increased fatigue-resistance of the quadriceps muscle (Fornusek et al. 2013). This result was supported by earlier recommendation that the health effects from FES-CE should be maximized for SCI survivors (Duffell et al. 2008; Liu et al. 2007). As evident from the reviewed literature, most FES-CE cycle ergometers are designed to operate at a cadence of 50 rpm. However, fatigue develops more slowly during FES-CE at lower pedal cadences, allowing greater muscle forces to be maintained (Johnston et al. 2015). Therefore, continuous stimulation in muscle's normal physiological frequency range (30-35 Hz) (Demchak et al. 2005; Duffell et al. 2008; Sadowsky et al. 2013) and pulse width (250-300 μ sec) (Baldi et al. 1998; Johnston et al. 2015; Liu et al. 2007) may offer a better electrical stimulus effect that could promote adaptation of a muscle to be more fatigue resistant. In a total of 10 studies, not all studies stated the resistance used while cycling with FES. However, it seemed that by increasing the resistance every \sim 3W- \sim 6W if participants able to maintain rotation per minutes of the cadence by three consecutive 30 min training sessions it could give more impact to the strength of the muscle (Demchak et al. 2005; Liu et al. 2007; Thrasher et al. 2013).

CONCLUSION

Although the available evidence is compelling, there is insufficient quantity and quality evidence to draw conclusions regarding the specific parameter of FES-CE that may optimally increase muscle strength, mass, and circumference. Nevertheless, based on Table 1, it can be safely concluded that an effective training session would spend between 45 and 60 min, 3 times a week for at least 4 weeks to see changes in muscle size and strength. The FES stimulation parameters of 250-300 microseconds pulse width, current amplitude 40mA-140mA, 30-35Hz frequency are optimally administered for each participant. Studies on leg FES-cycling have only been carried out in persons with complete SCI or those with a combination of complete and incomplete SCI. However, a lesser attention has been paid to the effects of FES-cycling on persons with incomplete SCI. Although FES-cycling in persons with incomplete SCI may give a greater improvement in muscle strength, the affected population may not be able to tolerate electrical stimulation of a similar intensity as administered to complete SCI survivors. Thus, further quality research is needed to examine the most suitable parameters that can be used to maximize its effectiveness. For clinical rehabilitation applications, a standardized FES-evoked cycling protocol will be guiding especially for clinicians

and other allied health professional administering FES-cycling for strength training.

ACKNOWLEDGEMENTS

The authors would like to thank Dr Morufu Olusola Ibitoye, for assisting with the preparation of the manuscript. We also would like to thank Universiti Malaya for the High Impact Research (HIR) Grant (Ref No: UM.C/HIR/MOHE/ENG/39) and Fundamental Research Grant Scheme (FRGS) awarded by the Ministry of Higher Education, Malaysia (Ref No: 600-RMI/FRGS 5/3 (0008/2016)) for funding the manuscript process.

REFERENCES

- American College of Sports Medicine (ACSM). 2010. *ACSM's Resources for Clinical Exercise Physiology: Musculoskeletal, Neuromuscular, Neoplastic, Immunologic and Hematologic Conditions (ACSM's Resources for the Clinical Exercise Physiology)*. 2nd edition. Philadelphia: Lippincott, Williams and Wilkins.
- Baldi, J.C., Jackson, R.D., Moraille, R. & Mysiw, W.J. 1998. Muscle atrophy is prevented in patients with acute spinal cord injury using functional electrical stimulation. *Spinal Cord* 36: 463-469.
- Demchak, T.J., Linderman, J.K., Mysiw, W.J., Jackson, R., Suun, J. & Devor, S.T. 2005. Effects of functional electric stimulation cycle ergometry training on lower limb musculature in acute SCI individuals. *Journal of Sports Science and Medicine* 4(3): 263-271.
- Dionyssiotsis, Y., Stathopoulos, K., Trovas, G., Papaioannou, N., Skarantavos, G. & Papagelopoulos, P. 2015. Impact on bone and muscle area after spinal cord injury. *BoneKEY Reports* 4(1): 1-8.
- Duffell, L.D., Donaldson, N.D.N., Perkins, T.I.M.A. & Ms, C. 2008. Long-term intensive electrically stimulated cycling by spinal cord-injured people: Effect on muscle properties and their relation to power output. *Muscle Nerve* 38(4): 1304-1311.
- Estigoni, E.H., Fornusek, C., Smith, R.M. & Davis, G.M. 2011. Evoked EMG and muscle fatigue during isokinetic FES-cycling in individuals with SCI. *Neuromodulation: Journal of the International Neuromodulation Society* 14(4): 349-355.
- Fornusek, C. & Davis, G. 2004. Maximizing muscle force via low-cadence functional electrical stimulation cycling. *Journal of Rehabilitation Medicine* 36(5): 232-237.
- Fornusek, C., Davis, G.M. & Russold, M.F. 2013. Pilot study of the effect of low-cadence functional electrical stimulation cycling after spinal cord injury on thigh girth and strength. *Archives of Physical Medicine and Rehabilitation* 94(5): 990-993.
- Gianguregorio, L.M., Hicks, A.L., Webber, C.E., Phillips, S.M., Craven, B.C., Bugaresti, J.M. & McCartney, N. 2005. Body weight supported treadmill training in acute spinal cord injury: Impact on muscle and bone. *Spinal Cord: The Official Journal of the International Medical Society of Paraplegia* 43(11): 649-657.
- Hamzaid, N.A. & Davis, G.M. 2009. Health and fitness benefits of functional electrical stimulation-evoked leg exercise for spinal cord-injured individuals. *Topics in Spinal Cord Injury Rehabilitation* 14(4): 88-121.
- Hasnan, N., Ektas, N., Tanhoffer, A.I.P., Tanhoffer, R., Fornusek, C., Middleton, J.W., Husain, R. & Davis, G.M. 2013. Exercise responses during functional electrical stimulation cycling in individuals with spinal cord injury. *Medicine and Science in Sports and Exercise* 45(6): 1131-1138.
- Hicks, A.L., Ginis, K.A.M., Pelletier, C.A., Ditor, D.S., Foulon, B. & Wolfe, D.L. 2011. The effects of exercise training on physical capacity, strength, body composition and functional performance among adults with spinal cord injury: A systematic review. *Spinal Cord*. 49(11): 1103-1127.
- Johnston, T.E., Schmidt-Read, M., Marino, R., Oleson, C., Leiby, B. & Modlesky, C. 2015. Musculoskeletal effects of two functional electrical stimulation cycling paradigms for people with spinal cord injury. *Archives of Physical Medicine and Rehabilitation* 97(9): 1413-1422.
- Kuhn, D., Leichtfried, V. & Schobersberger, W. 2014. Four weeks of functional electrical stimulated cycling after spinal cord injury: A clinical cohort study. *International Journal of Rehabilitation Research* 37(3): 243-250.
- Liu, C., Chen, S., Chen, C., Chen, T. & Chen, J.J. 2007. Effects of functional electrical stimulation on peak torque and body composition in patient with spinal cord injury. *The Kaohsiung Journal of Medical Sciences* 23(5): 232-240.
- LiVecchi, M.A. 2011. Spinal cord injury. *Continuum: Lifelong Learning Neurology* 17(3): 568-583.
- Lynch, C.L., Popovic, M.R. & Rushton, D. 2008. Functional electrical stimulation. *IEEE Control Systems Magazine* 28(2): 40-50.
- National Spinal Cord Injury Database. 2015. www.nscisc.uab.edu/nscisc-database.aspx.
- Peckham, P.H. & Knutson, J.S. 2005. Functional electrical stimulation for neuromuscular applications. *Annual Review of Biomedical Engineering* 7(1): 327-360.
- Ragnarsson, K.T. 2008. Functional electrical stimulation after spinal cord injury: Current use, therapeutic effects and future directions. *Spinal Cord* 46(4): 255-274.
- Sadowsky, C.L., Hammond, E.R., Strohl, A.B., Commean, K., Eby, S.A., Damiano, D.L., Wingert, J.R., Bae, K.T. & McDonald, J.W. 2013. Lower extremity functional electrical stimulation cycling promotes physical and functional recovery in chronic spinal cord injury. *J. Spinal Cord. Med.* 36(6): 623-631.
- Sköld, C., Lönn, L., Harms-Ringdahl, K., Hultling, C., Levi, R., Nash, M. & Seiger, Å. 2002. Effects of functional electrical stimulation training for six months on body composition and spasticity in motor complete tetraplegic spinal cord-injured individuals. *Journal of Rehabilitation Medicine* 34(1): 25-32.
- Thrasher, T.A., Ward, J.S. & Fisher, S. 2013. Strength and endurance adaptations to functional electrical stimulation leg cycle ergometry in spinal cord injury. *NeuroRehabilitation* 33(1): 133-138.
- Thrasher, T.A., Flett, H.M. & Popovic, M.R. 2006. Gait training regimen for incomplete spinal cord injury using functional electrical stimulation. *Spinal Cord* 44(6): 357-361.
- Westcott, W.L. & Rosa, S.A. 2010. Spinal cord injury. *Strength and Conditioning Journal* 32(6): 16-18.

Nurhaida Rosley & Haidzir Manaf*
Centre of Physiotherapy
Faculty of Health Sciences
Universiti Teknologi MARA (Puncak Alam Campus)

42300 Puncak Alam, Selangor Darul Ehsan
Malaysia

Nur Azah Hamzaid
Department of Biomedical Engineering
Faculty of Engineering, University Malaya
50603 Kuala Lumpur, Federal Territory
Malaysia

Nazirah Hasnan
Department of Rehabilitation Medicine
Faculty of Medicine, University Malaya
50603 Kuala Lumpur, Federal Territory
Malaysia

Glen M Davis
Department of Clinical Exercise and Rehabilitation Unit
Faculty of Health Science, University of Sydney
Camperdown NSW 2006
Australia

*Corresponding author; email: haidzir5894@uitm.edu.my

Received: 19 July 2017
Accepted: 9 January 2019