

University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

West Central Research and Extension Center, North
Platte

Agricultural Research Division of IANR

2018

Effects of red clover isoflavones on tall fescue seed fermentation and microbial populations *in vitro*

Emily A. Melchior
University of Tennessee, Knoxville

Jason K. Smith
University of Tennessee, Knoxville, Jason.Smith@utk.edu


Liesel G. Schneider
University of Tennessee, Knoxville, lschneider@utk.edu

Travis Mulliniks
West Central Research and Extension Center, University of Nebraska-Lincoln, North Platte, travis.mulliniks@unl.edu

Gary E. Bates
University of Tennessee, Knoxville, gbates@utk.edu

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/westcentresext>

 Part of the [Agriculture Commons](#), [Ecology and Evolutionary Biology Commons](#), and the [Plant Sciences Commons](#)

Melchior, Emily A.; Smith, Jason K.; Schneider, Liesel G.; Mulliniks, Travis; Bates, Gary E.; McFarlane, Zachary D.; Flythe, Michael D.; Klotz, James L.; Goodman, Jack P.; Ji, Huihua; and Meyer, Phillip R., "Effects of red clover isoflavones on tall fescue seed fermentation and microbial populations *in vitro*" (2018). *West Central Research and Extension Center, North Platte*. 109.
<https://digitalcommons.unl.edu/westcentresext/109>

This Article is brought to you for free and open access by the Agricultural Research Division of IANR at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in West Central Research and Extension Center, North Platte by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, Travis Mulliniks, Gary E. Bates, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, and Phillip R. Meyer

RESEARCH ARTICLE

Effects of red clover isoflavones on tall fescue seed fermentation and microbial populations *in vitro*


Emily A. Melchior¹, Jason K. Smith¹, Liesel G. Schneider¹, J. Travis Mulliniks^{1ma}, Gary E. Bates², Zachary D. McFarlane^{1mb}, Michael D. Flythe³, James L. Klotz³, Jack P. Goodman⁴, Huihua Ji⁵, Phillip R. Myer^{1*}

1 Department of Animal Sciences, University of Tennessee, Knoxville, TN, United States of America, **2** Department of Plant Sciences, University of Tennessee, Knoxville, TN, United States of America, **3** USDA-ARS, Forage-Animal Production Research Unit, Lexington, KY, United States of America, **4** Department of Plant and Soil Sciences, University of Kentucky, Lexington, KY, United States of America, **5** College of Agriculture, Food and Environment, University of Kentucky, Lexington, KY, United States of America

^{ma} Current address: West Central Research and Extension Center, University of Nebraska-Lincoln, North Platte, United States of America

^{mb} Current address: California Polytechnic State University Department of Animal Sciences, San Luis Obispo, United States of America

* pmyer@utk.edu



OPEN ACCESS

Citation: Melchior EA, Smith JK, Schneider LG, Mulliniks JT, Bates GE, McFarlane ZD, et al. (2018) Effects of red clover isoflavones on tall fescue seed fermentation and microbial populations *in vitro*. PLoS ONE 13(10): e0201866. <https://doi.org/10.1371/journal.pone.0201866>

Editor: Alex V. Chaves, The University of Sydney, AUSTRALIA

Received: June 18, 2018

Accepted: October 7, 2018

Published: October 18, 2018

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: Sequence files are available from the NCBI Sequence Read Archive (SRA Accession SRP150716). Additional descriptive information is associated with NCBI BioProject PRJNA476124.

Funding: This work was supported by a grant from the UT Center of Excellence in Livestock Diseases and Human Health (COE). The authors acknowledge PharmaCare Inc. for providing the Promensil® product, as well as the USDA-NIFA Hatch/Multistate Project W2010- TEN00493—

Abstract

Negative impacts of endophyte-infected *Lolium arundinaceum* (*Darbyshire*) (tall fescue) are responsible for over \$2 billion in losses to livestock producers annually. While the influence of endophyte-infected tall fescue has been studied for decades, mitigation methods have not been clearly elucidated. Isoflavones found in *Trifolium pratense* (red clover) have been the subject of recent research regarding tall fescue toxicosis mitigation. Therefore, the aim of this study was to determine the effect of ergovaline and red clover isoflavones on rumen microbial populations, fiber degradation, and volatile fatty acids (VFA) in an *in vitro* system. Using a dose of 1.10 mg × L⁻¹, endophyte-infected or endophyte-free tall fescue seed was added to ANKOM fiber bags with or without 2.19 mg of isoflavones in the form of a control, powder, or pulverized tablet, resulting in a 2 × 3 factorial arrangements of treatments. Measurements of pH, VFA, bacterial taxa, as well as the disappearance of neutral detergent fiber (aNDF), acid detergent fiber (ADF), and crude protein (CP) were taken after 48 h of incubation. aNDF disappearance values were significantly altered by seed type ($P = 0.003$) and isoflavone treatment ($P = 0.005$), and ADF disappearance values were significantly different in a seed × isoflavone treatment interaction ($P \leq 0.05$). A seed × isoflavone treatment interaction was also observed with respect to CP disappearance ($P \leq 0.05$). Eighteen bacterial taxa were significantly altered by seed × isoflavone treatment interaction groups ($P \leq 0.05$), eight bacterial taxa were increased by isoflavones ($P \leq 0.05$), and ten bacterial taxa were altered by seed type ($P \leq 0.05$). Due to the beneficial effect of isoflavones on tall fescue seed fiber degradation, these compounds may be viable options for mitigating fescue toxicosis. Further research should be conducted to determine physiological implications as well as microbiological changes *in vivo*.

Integrated Approach to Enhance Efficiency of Feed Utilization in Beef Production Systems. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors declare funding from a commercial source: PharmaCare Inc. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Ergot alkaloids produced by the endophytic fungus *Epichloë coenophiala* found in the endophyte-infected tall fescue plant (*Lolium arundinaceum* Schreb.) contribute to drought, heat, and disease tolerance of the plant, but are implicated in expensive production losses when grazed by livestock [1]. Concerns of the alkaloids manifest with reduced blood flow to the periphery, reduced reproductive efficiency, and reductions in growth performance [2–4]. Of the alkaloids associated with endophyte-infected tall fescue, ergovaline is the most predominant of concern in the plant, as it imposes the greatest detrimental effects on livestock [5, 6]. Reducing the consumption of endophyte-infected tall fescue has proven to be a challenge for livestock producers, as it occupies roughly 15 million hectares in the United States [7], with the majority in the east and southeast regions of the country. Replacing endophyte-infected varieties with novel and endophyte-free varieties of tall fescue can abate toxicosis, but may not be a cost-effective long-term solution [8]. Alternatively, the incorporation of cool-season legumes into endophyte-infected tall fescue pastures has been shown to mitigate a portion of tall fescue toxicosis symptoms [9, 10]. Phytoestrogenic compounds found in red and white clover have been reported to improve overall growth performance of livestock grazing endophyte-infected tall fescue throughout the spring and summer months [11–13]. Various isoflavones have been targeted for use in human medicine due to their beneficial reduction in breast cancer symptoms, improvements in cardiovascular health, as well as reductions in menopausal symptoms [14–16]. As such, determining additional benefits of isoflavones, such as the improvement of fiber utilization and rumen fermentation in livestock, is timely.

Ergot alkaloid pressure, or the physiological burden, is reduced in the presence of ruminal fluid [17, 18], indicating that ruminal microorganisms may be responsible for degradation of ergot alkaloids to less harmful compounds such as lysergic acid. Harlow, Goodman (19) identified several hyper-ammonia producing and tryptophan-utilizing bacteria capable of degrading ergovaline. Tryptophan is essential for ergovaline synthesis and formation of the ergoline ring structure. It has been hypothesized that bacterial species able to degrade tryptophan may also degrade ergovaline in the rumen [19, 20], proving essential for reducing the effects of tall fescue toxicosis. What is not as clear is the effect of ergovaline on the ruminal microbiome. As diet and toxins effect ruminal fermentation and production of glucogenic precursors required for the nutrition of the animal [21, 22], it is critical to determine the extent to which endophyte toxins in fescue may impact microbial numbers, alter microbial profiles, or influence volatile fatty acid (VFA) production and concentrations. While a phylogenetically [23] and functionally diverse rumen microbiome is essential to the health and nutritional status of ruminant animals, feed additives or plant-secondary compounds, such as isoflavones, may further improve rumen functionality in tandem with microbial populations. Isoflavones may enhance rumen function by improving fiber degradation and by reducing several microorganisms responsible for lowered ruminal pH and increased lactate production that can disrupt normal functional microflora [24, 25].

This study examined the effect of endophyte-infected or endophyte-free tall fescue seed with or without an isoflavone source on *in vitro* rumen fermentation, fiber degradation, and rumen bacterial communities. The objectives of this study were to 1) determine if isoflavones improve fiber degradation and fermentation of tall fescue seed *in vitro*, and 2) determine if *in vitro* rumen bacterial populations are affected by seed type or the addition of isoflavones. We hypothesized that the addition of isoflavones to endophyte-infected tall fescue seed would alter *in vitro* fermentation and rumen bacterial populations.

Materials and methods

Study design

This study was approved and carried out in accordance with the recommendations of the Institutional Animal Care and Use Committee at the University of Tennessee, Knoxville. Rumen content for the study was acquired from two fistulated Holstein heifers. The cattle are maintained and managed by the standard operating procedures of the Cherokee Farm (Knoxville, TN) as part of the University of Tennessee Institute of Agriculture. Heifers are continually maintained on mixed cool-season grass pastures.

This experiment was conducted as a randomized incomplete block design, and was blocked by each run. Eight separate fermentation runs were performed with the Daisy^{II} incubator (ANKOM Technology Co. Ltd. NY, USA). The first seven runs included four ANKOM fermentation jars (ANKOM Technology Co. Ltd. NY, USA) to which treatments were randomly allocated, and the eighth run consisted of two fermentation jars, for a total of five replications of six treatment combinations. Treatments were arranged as a 3×2 factorial design that included a commercial mix of isoflavones: (1) Promensil 80 mg (net) isoflavone tablet, (2) Promensil 80 mg (net) equivalent isoflavone powder, or (3) control, receiving no isoflavones, and two seed types: (1) endophyte-infected tall fescue seed (KY 31) or (2) endophyte-free tall fescue seed (KY 32). All seed was purchased from Turner Seed Inc. (Winchester, KY). Isoflavones in the form of the Promensil product were supplied by PharmaCare Laboratories (Warriewood, NSW 2102, Australia). As the Promensil tablet is processed and bound, the unprocessed powder was also examined.

For the responses of aNDF and ADF disappearance, ten fiber bags per fermentation jar were utilized as the sampling units. This yielded a total number of aNDF and ADF disappearance observations of 300. For the response of CP disappearance, five bags per fermentation jar were composited to a total of two weigh boats per fermentation jar as the sampling unit. This yielded a total number of CP disappearance observations of 60. Similarly, for the responses of ruminal pH and volatile fatty acids (VFA), two samples per fermentation jar were obtained using a 15-mL conical tube, yielding 60 ruminal pH and VFA observations.

Quantification of ergot alkaloids

Fescue seeds have been utilized in this study to eliminate any issues pertaining to volume and fiber that may impede or confound the interpretation of isoflavone effects on endophyte-infected tall fescue seed fermentation. This method has been utilized by numerous studies under this rationale [26, 27]. Prior to the study, concentrations of ergovaline and its epimer ergovalinine in fescue seed were determined using HPLC with fluorescence detection as described in Aiken et al. [27] with modifications described by Koontz et al. [28]. The endophyte-infected fescue seed contained 2.94 ppm of ergovaline and ergovalinine (1.85 and 1.09 ppm, respectively), and the endophyte-free tall fescue seed contained a total of 0 ppm of ergot alkaloids. Additionally, both seed varieties tested negative for the presence of the alkaloid ergotamine and its epimer ergotaminine. Seed samples were ground through a 5-mm screen using a Wiley Mill before 0.4425g of seed (1.3865×10^{-6} g ergot alkaloids) was added to each ANKOM fiber bag.

Quantification of isoflavones

Isoflavones were procured by finely grinding the product Promensil (PharmaCare Laboratories, Warriewood, NSW 2102, Australia), an over-the-counter isoflavone supplement isolated from red clover, with mortar and pestle to pass through a 1 mm screen. Quantification of

isoflavones in Promensil, including biochanin A, formononetin, genistein and daidzein, was performed using methods similar to those previously described by Aiken et al. (2016), using LC-MS rather than UV for detection. Briefly, isoflavone extracts were prepared by adding 7 mL of 85% methanol in 0.5% acetic acid to ground samples in 50-mL conical polypropylene tubes. Samples were briefly vortexed and sonicated for 30 min at ambient temperature. Three mL of deionized water was added to each sample prior to being vortexed and centrifuged for 8 min at $2200 \times g$. The resulting supernatant was filtered through a $0.45 \mu\text{m}$ GHP membrane syringe filter. Extracts were then diluted, and flavone added as internal standard. One portion of each sample was analyzed as-extracted and a second portion was heated at 85°C for 5 h to hydrolyze isoflavone malonyl-glucosides to their corresponding isoflavone glucosides. Concentrations of biochanin a-malonyl-glucoside and formononetin-malonyl-glucoside were determined by difference between hydrolyzed and un-hydrolyzed portions. Isoflavone extracts were analyzed by LC-MS on a Waters Acquity UPLC coupled to a Waters Synapt G2 (q-ToF) high resolution mass spectrometer. Chromatographic separation was obtained using a Waters BEH C18 UPLC column ($1.7 \mu\text{m}$, $2.1 \text{ mm} \times 150 \text{ mm}$). The mobile phase employed a mixture of water containing 0.1% formic acid (solvent A) and acetonitrile containing 0.1% formic acid (solvent B) in a linear gradient from 20% B to 80% B at a flow rate of $0.35 \text{ mL} \times \text{min}^{-1}$. The high resolution mass spectrometer was operated in positive ion electrospray mode with a resolving power of $\sim 14,000$ and scanned from 100 to 1000 Da in 0.3 s. Leucine enkephalin was used to provide a lock mass (m/z 554.2615). Quantification of isoflavones was performed using QuanLynx software with a linear calibration curve and internal standard method. Extracted ion chromatograms with a mass window of 0.02 Da around the accurate mass of each analyte were used to calculate peak areas. A total of 2.19 mg of total isoflavones were provided per bag (0.92 mg biochanin A, 0.89 mg formononetin, 0.0038 mg genistein and 0.0016 mg daidzein), for a concentration of $1.10 \text{ mg} \times \text{L}^{-1}$ of rumen fluid and buffer.

***In vitro* fiber and crude protein disappearance**

The Daisy^{II} *in vitro* incubation system (ANKOM Corp., Fairport, NY) was utilized to determine the rate and extent of fiber disappearance of the endophyte-infected or endophyte-free tall fescue seed with or without the addition of isoflavones from Promensil. Each substrate ($500 \pm 40 \text{ mg}$) was weighed into artificial fiber bags (F57 fiber bags, ANKOM Corp.), which were then heat sealed. Content included 442.5 mg of either endophyte-infected (KY 31) or endophyte-free tall fescue (KY 32) seed (dry matter basis), and 57.5 mg of Promensil (tablet or powder, dry matter basis). The fiber bags were separated into four groups of 12 bags within a seed type \times isoflavone treatment combination, including two empty bags for correction, and placed into separate upright plastic containers. A total of 1600 mL of rumen fluid was procured via aspiration from two fistulated Holstein heifers (Cherokee Farm, Knoxville, TN), and buffered using a 1:4 dilution of rumen fluid to McDougall's artificial saliva buffer [29]. A total of 400 mL were added to each ANKOM fermentation jar with 1600 mL of buffer, and adjusted to a pH of 6.8 with CO_2 . The fiber bags and diluted/buffered rumen fluid were added to the incubation containers. Incubation then proceeded for 48 h at $39 \pm 0.5^\circ\text{C}$.

Upon completion of the 48-h incubation, fermented rumen fluid pH was measured using a pH meter (Denver Instruments, Bohemia, NY) and 15 mL of fermented rumen content was sampled and stored at -80°C until further processing for bacterial DNA [30, 31]. All fiber bags were removed from fermentation jars, and washed under tap water before further analyses. Analysis of aNDF was obtained using α -amylase and sodium sulfite [32], and ADF was determined using sulfuric acid-based detergent [33]. Both procedures were conducted using the ANKOM200 fiber analysis system (ANKOM Corp., Fairport, NY). Crude protein was

determined by total nitrogen combustion analysis (LECO Instruments, Inc., St. Joseph, MI). Difference between pre- and post-fermentation aNDF, ADF, and CP concentrations were used to calculate disappearance.

VFA analysis

A subsample of rumen fermentation fluid was aliquoted from each fermentation jar for VFA analysis using HPLC, similar to the procedures previously described by Harlow et al. (2017). Briefly, samples were analyzed for concentrations of acetate, propionate, butyrate, valerate, and isovalerate/methylbutyrate (IVMB) using a Summit HPLC (Dionex; Sunnyvale, CA, USA) equipped with an anion exchange column (Aminex HP-87H; Bio-Rad, Hercules, CA, USA) and UV detector. Column parameters include a 300 x 7.8 mm, prepacked HPLC carbohydrate analysis column, hydrogen form, 9 μm particle size, 8% cross linkage, with a flow rate of 0.4 mL/min and an injection volume of 100 μL . The eluting compounds were separated isocratically with an aqueous sulfuric acid solution (5 mM). The parameters included an injection volume of 0.1 mL, flow rate of 0.4 mL \times min⁻¹, and column temperature of 50°C. Isovalerate and methyl butyrate (IVMB) resolve together in a single peak, and consequently, are reported together. A total VFA concentration (mMol) was calculated for each sample by summing the concentrations of all VFAs. A proportion of each specific VFA was calculated from the total by dividing the specific VFA by the total VFA and multiplying by 100 (units = mmol/100mmol total).

DNA extraction, PCR, and sequencing

DNA was extracted from ruminal fluid post-fermentation. The procedure of the DNA extraction method was similar to that previously described by Yu and Morrison [34]. After the chemical/mechanical cell lysis and isopropanol precipitation of nucleic acids, metagenomic DNA was purified with Rnase and proteinase K treatment, followed by the use of QIAamp columns from the Qiagen DNA Stool Mini Kit (Qiagen, Hilden, Germany). Genomic DNA concentration was determined using a Nanodrop 1000 spectrophotometer (Thermo Scientific, Wilmington, DE), and verified using Invitrogen Qubit fluorometer with PicoGreen (Thermo-Fisher Scientific, Wilmington, DE). Extractions were stored at -20°C until sequencing library preparation. Bacterial 16S rRNA genes were PCR-amplified with dual-barcoded primers targeting the V4 region, as per the protocol of Kozich [35]. Amplicons were sequenced with an Illumina MiSeq using the 250-bp paired-end kit (v.2). Sequences were denoised, taxonomically classified using Greengenes (v. 13_8) as the reference database, and clustered into 97% similarity operational taxonomic units (OTUs) with the mothur software package (v. 1.39.5), as previously described by Schloss, Westcott [36], and following the manufacturer-recommended procedure (https://www.mothur.org/wiki/MiSeq_SOP; accessed November 2017). Sequence files are available from the NCBI Sequence Read Archive (SRA Accession SRP150716). Additional descriptive information is associated with NCBI BioProject PRJNA476124.

Statistical analyses

The efficiency of the incomplete block design [37] was determined using PROC OPTEX in SAS 9.4 (SAS Inst. Inc., Cary, NC). Study analyses included responses of aNDF, ADF, CP disappearance, rumen pH, total VFA concentrations, specific VFA concentration in relation to total, and ruminal bacterial communities. Separate mixed model analyses of variance (ANOVA) were performed using the MIXED procedure of SAS to determine the fixed effects of isoflavone treatment, seed type, and seed type \times isoflavone treatment interaction. The ANOVA models also included random effects of run and run \times seed type \times isoflavone

treatment. Differences in least square means were separated using Fisher’s least significant differences. When an interaction between main effects was detected, multiple comparisons between all treatment combinations were adjusted using Tukey’s adjustments. Effects were considered significant at $P \leq 0.05$, and tendencies declared at $P > 0.05$ and ≤ 0.10 .

For rumen bacterial communities, analysis was conducted in the R environment. Alpha diversity was estimated with the Shannon index on raw OTU abundance tables. The significance of diversity differences was tested with an ANOVA. To estimate beta diversity across samples, OTUs were excluded if occurring in fewer than 10% of the samples with a count of less than three and computed Bray-Curtis indices. Beta diversity, emphasizing differences across samples, was visualized using non-metric multidimensional (NMDS) ordination. Variation in community structure was assessed with permutational multivariate analyses of variance (PERMANOVA) with treatment group as the main fixed factor and using 4,999 permutations for significance testing.

Results

Efficiency of Incomplete block design

Efficiency is a measurement of goodness of an experimental design [32]. D-efficiency is calculated based on the geometric mean of eigenvalues of the variance matrix [32]. In a perfectly balanced design, D-efficiency will equal 100%. In our study, the treatment D-efficiency was 87.6% for our allocation of treatments to runs.

aNDF, ADF, and CP disappearance

There was a main effect of seed type on aNDF disappearance ($P = 0.003$, Table 1). On average, there was reduced aNDF disappearance in E+ seed compared to that of E- seed ($11.53 \pm 0.97\%$ vs. $14.02 \pm 0.96\%$, respectively). Additionally, there was an effect of isoflavone treatment on aNDF disappearance ($P = 0.0053$). Isoflavone powder treatment, isoflavone tablet treatment, and control treatment resulted in $14.28 \pm 0.93\%$, $13.21 \pm 0.93\%$, and $10.82 \pm 0.93\%$ disappearance, respectively. There was no difference ($P = 0.49$) in aNDF disappearance between powder or tablet forms of isoflavone, and both forms of isoflavone resulted in significantly increased aNDF disappearance compared to the control.

ADF disappearance was associated with seed \times treatment type interaction ($P \leq 0.05$, Table 1). Effects of seed type on ADF disappearance was dependent upon isoflavone treatment. Samples that received the powdered isoflavone had greater ($P = 0.0007$) ADF disappearance

Table 1. Effects of isoflavones and fescue seed type on neutral detergent fiber (aNDFD), acid detergent fiber (ADF) and crude protein (CP) disappearance after rumen fluid incubation (48 h).

Measurement (%)	Isoflavone Treatment ¹						SEM	P-value		
	Control		Powder		Tablet			Seed	Isoflavone Treatment	Seed \times Isoflavone Treatment
	Fescue Seed ²									
	E+	E-	E+	E-	E+	E-				
aNDF	9.11	12.52	12.72	15.84	12.74	13.68	1.24	0.003	0.005	0.432
ADF	1.72 ^c	3.21 ^{bc}	7.01 ^a	2.19 ^c	1.27 ^c	5.91 ^b	1.03	0.36	0.009	< 0.001
CP	2.44 ^d	12.23 ^a	6.98 ^{bc}	7.33 ^{bc}	4.01 ^{cd}	9.08 ^{ab}	0.93	< 0.001	0.689	0.0002

¹Treatment: control, isoflavone powder, or isoflavone pulverized tablet.

²Seed type: KY 31 endophyte-infected (E+) tall fescue seed, KY 32 endophyte-free (E-) tall fescue seed.

^{abcd} Within an isoflavone treatment, seed types with like superscripts are not different at $P = 0.05$

<https://doi.org/10.1371/journal.pone.0201866.t001>

from endophyte-infected tall fescue seed ($7.01 \pm 1.04\%$) when compared to endophyte-free tall fescue seed ($2.19 \pm 1.04\%$). However, samples that received the tablet isoflavones had lesser ($P = 0.001$) ADF disappearance from endophyte-infected tall fescue seed ($1.27 \pm 1.04\%$) when compared to endophyte-free tall fescue seed ($5.91 \pm 1.04\%$). Within the control isoflavone treatment, there was no effect of seed type ($P = 0.52$).

There was a seed \times treatment interaction on overall CP disappearance ($P = 0.0002$, Table 1). In the control isoflavone group, reduced CP disappearance was observed with endophyte-infected tall fescue seed type ($2.44 \pm 0.93\%$) when compared to endophyte-free fescue seed ($12.23 \pm 0.93\%$, $P < 0.0001$). There was no CP disappearance difference between endophyte-infected and endophyte-free seed when the powder isoflavone was used ($P = 0.99$); however, when the tablet isoflavone form was supplemented, endophyte-infected tall fescue seed yielded decreased CP disappearance ($4.01 \pm 0.99\%$) compared to endophyte free tall fescue seed ($9.08 \pm 0.93\%$, $P = 0.01$, Table 1).

Rumen fermentation fluid VFA concentrations, bacterial populations, and pH

There was a tendency for an effect of seed type on total VFA concentrations. Total VFA was slightly increased in endophyte infected seed (21.68 ± 2.18) compared to endophyte-free seed (18.63 ± 2.1 , $P = 0.06$). Propionate concentrations tended to be reduced for endophyte-free tall fescue seed (31.11 ± 1.96) when compared to endophyte-infected seed (33.54 ± 1.96 , $P = 0.075$; Table 2), but no effect of treatment or treatment \times seed interaction was observed ($P > 0.1$). The acetate: propionate ratio tended to be reduced ($P = 0.09$; Table 2) with endophyte-infected tall fescue seed (1.74 ± 0.15) when compared to endophyte-free tall fescue seed (1.98 ± 0.15). No effects of seed type or isoflavone treatment were observed among acetate, butyrate, valerate or IVMB concentrations ($P > 0.1$).

After stringent sequence processing, a total of 408,813 high quality reads were obtained and averaged $17,034 \pm 3,180$ reads per sample, which is a consistent depth for sequencing analyses from ruminal samples [38]. Number of observed OTUs totaled 48,867 and averaged $2,037 \pm 169$ per sample. PERMANOVA global pairwise comparisons indicated significant

Table 2. Effects of isoflavones and fescue seed type on volatile fatty acid concentrations in the fluid phase of an *in vitro* rumen fluid digestion (48 h).

VFA	Isoflavone Treatment ¹						SEM	P-value		
	Control		Powder		Tablet			Seed	Isoflavone Treatment	Seed \times Isoflavone Treatment
	Fescue Seed ²									
	E+	E-	E+	E-	E+	E-				
Total VFA ³ VFA ⁴	23.55	20.00	19.66	17.98	21.84	17.90	2.73	0.06	0.33	0.86
Acetate	55.22	59.29	57.85	58.85	56.38	58.39	2.08	0.14	0.82	0.73
Propionate	35.24	31.23	31.15	30.78	34.23	31.30	2.38	0.07	0.34	0.56
Butyrate	7.94	7.88	9.58	8.06	7.07	8.81	1.12	0.95	0.52	0.28
Valerate	1.48	0.52	0.71	1.16	0.91	1.30	0.50	0.91	0.92	0.24
IVMB ⁵	0.81	1.05	0.76	0.81	0.75	0.84	0.24	0.15	0.35	0.68
Acetate:Propionate	1.60	1.97	1.90	1.99	1.73	1.98	0.20	0.09	0.62	0.73

¹Treatment: control, isoflavone powder, or isoflavone pulverized tablet.

²Seed type: KY 31 endophyte-infected (E+) tall fescue seed, KY 32 endophyte-free (E-) tall fescue seed.

³Concentration in mMol

⁴Concentration in mMol \times 100mMol⁻¹.

⁵IVMB: Isovalerate/methylbutyrate




Fig 1. Shannon's Diversity Index box plot of bacterial species diversity across treatment groups. Group 1: endophyte-infected tall fescue seed. Group 2: endophyte-free tall fescue seed. Group 3: endophyte-infected tall fescue seed with isoflavone powder. Group 4: endophyte-infected tall fescue seed with isoflavone tablet. Group 5: endophyte-free tall fescue seed with isoflavone powder. Group 6: endophyte-free tall fescue seed with isoflavone tablet.

<https://doi.org/10.1371/journal.pone.0201866.g001>

differences among treatment groups ($P = 0.03$, $R^2 = 0.32$), however Shannon's Diversity Index was not affected by treatment group ($P > 0.1$, Fig 1). Non-metric multidimensional scaling (NMDS) was utilized to analyze beta diversity (Figs 2 and 3), where clusters of samples represent similarity of 16S rRNA bacterial genera by group based on rank. While treatment groups may have appeared to cluster similarly, there was no observable pattern between groups.




Fig 2. NMDS ordination plot grouped by treatment. Points represent taxa within treatment groups. Similar clustering indicates similar taxa within treatment groups. Group 1: endophyte-infected tall fescue seed. Group 2: endophyte-free tall fescue seed. Group 3: endophyte-infected tall fescue seed with isoflavone powder. Group 4: endophyte-infected tall fescue seed with isoflavone tablet. Group 5: endophyte-free tall fescue seed with isoflavone powder. Group 6: endophyte-free tall fescue seed with isoflavone tablet.

<https://doi.org/10.1371/journal.pone.0201866.g002>




Fig 3. NMDS ordination plot indicating significant taxa at the genus level that influenced major bacterial shifts among treatments.

<https://doi.org/10.1371/journal.pone.0201866.g003>

Relative proportions of the genus level diversity between groups are represented in Fig 4. No differences in taxonomic profile were observed among groups ($P > 0.1$). However, finer changes in OTU abundances were identified. Numerous taxa were significantly different by seed, treatment, or seed \times treatment interaction ($P \leq 0.05$; Table 3). Fermented rumen fluid pH was not significantly affected by seed type, treatment or seed \times treatment interaction ($P > 0.1$), and pH ranged from 6.97 to 7.57.

Discussion

Previous research conducted by Flythe and Kagan [39] and Harlow et al. [25] that noted improved fiber degradation and ammonia reduction employed biochanin A as the primary isoflavone ($30 \text{ mg} \times \text{L}^{-1}$). The present study observed similar improvements in fiber degradation, including only $920 \text{ } \mu\text{g}$ biochanin A per fiber bag ($0.46 \text{ mg} \times \text{L}^{-1}$) and a total isoflavone concentration of $1.10 \text{ mg} \times \text{L}^{-1}$. The results from this experiment were consistent with those achieved by Harlow et al [25], however the amount of total isoflavones ($1.10 \text{ mg} \times \text{L}^{-1}$) in this experiment may have been insufficient to elucidate all of the beneficial effects on fiber degradation and ammonia reduction that were previously reported.

The increase in ADF disappearance observed in the present study is consistent with the observations of Harlow [25], when the inclusion of the isoflavone biochanin A reduced overall ADF values post fermentation compared to the controls. As acid detergent fiber measures the




Fig 4. Taxonomic profiles of the relative proportions of bacterial communities by genus, grouped by treatment. Group 1: endophyte-infected tall fescue seed. Group 2: endophyte-free tall fescue seed. Group 3: endophyte-infected tall fescue seed with isoflavone powder. Group 4: endophyte-infected tall fescue seed with isoflavone tablet. Group 5: endophyte-free tall fescue seed with isoflavone powder. Group 6: endophyte-free tall fescue seed with isoflavone tablet. No significant differences among groups ($P > 0.1$).

<https://doi.org/10.1371/journal.pone.0201866.g004>

amount of cellulose, which is a relatively indigestible fiber component, the inclusion of isoflavones with tall fescue seed in the present study increased indigestible fiber disappearance. Acid detergent fiber has an inverse relationship with digestible energy, and thus a high ADF component indicates less digestible energy available to the animal. In the present study, isoflavones resulted in a higher ADF disappearance, indicating possible improvement in available energy from tall fescue seed.

The results of CP disappearance in the present study are inconsistent with most tall fescue and CP literature, where proteolytic bacteria, including several species of hyper-ammonia producing bacteria (HAB), are more readily able to degrade ergot alkaloid compounds [19] such as those found in endophyte-infected tall fescue. While these hyper-ammonia bacteria are able to degrade ergovaline, they are a source of inefficiency in the animal when dietary nitrogen is lost as ammonia. Antimicrobials, including ionophores and similar compounds, may be employed in the diet to combat this loss of nitrogen, selectively inhibiting some of the microbiota and reducing wasteful byproducts of digestion [40–42]. Flythe et al. [39, 43] examined red clover isoflavones as antimicrobials due to their resistance to amino-acid degradation and action on HAB. They determined that biochanin A inhibited growth of *Clostridium stricklandii*, a member of HAB, and may prevent amino acid fermentation. A seed \times treatment interaction affected CP disappearance, and in general, increased disappearance was observed when endophyte-free tall fescue seed was fed while either no isoflavones were supplemented or when isoflavone was given in tablet form. However, the isoflavone concentration used in the present study may not have provided a sufficient amount of isoflavones to elicit a sizable reduction in ruminal amino acid degradation.

The donor animals ($n = 2$) used for rumen fluid procurement were not tall fescue-naïve animals, and consumed a forage-based diet. Thus, rumen fluid obtained for this experiment may have been previously exposed to ergot alkaloids, which may have affected the results of ruminal VFA concentrations. Additionally, ruminal microbial populations may have adapted or shifted from previous exposure to ergot alkaloids. The lack of significant differences in overall VFA concentrations (Table 2) provides support that the treatments did not inhibit or improve *in vitro* rumen fermentation. Richards and colleagues [44] similarly described no differences in rumen pH or VFA concentrations when steers consumed endophyte-infected tall fescue with or without soybean hulls. Rumen fermentation fluid pH samples were not below 6.0 across replications of the experiment indicating fiber digestion and cellulolytic bacteria function [45] was not reduced, as expected in the buffered system.

Overall, fiber degradability and VFA concentrations were notably reduced in the study in contrast to forage fermentation studies [46]. This was due to the fermentable substrate, as this study aimed to interrogate the effects of isoflavone supplementation on endophyte-infected tall fescue seed fermentation. Other studies have noted the benefits of utilizing ground tall fescue seed as a means of specifically examining the effects of ergot alkaloid exposure [26, 47–49]. Consistent with the previous studies, the current experiment reflects similar biologically significant results, including those regarding fiber degradation [25, 39], which support the conclusion that isoflavones may be effective at improving fiber utilization when supplemented. Although VFA concentrations were unexpectedly lower contrasted to *in vivo* studies [50, 51], several studies using *in vitro* fermentation have noted reduced concentrations using similar methodologies [24, 52].

Although diet imposes great variability on the bacterial community in the rumen [53], no large changes in bacterial community composition or phylogenetic diversity were identified among treatments. Diet similarity may contribute to these observations, as tall fescue seed-types were not drastically different among treatments and endophyte inclusion may induce smaller population changes. Indeed, rather than large changes in community composition,

Table 3. Effects of isoflavones and fescue seed type on relative abundance of significant *in vitro* rumen bacterial taxa.

Classification	Percentage of total sequences ¹						Effect	SEM	P-value ⁴
	Treatment ²								
	Control		Powder		Tablet				
	Fescue Seed ³								
	E+	E-	E+	E-	E+	E-			
<i>Actinomycetales_unclassified</i>	0.0011 ^b	ND ^b	0.013 ^{ab}	0.0027 ^b	0.012 ^{ab}	0.019 ^a	Treatment	0.0047	0.018
<i>Alphaproteobacteria_unclassified</i>	0.012 ^{bc}	0.017 ^{abc}	0.0255 ^{ab}	ND ^c	0.0203 ^{abc}	0.037 ^a	Seed × Treatment	0.0071	0.020
<i>Anaeroplasm</i>	0.054 ^{ab}	0.067 ^{ab}	0.0233 ^b	0.075 ^{ab}	0.044 ^b	0.108 ^a	Seed × Treatment	0.019	0.037
<i>Anaerovibrio</i>	0.074 ^{bc}	0.014 ^c	0.244 ^a	0.104 ^{bc}	0.158 ^{ab}	0.096 ^{bc}	Seed	0.0447	0.027
<i>Arcobacter</i>	0.928 ^{bc}	0.277 ^c	1.44 ^b	2.34 ^a	1.39 ^b	0.574 ^{bc}	Seed × Treatment	0.2937	0.015
<i>Bacteria_unclassified</i>	4.06 ^b	5.56 ^a	5.54 ^a	4.16 ^{ab}	4.23 ^{ab}	5.49 ^{ab}	Seed × Treatment	0.4920	0.015
<i>Bacteroidetes_unclassified</i>	8.77 ^d	13.96 ^a	12.62 ^{ab}	10.11 ^{cd}	11.26 ^{bc}	11.58 ^{abc}	Seed × Treatment	0.8247	0.001
<i>Bulleidia</i>	0.079 ^b	0.21 ^a	0.11 ^b	0.12 ^b	0.086 ^b	0.096 ^b	Seed × Treatment	0.0219	0.019
<i>Christensenellaceae_unclassified</i>	0.81 ^b	1.61 ^a	0.742 ^b	0.801 ^b	0.721 ^b	0.941 ^b	Seed	0.1092	0.031
<i>Clostridiales_unclassified</i>	3.33 ^b	5.60 ^a	4.18 ^{ab}	3.92 ^b	3.22 ^b	3.98 ^b	Seed × Treatment	0.2961	0.039
<i>Corynebacterium</i>	0.041 ^c	0.065 ^{bc}	0.131 ^a	0.109 ^{ab}	0.066 ^{bc}	0.076 ^{bc}	Seed	0.0014	0.009
<i>Endomicrobia_unclassified</i>	ND ^b	0.0115 ^a	ND ^b	0.0015 ^b	0.0015 ^b	0.0056 ^{ab}	Seed × Treatment	0.0024	0.007
<i>Gammaproteobacteria_unclassified</i>	0.018 ^{cd}	0.06 ^{ab}	0.056 ^{abc}	0.013 ^d	0.043 ^{bcd}	0.083 ^a	Seed × Treatment	0.0152	0.006
<i>Megasphaera</i>	0.117 ^a	0.018 ^b	0.0411 ^b	0.0431 ^b	0.0344 ^b	0.0286 ^b	Seed × Treatment	0.0502	0.008
<i>Methylobacillus</i>	ND	0.0014	0.0043	0.0042	ND	ND	Treatment	0.0051	0.020
<i>ML615J.28_unclassified</i>	0.027 ^b	0.065 ^b	0.109 ^b	0.123 ^b	0.123 ^b	0.456 ^a	Treatment	0.3030	0.009
<i>Moraxella</i>	0.0092 ^{ab}	0.0012 ^b	0.0142 ^{ab}	0.0029 ^b	0.027 ^a	0.0088 ^{ab}	Seed	0.0038	0.030
<i>Olsenella</i>	0.0292 ^b	0.148 ^a	0.053 ^b	0.038 ^b	0.034 ^b	0.039 ^b	Seed × Treatment	0.0233	0.021
<i>p.75.a5</i>	0.028 ^c	0.103 ^a	0.047 ^{bc}	0.071 ^{ab}	0.063 ^{abc}	0.071 ^{ab}	Seed	0.0134	0.004
<i>Peptoniphilus</i>	0.0069 ^{ab}	ND ^b	ND ^b	0.0146 ^a	0.0063 ^{ab}	0.0044 ^b	Seed × Treatment	0.0033	0.011
<i>Peptostreptococcaceae_unclassified</i>	0.0578 ^a	0.0088 ^c	0.0024 ^c	0.0278 ^{bc}	0.0373 ^{ab}	0.0162 ^{bc}	Seed × Treatment	0.0094	0.003
<i>Pirellulaceae_unclassified</i>	0.066 ^b	0.196 ^a	0.0811 ^b	0.125 ^{ab}	0.0677 ^b	0.103 ^{ab}	Seed	0.0192	0.019
<i>Planococcaceae_unclassified</i>	0.095 ^a	0.0014 ^b	0.0044 ^b	0.0118 ^b	0.007 ^b	0.0136 ^b	Seed × Treatment	0.0168	0.010
<i>Proteinclasticum</i>	1.45 ^{ab}	0.223 ^c	2.25 ^a	1.29 ^b	1.27 ^b	1.35 ^{ab}	Seed	0.1770	0.011
<i>Proteiniphilum</i>	0.097 ^{ab}	0.027 ^{ab}	0.0203 ^{ab}	0.0069 ^b	0.124 ^a	0.0148 ^b	Seed	0.0203	0.037
<i>Proteobacteria_unclassified</i>	0.157 ^b	0.196 ^b	0.272 ^{ab}	0.192 ^b	0.263 ^{ab}	0.378 ^a	Treatment	0.0332	0.021
<i>Pseudobutyrvibrio</i>	0.496 ^{bc}	0.237 ^c	1.11 ^{ab}	0.831 ^{abc}	1.24 ^a	0.681 ^{abc}	Treatment	0.1568	0.020
<i>Ruminococcaceae_unclassified</i>	2.18 ^{cd}	3.45 ^a	2.75 ^b	2.04 ^d	2.20 ^{cd}	2.66 ^{bc}	Seed × Treatment	0.1852	0.002
<i>Schwartzia</i>	0.0334 ^{bc}	0.0087 ^c	0.0766 ^{ab}	0.079 ^a	0.065 ^{ab}	0.060 ^{ab}	Treatment	0.0108	0.004
<i>Selenomonas</i>	0.219 ^{bc}	0.159 ^c	0.459 ^{abc}	0.629 ^a	0.543 ^{ab}	0.514 ^{ab}	Treatment	0.0842	0.012
<i>SRI_unclassified</i>	0.230 ^{bc}	0.0901 ^c	0.903 ^{ab}	0.950 ^a	0.954 ^a	0.607 ^{abc}	Treatment	0.236	0.0104
<i>Succiniclasticum</i>	0.672 ^b	1.67 ^a	0.662 ^b	0.774 ^b	0.627 ^b	0.632 ^b	Seed × Treatment	0.191	0.0349
<i>Succinivibrio</i>	0.002 ^b	0.014 ^a	ND ^b	0.004 ^{ab}	0.0015 ^b	0.0088 ^{ab}	Seed	0.1670	0.027
<i>Tenericutes_unclassified</i>	1.21 ^{cd}	1.01 ^d	2.43 ^a	1.32 ^{bcd}	2.14 ^{ab}	2.07 ^{abc}	Seed	0.2967	0.008
<i>Treponema</i>	0.0078 ^a	ND ^b	ND ^b	0.0028 ^b	ND ^b	0.0031 ^b	Seed × Treatment	0.0011	< 0.001
<i>Wautersiella</i>	0.445 ^b	0.797 ^b	0.865 ^{ab}	0.709 ^b	0.722 ^b	1.31 ^a	Seed × Treatment	0.153	< 0.001

¹ Data reported as LSMeans

² Treatment: control, isoflavone powder, or isoflavone pulverized tablet

³ Seed type: KY 31 endophyte infected (E+) tall fescue seed, KY 32 endophyte-free (E-) tall fescue seed

⁴ Within a row, differences between treatment groups are considered significant at $P \leq 0.05$, and trending toward significance at $P \leq 0.10$

^{abcd} Different means separation letters within a row indicate significant differences among LSMeans

<https://doi.org/10.1371/journal.pone.0201866.t003>

changes in specific taxa occurred, which may impact ruminal function and/or be impacted by supplementation of isoflavones. Overall, 36 taxa were significantly different by seed, treatment, or seed \times treatment interaction (Table 3). Studies by Harlow and colleagues [24] observed several species of amylolytic microbes that may be inhibited by biochanin A: *Streptococcus bovis* JB1, *Streptococcus bovis* HC5, *Lactobacillus reuteri*, and *Selenomonas ruminatum*. Additionally Harlow et al. [25] indicated several species of cellulolytic bacteria that are sensitive to biochanin A: *Fibrobacter succinogenes* S85, *Ruminococcus flavefaciens* 8, and *Ruminococcus albus* 8. In the current study, isoflavone supplementation was demonstrated to reduce Clostridiales with the E- seed, supporting results by Flythe et al. [39] where biochanin A inhibited growth of specific HAB Clostridium species, potentially preventing amino acid fermentation. Other cellulolytic and proteolytic bacteria also significantly varied among the treatments, including the genus *Megasphaera*, which was reduced with isoflavone supplementation. As a major lactic acid utilizer in the rumen, *Megasphaera elsdenii* has positive relationship with *S. bovis*, due to lactate production from *S. bovis* [54, 55]. Although no *S. bovis* variations were identified in the current study, the reduction in *M. elsdenii* hints at similar inhibitions noted by other researchers [24]. In contrast, the same species shifts that occurred in studies conducted by Harlow et al. [24] were not observed in the present study. Rather than being inhibited, *Selenomonas* proportions were increased when supplemented with isoflavones. However, in both studies conducted by Harlow et al. [24, 25], biochanin A was the only isoflavone supplemented and thus subsequent studies should be completed to verify population shifts with other isoflavones.

The combined results of this study indicated fermentation modifications which occurred due to treatments, specifically in aNDF, ADF, and CP disappearance, but these results may have been affected by several factors. Where both fiber and crude protein disappearance results were not consistent with previous *in vitro* studies, this may be the result of a lower quantity of isoflavone administration to the *in vitro* fermentation system. As previous studies utilized a minimum of $30 \text{ mg} \times \text{L}^{-1}$ of biochanin A to elucidate responses, the present study utilized $0.63 \text{ mg} \times \text{L}^{-1}$ of biochanin A, which may have not been enough to elicit a robust response. However, the current study was designed to examine effects of isoflavone administration on fermentation of ergot alkaloid-containing tall fescue, rather than effects on HAB and amino acid fermentation. Ergot alkaloid pressure on rumen fermentation was not clearly elucidated in the present study. VFA concentrations are indicative of rumen fermentation, which were not affected by each treatment. It is noteworthy that isoflavone administration did not alter VFA concentrations, and that seed type was the driving factor in any of the changes in VFA concentrations. Foote et al. [56] noted significant reductions in VFA flux and absorption in animals administered endophyte-infected tall fescue seed, lending support that nutrient utilization and energy absorption may be further delayed when cattle experience tall fescue toxicosis. As propionate supplies up to 65% of the glucose carbon in cattle [57], and livestock consuming endophyte-infected tall fescue produce reduced concentrations of VFAs [49], it is interesting that the acetate-to-propionate ratio tended to be reduced by endophyte-infected seed in the present study, as the combined effects of nutrient production, such as production of glucogenic precursors, and nutrient absorption will have significant impacts on animal production *in vivo*. The variability of this ratio may have been impacted by organisms identified in the current study that produce acetate, such as genera *Corynebacterium* and *Succinivibrio* [58, 59], as their respective abundances were similarly reduced with endophyte-infected seed. These microbiome-fermentation parameter associations hint at further evidence pertaining to reduced production and efficiency of cattle on endophyte-infected tall fescue. Further, Foote and colleagues reported similarly reduced VFA concentrations, when rumen buffers containing extracts of ergovaline were applied to heat stressed, ruminally cannulated steers [56]. The tendency for reduction of certain VFA concentrations observed in the present study, as well as

reduced total VFAs described by Foote and colleagues [56], may provide further evidence that consumption of endophyte-infected tall fescue impacts nutrient production and absorption.

While rumen fermentation fluid bacterial populations varied among groups, the rumen microbiome is a dynamic ecosystem that is affected by various environmental and dietary factors, and thus changes in the bacterial taxa among groups may not be solely indicative of treatment with fescue seed or isoflavone treatment. As other studies have noted specific bacterial species capable of ergovaline degradation and benefits from isoflavones, further research should be conducted to validate consistent shifts in the rumen microbiome from ergot alkaloid pressure. It must also be noted that isoflavones are estrogenic. In finishing steers or cattle intended for harvest, this is of little issue and poses no risk to human health. However, the reproductive effects of isoflavone supplementation must be accounted for when administered in studies and systems where reproductive status is critical.

Conclusions

In the current study, aNDF, ADF, and CP disappearance, as well as the abundances of 36 bacterial taxa, were impacted by isoflavone administration *in vitro*. Future research should be conducted to determine the impact of isoflavones on beef production *in vivo*, as the combined effects on ruminal feedstuff fermentation and HAB provide opportunities to naturally improve ruminal fermentation and enhance the efficiency of nutrient use in cattle.

Acknowledgments

The authors would like to thank Gloria Gellin, USDA-ARS, for HPLC analyses.

Author Contributions

Conceptualization: Emily A. Melchior, Jason K. Smith, J. Travis Mulliniks, Gary E. Bates, Michael D. Flythe, James L. Klotz, Phillip R. Myer.

Data curation: Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, J. Travis Mulliniks, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, Phillip R. Myer.

Formal analysis: Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, Gary E. Bates, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, Phillip R. Myer.

Funding acquisition: Emily A. Melchior, Gary E. Bates, Phillip R. Myer.

Investigation: Emily A. Melchior, Jason K. Smith, Gary E. Bates, Michael D. Flythe, Phillip R. Myer.

Methodology: Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, J. Travis Mulliniks, Gary E. Bates, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, Phillip R. Myer.

Project administration: Emily A. Melchior, J. Travis Mulliniks, Gary E. Bates, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Phillip R. Myer.

Resources: Jason K. Smith, Gary E. Bates, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, Phillip R. Myer.

Software: Phillip R. Myer.

Supervision: J. Travis Mulliniks, Gary E. Bates, Phillip R. Myer.

Validation: Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, J. Travis Mulliniks, Gary E. Bates, Michael D. Flythe, James L. Klotz, Phillip R. Myer.

Visualization: Emily A. Melchior, Jason K. Smith, J. Travis Mulliniks, Gary E. Bates, Phillip R. Myer.

Writing – original draft: Emily A. Melchior, Michael D. Flythe, Phillip R. Myer.

Writing – review & editing: Emily A. Melchior, Jason K. Smith, Liesel G. Schneider, J. Travis Mulliniks, Gary E. Bates, Zachary D. McFarlane, Michael D. Flythe, James L. Klotz, Jack P. Goodman, Huihua Ji, Phillip R. Myer.

References

1. Kallenbach RL, BILL E. KUNKLE INTERDISCIPLINARY BEEF SYMPOSIUM: Coping with tall fescue toxicosis: Solutions and realities12. *J Anim Sci.* 2015; 93(12):5487–95. <https://doi.org/10.2527/jas.2015-9229> PMID: 26641158
2. Bush L, Buckner RC. Tall Fescue Toxicity. In: Matches AG, editor. *Anti-Quality Components of Forages.* CSSA Special Publication. Madison, WI: Crop Science Society of America; 1973. p. 99–112.
3. Ball D, Lacefield GD, Hoveland CS. The tall fescue endophyte. 1991.
4. Bush L, Streeter C, Buckner R. Peroline Inhibition of *in vitro* Ruminal Cellulose Digestion 1. *Crop Science.* 1970; 10(1):108–9.
5. Klotz JL, Nicol AM. Ergovaline, an endophytic alkaloid. 1. Animal physiology and metabolism. *Animal Production Science.* 2016; 56(11):1761–74.
6. Strickland JR, Looper ML, Matthews J, Rosenkrans C, Flythe M, Brown K. Board-invited review: St. Anthony's Fire in livestock: causes, mechanisms, and potential solutions. *J Anim Sci.* 2011; 89(5):1603–26. <https://doi.org/10.2527/jas.2010-3478> PMID: 21521821
7. Kim Abney L, Oliver JW, Reinemeyer CR. Vasoconstrictive effects of tall fescue alkaloids on equine vasculature. *J Equine Vet Sci.* 1993; 13(6):334–40.
8. Zhuang J, Marchant MA, Schardl CL, Butler CM. Economic Analysis of Replacing Endophyte-Infected with Endophyte-Free Tall Fescue Pastures. *Agron J.* 2005; 97(3):711–6.
9. Lusby KS, McMurphy WE, Strasia CA, Smith SC, Muntz SH. Effects of Fescue Endophyte and Interseeded Clovers on Subsequent Finishing Performance of Steers. *Journal of Production Agriculture.* 1990; 3(1):103–5.
10. McMurphy WE, Lusby KS, Smith SC, Muntz SH, Strasia CA. Steer Performance on Tall Fescue Pasture. *Journal of Production Agriculture.* 1990; 3(1):100–2.
11. Aiken GE, Flythe MD, Kagan IA, Ji H, Bush LP. Mitigation of Ergot Vasoconstriction by Clover Isoflavones in Goats (*Capra hircus*). *Frontiers in Veterinary Science.* 2016; 3(17).
12. Harlow BE, Flythe MD, Kagan IA, Aiken GE. Biochanin A (an Isoflavone Produced by Red Clover) Promotes Weight Gain of Steers Grazed in Mixed Grass Pastures and Fed Dried-Distillers' Grains. *Crop Science.* 2017; 57(1):506–14.
13. Shappell N, Flythe M, Aiken G. The effects of steroid implant and dietary soybean hulls on estrogenic activity of sera of steers grazing toxic endophyte-infected tall fescue pasture. *Frontiers in Veterinary Science.* 2015; 2(30).
14. van de Weijer PHM, Barentsen R. Isoflavones from red clover (Promensil) significantly reduce menopausal hot flush symptoms compared with placebo. *Maturitas.* 2002; 42(3):187–93. PMID: 12161042
15. Beck V, Rohr U, Jungbauer A. Phytoestrogens derived from red clover: an alternative to estrogen replacement therapy? *The Journal of steroid biochemistry and molecular biology.* 2005; 94(5):499–518. <https://doi.org/10.1016/j.jsbmb.2004.12.038> PMID: 15876415
16. Cruz MN, Luksha L, Logman H, Poston L, Agewall S, Kublickiene K. Acute responses to phytoestrogens in small arteries from men with coronary heart disease. *American Journal of Physiology-Heart and Circulatory Physiology.* 2006; 290(5):H1969–H75. <https://doi.org/10.1152/ajpheart.01065.2005> PMID: 16361368
17. Westendorf M G. Mitchell, Tucker. R. Influence of rumen fermentation on response to endophyte infected tall fescue seed measured by a rat bioassay. *Drug Chem Toxicology.* 1992; 15:351–64.

18. De Lorme MJM, Lodge-Ivey SL, Craig AM. Physiological and digestive effects of *Neotyphodium coenophialum*-infected tall fescue fed to lambs¹. *J Anim Sci*. 2007; 85:1199–206. <https://doi.org/10.2527/jas.2005-430> PMID: 17296774
19. Harlow BE, Goodman JP, Lynn BC, Flythe MD, Ji H, Aiken GE. Ruminal tryptophan-utilizing bacteria degrade ergovaline from tall fescue seed extract¹. *J Anim Sci*. 2017.
20. Mantle PG. The role of tryptophan as a biosynthetic precursor of indole-diterpenoid fungal metabolites: Continuing a debate. *Phytochemistry*. 2009; 70(1):7–10. <https://doi.org/10.1016/j.phytochem.2008.11.004> PMID: 19136126
21. Weimer PJ. Redundancy, resilience, and host specificity of the ruminal microbiota: implications for engineering improved ruminal fermentations. *Front Microbiol*. 2015; 6:296. <https://doi.org/10.3389/fmicb.2015.00296> PMID: 25914693
22. Allison MJ, Dawson KA, Mayberry WR, Foss JG. *Oxalobacter formigenes* gen. nov., sp. nov.: oxalate-degrading anaerobes that inhabit the gastrointestinal tract. *Arch Microbiol*. 1985; 141(1):1–7. PMID: 3994481
23. Myer PR, Freetly HC, Wells JE, Smith TPL, Kuehn LA. Analysis of the gut bacterial communities in beef cattle and their association with feed intake, growth, and efficiency. *J Anim Sci*. 2017; 95(7):3215–24. <https://doi.org/10.2527/jas.2016.1059> PMID: 28727105
24. Harlow BE, Flythe M.D., Aiken G.E. Effect of biochanin A on corn grain (*Zea mays*) fermentation by bovine rumen amylolytic bacteria. *J Appl Microbiol*. 2017; 122:870–80. <https://doi.org/10.1111/jam.13397> PMID: 28055130
25. Harlow BE, Flythe M. D. and Aiken G. E. Biochanin A improves fibre fermentation by cellulolytic bacteria. *J Appl Microbiol*. 2018; 124:58–66. <https://doi.org/10.1111/jam.13632> PMID: 29112792
26. Green MA, Whitlock BK, Edwards JL, Scholljegerdes EJ, Mulliniks JT. Rumen-protected arginine alters blood flow parameters and luteinizing hormone concentration in cyclic beef cows consuming toxic endophyte-infected tall fescue seed. *J Anim Sci*. 2017; 95(4):1537–44. <https://doi.org/10.2527/jas.2016.1309> PMID: 28464107
27. Aiken GE, Strickland J. R., Looper M. L, Bush L. P, and Schrick F. N. Hemodynamics are altered in the caudal artery of beef heifers fed different ergot alkaloid concentrations. *J Anim Sci*. 2009; 87:2142–50. <https://doi.org/10.2527/jas.2008-1562> PMID: 19251925
28. Koontz AF, Bush LP, Klotz JL, McLeod KR, Schrick FN, Harmon DL. Evaluation of a ruminally dosed tall fescue seed extract as a model for fescue toxicosis in steers¹. *J Anim Sci*. 2012; 90(3):914–21. <https://doi.org/10.2527/jas.2011-4292> PMID: 22064740
29. McDougall EI. Studies on ruminant saliva. 1. The composition and output of sheep's saliva. *Biochem J*. 1948; 43(1):99–109.
30. Contreras-Govea F, Muck R, Weimer P, Hymes-Fecht U. *In vitro* ruminal fermentation of treated alfalfa silage using ruminal inocula from high and low feed-efficient lactating cows. *J Appl Microbiol*. 2016; 121(2):333–40. <https://doi.org/10.1111/jam.13103> PMID: 27241014
31. Jeyanathan J, Martin C, Morgavi D. Screening of bacterial direct-fed microbials for their antimethanogenic potential *in vitro* and assessment of their effect on ruminal fermentation and microbial profiles in sheep. *J Anim Sci*. 2016; 94(2):739–50. <https://doi.org/10.2527/jas.2015-9682> PMID: 27065144
32. Mertens DR. Gravimetric Determination of Amylase-Treated Neutral Detergent Fiber in Feeds with Refluxing in Beakers or Crucibles: Collaborative Study. *J AOAC Int*. 2002; 85(6):1217–40. PMID: 12477183
33. Vogel KP, Pedersen JF, Masterson SD, Toy JJ. Evaluation of a Filter Bag System for NDF, ADF, and IVDMD Forage Analysis. *Crop Science*. 1999; 39(1):276–9.
34. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques*. 2004; 36(5):808–13. <https://doi.org/10.2144/04365ST04> PMID: 15152600
35. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Appl Environ Microbiol*. 2013; 79(17):5112–20. <https://doi.org/10.1128/AEM.01043-13> PMID: 23793624
36. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol*. 2009; 75(23):7537–41. <https://doi.org/10.1128/AEM.01541-09> PMID: 19801464
37. Kuhfeld W. Experimental design: efficiency, coding, and choice designs. 2010. MR-2010C; 2010.
38. Myer PR, Smith TPL, Wells JE, Kuehn LA, Freetly HC. Rumen microbiome from steers differing in feed efficiency. *PLoS One*. 2015; 10.

39. Flythe M, Kagan I. Antimicrobial Effect of Red Clover (*Trifolium pratense*) Phenolic Extract on the Ruminal Hyper Ammonia-Producing Bacterium, *Clostridium sticklandii*. *Curr Microbiol*. 2010; 61(2):125–31. <https://doi.org/10.1007/s00284-010-9586-5> PMID: 20087740
40. Chen G, Russell J. More monensin-sensitive, ammonia-producing bacteria from the rumen. *Appl Environ Microbiol*. 1989; 55(5):1052–7. PMID: 2757371
41. Tedeschi LO, Fox DG, Tylutki TP. Potential Environmental Benefits of Ionophores in Ruminant Diets Mention of trade names, proprietary products, or specific equipment does not constitute a guarantee or warranty of the product by the authors and does not imply its approval to the exclusion of other products that also may be suitable. *J Environ Qual*. 2003; 32(5):1591–602. PMID: 14535299
42. Reynolds CK, Humphries DJ, Kirton P, Kindermann M, Duval S, Steinberg W. Effects of 3-nitrooxypropanol on methane emission, digestion, and energy and nitrogen balance of lactating dairy cows. *J Dairy Sci*. 2014;97.
43. Flythe M, Harrison B, Kagan I, Klotz J, Gellin G, Goff B, et al. Antimicrobial activity of red clover (*Trifolium pratense* L.) extract on caprine hyper ammonia-producing bacteria. *Agric Food Anal Bacteriol*. 2013; 3:176–85.
44. Richards CJ, Pugh RB, Waller JC. Influence of soybean hull supplementation on rumen fermentation and digestibility in steers consuming freshly clipped, endophyte-infected tall fescue1. *J Anim Sci*. 2006; 84(3):678–85. PMID: 16478960
45. Orskov ER. Protein nutrition in ruminants. London: Academic Press Inc. (London) Ltd.; 1982. viii + 160pp. p.
46. Sari M, Ferret A, Calsamiglia S. Effect of pH on *in vitro* microbial fermentation and nutrient flow in diets containing barley straw or non-forage fiber sources. *Animal Feed Science and Technology*. 2015; 200:17–24.
47. Aiken GE, Kirch BH, Strickland JR, Bush LP, Looper ML, Schrick FN. Hemodynamic responses of the caudal artery to toxic tall fescue in beef heifers1. *J Anim Sci*. 2007; 85(9):2337–45. <https://doi.org/10.2527/jas.2006-821> PMID: 17526671
48. Jackson JA, Hemken R, Boling J, Harmon R, Buckner R, Bush L. Summer Fescue Toxicity in Dairy Steers Fed Tall Fescue Seed 1. *J Anim Sci*. 1984; 58(5):1057–61. PMID: 6735937
49. Hannah S, Paterson J, Williams J, Kerley M, Miner J. Effects of increasing dietary levels of endophyte-infected tall fescue seed on diet digestibility and ruminal kinetics in sheep. *J Anim Sci*. 1990; 68(6):1693–701. PMID: 2117003
50. Van Gastelen S, Antunes-Fernandes E, Hettinga K, Klop G, Alferink S, Hendriks W, et al. Enteric methane production, rumen volatile fatty acid concentrations, and milk fatty acid composition in lactating Holstein-Friesian cows fed grass silage- or corn silage-based diets. *J Dairy Sci*. 2015; 98(3):1915–27. <https://doi.org/10.3168/jds.2014-8552> PMID: 25582590
51. Weimer P, Cox M, Vieira de Paula T, Lin M, Suen G. 1657 Changes in milk production efficiency and ruminal bacterial community composition following near-total exchange of ruminal contents between high- and low-efficiency Holstein cows. *J Anim Sci*. 2016; 94(suppl_5):807–8.
52. Flythe M, Aiken G. Effects of hops (*Humulus lupulus* L.) extract on volatile fatty acid production by rumen bacteria. *J Appl Microbiol*. 2010; 109(4):1169–76. <https://doi.org/10.1111/j.1365-2672.2010.04739.x> PMID: 20456526
53. Henderson G, Cox F, Ganesh S, Jonker A, Young W, Collaborators GRC, et al. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci Rep*. 2015; 5:14567. <https://doi.org/10.1038/srep14567> PMID: 26449758
54. Kung L Jr, Hession AO. Preventing *in vitro* lactate accumulation in ruminal fermentations by inoculation with *Megasphaera elsdenii*. *J Anim Sci*. 1995; 73(1):250–6. PMID: 7601741
55. Luo J, Ranadheera CS, King S, Evans CA, Baines S. Dairy propionibacteria as direct-fed microbials: *in vitro* effect on acid metabolism of *Streptococcus bovis* and *Megasphaera elsdenii*. *Ann Microbiol*. 2018; 68(3):153–8.
56. Foote AP, Kristensen NB, Klotz JL, Kim DH, Koontz AF, McLeod KR, et al. Ergot alkaloids from endophyte-infected tall fescue decrease reticuloruminal epithelial blood flow and volatile fatty acid absorption from the washed reticulorumen1,2. *J Anim Sci*. 2013; 91(11):5366–78. <https://doi.org/10.2527/jas.2013-6517> PMID: 23989869
57. Herbein JH, Van Maanen RW, McGilliard AD, Young JW. Rumen Propionate and Blood Glucose Kinetics in Growing Cattle Fed Isoenergetic Diets. *The Journal of Nutrition*. 1978; 108(6):994–1001. <https://doi.org/10.1093/jn/108.6.994> PMID: 650302
58. Pan X, Xue F, Nan X, Tang Z, Wang K, Beckers Y, et al. Illumina sequencing approach to characterize thiamine metabolism related bacteria and the impacts of thiamine supplementation on ruminal

microbiota in dairy cows fed high-grain diets. *Front Microbiol.* 2017; 8:1818. <https://doi.org/10.3389/fmicb.2017.01818> PMID: 28979254

59. Reddy CA, Kao M. Value of acid metabolic products in identification of certain corynebacteria. *J Clin Microbiol.* 1978; 7(5):428–33. PMID: 96126