



**UNIVERSITI PUTRA MALAYSIA**

***MODULATION OF NMDA AND AMPA/KAINATE RECEPTORS BY  
TOCOTRIENOL-RICH FRACTION AND  $\alpha$ -TOCOPHEROL IN  
GLUTAMATE- INDUCED INJURY OF PRIMARY ASTROCYTES***

**ZAHRA ABEDI**

**FPSK(M) 2017 17**



**MODULATION OF NMDA AND AMPA/KAINATE RECEPTORS BY  
TOCOTRIENOL-RICH FRACTION AND  $\alpha$ -TOCOPHEROL IN  
GLUTAMATE- INDUCED INJURY OF PRIMARY ASTROCYTES**

By

**ZAHRA ABEDI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Master of Science**

**March 2017**

## COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**MODULATION OF NMDA & AMPA/KAINATE RECEPTORS BY  
TOCOTRIENOL-RICH FRACTION AND  $\alpha$ -TOCOPHEROL  
IN GLUTAMATE INDUCED INJURY OF PRIMARY ASTROCYTES**

By

**ZAHRA ABEDI**

**March 2017**

**Chairman : Huzwah Binti Khaza'ai, PhD**  
**Faculty : Medicine and Health Sciences**

Neurodegenerative diseases such as Huntington's, Alzheimer's, Parkinson's disease and stroke are the most common diseases suffered by the aged population. Glutamate is considered as a main excitatory amino acid neurotransmitter in the mammalian central nervous system which can be excitotoxic, playing a key role in series of chronic neurodegenerative diseases. The Vitamin E which consist of tocopherol and tocotrienols, are different in their side chain either in saturated or unsaturated phytol tail. Previous studies have demonstrated that tocopherol and tocotrienol have protective effects against glutamate toxicity in an astrocytic cell line. The aim of current study is to demonstrate the potential of tocopherol and tocotrienol in protecting glutamate injured primary astrocytes. For this purpose, the primary astrocytes were isolated from mixed glial cells of C57BL/6 mice by using the Easysep Mouse CD11b positive selection kit and cultured in supplemented DMEM. Mixed glial cultures were treated with 50-75 mM L-leucine methyl ester (LME) for 60-90 minutes to improve purity of cultures. The purity of primary astrocytes was measured by flow-cytometer and is approximately 79.4%. The  $IC_{20}$  and  $IC_{50}$  values of glutamate were determined by MTT assay at 10 mM and 100 mM respectively. Cell were induced injury at  $IC_{20}$  and  $IC_{50}$  of glutamate and the effects of tocopherol and tocotrienol rich fraction (TRF) was determined.in pre and post-treatment study. For the high yield of RNA, the  $IC_{20}$  of glutamate was used in the experiment. Exposure to 100 mM of glutamate in primary astrocytes reduced cell viability by approximately 64.75 % and 61.10 % in pre and post treatment study respectively. The mitochondrial membrane potential (MMP) detected in primary astrocytes were assessed with 100, 200 and 300 ng/ml concentration of TRF and  $\alpha$ -Tocopherol. The results depicted that pre-treatment with TRF and  $\alpha$ -Tocopherol caused the mitochondrial activity to achieve 88.46%, 82.42%, 80.74% and 93.31%, 87.51%, 83.70%, respectively. In post-treatment study, with increase of TRF (100, 200 and 300 ng/ml) concentration causes the increase to 61.21%, 73.01%, 78.43% of MMP value. Similarly, increase of MMP value from 66.12%, 76.46%, and 81.22% was observed with increasing

concentration of  $\alpha$ -Tocopherol. Then the expression of ionotropic glutamate receptors genes was elucidated using Real-time PCR. The gene of interest consists of the Gria2 (Glutamate Receptor, Ionotropic AMPA), GRIK1 (Glutamate Receptor, Ionotropic, kainate1) and Grin2A (Glutamate Receptor, Ionotropic, N-Methyl D-Aspartate 2A). The results showed that in both pre and post studies, the ionotropic glutamate receptors genes were down regulated after the treatment and  $\alpha$ -tocopherol played an important role in down regulating the genes. The most affected genes were Gria2, GriK1 and Grin2A respectively in both pre and post studies. Decreased intracellular calcium concentration also was observed indicating the present of vitamin E altered the polarization of astrocytes. As a conclusion, this study shown that  $\alpha$ -tocopherol is more effective and only required low concentration of  $\alpha$ -tocopherol for prophylactic purposes compared to post-treatment in primary astrocytes cells.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**MODULASI RESEPTOR-RESEPTOR NMDA & AMPA/ KAINATE OLEH FRAKSI-KAYA TOKOTRIENOL DAN TOKOFEROL DI DALAM SEL PRIMER ASTROSIT YANG DIRANGSANG KECEDEeraan OLEH GLUTAMAT**

Oleh

**ZAHRA ABEDI**

**Mac 2017**

**Pengerusi : Huzwah Khaza'ai, PhD**  
**Fakulti : Perubatan dan Sains kesihatan**

Penyakit neurodegeneratif seperti Huntington's, Alzheimer's dan Parkinson's dan strok adalah penyakit yang sering dialami oleh populasi golongan warga tua. Glutamate merupakan neurotransmitter utama asid amino bagi sistem saraf pusat mamalia yang boleh menyebabkan eksitotoksik dan ia memainkan peranan penting dalam siri penyakit kronik neurodegeneratif. Vitamin E terdiri daripada tokoferol dan tokotrienol yang mempunyai perbezaan dari segi struktur rantai sisi lemak tidak tepu dan lemak tepu. Berdasarkan kajian sebelum ini, tokoferol dan tokotrienol didapati mempunyai kesan pelindung terhadap toksisiti glutamat di dalam sel astrosit. Tujuan kajian ini dilakukan adalah bagi melihat potensi Vitamin E dalam melindungi sel astrosit primer daripada kecederaan yang disebabkan oleh glutamate. Dengan ini, sel primer astrosit telah dipencilkan daripada campuran sel glial tikus C57BL/6 dengan menggunakan kit pemilihan positif Easysep Mouse CD116 dan dikulturkan di dalam media *DMEM* yang mengandungi supplemen. Kultur campuran glial dirawat selama 60-90 minit dengan 50-75 mM L-leucine methyl ester (LME) bagi tujuan memperbaiki ketulenan kultur tersebut. Ketulenan sel astrosit yang diperolehi adalah 79.4% dengan menggunakan "flow-sitometer". Nilai  $IC_{20}$  dan  $IC_{50}$  glutamate ditentukan dengan menggunakan MTT asai iaitu 10 mM dan 100 Mm. Sel diceredakan dengan glutamat pada aras  $IC_{20}$  dan  $IC_{50}$  untuk kesan tokoferol dan fraksi kaya tokotrienol (TRF) dan ditentukan di dalam pra dan pasca rawatan. Bagi tujuan menghasilkan RNA yang tinggi, eksperimen dijalankan dengan menggunakan glutamat pada  $IC_{20}$ . Pada kepekatan 100 mM glutamat di dalam sel astrosit utama menyebabkan pengurangan sel hidup dengan anggaran 64.75% dan 61.10% di dalam pra dan pasca rawatan. Potensi membran mitokondria (MMP) mengesan sel utama astrosit pada kepekatan 100, 200 dan 300 ng/ml untuk fraksi kaya tokotrienol (TRF) dan  $\alpha$ -tokoferol. Hasil kajian pra-rawatan dengan menggunakan TRF dan  $\alpha$ -tokoferol menyebabkan aktiviti mitokondria telah mencapai sebanyak 88.46%, 82.42%, 80.47% bagi TRF dan 93.31%, 87.51%, 83.70% untuk tokoferol. Dalam kajian pasca-rawatan

pula, pertambahan kepekatan TRF pada 100, 200 dan 300 ng/mL menyebabkan kenaikan nilai MMP kepada 61.21%, 73.01% dan 78.43%. Kenaikan yang serupa bagi nilai MMP daripada 66.12%, 76.46% ke 81.22% diperhatikan apabila kepekatan  $\alpha$ -tokoferol meningkat. Ekspresi reseptor gen ionotropik glutamat juga telah dikenalpasti menggunakan "Real-time PCR". Gen-gen tersebut ialah Gria2 (Reseptor glutamate, Ionotropik AMPA), GRIK1 (Reseptor Glutamate, Ionotropik, Kinase 1) dan Grin2A (Reseptor Glutamate, Ionotropik, N-Methyl-D-Aspartate 2A). Kajian didapati, kedua-dua pra dan pasca rawatan menunjukkan reseptor ionotropik glutamate menurun selepas rawatan, dan ini menunjukkan keberkesanan  $\alpha$ -tokoferol dalam mengawal selia gen-gen tersebut. Gen yang paling terkesan dalam pra dan pasca rawatan kajian, adalah Gria2, Grik1 dan Grin2A. Penurunan kepekatan kalsium pada intraselular menyebabkan perubahan polarisasi pada sel astrosit dengan kehadiran vitamin E. Kesimpulannya, kajian ini menunjukkan  $\alpha$ -tokoferol adalah lebih berkesan dan memerlukan hanya kepekatan yang rendah bagi tujuan profilaktik berbanding penggunaannya dalam pasca rawatan di dalam sel primer astrosit

## ACKNOWLEDGEMENTS

I humbly thank Allah Almighty, the Beneficent and the Merciful, who gave me health, thoughts and co-operative people to enable me to achieve this goal.

This work would not be accomplished without the help of so many people. In the following lines is a brief account of some but not all who deserve my thanks.

First, I offer my sincerest gratitude to my supervisor, Dr. Huzwah Khaza' ai for taking the burden of supervising this research. Her theoretical insight and her relentless enthusiasm assisting in creating such inspiration to make this direction and finally, to succeed in this study. Without her, this thesis would not have been completed.

I am also delighted to convey my appreciations to my co-supervisor, Assoc. Prof. Dr. Sharmili Vidyadaran, for her unwavering efforts in reviewing this thesis and other support. Appreciate from Miss Tan Shi wei for her valuable assist to complete the section 3.2.2.1 in Immunology lab.

My warmest gratitude goes to all my friends: Dr. Atabak Kheirkhah, Dr. Hamid Reza Abolkheir, Dr. Kambiz Yousefi Talooki and Dr. Thilaga Rati Selvaraju for their endless patient, support, understanding and encouragement.

My heartfelt thanks go to all my family member, especially my parents Mr.Hossein Abedi and Mrs.Fatima Abedi for their endless love and support in all my life, with all nice memory that I could never forget.



I certify that a Thesis Examination Committee has met on 2 March 2017 to conduct the final examination of Zahra Abedi on her thesis entitled "Modulation of NMDA and AMPA/Kainate Receptors by Tocotrienol-Rich Fraction and  $\alpha$ -Tocopherol in Glutamate-Induced Injury of Primary Astrocytes" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Suhaili binti Abu Bakar @ Jamaludin, PhD**

Senior Lecturer  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Chairman)

**Abdah binti Md Akim, PhD**

Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Junedah binti Sanusi, PhD**

Associate Professor  
Universiti Malaya  
Malaysia  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 2 June 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Huzwah Khaza'ai, PhD**

Senior Lecturer  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Chairman)

**Sharmili Vidyadaran, PhD**

Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice- Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in
- electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Zahra Abedi, GS35289

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature:  
Name of  
Chairman of  
Supervisory  
Committee:



DR. HUZWAH KHA'AI

**DR. HUZWAH KHAZA'AI**  
DEPARTMENT OF BIOMEDICAL SCIENCES  
FACULTY OF MEDICINE AND HEALTH SCIENCES  
UNIVERSITI PUTRA MALAYSIA  
43400 SERDANG SELANGOR  
(PHD, UNIVERSITY OF ABERDEEN)  
TEL: 03-8947 2436 FAX: 03-8943 6178

Signature:  
Name of  
Member of  
Supervisory  
Committee:



Sharmili Vidyadaran

**DR. SHARMILI VIDYADARAN**  
Associate Professor  
Immunology Unit, Department of Pathology  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
43400 Serdang, Malaysia.

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF APPENDICES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Background of the study	1
1.2 Research Problem	3
1.3 Research Objectives	3
1.3.1 General Objective	3
1.3.2 Specific Objectives	3
1.4 Hypotheses	4
<b>2 LITERATURE REVIEW</b>	<b>5</b>
2.1 Vitamin E	5
2.1.1 Tocopherols	5
2.1.2 Tocotrienols	7
2.2 Neurodegenerative diseases	12
2.2.1 Alzheimer's disease (AD)	12
2.2.2 Parkinson's disease (PD)	13
2.2.3 Other neurodegenerative disorders	14
2.3 Glial cells	14
2.3.1 Importance of interaction between neurons and astrocytes	15
2.4 Glutamate	17
2.4.1 Glutamate receptors	17
2.4.2 Inotropic glutamate receptors	18
2.4.3 Glutamate receptors gene	20
2.4.4 Glutamate transporters	21
2.4.5 Glutamate excitotoxicity	22
2.4.6 Glutamate homeostasis	23
2.4.7 Glutamate metabolism	24
2.5 Calcium ion (Ca <sup>2+</sup> ) influx and mitochondrial dysfunction	26
2.6 Modes of cell death in astrocytes cells: Apoptosis and necrosis	27
2.7 Biological markers in astrocytes cells	28

<b>3</b>	<b>MATERIALS AND METHODS</b>	<b>30</b>
3.1	Materials	30
3.1.1	Cell culture	30
3.2	Methods	31
3.2.1	Chemical and solution preparation	31
3.2.2	Isolation and purification of primary cells	32
3.2.3	Cells culture	34
3.2.4	Flow Cytometry	37
3.2.5	Culturing of astrocytes in 96-multiwell plate	37
3.2.6	MTT assay	38
3.2.7	MMP assay	39
3.2.8	Real-time PCR	40
3.2.9	Calcium assay Kit (colorimetric)	44
<b>4</b>	<b>RESULTS</b>	<b>46</b>
4.1	Isolation and purification of primary astrocytes	46
4.2	Cell viability study (MTT assay)	48
4.2.1	Dose response of glutamate in astrocytes	48
4.2.2	Vitamin E pre-treatment	48
4.2.3	Vitamin E post-treatment	49
4.3	Mitochondrial membrane potential study (MMP assay)	50
4.3.1	Vitamin E pre-treatment	51
4.3.2	Vitamin E post-treatment	51
4.4	Effect of vitamin E on the expression of ionotropic glutamate receptors genes	52
4.4.1	RT-PCR standard curve	52
4.4.2	GRIA2 gene (AMPA glutamate receptor gene)	53
4.4.3	GRIK1 gene (Kainat glutamate receptor gene)	55
4.4.4	GRIN2A gene (NMDA glutamate receptor gene)	57
4.5	Ca <sup>2+</sup> ion assay	59
4.5.1	Vitamin E pre-treatment	59
4.5.2	Vitamin E post-treatment	60
<b>5</b>	<b>DISCUSSION</b>	<b>62</b>
5.1	Isolation and purification of primary astrocytes	62
5.2	Cell viability upon glutamate toxicity	62
5.2.1	Protective effect of vitamin E on glutamate-induced injury	63
5.3	Survivability of glutamate-induced injury cells by Vitamin E	65
5.4	Expression of glutamate receptor gene upon glutamate toxicity	66
5.5	Calcium (Ca <sup>2+</sup> ) influx in primary astrocytes upon glutamate toxicity	69
<b>6</b>	<b>CONCLUSION</b>	<b>72</b>
6.1	Recommendation for future study	72

<b>REFERENCES</b>	74
<b>APPENDICES</b>	103
<b>BIODATA OF STUDENT</b>	114
<b>PUBLICATION</b>	115



## LIST OF TABLES

Table		Page
2.1	The term vitamin E includes four tocopherols ( $\alpha$ , $\beta$ , $\gamma$ , $\delta$ ) and four tocotrienols ( $\alpha$ , $\beta$ , $\gamma$ , $\delta$ ). Source: (Sen et al., 2006)	6
3.1	Experimental design of vitamin E treatments against glutamate injury	38
3.2	Amount of Master Mix for 1 Reaction of Real-time PCR	42





## LIST OF FIGURES

Figure		Page
2.1	Chemical structures of vitamin E analogs. Source: (Sen et al., 2006)	6
2.2	Glutamate homeostasis between Astrocytes and Neuron.	25
3.1	Appearance of the hemacytometer grid visualized under microscope	36
4.1	Phase contrast micrographs of different culture.	46
4.2	GFAP immunophenotyping to determining purity of astrocyte culture.	47
4.3	Effect of various concentration of glutamate on primary astrocytes cells.	48
4.4	Effects of TRF and $\alpha$ -TCP pre-treatment against 100 mM glutamate on primary astrocytes viability.	49
4.5	Effects of TRF and $\alpha$ -TCP post-treatment against 100 mM glutamate on primary astrocyte viability.	50
4.6	Effects of TRF and $\alpha$ -TCP pre-treatment against 100 mM glutamate on primary astrocytes viability in terms of MMP.	51
4.7	Effects of TRF and $\alpha$ -TCP post-treatment against 100 mM glutamate on primary astrocytes viability in terms of MMP.	52
4.8	Representative standard curve for ACTB plotted based on serial dilution of cDNA against Cq value obtained during amplification of each dilution series.	53
4.9	GRIA2 gene expression in primary astrocytes with TRF and $\alpha$ -TCP pre-treatment against glutamate insult.	54
4.10	GRIA2 expression in primary astrocytes with TRF and $\alpha$ -TCP post-treatment against glutamate insult.	55
4.11	GRIK1 gene expression in primary astrocytes with TRF and $\alpha$ -TCP pre-treatment against glutamate insult.	56
4.12	GRIK1 expression in primary astrocytes with TRF and $\alpha$ -TCP post-treatment against glutamate insult.	57
4.13	GRIN2A gene expression in primary astrocytes with TRF and $\alpha$ -TCP pre-treatment against glutamate insult.	58
4.14	GRIN2A expression in primary astrocytes with TRF and $\alpha$ -TCP post-treatment against glutamate insult.	59
4.15	Effects of TRF and $\alpha$ -TCP pre-treatment against 100 mM glutamate on primary astrocytes viability in terms of $\text{Ca}^{2+}$ ion assay.	60
4.16	Effects of TRF and $\alpha$ -TCP post-treatment against 100 mM glutamate on primary astrocytes viability in terms of $\text{Ca}^{2+}$ ion assay.	61
5.1	Summary of process involved glutamate-induce cell death. The figure has been created based on this study.	70

- 5.2 Schematic representation of role of vitamin E in prevention of Cell death induced by elevated concentration. The figure has been created based on this study.

71



## LIST OF APPENDICES

<b>Appendix</b>		<b>Page</b>
A	Dose response of glutamate in primary astrocytes	103
B	Protective effect of vitamin E on glutamate-induced injury	104
C	Survivability of glutamate-induced injury cells by vitamin E	105
D	Expression of glutamate receptors gene upon glutamate toxicity	106
E	Calcium influx in primary astrocyte upon glutamate toxicity	109
F	GFAP Immunophenotyping to determine purity of astrocyte culture	110
G	Real-time PCR supplementary data	111

## LIST OF ABBREVIATIONS

AB	Amyloid beta
ROS	Reactive oxygen species
NOS	Nitric oxide synthetase
PARP	Poly ADP ribose polymerase
AIF	Apoptosis- inducing factor
TRF	Palm tocotrienol-rich fraction
$\alpha$ -TCP	A-tocopherol
HGA	Homogentisate
HPT	HGA phytyltransferase
GSH	Glutathione
DMBA	7, 12-dimethyl benz[ $\alpha$ ] anthracene
EBV	Epstein-Barr virus
LDL	Low-density lipoprotein
HMG-CoA	3-hydroxy-3-methylglutaryl coA
RNS	Reactive nitrogen species
ALS	Amyotropic lateral sclerosis
HUVEC	Human umbilical vein endothelial cells
TBARS	Thiobarbituric acid reactive substances
12-Lox	12-lipoxygenase
AD	Alzheimer's disease
NFT	Neurofibrillary tangles
APP	Amyloid precursor protein
BBB	The blood-brain barrier
CNS	Central nervous system
NSAIDS	Non-steroidal anti-inflammatory drugs
SNpc	Substantia nigra pars compact
HD	Huntington's disease
ALS	Amyotropic lateral sclerosis
PNS	Peripheral nervous system
GABA	$\Gamma$ -aminobutyric acid
ATP	Adenosine triphosphate
AMPA	A-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
NMDA	N-methyl-D-aspartate
GSH	Glutathione
TAC	Tricarboxylic acid
MPT	Mitochondrial permeability transition
NSE	Neuron specific enolase
GFAP	Glialfibrillary acidic protein
MBP	Myelin basic protein
TBI	Traumatic brain injury
DMEM	Dulbecco's modified eagle medium
DMSO	Dimethyl sulfoxide
Rlio 123	Dye rhodamine 123
Day in vitro	Div
GFAP	Glial fibrillary acidic protein
BSO	Buthionine sulfoximine
AHS	Ammon's horn sclerosis
iGluR	Ionotropic glutamate receptor

# CHAPTER 1

## INTRODUCTION

### 1.1 Background of the study

Neurodegenerative diseases such as Alzheimer's disease and Parkinson as well as stroke are some of the common diseases among the aged individuals. In neurodegenerative diseases in mammals, cell death occurs in the brain due to two reasons: either the accumulation of amyloid beta (AB) protein or oxidative glutamate toxicity that causes neuronal cell death (Murphy et al., 1990).

The endogenous chemicals which transmit a signal from one neuron to another neuron across a synapse are referred to as neurotransmitters. These transmitters bind to a receptor in postsynaptic membrane where depolarization is caused by an excitatory neurotransmitter. In contrast, hyper-polarization in the postsynaptic membrane occurs as a result of an inhibitory neurotransmitter. It is assumed that glutamate, which is the main amino-acid neurotransmitter stimulant in the mammals' central nervous system can be excitotoxic at higher concentration, which mainly responsible for neurodegenerative diseases.

Glutamate which is considered as an important excitatory amino acid neurotransmitter plays a key role in the mammals' central nervous system. It can be excitotoxic and is assumed to be a cause of most types of neurodegenerative diseases (Zou & Crews, 2005). According to (Coyle & Puttfarcken, 1993), glutamate toxicity is a kind of brain-cell death which is associated with many types of chronic neurodegenerative diseases such as Alzheimer's, Parkinson's, and Huntington's diseases.

There are two reasons for the occurrence of glutamate toxicity: receptor-originated excitotoxicity and non-receptor mediated toxicity. In the former type, excitotoxicity is caused by the activation of glutamate receptors, while in the latter it is the result of a series of disturbances, which reduce and oxide homeostasis in the cells. There are two main types of glutamate receptors: ionotropic receptors that are ligand-gated ion channels and metabotropic receptors, which are coupled with second messenger systems by G proteins (Hollmann & Heinemann, 1994). Cell death caused by excitotoxicity is due to the lack of three functional families of ionotropic glutamate receptors known as N-Methyl-D- Aspartate (NMDA) receptor,  $\alpha$ -Amino-3-Hydroxy-5-Methyl-4- isoxazolepropionic acid (AMPA) receptor and Kainate receptor. These receptors are considerably penetrable to  $\text{Ca}^{2+}$  ions as well as  $\text{Na}^+$  and  $\text{K}^+$ . Thus, the increase in  $[\text{Ca}^{2+}]$  through this receptor-gated channels is essential for physiological function and Excitotoxicity (Choi et al., 2005; Singh et al., 2003).

When a neuron and glial cell injury occur, as a result of their deprivation of oxygen and glucose, ATP decreases rapidly and becomes depolarized which leads to the release of glutamate from nerve terminals and eventually caused reduction of glutamate uptake by glial cells (SIESJÖ et al., 1995).

Additionally, glutamate accumulation over-excites the post-synaptic NMDA and AMPA receptors leading to an intracellular overload of  $\text{Ca}^{2+}$  (Sattler & Tymianski, 2001). Furthermore, increased of  $\text{Ca}^{2+}$  uptake into mitochondria due to strong loading, which can disrupt their function and lead to the release of damaging reactive oxygen species (ROS)(Dowling, 2001).  $\text{Ca}^{2+}$  loading may also lead to the generation of ROS through other mechanisms, such as the activation of nitric oxide synthetase (NOS), resulting in nitric oxide production, and the activation of NADPH-oxidase producing superoxide in mitochondria (Brennan et al., 2009). The generated ROS can induce DNA damage, which causes the activation of Poly ADP ribose polymerase (PARP) enzyme, capable of injuring neurons via  $\text{NAD}^+$  depletion, causing glycolytic block, ATP depletion, and induction of apoptotic signaling pathway through the release of apoptosis- inducing factor (AIF) from mitochondria (Alano et al., 2010).

In the case of a cerebral insult, the role of astrocytes in improving neuronal survival and recovery is increasingly recognized. The large body of recent studies on neuroprotection has focused on improving the neuron survival (Chang et al., 2009); however, a main simultaneous effect of ischemic infarction is the death of glia, especially astrocytes.

When an astrocyte is damaged, reduction of astrocytes function can lead to loss of CNS and the overall pathology related to the excitatory amino acids (Chen et al., 2000). Thus, it is essential to identify the responsible mechanisms in astrocytes death after insults and it should result in developing strategies with the specific aim of promoting astrocytes survival. Many studies have shown astrocytes damage after neurodegenerative disorders (Landis, 1994; Liu et al., 1999). For example, a study by Chen (2000)(Chen et al., 2000) showed that stimulating astrocytes with glutamate cause cell swelling and cell death.

With regard to the significance of the astrocyte survival in the occurrence of glutamate insult, it is highly desirable to find treatments to prevent excitotoxicity that causes cell death. It has been shown that astrocytes treated with vitamin E were able to resist glutamate excitotoxicity (Chen et al., 2001). The neuroprotective properties of Vitamin E composed of eight different isoforms, four tocopherols ( $\alpha$ -, $\beta$ -, $\gamma$ -, $\delta$ -) and four tocotrienols ( $\alpha$ -, $\beta$ -, $\gamma$ -, $\delta$ -) have been identified (Aggarwal et al., 2010).

Structurally, tocopherols and tocotrienols are different, in which the former possesses a saturated phytol tail, while the latter has an unsaturated isoprenoid side chain. Palm tocotrienol-rich fraction (TRF) which is extracted from palm oil contains 75% tocotrienols and 25%  $\alpha$ -tocopherol. It was illustrated that TRF has potent antioxidant (Maniam et al., 2008; Serbinova & Packer, 1994), anti-inflammatory (Wu et al., 2008), anticancer (Goh et al., 1994; Takahashi & Loo, 2004; WU & Ng, 2007),

neuroprotection (Osakada et al., 2004; Sen et al., 2000) and cholesterol-lowering effects (Minhajuddin et al., 2005; Mutalib et al., 2003; Qureshi et al., 1995).

Most of the studies have focused on the neuronal cell protection rather than astrocytes (Pettmann & Henderson, 1998; Wang et al., 2005). Hence, the present study attempts to investigate the effects of tocotrienols and tocopherols in preventing glutamate excitotoxicity in astrocytes. Besides promoting astrocytes survival, advanced neuroprotection would be expected. The prophylactic and preventive properties of tocotrienols and tocopherol in neurodegeneration are expected to be achieved and an alternative therapeutics based on nutrition would be offered.

## **1.2 Research Problem**

Glutamate excitotoxicity in astrocytes cells leads to the release of  $\text{Ca}^{2+}$  and reactivation of free radicals by over-activation of glutamate receptors that eventually cause neuronal neurodegeneration and cell death. Controlling the receptor activity and inhibition of  $\text{Ca}^{2+}$  release of glutamate and free radicals from astrocyte cells may be an effective method for preventing neurodegeneration. Does neuroprotective action of tocotrienol and tocopherols prevent astrocyte cells death induced glutamate?

## **1.3 Research Objectives**

### **1.3.1 General Objective**

To determine the role of tocotrienols and tocopherols in the prevention of glutamate toxicity in injured primary astrocytes.

### **1.3.2 Specific Objectives**

- To determine the  $\text{IC}_{20}$  and  $\text{IC}_{50}$  values of glutamate in primary astrocyte cells.
- To determine the effect of tocotrienols and tocopherols on cell viability in primary astrocytes injured by glutamate.
- To determine the effect of tocotrienols and tocopherols on NMDA receptors in primary astrocytes injured by glutamate.
- To determine the effect of tocotrienol and tocopherols on the activation of  $\text{Ca}^{2+}$  permeability in AMPA/Kainate receptors in primary astrocytes injured by glutamate.
- To determine the effect of tocotrienol and tocopherols on  $\text{Ca}^{2+}$  ion influx in primary astrocytes injured by glutamate.

#### 1.4 Hypotheses

Tocotrienols and tocopherols are capable of modulating glutamate receptors and  $\text{Ca}^{2+}$  influx in increasing survivability of primary astrocytes.





## REFERENCES

- Abramov, A. Y., & Duchen, M. R. (2008). Mechanisms underlying the loss of mitochondrial membrane potential in glutamate excitotoxicity. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1777(7), 953-964.
- Aggarwal, B. B., Sundaram, C., Prasad, S., & Kannappan, R. (2010). Tocotrienols, the vitamin E of the 21st century: its potential against cancer and other chronic diseases. *Biochemical pharmacology*, 80(11), 1613-1631.
- Agulhon, C., Petravicz, J., McMullen, A. B., Sweger, E. J., Minton, S. K., Taves, S. R., . . . McCarthy, K. D. (2008). What is the role of astrocyte calcium in neurophysiology? *Neuron*, 59(6), 932-946.
- Akopian, G., Kuprijanova, E., Kressin, K., & Steinhäuser, C. (1997). Analysis of ion channel expression by astrocytes in red nucleus brain stem slices of the rat. *Glia*, 19(3), 234-246.
- Al-Hallaq, R. A., Jarabek, B. R., Fu, Z., Vicini, S., Wolfe, B. B., & Yasuda, R. P. (2002). Association of NR3A with the N-Methyl-D-aspartate Receptor NR1 and NR2 Subunits. *Molecular pharmacology*, 62(5), 1119-1127.
- Alano, C. C., Garnier, P., Ying, W., Higashi, Y., Kauppinen, T. M., & Swanson, R. A. (2010). NAD<sup>+</sup> Depletion Is Necessary and Sufficient for Poly (ADP-Ribose) Polymerase-1-Mediated Neuronal Death. *The Journal of Neuroscience*, 30(8), 2967-2978.
- Alberdi, E., Sánchez-Gómez, M. a. V., Marino, A., & Matute, C. (2002). Ca<sup>2+</sup> influx through AMPA or kainate receptors alone is sufficient to initiate excitotoxicity in cultured oligodendrocytes. *Neurobiology of disease*, 9(2), 234-243.
- Allen, N. J., & Barres, B. A. (2005). Signaling between glia and neurons: focus on synaptic plasticity. *Current opinion in neurobiology*, 15(5), 542-548.
- Ames, A. (2000). CNS energy metabolism as related to function. *Brain Research Reviews*, 34(1), 42-68.
- Anderson, C. M., & Swanson, R. A. (2000). Astrocyte glutamate transport: review of properties, regulation, and physiological functions. *Glia*, 32(1), 1-14.
- Ankarcrona, M., Dypbukt, J. M., Bonfoco, E., Zhivotovsky, B., Orrenius, S., Lipton, S. A., & Nicotera, P. (1995). Glutamate-induced neuronal death: a succession of necrosis or apoptosis depending on mitochondrial function. *Neuron*, 15(4), 961-973.
- Aoki, C., Venkatesan, C., Go, C., Mong, J. A., & Dawson, T. M. (1994). Cellular and subcellular localization of NMDA-R1 subunit immunoreactivity in the visual

cortex of adult and neonatal rats. *The Journal of neuroscience*, 14(9), 5202-5222.

- Araque, A., Parpura, V., Sanzgiri, R. P., & Haydon, P. G. (1999). Tripartite synapses: glia, the unacknowledged partner. *Trends in neurosciences*, 22(5), 208-215.
- Arriza, J. L., Fairman, W. A., Wadiche, J. I., Murdoch, G. H., Kavanaugh, M. P., & Amara, S. G. (1994). Functional comparisons of three glutamate transporter subtypes cloned from human motor cortex. *The Journal of neuroscience*, 14(9), 5559-5569.
- Ashfaq, M., Zuberi, H. S., & Waqar, M. A. (2000). Vitamin E and  $\beta$ -carotene affect natural killer cell function. *International journal of food sciences and nutrition*, 51(sup1), s13-s20.
- Atlante, A., Calissano, P., Bobba, A., Giannattasio, S., Marra, E., & Passarella, S. (2001). Glutamate neurotoxicity, oxidative stress and mitochondria. *FEBS letters*, 497(1), 1-5.
- Auger, C., & Attwell, D. (2000). Fast removal of synaptic glutamate by postsynaptic transporters. *Neuron*, 28(2), 547-558.
- Aurell, A., Rosengren, L. E., Karlsson, B., Olsson, J.-E., Zbornikova, V., & Haglid, K. G. (1991). Determination of S-100 and glial fibrillary acidic protein concentrations in cerebrospinal fluid after brain infarction. *Stroke*, 22(10), 1254-1258.
- Backus, K. H., Elsen, F. P., & Schulze, C. H. (1998). Effect of kainate on the membrane conductance of hilar glial precursor cells recorded in the perforated-patch configuration. *Glia*, 23(1), 35-44.
- Baltrons, M. A., & Garcia, A. (1997). AMPA receptors are coupled to the nitric oxide/cyclic GMP pathway in cerebellar astroglial cells. *European Journal of Neuroscience*, 9(11), 2497-2501.
- Bandyopadhyay, S., Hennes, H., Gorelick, M. H., Wells, R. G., & Walsh-Kelly, C. M. (2005). Serum Neuron-specific Enolase as a Predictor of Short-term Outcome in Children with Closed Traumatic Brain Injury. *Academic Emergency Medicine*, 12(8), 732-738.
- Bartłomiej, S., Justyna, R.-K., & Ewa, N. (2012). Bioactive compounds in cereal grains—occurrence, structure, technological significance and nutritional benefits—a review. *Food science and technology international*, 18(6), 559-568.
- Beal, M. F. (1996). Mitochondria, free radicals, and neurodegeneration. *Current opinion in neurobiology*, 6(5), 661-666.
- Beems, T., Simons, K., Van Geel, W., De Reus, H., Vos, P., & Verbeek, M. (2003). Serum-and CSF-concentrations of brain specific proteins in hydrocephalus. *Acta neurochirurgica*, 145(1), 37-43.

- Behl, C., Davis, J., Cole, G. M., & Schubert, D. (1992). Vitamin E protects nerve cells from amyloid  $\beta$ protein toxicity. *Biochemical and biophysical research communications*, 186(2), 944-950.
- Bellucci, A., Zaltieri, M., Navarria, L., Grigoletto, J., Missale, C., & Spano, P. (2012). From  $\alpha$ -synuclein to synaptic dysfunctions: new insights into the pathophysiology of Parkinson's disease. *Brain research*, 1476, 183-202.
- Benarroch, E. E. (2005). *Neuron-astrocyte interactions: partnership for normal function and disease in the central nervous system*. Paper presented at the Mayo Clinic Proceedings.
- Berger, U. V., DeSilva, T. M., Chen, W., & Rosenberg, P. A. (2005). Cellular and subcellular mRNA localization of glutamate transporter isoforms GLT1a and GLT1b in rat brain by in situ hybridization. *Journal of Comparative Neurology*, 492(1), 78-89.
- Bergles, D. E., Dzubay, J. A., & Jahr, C. E. (1997). Glutamate transporter currents in bergmann glial cells follow the time course of extrasynaptic glutamate. *Proceedings of the National Academy of Sciences*, 94(26), 14821-14825.
- Bergles, D. E., Roberts, J. D. B., Somogyi, P., & Jahr, C. E. (2000). Glutamatergic synapses on oligodendrocyte precursor cells in the hippocampus. *Nature*, 405(6783), 187-191.
- Bernard, A., Ferhat, L., Dessi, F., Charton, G., Represa, A., Ben-Ari, Y., & Khrestchatsky, M. (1999). Q/R editing of the rat GluR5 and GluR6 kainate receptors in vivo and in vitro: evidence for independent developmental, pathological and cellular regulation. *European Journal of Neuroscience*, 11(2), 604-616.
- Bezzi, P., Carmignoto, G., Pasti, L., Vesce, S., Rossi, D., Rizzini, B. L., . . . Volterra, A. (1998). Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature*, 391(6664), 281-285.
- Bignami, A., & Dahl, D. (1974). Astrocyte-specific protein and radial glia in the cerebral cortex of newborn rat. *Nature*, 252, 55-56.
- Bignami, A., Eng, L., Dahl, D., & Uyeda, C. (1972). Localization of the glial fibrillary acidic protein in astrocytes by immunofluorescence. *Brain research*, 43(2), 429-435.
- Bonfoco, E., Krainc, D., Ankarcrona, M., Nicotera, P., & Lipton, S. A. (1995). Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with N-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proceedings of the National Academy of Sciences*, 92(16), 7162-7166.
- Borlongan, C. V., Nishino, H., & Sanberg, P. R. (1997). Systemic, but not intraparenchymal, administration of 3-nitropropionic acid mimics the

neuropathology of Huntington's disease: a speculative explanation. *Neuroscience research*, 28(3), 185-189.

Boyce-Rustay, J. M., & Holmes, A. (2006). Genetic inactivation of the NMDA receptor NR2A subunit has anxiolytic-and antidepressant-like effects in mice. *Neuropsychopharmacology*, 31(11), 2405-2414.

Brennan, A. M., Suh, S. W., Won, S. J., Narasimhan, P., Kauppinen, T. M., Lee, H., . . . Swanson, R. A. (2009). NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation. *Nature neuroscience*, 12(7), 857-863.

Brigelius-Flohe, R., & Traber, M. G. (1999). Vitamin E: function and metabolism. *The FASEB Journal*, 13(10), 1145-1155.

Brodal, P. (2004). *The central nervous system: structure and function*: Oxford University Press.

Brown, A. M., & Ransom, B. R. (2007). Astrocyte glycogen and brain energy metabolism. *Glia*, 55(12), 1263-1271.

Brugge, J. S., Cotton, P. C., Queral, A., Barrett, J. N., Nonner, D., & Keane, R. W. (1985). Neurones express high levels of a structurally modified, activated form of pp60c-src.

Bruno, R. S., Leonard, S. W., Park, S.-i., Zhao, Y., & Traber, M. G. (2006). Human vitamin E requirements assessed with the use of apples fortified with deuterium-labeled  $\alpha$ -tocopheryl acetate. *The American journal of clinical nutrition*, 83(2), 299-304.

Burnashev, N. (1996). Calcium permeability of glutamate-gated channels in the central nervous system. *Current opinion in neurobiology*, 6(3), 311-317.

Burnashev, N., Monyer, H., Seeburg, P. H., & Sakmann, B. (1992). Divalent ion permeability of AMPA receptor channels is dominated by the edited form of a single subunit. *Neuron*, 8(1), 189-198.

Busciglio, J., & Yankner, B. A. (1994). Apoptosis and increased generation of reactive oxygen species in Down's syndrome neurons in vitro. *Nature*, 378(6559), 776-779.

Butcher, S. P., & Hamberger, A. (1987). In Vivo Studies on the Extracellular, and Veratrine-Releasable, Pools of Endogenous Amino Acids in the Rat Striatum: Effects of Corticostriatal Deafferentation and Kainic Acid Lesion. *Journal of neurochemistry*, 48(3), 713-721.

Butterfield, D. A., & Sultana, R. (2007). Redox proteomics identification of oxidatively modified brain proteins in Alzheimer's disease and mild cognitive impairment: insights into the progression of this dementing disorder. *Journal of Alzheimer's Disease*, 12(1), 61-72.



- Byeon, S. E., Yi, Y.-S., Oh, J., Yoo, B. C., Hong, S., & Cho, J. Y. (2012). The role of Src kinase in macrophage-mediated inflammatory responses. *Mediators of inflammation*, 2012.
- Cao, G., Minami, M., Pei, W., Yan, C., Chen, D., O'Horo, C., . . . Chen, J. (2001). Intracellular Bax Translocation After Transient Cerebral Ischemia&colon; Implications for a Role of the Mitochondrial Apoptotic Signaling Pathway in Ischemic Neuronal Death. *Journal of Cerebral Blood Flow & Metabolism*, 21(4), 321-333.
- Cauley, K., Kukekov, V., & Young, D. (1997). Kainate/AMPA receptors expressed on human fetal astrocytes in long-term culture. *Journal of neuroscience research*, 47(3), 311-321.
- Chang, P. N., Yap, W. N., Wing Lee, D. T., Ling, M., Wong, Y. C., & Yap, Y. L. (2009). Evidence of  $\gamma$ -tocotrienol as an apoptosis-inducing, invasion-suppressing, and chemotherapy drug-sensitizing agent in human melanoma cells. *Nutrition and cancer*, 61(3), 357-366.
- Chaudhry, F. A., Lehre, K. P., van Lookeren Campagne, M., Ottersen, O. P., Danbolt, N. C., & Storm-Mathisen, J. (1995). Glutamate transporters in glial plasma membranes: highly differentiated localizations revealed by quantitative ultrastructural immunocytochemistry. *Neuron*, 15(3), 711-720.
- Chen, C. J., Liao, S. L., & Kuo, J. S. (2000). Gliotoxic action of glutamate on cultured astrocytes. *Journal of neurochemistry*, 75(4), 1557-1565.
- Chen, H., Zhang, S., Schwarzschild, M., Hernan, M., & Ascherio, A. (2005). Physical activity and the risk of Parkinson disease. *Neurology*, 64(4), 664-669.
- Chen, H., Zhang, S. M., Hernán, M. A., Schwarzschild, M. A., Willett, W. C., Colditz, G. A., . . . Ascherio, A. (2003). Nonsteroidal anti-inflammatory drugs and the risk of Parkinson disease. *Archives of Neurology*, 60(8), 1059-1064.
- Chen, W., Mahadomrongkul, V., Berger, U. V., Bassan, M., DeSilva, T., Tanaka, K., . . . Rosenberg, P. A. (2004). The glutamate transporter GLT1a is expressed in excitatory axon terminals of mature hippocampal neurons. *The Journal of neuroscience*, 24(5), 1136-1148.
- Chen, Y., Vartiainen, N. E., Ying, W., Chan, P. H., Koistinaho, J., & Swanson, R. A. (2001). Astrocytes protect neurons from nitric oxide toxicity by a glutathione-dependent mechanism. *Journal of neurochemistry*, 77(6), 1601-1610.
- Choi, D.-K., Pennathur, S., Perier, C., Tieu, K., Teismann, P., Wu, D.-C., . . . Heinecke, J. W. (2005). Ablation of the inflammatory enzyme myeloperoxidase mitigates features of Parkinson's disease in mice. *The Journal of neuroscience*, 25(28), 6594-6600.
- Choi, D. W. (1992). Excitotoxic cell death. *Journal of neurobiology*, 23(9), 1261-1276.

- Ciani, E., Grøneng, L., Voltattorni, M., Rolseth, V., Contestabile, A., & Paulsen, R. E. (1996). Inhibition of free radical production or free radical scavenging protects from the excitotoxic cell death mediated by glutamate in cultures of cerebellar granule neurons. *Brain research*, 728(1), 1-6.
- Condorelli, D., Dell'Albani, P., Corsaro, M., Barresi, V., & Stella, A. (1993). AMPA-Selective glutamate receptor subunits in astroglial cultures. *Journal of neuroscience research*, 36(3), 344-356.
- Conti, F., Barbaresi, P., Melone, M., & Ducati, A. (1999). Neuronal and glial localization of NR1 and NR2A/B subunits of the NMDA receptor in the human cerebral cortex. *Cerebral cortex*, 9(2), 110-120.
- Conti, F., Minelli, A., & Brecha, N. C. (1994). Cellular localization and laminar distribution of AMPA glutamate receptor subunits mRNAs and proteins in the rat cerebral cortex. *Journal of Comparative Neurology*, 350(2), 241-259.
- Conti, F., & Weinberg, R. J. (1999). Shaping excitation at glutamatergic synapses. *Trends in neurosciences*, 22(10), 451-458.
- Cornell-Bell, A. H., Thomas, P. G., & Smith, S. J. (1990). The excitatory neurotransmitter glutamate causes filopodia formation in cultured hippocampal astrocytes. *Glia*, 3(5), 322-334.
- Coyle, J. T., & Puttfarcken, P. (1993). Oxidative stress, glutamate, and neurodegenerative disorders. *Science*, 262(5134), 689-695.
- Cui, X., Wang, L., Zuo, P., Han, Z., Fang, Z., Li, W., & Liu, J. (2004). D-galactose-caused life shortening in *Drosophila melanogaster* and *Musca domestica* is associated with oxidative stress. *Biogerontology*, 5(5), 317-325.
- Daikhin, Y., & Yudkoff, M. (2000). Compartmentation of brain glutamate metabolism in neurons and glia. *The Journal of nutrition*, 130(4), 1026S-1031S.
- Danbolt, N. C. (2001). Glutamate uptake. *Progress in neurobiology*, 65(1), 1-105.
- Danial, N. N., & Korsmeyer, S. J. (2004). Cell death: critical control points. *Cell*, 116(2), 205-219.
- Daniels, B. A. (2011). *Modulation of ionotropic glutamate receptors in retinal neurons by the amino acid D-serine*. Dalhousie University Halifax.
- Dauer, W., & Przedborski, S. (2003). Parkinson's disease: mechanisms and models. *Neuron*, 39(6), 889-909.
- David, J. C., Yamada, K. A., Bagwe, M. R., & Goldberg, M. P. (1996). AMPA receptor activation is rapidly toxic to cortical astrocytes when desensitization is blocked. *The Journal of neuroscience*, 16(1), 200-209.

- Davis, J. B., & Maher, P. (1994). Protein kinase C activation inhibits glutamate-induced cytotoxicity in a neuronal cell line. *Brain research*, 652(1), 169-173.
- Dawson, T. M., & Dawson, V. L. (2003). Molecular pathways of neurodegeneration in Parkinson's disease. *Science*, 302(5646), 819-822.
- Day, N. C., Williams, T. L., Ince, P. G., Kamboj, R. K., Lodged, D., & Shaw, P. J. (1995). Distribution of AMPA-selective glutamate receptor subunits in the human hippocampus and cerebellum. *Molecular brain research*, 31(1), 17-32.
- Dingledine, R., Borges, K., Bowie, D., & Traynelis, S. F. (1999). The glutamate receptor ion channels. *Pharmacological reviews*, 51(1), 7-62.
- DOEGE, H., BOCIANSKI, A., SCHEEPERS, A., Hubertus, A., ECKEL, J., JOOST, H.-G., & SCHÜRMAN, A. (2001). Characterization of human glucose transporter (GLUT) 11 (encoded by SLC2A11), a novel sugar-transport facilitator specifically expressed in heart and skeletal muscle. *Biochemical Journal*, 359(2), 443-449.
- Doetsch, F. (2003). The glial identity of neural stem cells. *Nature neuroscience*, 6(11), 1127-1134.
- Domercq, M., Sánchez-Gómez, M. V., Areso, P., & Matute, C. (1999). Expression of glutamate transporters in rat optic nerve oligodendrocytes. *European Journal of Neuroscience*, 11(7), 2226-2236.
- Dowling, J. E. (2001). *Neurons and networks: an introduction to behavioral neuroscience*: Harvard University Press.
- Dringen, R., Gutterer, J. M., & Hirrlinger, J. (2000). Glutathione metabolism in brain. *European Journal of Biochemistry*, 267(16), 4912-4916.
- Du, Y., Ma, Z., Lin, S., Dodel, R. C., Gao, F., Bales, K. R., . . . Nelson, D. L. (2001). Minocycline prevents nigrostriatal dopaminergic neurodegeneration in the MPTP model of Parkinson's disease. *Proceedings of the National Academy of Sciences*, 98(25), 14669-14674.
- Echols, H., & Goodman, M. F. (1991). Fidelity mechanisms in DNA replication. *Annual review of biochemistry*, 60(1), 477-511.
- Edinger, A. L., & Thompson, C. B. (2004). Death by design: apoptosis, necrosis and autophagy. *Current opinion in cell biology*, 16(6), 663-669.
- Ehlers, S., Kyllerman, M., & Rosengren, L. (1994). Analysis of glial fibrillary acidic protein in the cerebrospinal fluid of children investigated for encephalopathy. *Neuropediatrics*, 25(3), 129-133.
- Emerit, J., Edeas, M., & Bricaire, F. (2004). Neurodegenerative diseases and oxidative stress. *Biomedicine & pharmacotherapy*, 58(1), 39-46.

- Endele, S., Rosenberger, G., Geider, K., Popp, B., Tamer, C., Stefanova, I., . . . Pientka, F. K. (2010). Mutations in GRIN2A and GRIN2B encoding regulatory subunits of NMDA receptors cause variable neurodevelopmental phenotypes. *Nature genetics*, 42(11), 1021-1026.
- Engelman, H. S., & MacDermott, A. B. (2004). Presynaptic ionotropic receptors and control of transmitter release. *Nature Reviews Neuroscience*, 5(2), 135-145.
- Erecińska, M., & Silver, I. A. (1990). Metabolism and role of glutamate in mammalian brain. *Progress in neurobiology*, 35(4), 245-296.
- Fan, D., Grooms, S. Y., Araneda, R. C., Johnson, A. B., Dobrenis, K., Kessler, J. A., & Zukin, R. S. (1999). AMPA receptor protein expression and function in astrocytes cultured from hippocampus. *Journal of neuroscience research*, 57(4), 557-571.
- Farb, C., Aoki, C., & Ledoux, J. (1995). Differential localization of NMDA and AMPA receptor subunits in the lateral and basal nuclei of the amygdala: a light and electron microscopic study. *Journal of Comparative Neurology*, 362(1), 86-108.
- Fields, R. D., & Stevens-Graham, B. (2002). New insights into neuron-glia communication. *Science*, 298(5593), 556-562.
- Finiels, F., y Ribotta, M. G., Barkats, M., Samolyk, M.-L., Robert, J.-J., Privat, A., . . . Mallet, J. (1995). Specific and efficient gene transfer strategy offers new potentialities for the treatment of motor neurone diseases. *Neuroreport*, 7(1), 373-378.
- Floyd, R. A., & Hensley, K. (2002). Oxidative stress in brain aging: implications for therapeutics of neurodegenerative diseases. *Neurobiology of aging*, 23(5), 795-807.
- Forman, M. S., Trojanowski, J. Q., & Lee, V. M. (2004). Neurodegenerative diseases: a decade of discoveries paves the way for therapeutic breakthroughs. *Nature medicine*, 10(10), 1055-1063.
- Fraga, C. G., Shigenaga, M. K., Park, J.-W., Degan, P., & Ames, B. N. (1990). Oxidative damage to DNA during aging: 8-hydroxy-2'-deoxyguanosine in rat organ DNA and urine. *Proceedings of the National Academy of Sciences*, 87(12), 4533-4537.
- Froissard, P., Monrocq, H., & Duval, D. (1997). Role of glutathione metabolism in the glutamate-induced programmed cell death of neuronal-like PC12 cells. *European journal of pharmacology*, 326(1), 93-99.
- Fujimura, M., Morita-Fujimura, Y., Murakami, K., Kawase, M., & Chan, P. H. (1998). Cytosolic redistribution of cytochrome c after transient focal cerebral ischemia in rats. *Journal of Cerebral Blood Flow & Metabolism*, 18(11), 1239-1247.



- Fukamachi, S., Furuta, A., Ikeda, T., Ikenoue, T., Kaneoka, T., Rothstein, J. D., & Iwaki, T. (2001). Altered expressions of glutamate transporter subtypes in rat model of neonatal cerebral hypoxia–ischemia. *Developmental brain research*, *132*(2), 131-139.
- Furuta, A., Martin, L., Lin, C.-L., Dykes-Hoberg, M., & Rothstein, J. (1997). Cellular and synaptic localization of the neuronal glutamate transporters excitatory amino acid transporter 3 and 4. *Neuroscience*, *81*(4), 1031-1042.
- Gallo, V., Patneau, D. K., Mayer, M. L., & Vaccarino, F. M. (1994). Excitatory amino acid receptors in glial progenitor cells: molecular and functional properties. *Glia*, *11*(2), 94-101.
- Garcia-Barcina, J. M., & Matute, C. (1996). Expression of Kainate-selective Glutamate Receptor Subunits in Glial Cells of the Adult Bovine White Matter. *European Journal of Neuroscience*, *8*(11), 2379-2387.
- Geiger, J. R., Melcher, T., Koh, D.-S., Sakmann, B., Seeburg, P. H., Jonas, P., & Monyer, H. (1995). Relative abundance of subunit mRNAs determines gating and Ca<sup>2+</sup> permeability of AMPA receptors in principal neurons and interneurons in rat CNS. *Neuron*, *15*(1), 193-204.
- Gill, R., & Lodge, D. (1996). Pharmacology of AMPA antagonists and their role in neuroprotection. *Neuroprotective Agents and Cerebral Ischemia*, 197-232.
- Goh, S., Hew, N., Norhanom, A., & Yadav, M. (1994). Inhibition of tumour promotion by various palm-oil tocotrienols. *International Journal of Cancer*, *57*(4), 529-531.
- Goldberg, Y. P., Andrew, S. E., Clarke, L. A., & Hayden, M. R. (1993). A PCR method for accurate assessment of trinucleotide repeat expansion in Huntington disease. *Human molecular genetics*, *2*(6), 635-636.
- Goodman, Y., & Mattson, M. P. (1994). Secreted forms of  $\beta$ -amyloid precursor protein protect hippocampal neurons against amyloid  $\beta$ -peptide-induced oxidative injury. *Experimental neurology*, *128*(1), 1-12.
- Goodwin, B., Moore, L. B., Stoltz, C. M., McKee, D. D., & Kliewer, S. A. (2001). Regulation of the human CYP2B6 gene by the nuclear pregnane X receptor. *Molecular pharmacology*, *60*(3), 427-431.
- Gottlieb, M., & Matute, C. (1997). Expression of ionotropic glutamate receptor subunits in glial cells of the hippocampal CA1 area following transient forebrain ischemia. *Journal of Cerebral Blood Flow & Metabolism*, *17*(3), 290-300.
- Gras, G., Chrétien, F., Vallat-Decouvelaere, A. V., Le Pavec, G., Porcheray, F., Bossuet, C., . . . Clayette, P. (2003). Regulated expression of sodium-dependent glutamate transporters and synthetase: a neuroprotective role for

activated microglia and macrophages in HIV infection? *Brain pathology*, 13(2), 211-222.

Green, D. R., & Reed, J. C. (1998). Mitochondria and apoptosis. *Science*, 281(5381), 1309.

Grosche, J., Matyash, V., Möller, T., Verkhratsky, A., Reichenbach, A., & Kettenmann, H. (1999). Microdomains for neuron–glia interaction: parallel fiber signaling to Bergmann glial cells. *Nature neuroscience*, 2(2), 139-143.

Gu, J., Liu, Y., Wen, Y., Natarajan, R., Lanting, L., & Nadler, J. L. (2001). Evidence that increased 12-lipoxygenase activity induces apoptosis in fibroblasts. *Journal of cellular physiology*, 186(3), 357-365.

Had-Aissouni, L., Ré, D.-B., Nieoullon, A., & Kerkerian-Le Goff, L. (2002). Importance of astrocytic inactivation of synaptically released glutamate for cell survival in the central nervous system—are astrocytes vulnerable to low intracellular glutamate concentrations? *Journal of Physiology-Paris*, 96(3), 317-322.

Hagberg, H., Lehmann, A., Sandberg, M., Nyström, B., Jacobson, I., & Hamberger, A. (1985). Ischemia-induced shift of inhibitory and excitatory amino acids from intra-to extracellular compartments. *Journal of Cerebral Blood Flow & Metabolism*, 5(3), 413-419.

Hamby, M. E., Uliasz, T. F., Hewett, S. J., & Hewett, J. A. (2006). Characterization of an improved procedure for the removal of microglia from confluent monolayers of primary astrocytes. *Journal of neuroscience methods*, 150(1), 128-137.

Han, D., Sen, C. K., Roy, S., Kobayashi, M. S., Tritschler, H. J., & Packer, L. (1997). Protection against glutamate-induced cytotoxicity in C6 glial cells by thiol antioxidants. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 273(5), R1771-R1778.

Hanley, L., Hilary Jackson, T., Chater, P. H., & Henley, J. (2010). Glutamate receptors. *Ion Channels: From Structure to Function*, 1, 277.

He, L., Mo, H., Hadisusilo, S., Qureshi, A. A., & Elson, C. E. (1997). Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo. *The Journal of nutrition*, 127(5), 668-674.

Herrmann, M., Ebert, A. D., Galazky, I., Wunderlich, M. T., Kunz, W. S., & Huth, C. (2000). Neurobehavioral outcome prediction after cardiac surgery role of neurobiochemical markers of damage to neuronal and glial brain tissue. *Stroke*, 31(3), 645-650.

Hersch, S. M., & Rosas, H. D. (2008). Neuroprotection for Huntington's disease: ready, set, slow. *Neurotherapeutics*, 5(2), 226-236.

- Hiura, Y., Tachibana, H., Arakawa, R., Aoyama, N., Okabe, M., Sakai, M., & Yamada, K. (2009). Specific accumulation of  $\gamma$ - and  $\delta$ -tocotrienols in tumor and their antitumor effect in vivo. *The Journal of nutritional biochemistry*, 20(8), 607-613.
- Hollensworth, S. B., Shen, C.-C., Sim, J. E., Spitz, D. R., Wilson, G. L., & LeDoux, S. P. (2000). Glial cell type-specific responses to menadione-induced oxidative stress. *Free Radical Biology and Medicine*, 28(8), 1161-1174.
- Hollmann, M., & Heinemann, S. (1994). Cloned glutamate receptors. *Annual review of neuroscience*, 17(1), 31-108.
- Holzwarth, J. A., Gibbons, S. J., Brorson, J. R., Philipson, L. H., & Miller, R. J. (1994). Glutamate receptor agonists stimulate diverse calcium responses in different types of cultured rat cortical glial cells. *The Journal of neuroscience*, 14(4), 1879-1891.
- Hosein, M. (2015). Biomarkers Of Mild Traumatic Brain Injury In Cerebrospinal Fluid And Blood. *Clinical Chemistry and Laboratory Medicine*, 53, S176.
- Hosomi, A., Arita, M., Sato, Y., Kiyose, C., Ueda, T., Igarashi, O., . . . Inoue, K. (1997). Affinity for  $\alpha$ -tocopherol transfer protein as a determinant of the biological activities of vitamin E analogs. *FEBS letters*, 409(1), 105-108.
- Inage, Y., Itoh, M., Wada, K., & Takashima, S. (1998). Expression of Two Glutamate Transporters, GLAST and EAAT4, in the Human Cerebellum: Their Correlation in Development and Neonatal Hypoxic-Ischemic Damage. *Journal of Neuropathology & Experimental Neurology*, 57(6), 554-562.
- Iqbal, J., Minhajuddin, M., & Beg, Z. (2003). Suppression of 7, 12-dimethylbenz [ $\alpha$ ] anthracene-induced carcinogenesis and hypercholesterolaemia in rats by tocotrienol-rich fraction isolated from rice bran oil. *European journal of cancer prevention*, 12(6), 447-453.
- Jabs, R., Kirchhoff, F., Kettenmann, H., & Steinhäuser, C. (1994). Kainate activates  $Ca^{2+}$ -permeable glutamate receptors and blocks voltage-gated  $K^{+}$  currents in glial cells of mouse hippocampal slices. *Pflügers Archiv*, 426(3-4), 310-319.
- Jackson-Lewis, V., Vila, M., Tieu, K., Teismann, P., Vadseth, C., Choi, D.-K., . . . Przedborski, S. (2002). Blockade of microglial activation is neuroprotective in the 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine mouse model of Parkinson disease. *The Journal of Neuroscience*, 22(5), 1763-1771.
- Jose, S., Tan, S. W., Tong, C. K., & Vidyadaran, S. (2015). Isolation and characterization of primary microglia from post-natal murine brain tissues: a comparison of two methods. *Cell biology international*, 39(12), 1355-1363.
- Kalia, L. V., Kalia, S. K., & Salter, M. W. (2008). NMDA receptors in clinical neurology: excitatory times ahead. *The Lancet Neurology*, 7(8), 742-755.

- Kamat, J., & Devasagayam, T. (1995). Tocotrienols from palm oil as potent inhibitors of lipid peroxidation and protein oxidation in rat brain mitochondria. *Neuroscience letters*, 195(3), 179-182.
- Kamat, J., Sarma, H., Devasagayam, T., Nesaretnam, K., & Basiron, Y. (1997). Tocotrienols from palm oil as effective inhibitors of protein oxidation and lipid peroxidation in rat liver microsomes. *Molecular and cellular biochemistry*, 170(1-2), 131-138.
- Kato, S., Negishi, K., Mawatari, K., & Kuo, C.-H. (1992). A mechanism for glutamate toxicity in the C6 glioma cells involving inhibition of cystine uptake leading to glutathione depletion. *Neuroscience*, 48(4), 903-914.
- Khanna, S., Roy, S., Parinandi, N. L., Maurer, M., & Sen, C. K. (2006). Characterization of the potent neuroprotective properties of the natural vitamin E  $\alpha$ -tocotrienol. *Journal of neurochemistry*, 98(5), 1474-1486.
- Khanna, S., Roy, S., Park, H.-A., & Sen, C. K. (2007). Regulation of c-Src activity in glutamate-induced neurodegeneration. *Journal of Biological Chemistry*, 282(32), 23482-23490.
- Khanna, S., Roy, S., Ryu, H., Bahadduri, P., Swaan, P. W., Ratan, R. R., & Sen, C. K. (2003). Molecular Basis of Vitamin E Action TOCOTRIENOL MODULATES 12-LIPOXYGENASE, A KEY MEDIATOR OF GLUTAMATE-INDUCED NEURODEGENERATION. *Journal of Biological Chemistry*, 278(44), 43508-43515.
- Khanna, S., Roy, S., Slivka, A., Craft, T. K., Chaki, S., Rink, C., . . . Sen, C. K. (2005). Neuroprotective properties of the natural vitamin E  $\alpha$ -tocotrienol. *Stroke*, 36(10), e144-e152.
- Khor, H., Chieng, D., & Ong, K. (1995). Tocotrienols inhibit liver HMG CoA reductase activity in the guinea pig. *Nutrition Research*, 15(4), 537-544.
- Kilminster, S., Treasure, T., McMillan, T., & Holt, D. W. (1999). Neuropsychological change and S-100 protein release in 130 unselected patients undergoing cardiac surgery. *Stroke*, 30(9), 1869-1874.
- Kirchhoff, F., Blümcke, I., Celio, M., & Kettenmann, H. (1993). *Patch-clamp analysis of glial cells in brain slices of murine neocortex*. Paper presented at the Soc. Neurosci. Abstr.
- Koch, H. P., & Larsson, H. P. (2005). Small-scale molecular motions accomplish glutamate uptake in human glutamate transporters. *The Journal of neuroscience*, 25(7), 1730-1736.
- Kohutnicka, M., Lewandowska, E., Kurkowska-Jastrzebska, I., Członkowski, A., & Członkowska, A. (1998). Microglial and astrocytic involvement in a murine model of Parkinson's disease induced by 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP). *Immunopharmacology*, 39(3), 167-180.



- Komiyama, K., Iizuka, K., Yamaoka, M., Watanabe, H., Tsuchiya, N., & Umezawa, I. (1989). Studies on the biological activity of tocotrienols. *Chemical & pharmaceutical bulletin*, 37(5), 1369-1371.
- Kono, N., & Arai, H. (2015). Intracellular transport of fat-soluble vitamins A and E. *Traffic*, 16(1), 19-34.
- Kroemer, G., & Reed, J. C. (2000). Mitochondrial control of cell death. *Nature medicine*, 6(5).
- Kurkowska-Jastrzębska, I., Wrońska, A., Kohutnicka, M., Członkowski, A., & Członkowska, A. (1999). The inflammatory reaction following 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine intoxication in mouse. *Experimental neurology*, 156(1), 50-61.
- Kuryatov, A., Laube, B., Betz, H., & Kuhse, J. (1994). Mutational analysis of the glycine-binding site of the NMDA receptor: structural similarity with bacterial amino acid-binding proteins. *Neuron*, 12(6), 1291-1300.
- Lamers, K., Van Engelen, B., Gabreels, F., Hommes, O., Borm, G., & Wevers, R. (1995). Cerebrospinal neuron-specific enolase, S-100 and myelin basic protein in neurological disorders. *Acta neurologica scandinavica*, 92(3), 247-251.
- Landis, D. M. (1994). The early reactions of non-neuronal cells to brain injury. *Annual review of neuroscience*, 17(1), 133-151.
- Lawrence, M. S., Ho, D. Y., Sun, G. H., Steinberg, G. K., & Sapolsky, R. M. (1996). Overexpression of Bcl-2 with herpes simplex virus vectors protects CNS neurons against neurological insults in vitro and in vivo. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 16(2), 486-496.
- Lee, J.-M., Zipfel, G. J., & Choi, D. W. (1999). The changing landscape of ischaemic brain injury mechanisms. *Nature*, 399, A7-A14.
- Lehre, K. P., Levy, L. M., Ottersen, O. P., Storm-Mathisen, J., & Danbolt, N. C. (1995). Differential expression of two glial glutamate transporters in the rat brain: quantitative and immunocytochemical observations. *The Journal of neuroscience*, 15(3), 1835-1853.
- Levi, G., & Patrizio, M. (1992). Astrocyte Heterogeneity: Endogenous Amino Acid Levels and Release Evoked by Non-N-Methyl-D-Aspartate Receptor Agonists and by Potassium-Induced Swelling in Type-1 and Type-2 Astrocytes. *Journal of neurochemistry*, 58(5), 1943-1952.
- Lewerenz, J., Klein, M., & Methner, A. (2006). Cooperative action of glutamate transporters and cystine/glutamate antiporter system Xc<sup>-</sup> protects from oxidative glutamate toxicity. *Journal of neurochemistry*, 98(3), 916-925.

- Li, S., & Stys, P. K. (2000). Mechanisms of ionotropic glutamate receptor-mediated excitotoxicity in isolated spinal cord white matter. *The Journal of Neuroscience*, 20(3), 1190-1198.
- Li, Y., Maher, P., & Schubert, D. (1997). A role for 12-lipoxygenase in nerve cell death caused by glutathione depletion. *Neuron*, 19(2), 453-463.
- Liu, D., Smith, C. L., Barone, F. C., Ellison, J. A., Lysko, P. G., Li, K., & Simpson, I. A. (1999). Astrocytic demise precedes delayed neuronal death in focal ischemic rat brain. *Molecular brain research*, 68(1), 29-41.
- Loo, D. T., Copani, A., Pike, C. J., Whittemore, E. R., Walencewicz, A. J., & Cotman, C. W. (1993). Apoptosis is induced by beta-amyloid in cultured central nervous system neurons. *Proceedings of the National Academy of Sciences*, 90(17), 7951-7955.
- Lopez-Torres, M., Thiele, J., Shindo, Y., Han, D., & Packer, L. (1998). Topical application of -tocopherol modulates the antioxidant network and diminishes ultraviolet-induced oxidative damage in murine skin. *British Journal of Dermatology*, 138(2), 207-215.
- Lowe, D. L., Jahn, K., & Smith, D. O. (1997). Glutamate receptor editing in the mammalian hippocampus and avian neurons. *Molecular brain research*, 48(1), 37-44.
- Luetjens, C. M., Bui, N. T., Sengpiel, B., Münstermann, G., Poppe, M., Krohn, A. J., . . . Prehn, J. H. (2000). Delayed mitochondrial dysfunction in excitotoxic neuron death: cytochrome c release and a secondary increase in superoxide production. *The journal of neuroscience*, 20(15), 5715-5723.
- Luque, J. M., & Richards, J. G. (1995). Expression of NMDA 2B receptor subunit mRNA in Bergmann glia. *Glia*, 13(3), 228-232.
- MacAulay, N., Hamann, S., & Zeuthen, T. (2004). Water transport in the brain: role of cotransporters. *Neuroscience*, 129(4), 1029-1042.
- Mack, K. J., Kriegler, S., Chang, S., & Chiu, S.-Y. (1994). Transcription factor expression is induced by axonal stimulation and glutamate in the glia of the developing optic nerve. *Molecular brain research*, 23(1), 73-80.
- Madry, C., Mesic, I., Betz, H., & Laube, B. (2007). The N-terminal domains of both NR1 and NR2 subunits determine allosteric Zn<sup>2+</sup> inhibition and glycine affinity of N-methyl-D-aspartate receptors. *Molecular pharmacology*, 72(6), 1535-1544.
- Magistretti, P. (2004). Brain energy metabolism: Elsevier Academic Press.
- Magistretti, P. J. (2006). Neuron–glia metabolic coupling and plasticity. *Journal of Experimental Biology*, 209(12), 2304-2311.

- Magistretti, P. J., Pellerin, L., Rothman, D. L., & Shulman, R. G. (1999). Energy on demand. *Science*, 283(5401), 496.
- Maher, P., & Davis, J. B. (1996). The role of monoamine metabolism in oxidative glutamate toxicity. *The Journal of neuroscience*, 16(20), 6394-6401.
- Majno, G., & Joris, I. (1995). Apoptosis, oncosis, and necrosis. An overview of cell death. *The American journal of pathology*, 146(1), 3.
- Mandrekar, S., & Landreth, G. E. (2010). Microglia and inflammation in Alzheimer's disease. *CNS & neurological disorders drug targets*, 9(2), 156.
- Mangialasche, F., Kivipelto, M., Mecocci, P., Rizzuto, D., Palmer, K., Winblad, B., & Fratiglioni, L. (2010). High plasma levels of vitamin E forms and reduced Alzheimer's disease risk in advanced age. *Journal of Alzheimer's Disease*, 20(4), 1029-1037.
- Maniam, S., Mohamed, N., Shuid, A. N., & Soelaiman, I. N. (2008). Palm Tocotrienol Exerted Better Antioxidant Activities in Bone than  $\alpha$ -Tocopherol. *Basic & clinical pharmacology & toxicology*, 103(1), 55-60.
- Martin, L., Blackstone, C., Levey, A., Huganir, R., & Price, D. (1993). AMPA glutamate receptor subunits are differentially distributed in rat brain. *Neuroscience*, 53(2), 327-358.
- Martin, L. J., Brambrink, A. M., Lehmann, C., Portera-Cailliau, C., Koehler, R., Rothstein, J., & Traystman, R. J. (1997). Hypoxia—ischemia causes abnormalities in glutamate transporters and death of astroglia and neurons in newborn striatum. *Annals of neurology*, 42(3), 335-348.
- Matute, C., Alberdi, E., Domercq, M. a., Pérez-Cerdá, F., Pérez-Samartín, A., & Sánchez-Gómez, M. a. V. (2001). The link between excitotoxic oligodendroglial death and demyelinating diseases. *Trends in neurosciences*, 24(4), 224-230.
- Matute, C., Domercq, M., Fogarty, D. J., de Zulueta, M. P., & Sánchez-Gómez, M. V. (1999). On how altered glutamate homeostasis may contribute to demyelinating diseases of the CNS *The Functional Roles of Glial Cells in Health and Disease* (pp. 98-107): Springer.
- Matute, C., García-Barcina, J. M., & Miledi, R. (1994). Expression of neurotransmitter receptors and Ca<sup>2+</sup> channels in the adult fornix and optic nerve. *NeuroReport*, 5(12), 1457-1460.
- Maulik, G., Shrikhande, A., Kijima, T., Ma, P. C., Morrison, P. T., & Salgia, R. (2002). Role of the hepatocyte growth factor receptor, c-Met, in oncogenesis and potential for therapeutic inhibition. *Cytokine & growth factor reviews*, 13(1), 41-59.

- Mazlan, M., & Ngah, W. Z. W. (2010). Comparison of the Effects of  $\alpha$ -Tocopherol and  $\gamma$ -tocotrienol Against Oxidative Stress in Two Different Neuronal Cultures.
- Mazlan, M., Then, S.-M., Top, G. M., & Ngah, W. Z. W. (2006). Comparative effects of  $\alpha$ -tocopherol and  $\gamma$ -tocotrienol against hydrogen peroxide induced apoptosis on primary-cultured astrocytes. *Journal of the neurological sciences*, 243(1), 5-12.
- McGeer, P. L., & McGeer, E. G. (2002). Local neuroinflammation and the progression of Alzheimer's disease. *Journal of neurovirology*, 8(6), 529-538.
- McKeating, E. G., Andrews, P., & Mascia, L. (1998). *Relationship of neuron specific enolase and protein S-100 concentrations in systemic and jugular venous serum to injury severity and outcome after traumatic brain injury*: Springer.
- McNaughton, L., & Hunt, S. (1992). Regulation of gene expression in astrocytes by excitatory amino acids. *Molecular brain research*, 16(3), 261-266.
- Meng, S. Z., Ohyu, J., & Takashima, S. (1997). Changes in AMPA glutamate and dopamine D 2 receptors in hypoxic-ischemic basal ganglia necrosis. *Pediatric neurology*, 17(2), 139-143.
- Mennerick, S., Benz, A., & Zorumski, C. F. (1996). Components of glial responses to exogenous and synaptic glutamate in rat hippocampal microcultures. *The Journal of neuroscience*, 16(1), 55-64.
- Mennerick, S., Dhond, R. P., Benz, A., Xu, W., Rothstein, J. D., Danbolt, N. C., . . . Zorumski, C. F. (1998). Neuronal expression of the glutamate transporter GLT-1 in hippocampal microcultures. *The Journal of neuroscience*, 18(12), 4490-4499.
- Meucci, O., Fatatis, A., Holzwarth, J. A., & Miller, R. J. (1996). Developmental regulation of the toxin sensitivity of  $\text{Ca}^{2+}$ -permeable AMPA receptors in cortical glia. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 16(2), 519.
- Meydani, S. N., & Beharka, A. A. (1998). Recent developments in vitamin E and immune response. *Nutrition reviews*, 56(1), S49.
- Michaelis, E. K. (1998). Molecular biology of glutamate receptors in the central nervous system and their role in excitotoxicity, oxidative stress and aging. *Progress in neurobiology*, 54(4), 369-415.
- Minelli, A., Barbaresi, P., Reimer, R., Edwards, R., & Conti, F. (2001). The glial glutamate transporter GLT-1 is localized both in the vicinity of and at distance from axon terminals in the rat cerebral cortex. *Neuroscience*, 108(1), 51-59.
- Minhajuddin, M., Beg, Z. H., & Iqbal, J. (2005). Hypolipidemic and antioxidant properties of tocotrienol rich fraction isolated from rice bran oil in



experimentally induced hyperlipidemic rats. *Food and chemical toxicology*, 43(5), 747-753.

- Miyamoto, K., Shiozaki, M., Shibata, M., Koike, M., Uchiyama, Y., & Gotow, T. (2009). Very-high-dose  $\alpha$ -tocopherol supplementation increases blood pressure and causes possible adverse central nervous system effects in stroke-prone spontaneously hypertensive rats. *Journal of neuroscience research*, 87(2), 556-566.
- Miyamoto, M., Murphy, T. H., Schnaar, R. L., & Coyle, J. T. (1989). Antioxidants protect against glutamate-induced cytotoxicity in a neuronal cell line. *Journal of Pharmacology and Experimental Therapeutics*, 250(3), 1132-1140.
- Moreira, P. I., Smith, M. A., Zhu, X., Nunomura, A., Castellani, R. J., & Perry, G. (2005). Oxidative stress and neurodegeneration. *Annals of the New York Academy of Sciences*, 1043(1), 545-552.
- Müller, T., Grosche, J., Ohlemeyer, C., & Kettenmann, H. (1993). NMDA-activated currents in Bergmann glial cells. *Neuroreport*, 4(6), 671-674.
- Muller, T., Moller, T., Berger, T., Schnitzer, J., & Kettenmann, H. (1992). Calcium entry through kainate receptors and resulting potassium-channel blockade in Bergmann glial cells. *Science*, 256(5063), 1563-1566.
- Müller, T., Möller, T., Neuhaus, J., & Kettenmann, H. (1996). Electrical coupling among Bergmann glial cells and its modulation by glutamate receptor activation. *Glia*, 17(4), 274-284.
- Murphy, A. N., Fiskum, G., & Beal, M. F. (1999). Mitochondria in neurodegeneration: bioenergetic function in cell life and death. *Journal of Cerebral Blood Flow & Metabolism*, 19(3), 231-245.
- Murphy, M. P. (2009). How mitochondria produce reactive oxygen species. *Biochemical Journal*, 417(1), 1-13.
- Murphy, T., Schnaar, R., & Coyle, J. (1990). Immature cortical neurons are uniquely sensitive to glutamate toxicity by inhibition of cystine uptake. *The FASEB Journal*, 4(6), 1624-1633.
- Murphy, T. H., Miyamoto, M., Sastre, A., Schnaar, R. L., & Coyle, J. T. (1989). Glutamate toxicity in a neuronal cell line involves inhibition of cystine transport leading to oxidative stress. *Neuron*, 2(6), 1547-1558.
- Murphy, T. H., Schnaar, R. L., & Coyle, J. T. (1990). Immature cortical neurons are uniquely sensitive to glutamate toxicity by inhibition of cystine uptake. *FASEB J*, 4(6), 1624-1633.
- Mutalib, M. S. A., Khaza'ai, H., & Wahle, K. W. (2003). Palm-tocotrienol rich fraction (TRF) is a more effective inhibitor of LDL oxidation and endothelial

cell lipid peroxidation than  $\alpha$ -tocopherol in vitro. *Food research international*, 36(5), 405-413.

Myers, R., MacDonald, M., Koroshetz, W., Duyao, M., Ambrose, C., Taylor, S., . . . Whaley, W. (1993). De novo expansion of a (CAG) n repeat in sporadic Huntington's disease. *Nature genetics*, 5(2), 168-173.

Naderi, J., Hung, M., & Pandey, S. (2003). Oxidative stress-induced apoptosis in dividing fibroblasts involves activation of p38 MAP kinase and over-expression of Bax: resistance of quiescent cells to oxidative stress. *Apoptosis*, 8(1), 91-100.

Nakanishi, S., & Masu, M. (1994). Molecular diversity and functions of glutamate receptors. *Annual review of biophysics and biomolecular structure*, 23(1), 319-348.

Nedergaard, M., Ransom, B., & Goldman, S. A. (2003). New roles for astrocytes: redefining the functional architecture of the brain. *Trends in neurosciences*, 26(10), 523-530.

Nesaretnam, S. D., Philippa D. Darbre, Kalanithi. (2000). Tocotrienols inhibit growth of ZR-75-1 breast cancer cells. *International Journal of Food Sciences and Nutrition*, 51(1), 95-103.

Nicholls, D., & Attwell, D. (1990). The release and uptake of excitatory amino acids. *Trends in pharmacological sciences*, 11(11), 462-468.

Nicholls, D. G., & Budd, S. L. (1998). Mitochondria and neuronal glutamate excitotoxicity. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1366(1), 97-112.

Niki, E., & Traber, M. G. (2012). A history of vitamin E. *Annals of Nutrition and Metabolism*, 61(3), 207-212.

Oka, A., Belliveau, M. J., Rosenberg, P. A., & Volpe, J. J. (1993). Vulnerability of oligodendroglia to glutamate: pharmacology, mechanisms, and prevention. *The Journal of neuroscience*, 13(4), 1441-1453.

Olanow, C. (1992). An introduction to the free radical hypothesis in Parkinson's disease. *Annals of Neurology*, 32(S1), S2-S9.

Olanow, C. (1993). A radical hypothesis for neurodegeneration. *Trends in neurosciences*, 16(11), 439-444.

Olsen, R. W., Chang, C.-S. S., Li, G., Hanchar, H. J., & Wallner, M. (2004). Fishing for allosteric sites on GABA A receptors. *Biochemical pharmacology*, 68(8), 1675-1684.

- Omar, N. S., Zakaria, Z. A. C., Mian, T. S., Ngah, W. Z. W., & Mazlan, M. (2011). Centella asiatica modulates neuron cell survival by altering caspase-9 pathway. *Journal of Medicinal Plants Research*, 5(11), 2201-2109.
- Osakada, F., Hashino, A., Kume, T., Katsuki, H., Kaneko, S., & Akaike, A. (2003). Neuroprotective effects of  $\alpha$ -tocopherol on oxidative stress in rat striatal cultures. *European journal of pharmacology*, 465(1), 15-22.
- Osakada, F., Hashino, A., Kume, T., Katsuki, H., Kaneko, S., & Akaike, A. (2004).  $\alpha$ -Tocotrienol provides the most potent neuroprotection among vitamin E analogs on cultured striatal neurons. *Neuropharmacology*, 47(6), 904-915.
- Pannicke, T. (1999). *A single-cell PCR study of NMDA receptor mRNA in Müller glial cells*. Paper presented at the Göttingen Conference German Neurosci. Soc. Abstr II.
- Parker, R. A., Pearce, B. C., Clark, R. W., Gordon, D. A., & Wright, J. (1993). Tocotrienols regulate cholesterol production in mammalian cells by post-transcriptional suppression of 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *Journal of Biological Chemistry*, 268(15), 11230-11238.
- Partin, K., Patneau, D., Winters, C., Mayer, M., & Buonanno, A. (1993). Selective modulation of desensitization at AMPA versus kainate receptors by cyclothiazide and concanavalin A. *Neuron*, 11(6), 1069-1082.
- Pasti, L., Volterra, A., Pozzan, T., & Carmignoto, G. (1997). Intracellular calcium oscillations in astrocytes: a highly plastic, bidirectional form of communication between neurons and astrocytes in situ. *The Journal of Neuroscience*, 17(20), 7817-7830.
- Petegnief, V., Saura, J., Dewar, D., Cummins, D., Dragunow, M., & Mahy, N. (1999). Long-term effects of  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionate and 6-nitro-7-sulphamoylbenzo (f) quinoxaline-2, 3-dione in the rat basal ganglia: calcification, changes in glutamate receptors and glial reactions. *Neuroscience*, 94(1), 105-115.
- Petralia, R. S., & Wenthold, R. J. (1992). Light and electron immunocytochemical localization of AMPA-selective glutamate receptors in the rat brain. *Journal of Comparative Neurology*, 318(3), 329-354.
- Pettmann, B., & Henderson, C. E. (1998). Neuronal cell death. *Neuron*, 20(4), 633-647.
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research*, 29(9), e45-e45.
- Phillis, J., Ren, J., & O'Regan, M. (2000). Transporter reversal as a mechanism of glutamate release from the ischemic rat cerebral cortex: studies with DL-threo- $\beta$ -benzyloxyaspartate. *Brain research*, 868(1), 105-112.

- Pines, G., Danbolt, N. C., Bjørås, M., Zhang, Y., Bendahan, A., Eide, L., . . . Kanner, B. I. (1992). Cloning and expression of a rat brain L-glutamate transporter.
- Porter, J. T., & McCarthy, K. D. (1995). GFAP-positive hippocampal astrocytes in situ respond to glutamatergic neuroligands with increases in  $[Ca^{2+}]_i$ . *Glia*, *13*(2), 101-112.
- Porter, J. T., & McCarthy, K. D. (1996). Hippocampal astrocytes in situ respond to glutamate released from synaptic terminals. *The Journal of Neuroscience*, *16*(16), 5073-5081.
- Portt, L., Norman, G., Clapp, C., Greenwood, M., & Greenwood, M. T. (2011). Anti-apoptosis and cell survival: a review. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, *1813*(1), 238-259.
- Pratico, D., Clark, C. M., Liun, F., Lee, V. Y.-M., & Trojanowski, J. Q. (2002). Increase of brain oxidative stress in mild cognitive impairment: a possible predictor of Alzheimer disease. *Archives of Neurology*, *59*(6), 972-976.
- Prieto, M., & Alonso, G. (1999). Differential sensitivity of cultured tancytes and astrocytes to hydrogen peroxide toxicity. *Experimental neurology*, *155*(1), 118-127.
- Puro, D. G., Yuan, J. P., & Sucher, N. J. (1996). Activation of NMDA receptor-channels in human retinal Müller glial cells inhibits inward-rectifying potassium currents. *Visual neuroscience*, *13*(02), 319-326.
- Pyper, J., & Bolen, J. (1989). Neuron-specific splicing of C-SRC cDNA RNA in human brain. *Journal of neuroscience research*, *24*(1), 89-96.
- Queiroz, G., Gebicke-Haerter, P., Schobert, A., Starke, K., & Von Kügelgen, I. (1997). Release of ATP from cultured rat astrocytes elicited by glutamate receptor activation. *Neuroscience*, *78*(4), 1203-1208.
- Qureshi, A., Bradlow, B., Brace, L., Manganello, J., Peterson, D., Pearce, B., . . . Elson, C. (1995). Response of hypercholesterolemic subjects to administration of tocotrienols. *Lipids*, *30*(12), 1171-1177.
- Qureshi, A. A., Burger, W., Peterson, D., & Elson, C. (1986). The structure of an inhibitor of cholesterol biosynthesis isolated from barley. *Journal of Biological Chemistry*, *261*(23), 10544-10550.
- Qureshi, A. A., Qureshi, N., Wright, J., Shen, Z., Kramer, G., Gapor, A., . . . Peterson, D. (1991). Lowering of serum cholesterol in hypercholesterolemic humans by tocotrienols (palmvitee). *The American journal of clinical nutrition*, *53*(4), 1021S-1026S.
- Qureshi, A. A., Sami, S. A., Salser, W. A., & Khan, F. A. (2002). Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF 25) of rice bran in hypercholesterolemic humans. *Atherosclerosis*, *161*(1), 199-207.



- Rao, V. L. R., Rao, A. M., Dogan, A., Bowen, K. K., Hatcher, J., Rothstein, J. D., & Dempsey, R. J. (2000). Glial glutamate transporter GLT-1 down-regulation precedes delayed neuronal death in gerbil hippocampus following transient global cerebral ischemia. *Neurochemistry international*, 36(6), 531-537.
- Ratan, R. R., Murphy, T., & Baraban, J. (1994). Macromolecular synthesis inhibitors prevent oxidative stress-induced apoptosis in embryonic cortical neurons by shunting cysteine from protein synthesis to glutathione. *The Journal of neuroscience*, 14(7), 4385-4392.
- Rauen, T., Wießner, M., Sullivan, R., Lee, A., & Pow, D. V. (2004). A new GLT1 splice variant: cloning and immunolocalization of GLT1c in the mammalian retina and brain. *Neurochemistry international*, 45(7), 1095-1106.
- Rego, A. C., Ward, M. W., & Nicholls, D. G. (2001). Mitochondria control ampa/kainate receptor-induced cytoplasmic calcium deregulation in rat cerebellar granule cells. *The Journal of Neuroscience*, 21(6), 1893-1901.
- Reichert, S. A., Kim-Han, J. S., & Dugan, L. L. (2001). The mitochondrial permeability transition pore and nitric oxide synthase mediate early mitochondrial depolarization in astrocytes during oxygen-glucose deprivation. *The Journal of Neuroscience*, 21(17), 6608-6616.
- Reye, P., Sullivan, R., & Pow, D. V. (2002). Distribution of two splice variants of the glutamate transporter GLT-1 in the developing rat retina. *Journal of Comparative Neurology*, 447(4), 323-330.
- Richardson, J. S. (1993). Free radicals in the genesis of Alzheimer's disease. *Annals of the New York Academy of Sciences*, 695(1), 73-76.
- Ripellino, J., Neve, R., & Howe, J. (1997). Expression and heteromeric interactions of non-N-methyl-D-aspartate glutamate receptor subunits in the developing and adult cerebellum. *Neuroscience*, 82(2), 485-497.
- Robert, A., & Magistretti, P. J. (1997). AMPA/kainate receptor activation blocks K<sup>+</sup> currents via internal Na<sup>+</sup> increase in mouse cultured stellate astrocytes. *Glia*, 20(1), 38-50.
- Romner, B., Ingebrigtsen, T., Kongstad, P., & Børgesen, S. E. (2000). Traumatic brain damage: serum S-100 protein measurements related to neuroradiological findings. *Journal of neurotrauma*, 17(8), 641-647.
- Rosenthal, J. J., & Seeburg, P. H. (2012). A-to-I RNA editing: effects on proteins key to neural excitability. *Neuron*, 74(3), 432-439.
- Rossi, D. J., Oshima, T., & Attwell, D. (2000). Glutamate release in severe brain ischaemia is mainly by reversed uptake. *Nature*, 403(6767), 316-321.
- Rothstein, J. D., Dykes-Hoberg, M., Pardo, C. A., Bristol, L. A., Jin, L., Kuncl, R. W., . . . Schielke, J. P. (1996). Knockout of glutamate transporters reveals a major

role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron*, 16(3), 675-686.

Rothstein, J. D., Martin, L., Levey, A. I., Dykes-Hoberg, M., Jin, L., Wu, D., . . . Kuncl, R. W. (1994). Localization of neuronal and glial glutamate transporters. *Neuron*, 13(3), 713-725.

Sagara, Y., & Schubert, D. (1998). The activation of metabotropic glutamate receptors protects nerve cells from oxidative stress. *The Journal of neuroscience*, 18(17), 6662-6671.

Saito, Y., Nishio, K., Numakawa, Y., Ogawa, Y., Yoshida, Y., Noguchi, N., & Niki, E. (2007). Protective effects of 15-deoxy- $\Delta$ 12, 14-prostaglandin J2 against glutamate-induced cell death in primary cortical neuron cultures: induction of adaptive response and enhancement of cell tolerance primarily through up-regulation of cellular glutathione. *Journal of neurochemistry*, 102(5), 1625-1634.

Saito, Y., Yoshida, Y., Noguchi, N., & Niki, E. (2010). Cytoprotective Effect of Vitamin E Homologues Against Glutamate-induced Cell Death in Immature Primary Cortical Neuron Cultures: Tocopherols and Tocotrienols Exert Similar Effects by Antioxidant Function. *Free Radical Biology and Medicine*, 49, S196.

Sakahira, H., Enari, M., & Nagata, S. (1998). Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature*, 391(6662), 96-99.

Sánchez-Gómez, M., Alberdi, E., Ibarretxe, G., Torre, I., & Matute, C. (2002). Activation of AMPA and kainate receptors triggers apoptotic and necrotic oligodendroglial death in vitro. Paper presented at the GLIA.

Sánchez, G., & Ortega, A. (1994). AMPA/KA receptor induced AP-1 DNA binding activity in cultured Bergmann glia cells. *NeuroReport*, 5(16), 2109-2112.

Sato, H., Tamba, M., Ishii, T., & Bannai, S. (1999). Cloning and expression of a plasma membrane cystine/glutamate exchange transporter composed of two distinct proteins. *Journal of Biological Chemistry*, 274(17), 11455-11458.

Sattler, R., & Tymianski, M. (2001). Molecular mechanisms of glutamate receptor-mediated excitotoxic neuronal cell death. *Molecular neurobiology*, 24(1-3), 107-129.

Schmitt, A., Asan, E., Lesch, K.-P., & Kugler, P. (2002). A splice variant of glutamate transporter GLT1/EAAT2 expressed in neurons: cloning and localization in rat nervous system. *Neuroscience*, 109(1), 45-61.

Schröder, W., Seifert, G., Hüttmann, K., Hinterkeuser, S., & Steinhäuser, C. (2002). AMPA receptor-mediated modulation of inward rectifier K<sup>+</sup> channels in astrocytes of mouse hippocampus. *Molecular and Cellular Neuroscience*, 19(3), 447-458.



- Schubert, D., Kimura, H., & Maher, P. (1992). Growth factors and vitamin E modify neuronal glutamate toxicity. *Proceedings of the National Academy of Sciences*, 89(17), 8264-8267.
- Schubert, D., & Piasecki, D. (2001). Oxidative glutamate toxicity can be a component of the excitotoxicity cascade. *The Journal of Neuroscience*, 21(19), 7455-7462.
- Schulz, J. B., Lindenau, J., Seyfried, J., & Dichgans, J. (2000). Glutathione, oxidative stress and neurodegeneration. *European Journal of Biochemistry*, 267(16), 4904-4911.
- Seifert, G., Schilling, K., & Steinhäuser, C. (2006). Astrocyte dysfunction in neurological disorders: a molecular perspective. *Nature Reviews Neuroscience*, 7(3), 194-206.
- Seifert, G., Schumacher, T., Weber, M., Schramm, J., & Steinhäuser, C. (1999). *AMPA receptor properties of astrocytes in human epilepsy*. Paper presented at the JOURNAL OF NEUROCHEMISTRY.
- Seifert, G., & Steinhäuser, C. (2001). Ionotropic glutamate receptors in astrocytes *Glial Cell Function*: Elsevier Science BV, Amsterdam.
- Seifert, G., & Steinhäuser, C. (1995). Glial cells in the mouse hippocampus express AMPA receptors with an intermediate Ca<sup>2+</sup> permeability. *European Journal of Neuroscience*, 7(9), 1872-1881.
- Seifert, G., & Steinhäuser, C. (2001). Ionotropic glutamate receptors in astrocytes. *Progress in brain research*, 132, 287-299.
- Seifert, G., Zhou, M., & Steinhäuser, C. (1997). Analysis of AMPA receptor properties during postnatal development of mouse hippocampal astrocytes. *Journal of neurophysiology*, 78(6), 2916-2923.
- Selvaraju, T. R., Khaza'ai, H., Vidyadaran, S., Mutalib, M. S. A., Ramachandran, V., & Hamdan, Y. (2014). Cytoprotective effect of tocotrienol-rich fraction and  $\alpha$ -tocopherol vitamin E isoforms against glutamate-induced cell death in neuronal cells. *Int J Vitam Nutr Res*, 84, 140-151.
- Selvaraju, T. R., Khaza'ai, H., Vidyadaran, S., Mutalib, M. S. A., & Vasudevan, R. (2014). The neuroprotective effects of tocotrienol rich fraction and alpha tocopherol against glutamate injury in astrocytes. *Bosnian Journal of Basic Medical Sciences*, 14(4), 195.
- Sen, C. K., Khanna, S., & Roy, S. (2006). Tocotrienols: vitamin E beyond tocopherols. *Life sciences*, 78(18), 2088-2098.
- Sen, C. K., Khanna, S., & Roy, S. (2007). Tocotrienols in health and disease: the other half of the natural vitamin E family. *Molecular aspects of medicine*, 28(5), 692-728.

- Sen, C. K., Khanna, S., Roy, S., & Packer, L. (2000). Molecular Basis of Vitamin E Action Tocotrienol potently inhibits glutamate-induced pp60c-Src Kinase activation and death of ht4 neuronal cells. *Journal of Biological Chemistry*, 275(17), 13049-13055.
- Sen, C. K., Rink, C., & Khanna, S. (2010). Palm oil-derived natural vitamin E  $\alpha$ -tocotrienol in brain health and disease. *Journal of the American College of Nutrition*, 29(sup3), 314S-323S.
- Sensi, S. L., Yin, H. Z., & Weiss, J. H. (1999). Glutamate triggers preferential Zn<sup>2+</sup> flux through Ca<sup>2+</sup> permeable AMPA channels and consequent ROS production. *Neuroreport*, 10(8), 1723-1727.
- Serbinova, E., Kagan, V., Han, D., & Packer, L. (1991). Free radical recycling and intramembrane mobility in the antioxidant properties of alpha-tocopherol and alpha-tocotrienol. *Free Radical Biology and Medicine*, 10(5), 263-275.
- Serbinova, E. A., & Packer, L. (1994). Antioxidant properties of alpha-tocopherol and alpha-tocotrienol. *Methods in enzymology*, 234, 354-366.
- Sheldon, A. L., & Robinson, M. B. (2007). The role of glutamate transporters in neurodegenerative diseases and potential opportunities for intervention. *Neurochemistry international*, 51(6), 333-355.
- Shelton, M. K., & McCarthy, K. D. (1999). Mature hippocampal astrocytes exhibit functional metabotropic and ionotropic glutamate receptors in situ. *Glia*, 26(1), 1-11.
- Shukla, V., Mishra, S. K., & Pant, H. C. (2011). Oxidative stress in neurodegeneration. *Advances in pharmacological sciences*, 2011.
- SIESJÖ, B. K., KATSURA, K.-I., Zhao, Q., FOLBERGROVÁ, J., PAHLMARK, K., SIESJÖ, P., & SMITH, M.-L. (1995). Mechanisms of secondary brain damage in global and focal ischemia: a speculative synthesis. *Journal of neurotrauma*, 12(5), 943-956.
- Singh, R. P., Khanna, R., Kaw, J. L., Khanna, S. K., & Das, M. (2003). Comparative effect of benzantrone and 3-bromobenzanthrone on hepatic xenobiotic metabolism and anti-oxidative defense system in guinea pigs. *Archives of toxicology*, 77(2), 94-99.
- Soriano, P., Montgomery, C., Geske, R., & Bradley, A. (1991). Targeted disruption of the c-src proto-oncogene leads to osteopetrosis in mice. *Cell*, 64(4), 693-702.
- Spreafico, R., Frassoni, C., Arcelli, P., Battaglia, G., Wenthold, R., & De Biasi, S. (1994). Distribution of AMPA selective glutamate receptors in the thalamus of adult rats and during postnatal development. A light and ultrastructural immunocytochemical study. *Developmental brain research*, 82(1), 231-244.

- Srivastava, J. K., & Gupta, S. (2006). Tocotrienol-rich fraction of palm oil induces cell cycle arrest and apoptosis selectively in human prostate cancer cells. *Biochemical and biophysical research communications*, 346(2), 447-453.
- Steinhäser, C., Jabs, R., & Kettenmann, H. (1994). Properties of GABA and glutamate responses in identified glial cells of the mouse hippocampal slice. *Hippocampus*, 4(1), 19-35.
- Storck, T., Schulte, S., Hofmann, K., & Stoffel, W. (1992). Structure, expression, and functional analysis of a Na (+)-dependent glutamate/aspartate transporter from rat brain. *Proceedings of the National Academy of Sciences*, 89(22), 10955-10959.
- Stout, A. K., Raphael, H. M., Kanterewicz, B. I., Klann, E., & Reynolds, I. J. (1998). Glutamate-induced neuron death requires mitochondrial calcium uptake. *Nature neuroscience*, 1(5), 366-373.
- Suarez, I., Bodega, G., & Fernandez, B. (1997). Modulation of AMPA receptor subunits GluR1 and GluR2/3 in the rat cerebellum in an experimental hepatic encephalopathy model. *Brain research*, 778(2), 346-353.
- Sullivan, R., Rauen, T., Fischer, F., Wießner, M., Grewer, C., Bicho, A., & Pow, D. V. (2004). Cloning, transport properties, and differential localization of two splice variants of GLT-1 in the rat CNS: Implications for CNS glutamate homeostasis. *Glia*, 45(2), 155-169.
- Sun, Z.-W., Zhang, L., Zhu, S.-J., Chen, W.-C., & Mei, B. (2010). Excitotoxicity effects of glutamate on human neuroblastoma SH-SY5Y cells via oxidative damage. *Neuroscience bulletin*, 26(1), 8-16.
- Sundram, K., Khor, H., Ong, A. S., & Pathmanathan, R. (1989). Effect of dietary palm oils on mammary carcinogenesis in female rats induced by 7, 12-dimethylbenz (a) anthracene. *Cancer Research*, 49(6), 1447-1451.
- Suzuki, Y. J., Tsuchiya, M., Wassall, S. R., Choo, Y. M., Govil, G., Kagan, V. E., & Packer, L. (1993). Structural and dynamic membrane properties of Alpha-tocopherol and Alpha-tocotrienol: implication to the molecular mechanism of their antioxidant potency. *Biochemistry*, 32(40), 10692-10699.
- Takahashi, K., & Loo, G. (2004). Disruption of mitochondria during tocotrienol-induced apoptosis in MDA-MB-231 human breast cancer cells. *Biochemical pharmacology*, 67(2), 315-324.
- Tan, S., Schubert, D., & Maher, P. (2001). Oxytosis: a novel form of programmed cell death. *Current topics in medicinal chemistry*, 1(6), 497-506.
- Tapiero, H., Mathe, G., Couvreur, P., & Tew, K. (2002). II. Glutamine and glutamate. *Biomedicine & pharmacotherapy*, 56(9), 446-457.

- Then, S. M., Mazlan, M., Top, G. M., & Ngah, W. Z. W. (2009). Is vitamin E toxic to neuron cells? *Cellular and molecular neurobiology*, 29(4), 485-496.
- Thiele, D. L., & Lipsky, P. E. (1990). The action of leucyl-leucine methyl ester on cytotoxic lymphocytes requires uptake by a novel dipeptide-specific facilitated transport system and dipeptidyl peptidase I-mediated conversion to membranolytic products. *The Journal of experimental medicine*, 172(1), 183-194.
- Thiele, D. L., & Lipsky, P. E. (1992). Apoptosis is induced in cells with cytolytic potential by L-leucyl-L-leucine methyl ester. *The Journal of Immunology*, 148(12), 3950-3957.
- Tirosh, O., Sen, C. K., Roy, S., Kobayashi, M. S., & Packer, L. (1999). Neuroprotective effects of  $\alpha$ -lipoic acid and its positively charged amide analogue. *Free Radical Biology and Medicine*, 26(11), 1418-1426.
- Torp, R., Lekieffre, D., Levy, L., Haug, F., Danbolt, N., Meldrum, B., & Ottersen, O. (1995). Reduced postischemic expression of a glial glutamate transporter, GLT1, in the rat hippocampus. *Experimental brain research*, 103(1), 51-58.
- Tort, A. B., Portela, L. V., Rockenbach, I. C., Monte, T. L., Pereira, M. L., Souza, D. O., . . . Jardim, L. B. (2005). S100B and NSE serum concentrations in Machado Joseph disease. *Clinica chimica acta*, 351(1), 143-148.
- Tullberg, M., Rosengren, L., Blomsterwall, E., Karlsson, J.-E., & Wikkelso, C. (1998). CSF neurofilament and glial fibrillary acidic protein in normal pressure hydrocephalus. *Neurology*, 50(4), 1122-1127.
- Turner, M. R., Hardiman, O., Benatar, M., Brooks, B. R., Chio, A., De Carvalho, M., . . . Mitsumoto, H. (2013). Controversies and priorities in amyotrophic lateral sclerosis. *The Lancet Neurology*, 12(3), 310-322.
- Uchihori, Y., & Puro, D. G. (1993). Glutamate as a neuron-to-glial signal for mitogenesis: role of glial N-methyl-D-aspartate receptors. *Brain research*, 613(2), 212-220.
- Vos, P., & Verbeek, M. (2002). Brain specific proteins in serum: do they reliably reflect brain damage? *Shock*, 18(5), 481-482.
- Wang, J., Sun, P., Bao, Y., Dou, B., Song, D., & Li, Y. (2012). Vitamin E renders protection to PC12 cells against oxidative damage and apoptosis induced by single-walled carbon nanotubes. *Toxicology in Vitro*, 26(1), 32-41.
- Wang, J., Zhai, Q., Chen, Y., Lin, E., Gu, W., McBurney, M. W., & He, Z. (2005). A local mechanism mediates NAD-dependent protection of axon degeneration. *J Cell Biol*, 170(3), 349-355.



- Wang, K., & Walz, W. (2003). Unusual topographical pattern of proximal astrogliosis around a cortical devascularizing lesion. *Journal of neuroscience research*, 73(4), 497-506.
- Wang, L., Jing, W., & Hang, Y. N. (2008). Glutamate-induced c-Jun expression in neuronal PC12 cells: the effects of ketamine and propofol. *J Neurosurg Anesthesiol*, 20(2), 124-130. doi: 10.1097/ANA.0b013e3181667c27
- Wang, S., Yu, H., & Wickliffe, J. K. (2011). Limitation of the MTT and XTT assays for measuring cell viability due to superoxide formation induced by nano-scale TiO<sub>2</sub>. *Toxicology in Vitro*, 25(8), 2147-2151.
- Weimann, B. J., & Weiser, H. (1991). Functions of vitamin E in reproduction and in prostacyclin and immunoglobulin synthesis in rats. *The American journal of clinical nutrition*, 53(4), 1056S-1060S.
- Whittemore, E., Loo, D., Watt, J., & Cotmans, C. (1995). A detailed analysis of hydrogen peroxide-induced cell death in primary neuronal culture. *Neuroscience*, 67(4), 921-932.
- Winterbone, M. S., Sampson, M. J., Saha, S., Hughes, J. C., & Hughes, D. A. (2007). Pro-oxidant effect of alpha-tocopherol in patients with type 2 diabetes after an oral glucose tolerance test--a randomised controlled trial. *Cardiovasc Diabetol*, 6(8).
- Wirdefeldt, K., Adami, H.-O., Cole, P., Trichopoulos, D., & Mandel, J. (2011). Epidemiology and etiology of Parkinson's disease: a review of the evidence. *European journal of epidemiology*, 26(1), 1.
- Wong, A., & Cortopassi, G. A. (2002). High-throughput measurement of mitochondrial membrane potential in a neural cell line using a fluorescence plate reader. *Biochemical and biophysical research communications*, 298(5), 750-754.
- Wu, S. J., Liu, P. L., & Ng, L. T. (2008). Tocotrienol-rich fraction of palm oil exhibits anti-inflammatory property by suppressing the expression of inflammatory mediators in human monocytic cells. *Molecular nutrition & food research*, 52(8), 921-929.
- WU, S. J., & Ng, L. T. (2007). Antioxidant and antihepatoma activities of palm oil extract. *Journal of Food Lipids*, 14(2), 122-137.
- Wyllie, D., & Cull-Candy, S. G. (1994). A comparison of non-NMDA receptor channels in type-2 astrocytes and granule cells from rat cerebellum. *The Journal of physiology*, 475(1), 95.
- Xiang, H., Kinoshita, Y., Knudson, C. M., Korsmeyer, S. J., Schwartzkroin, P. A., & Morrison, R. S. (1998). Bax involvement in p53-mediated neuronal cell death. *The Journal of neuroscience*, 18(4), 1363-1373.

- Yano, Y., Satoh, H., Fukumoto, K., Kumadaki, I., Ichikawa, T., Yamada, K., . . . Yano, T. (2005). Induction of cytotoxicity in human lung adenocarcinoma cells by 6-O-carboxypropyl- $\alpha$ -tocotrienol, a redox-silent derivative of  $\alpha$ -tocotrienol. *International journal of cancer*, 115(5), 839-846.
- Yardan, T., Cevik, Y., Donderici, O., Kavalci, C., Yilmaz, F. M., Yilmaz, G., . . . Sezer, A. A. (2009). Elevated serum S100B protein and neuron-specific enolase levels in carbon monoxide poisoning. *The American journal of emergency medicine*, 27(7), 838-842.
- Yernool, D., Boudker, O., Folta-Stogniew, E., & Gouaux, E. (2003). Trimeric subunit stoichiometry of the glutamate transporters from *Bacillus caldotenax* and *Bacillus stearothermophilus*. *Biochemistry*, 42(44), 12981-12988.
- Yernool, D., Boudker, O., Jin, Y., & Gouaux, E. (2004). Structure of a glutamate transporter homologue from *Pyrococcus horikoshii*. *Nature*, 431(7010), 811-818.
- Ying, W., Sevigny, M. B., Chen, Y., & Swanson, R. A. (2001). Poly (ADP-ribose) glycohydrolase mediates oxidative and excitotoxic neuronal death. *Proceedings of the National Academy of Sciences*, 98(21), 12227-12232.
- Ying, W., & Swanson, R. A. (2000). The poly (ADP-ribose) glycohydrolase inhibitor gallotannin blocks oxidative astrocyte death. *Neuroreport*, 11(7), 1385-1388.
- Zamzami, N., Maise, C., Metivier, D., & Kroemer, G. (2001). Measurement of membrane permeability and permeability transition of mitochondria. *Methods in cell biology*, 147-158.
- Zamzami, N., Maise, C., Métivier, D., & Kroemer, G. (2001). Measurement of membrane permeability and permeability transition of mitochondria. *Methods in cell biology*, 65, 147-158.
- Zetterberg, H., Smith, D. H., & Blennow, K. (2013). Biomarkers of mild traumatic brain injury in cerebrospinal fluid and blood. *Nature Reviews Neurology*, 9(4), 201-210.
- Zhang, Y., Lu, X., & Bhavnani, B. R. (2003). Equine estrogens differentially inhibit DNA fragmentation induced by glutamate in neuronal cells by modulation of regulatory proteins involved in programmed cell death. *BMC neuroscience*, 4(1), 1.
- Žiak, D., Chvatal, A., & Sykova, E. (1998). Glutamate-, kainate- and NMDA-evoked membrane currents in identified glial cells in rat spinal cord slice. *Physiol Res*, 47, 365-375.
- Zou, H., Li, Y., Liu, X., & Wang, X. (1999). An APAF-1· cytochrome c multimeric complex is a functional apoptosome that activates procaspase-9. *Journal of Biological Chemistry*, 274(17), 11549-11556.



Zou, J. Y., & Crews, F. T. (2005). TNF $\alpha$  potentiates glutamate neurotoxicity by inhibiting glutamate uptake in organotypic brain slice cultures: neuroprotection by NF $\kappa$ B inhibition. *Brain research*, 1034(1), 11-24.

