



Tonks, J., Cloke, G., Lee-Kelland, R., Jary, S., Thoresen, M., Cowan, F. M., & Chakkarapani, E. (2019). Attention and visuo-spatial function in children without cerebral palsy who were cooled for neonatal encephalopathy: a case-control study. *Brain Injury*, 33(7), 894-898. <https://doi.org/10.1080/02699052.2019.1597163>

Peer reviewed version

License (if available):
Other

Link to published version (if available):
[10.1080/02699052.2019.1597163](https://doi.org/10.1080/02699052.2019.1597163)


[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Taylor & Francis at <https://doi.org/10.1080/02699052.2019.1597163> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



Attention and visuo-spatial function in children without cerebral palsy who were 'cooled' for neonatal encephalopathy: a case-control study.

Journal:	<i>Brain Injury</i>
Manuscript ID	Draft
Manuscript Type:	Original Paper
Keywords:	neonatal encephalopathy, therapeutic hypothermia, attention, visuo-spatial function, dorsal stream

SCHOLARONE™
Manuscripts

Abstract:

Objectives: Dorsal-stream functions are vulnerable to early brain injury associated with neonatal encephalopathy (NE) following perinatal asphyxia, even in children not developing cerebral palsy (CP). Since therapeutic hypothermia (TH) became the standard treatment for NE, the incidence of CP is reduced but the impact on dorsal-stream functions is unknown. We aimed to compare dorsal-stream functions in TH-treated survivors of NE, without CP, with those of matched controls.

Methods: We administered tests of dorsal-stream function to 29 case children aged 6-to-8 years treated with TH for NE and without CP, and 20 age, sex and social class matched controls. We used the Conner's Continuous Performance Test (CPT) 2nd Edition to assess attentiveness, based upon Hit Reaction Time (HRT) percentile score and HRT standard error percentile, the CPT HRT block change measure to assess sustained attention and the NEPSY-II block construction and arrows tests to assess visuo-spatial performance and mental rotation.

Results: Case children performed significantly worse than controls on measures of attention and visuo-spatial function.

Conclusions: Children given TH treatment for NE can have subtle attention difficulties with slower reaction times and reduced visuo-spatial processing. These findings illustrate the continued vulnerability of dorsal-stream functions following NE despite the use of TH.

INTRODUCTION

Neonatal encephalopathy (NE) secondary to intrapartum events affects nearly 1.15 million babies worldwide. (Lee et al., 2013). Children with NE are at an increased risk of developing motor, cognitive and visuo-spatial, sensorimotor, attention and executive impairments (Marlow et al., 2005). Clinical trials have shown that moderate cooling following NE,

1
2
3 (reducing the core temperature to 33-34°C within 6 hours of birth for three days - termed
4 therapeutic hypothermia (TH)) reduces death and disability at 18 months (Jacobs et al., 2013),
5
6 the incidence and severity of cerebral palsy (CP) (Jary et al., 2015) and furthermore, survival
7
8 with an IQ>85 is increased (Azzopardi et al., 2014). TH is now the standard care for infants
9
10 with NE (NICE, May 2010).
11
12
13
14

15 In one of the two clinical trials of TH known as ‘the Toby trial,’(Azzopardi et al., 2014) the
16
17 infants who received TH, compared to the non-cooled group, had higher attention and
18
19 executive function scores and comparable visuo-spatial function assessed using the
20
21 “NEuroPSYchological” Assessment second edition-NEPSY-II) (Holcomb and Davis, 2011).
22
23 In the other clinical trial, the NICHD trial (Shankaran et al., 2012), the proportion of children
24
25 with attention and executive function or visuo-spatial scores (below a standard score of 70) did
26
27 not differ between the TH and non-cooled groups. Both studies included children with and
28
29 without CP and compared the attention, executive and visuo-spatial scores to the test norm.
30
31 Although cooling reduced the occurrence of severe disabilities, it is not known whether subtle
32
33 impairments in attention and visuo-spatial function will remain in cooled children without
34
35 severe disabilities such as CP.
36
37
38
39
40

41 It is methodologically problematic that both of these studies reported only index scores (created
42
43 from amalgamating individual tests designed to assess separate sub-components of executive
44
45 function and attention). Executive functions include a range of abilities, such as assigning
46
47 priority to tasks, sequencing, planning, initiation, inhibition, fluency emotional regulation,
48
49 processing speed and managing the resources of attention. Separate tests are used to assess
50
51 these discrete components of executive function, but then they are commonly collapsed in
52
53 assessment tools (as index scores) to create a single index score of executive function and
54
55 attention. Lower-scores are then commonly ‘smoothed-out’ by the averaged, or amalgamated
56
57 index-score. Such an approach can inflate Type II error rates (Tonks et al., 2014). Thus in both
58
59
60

1
2
3 of the studies to date, reporting only index scores could have masked more subtle strengths and
4
5 weaknesses in the neuropsychological profiles of these groups. We further note that the
6
7 attention tasks used in the studies were of relatively short duration, and so placed low demands
8
9 upon sustained attention.
10

11
12
13 One theory of relevance in understanding how NE may affect attention and visuo-spatial
14
15 functions is the Dorsal Stream Vulnerability model (Atkinson et al., 2003, Atkinson &
16
17 Braddick, 2011). Visual processing is divided into two specialized extra-striate streams; a
18
19 ventral stream, or ‘what’ stream projecting from the visual cortex to temporal lobe structures,
20
21 which is integral to object processing; and a dorsal stream, or ‘where’ stream projecting from
22
23 the visual cortex to the parietal lobe, which is essential in spatial processing (Felleman and Van
24
25 Essen, 1991, Young, 1992, Milner and Goodale, 2006). Selective, rapid visual attention
26
27 processes in location and movement of objects, and visuo-spatial functions (such as spatial
28
29 transformation and visuo-spatial relations) are dependent on the “dorsal stream” pathway in
30
31 the brain (Spencer et al., 2000, Atkinson and Braddick, 2011). Previous studies have indicated
32
33 that abilities that we now associate with dorsal stream function do appear to be affected by
34
35 perinatal brain injury, and particularly hypoxic-ischemic injury (Mercuri et al., 1997, Mercuri
36
37 et al., 1999). It is therefore reasonable to hypothesize that children cooled for NE who do not
38
39 have problems severe enough to cause CP could still have effects on dorsal stream functions,
40
41 despite treatment. Children with early (though not neonatal) brain injury commonly have
42
43 problems with visuo-spatial relations, which often become more apparent at around 10 years
44
45 (Tonks et al., 2009a). That is, visual-spatial tests are sensitive in differentiating between healthy
46
47 children and children after brain injury. Similar findings have been reported in relation to
48
49 attention difficulties which can also become more evident from 10 years, with attention-
50
51 switching skills having some interdependence with later developing executive function (Tonks
52
53 et al., 2011).
54
55
56
57
58
59
60

1
2
3 In summary, little is known about subtle visual attention processing, and visuo-spatial abilities
4 associated with dorsal-stream function in children who were cooled as standard care for NE.
5
6 In this study we compared performance on tests of attention and visuo-spatial function,
7
8 between children aged 6-8 years who were cooled for NE and did not develop CP, and matched
9
10 'healthy' control children, in order to determine whether there is evidence of subtle, enduring
11
12 brain injury effects.
13
14
15
16
17
18
19
20

21 **METHOD**

22
23 This was a prospective case-control study conducted between October 2016 and October 2017
24 within the University of Bristol, UK with ethics approval from the North Bristol Research
25 Ethics Committee (REC ID:15. SW.0148) and the Health Research Authority.
26
27
28
29

30 **Participants:**

31
32
33 **Cases:** Eligibility criteria were as follows: gestation at birth ≥ 36 weeks, treatment with TH as
34 standard clinical care, based on TOBY trial eligibility criteria including signs of perinatal
35 asphyxia and moderate to severe encephalopathy confirmed by amplitude integrated
36 electroencephalogram (aEEG)(Azzopardi et al 2009). All the children had English as their
37 primary language and did not have an additional medical diagnosis other than NE.
38
39
40
41
42
43
44
45
46

47 Cases were selected from the cohort of children who received TH for NE between 2008 and
48 2011. These data are maintained by the Bristol Neonatal Neurosciences group at St Michael's
49 Hospital, Bristol, UK. A diagnosis of CP was ruled-out from the assessment at 2 years and
50 reconfirmed at 6-8 years assessment. Forty potential cases were identified after
51 inclusion/exclusion criteria were applied.
52
53
54
55
56
57
58
59
60

1
2
3 **Controls:** The study sample was compared to a control group of children matched for age, sex
4 and socio-economic status (SES). SES was defined based on the national readership survey
5 (NRS) A, B, C1, C2, D and E criteria (2015). Exclusion criteria included children born before
6 36 weeks gestation, any history of NE or other medical issues of a neurological nature, and
7 English not being the primary language. Children who were best matched to the case in terms
8 of age, sex and social class were invited to participate in the study.
9
10
11
12
13
14
15
16
17
18
19

20 **Procedure and measures**

21
22 Cases and controls attended for a single day of testing at the Clinical Research and Imaging
23 Centre (CRIC), Bristol. Written informed consent was obtained from the parents and the
24 participants. We administered selected, standardised tests that assess performance in visuo-
25 spatial function and attention (details of each test are provided below). Each participant was
26 tested individually in a quiet testing room by assessors who were unaware of case/control
27 status.
28
29
30
31
32
33
34
35
36
37
38

39 **Measures.**

40 ***Attention Measures:***

41
42 The Conners Continuous Performance Test (CPT) 2nd Edition (16) (Conners et al, 2004)
43 assesses attention-related problems in individuals from 6-years through to adulthood. During a
44 14-minute, 360 trial administration, children are required to respond by clicking a computer
45 mouse when any letter appears on a computer screen, except the non-target letter "X." By
46 assessing the participant's response times across the test (and comparing the participant's
47 mean-score to the mean score of the published normative sample), it is possible to understand
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 performance in areas of inattentiveness, and/or sustained attention. Three CPT measures that
4
5 provide percentile scores for response time were focused upon for this study. These were:
6
7

- 8
9 • ***Overall Hit Reaction Time (HRT)***: Each time a trial is presented to the participant and
10 they correctly respond, a response time is recorded. The overall HRT is the mean
11 correct response-time from the 360 trials, which is given as a percentile for comparison
12 against the population norm. It provides an indication of efficiency in selectively
13 processing targets and can therefore index inattentiveness. Higher scores indicate worse
14 performance.
15
- 16 • ***HRT Standard Error of the Mean***: this is a measure of response speed consistency for
17 correct items throughout the administration of the test. The higher the overall standard
18 error (SE) of mean percentile, the greater the inconsistency in the response speed.
19 Response speed inconsistency is indicative of inattentiveness and can indicate when a
20 participant is less engaged or processing stimuli less efficiently during parts of the test-
21 administration. The mean percentile for both overall HRT and HRT SE of the Mean is
22 50. One standard deviation (SD) is a score of 84, 2 SDs is 98.
23
- 24 • ***HRT by Block (HRT Block Change)*** HRT Block Change measures change in reaction
25 time across the duration of the test. High values of HRT Block Change indicate a
26 substantial slowing in reaction times. Low values indicate that responses got quicker as
27 the test progressed. If reaction times slow down, as indicated by a higher HRT Block
28 Change score, the participant's information processing efficiency declines, and a loss
29 of sustained attention is indicated.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Visuo-spatial mental rotation and visual discrimination:**

56
57
58
59
60

1
2
3 The NEPSY-II Block construction test (Korkman et al., 2007) was used to assess visual-spatial
4 performance in a task of mental rotation/transformation. Participants were presented with
5 images of block designs (towers and other forms) and blocks for design construction. Each
6 image depicts complex arrangements of various numbers of 3D cubes positioned in structures.
7
8 The task is to visualise and construct the designs based upon studying the image before them.
9
10 Scaled-scores were utilised.
11
12
13
14
15
16
17
18

19 The NEPSY-II Arrows' test (Korkman et al., 2007) was administered as a measure of visual
20 discriminatory skill. Participants were presented with images of arrows around a target. Only
21 two arrows were on a trajectory to directly hit the target. The task is to identify the arrows that
22 will hit the centre of the target. Scaled scores were utilised. Average scaled scores range from
23 8 to 12, scores <7 are 1SD below the mean, scores of 5 are considered borderline for
24 impairment and scores of 4 or less are considered to represent impairment.
25
26
27
28
29
30
31
32
33
34

35 **Study size**

36
37 Forty eligible case children were identified. Seven families could not be traced and 4 families
38 were not willing to participate in the study leaving 29 participants in the case group. One of the
39 case-participants did not complete the CPT-2 test leaving 28 case children for the analysis of
40 CPT performance. The control children were recruited from schools in and around the Bristol
41 area. We contacted 133 schools to ask if study information could be put in their school-
42 newsletter. In accordance with Ethics protocols, schools were excluded if any case participant
43 was a pupil. Fifty-one families volunteered to participate. We best-matched 20 control children
44 with the case group based on age, sex and social class. Two of the control-participants refused
45 to complete the CPT-2 test leaving 18 control children for the analysis of CPT performance.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The NEPSY-II Block construction and the Arrows test were completed by all 29 cases and 20
4
5 control children.
6
7
8
9

10 **Statistical methods**

11
12
13 Checks for *departure from normality* were undertaken on all data for analysis using Shapiro-
14
15 Wilk's test. Checks of *homogeneity of variance* were undertaken using Levene's test. We
16
17 used independent *t-tests* with group (case/controls) as the independent variable. Analysis 1
18
19 was undertaken with HRT Block Change as the dependent variable, analysis 2 was undertaken
20
21 with visuo-spatial (Block construction) results as the dependent variable, and analysis 3 was
22
23 undertaken with visual discrimination (NEPSY-II Arrows) as the dependent variable. Two
24
25 further analyses were undertaken (in considering Overall HRT and HRT SE of the mean)
26
27 using Mann-Whitney U testing on mean ranks due to unequal variance and departure from
28
29 normality. All analyses were performed using IBM SPSS version 24 (IBM Corp., Armonk,
30
31 NY, USA). All reported p values are two tailed and $p < 0.05$ was considered as significant.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

In case children the median (IQR) Apgar score at 10 mins was 6 (5, 8). Median (IQR) pH value within 1 hour of birth was 6.9 (6.8-7.1). The aEEG patterns were moderately abnormal in 93% (27/29) and severely abnormal in 7% (2/29).

Cases were matched with controls for age (mean (SD): 7.0 years(0.5) versus 7.0 years (0.6)), sex (males: 69% versus 60%) and social class (upper middle class: 14% versus 15%; middle class: 28% versus 30%; lower middle class 21% versus 20%; skilled working class 24% versus 25%; working class: 10% versus 10% and casual worker or unemployed: 3% versus 0%).

Differential Performance in attention:

Comparison of the mean rank of the overall HRT percentiles indicated significantly slower response-times in the case children (84.1, Mean Rank 27.0) compared to the control children, (67.3, Mean Rank 17.89), [U=151.5, p=.024]. Similarly, HRT SE of the mean indicated greater variability in performance in case children (Mean Rank 26.8) compared to the control children (Mean Rank 18.2), [U=157, p=.032]. Figure 1 shows the distribution of scores for cases and controls on these measures, where higher percentiles indicate slower response-times (and therefore worsening score). A larger number of cases compared to controls performed poorly on these tests; the proportion of case and control children below 2SD (98th percentile) were 32% versus 11% respectively for HRT, and 18% versus 11% respectively for HRT SE. Additionally, many more cases than controls performed below 1SD (84th percentile), specifically, 57% versus 33% for HRT percentile and 50% versus 33% for HRT SE.




Figure 1: Distribution of scores for cases and controls: Overall Hit Reaction Time (HRT), HRT standard error of the mean and HRT change across each block of the Conner's Continuous Performance Test. Dotted lines mark 1 and 2 SD cut-offs.

In considering HRT by block, the control group mean was marginally below the 50th percentile (M=49.1, SD 23.9) while the mean for the case children was higher (M=61.9, SD 18.4) indicating that they had worse (slowed) performance as the assessment progressed. This difference was significant [$t(44)=-2.04, p = .047$]. Figure 1 shows the distribution of scores for cases and controls for HRT by block, where a higher percentile indicates slower response-times (and therefore worsening score). Again, more cases (14%) than controls (0%) perform below 1SD (84th percentile).

DISCUSSION

We undertook tests of attention ability and visual-spatial processing in children treated with TH after NE who did not develop CP and compared them to matched controls. In summarising our findings in relation to attention, we note that the case children compared to the controls had higher response speed percentiles indicating slower response times, greater inconsistency in their response speed and worsening reaction times over the duration of assessment. Taken together these results are indicative of reduced efficiency in attention processing, and loss of sustained attention at a rate that is significantly different to age-matched control peers. Analysis of group differences in visual-spatial function, similarly showed that the case group performed significantly worse than same age peers.

Attention problems have been reported in the pre-cooling era for children who were defined as having moderate to severe NE based on a broad clinical definition and who did not develop CP. Of those children nearly 11% were reported to have attention problems assessed using the NEPSY-II and Child Behaviour Checklist at 5-16 years (Hayes et al., 2018). At 6.5-9 years children with medical histories of severe encephalopathy had significantly lower attention and executive scores assessed using the NEPSY than children with moderate encephalopathy and a control group. Additionally, the moderate encephalopathy group had lower scores than a matched control group (Marlow et al., 2005). At 15-19 years, children who have had moderate encephalopathy were compared to control children and were noted to score higher on the Inattention subscale of the Attention Deficit Hyperactivity Disorder (ADHD) Rating Scale-IV (Lindström et al., 2006).

The school age (6-7 years) follow-up data from the clinical trials of TH for NE (including children with CP) reported varying results, with the TOBY trial finding higher attention and executive scores in the cooled group than the non-cooled group (Azzopardi et al., 2014) and the NICHD trial reporting similar scores between the cooled and control group (Shankaran et

1
2
3 al., 2012). The clinical trials combined attention and executive function into a single measure
4
5 in order to understand the effects of TH as did the studies in the 'pre-hypothermia era'. Given
6
7 that executive functions develop across the span of childhood and adolescence, with most
8
9 significant advances arising with pre-frontal cortex development after 10-to-12 years in healthy
10
11 maturation (Tonks et al., 2017), it seems probable that any immediate, but subtle attention-
12
13 based difficulties could have been concealed by the methodological approach used, which did
14
15 not base test selection on early brain injury models of post-injury attention and visuo-spatial
16
17 development. We compared performance differences between healthy children and children
18
19 treated with TH after NE, focusing more specifically upon the dorsal-stream vulnerability
20
21 hypothesis in exploring outcome after TH. Our approach recognises a temporal (early vs late)
22
23 evaluation of the effects of NE, which considers the effect of brain growth and the changing
24
25 nature of neuropsychological abilities as children mature. Dorsal-stream functions of visuo-
26
27 spatial mental rotation and attention efficiency in terms of response-times were focussed upon
28
29 here, based upon the current age of the sample, and the patterns of neuropsychological
30
31 difficulty that might be predicted after NE. Mean HRT is shown to be associated with
32
33 hyperactivity and mean HRT SE is related to hyperactive-impulsive and inattention symptoms
34
35 in a cohort of children with attention deficit hyperactivity disorder (Epstein et al., 2003).
36
37
38
39
40
41
42

43 In non-cooled children without CP, visuo-spatial function was reported to be impaired in nearly
44
45 5% of children aged 3-4 years (Hayes et al., 2018), and children who have had severe NE had
46
47 significantly lower scores than children with moderate encephalopathy at 6.5-9 years (Marlow
48
49 et al., 2005). In children aged 6-7 years (including children with CP) who underwent TH the
50
51 visuo-spatial processing scores did not differ from the non-cooled group (Azzopardi et al.,
52
53 2014, Shankaran et al., 2012). Our findings support the dorsal-stream vulnerability model in
54
55 this age group and our findings indicate that sustained attention and visual-spatial processing
56
57
58
59
60

1
2
3 can be negatively affected in children with NE following TH, much as can be observed in
4
5 children who had NE and who have received no treatment.
6
7
8
9

10
11 Our study is limited by sample-size. Future research utilizing a longitudinal approach with
12
13 larger samples is needed in order to enhance understanding of outcome, particularly in the
14
15 context of age-related brain maturation.
16
17
18
19
20

21 **In conclusion**, based upon this preliminary study, despite therapeutic cooling, children who
22
23 have had NE can have problems with attention and visuo-spatial function even in the absence
24
25 developing severer problems such as CP. It, therefore remains important to maintain a temporal
26
27 approach in following-up children treated with TH after NE, seeking to remediate difficulties
28
29 that might emerge for individuals.
30
31
32
33
34

35 **Funding details**

36
37 This work was supported by the Baily Thomas Charitable Fund under grant 3548/6501 and
38
39 the David Telling Charitable Trust under grant 357.
40
41

42 **Disclosure statement**

43
44
45 The authors do not have any conflicts of interest to disclose.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Atkinson J, Braddick O. From genes to brain development to phenotypic behavior: “dorsal-stream vulnerability” in relation to spatial cognition, attention, and planning of actions in Williams syndrome (WS) and other developmental disorders. In *Progress in brain research* 2011 Jan 1 (Vol. 189, pp. 261-283). Elsevier.

Atkinson J, Braddick O, Anker S, Curran W, Andrew R, Wattam-Bell J, Braddick F. Neurobiological models of visuospatial cognition in children with Williams syndrome: measures of dorsal-stream and frontal function. *Developmental Neuropsychology*. 2003 Feb 1;23(1-2):139-72.

Azzopardi DV, Strohm B, Edwards AD, Dyet L, Halliday HL, Juszczak E, Kapellou O, Levene M, Marlow N, Porter E, Thoresen M. Moderate hypothermia to treat perinatal asphyxial encephalopathy. *New England Journal of Medicine*. 2009 Oct 1;361(14):1349-58.

Azzopardi D, Strohm B, Marlow N, Brocklehurst P, Deierl A, Eddama O, Goodwin J, Halliday HL, Juszczak E, Kapellou O, Levene M. Effects of hypothermia for perinatal asphyxia on childhood outcomes. *New England Journal of Medicine*. 2014 Jul 10;371(2):140-9.

Conners CK, Staff MH, Connelly V, Campbell S, MacLean M, Barnes J. Conners' continuous performance Test II (CPT II v. 5). Multi-Health Syst Inc. 2000; 29:175-96.

1
2
3 Epstein JN, Erkanli A, Conners CK, Klaric J, Costello JE, Angold A. Relations between
4 continuous performance test performance measures and ADHD behaviors. *Journal of*
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Abnormal Child Psychology. 2003 Oct 1;31(5):543-54.

Felleman DJ, Van DE. Distributed hierarchical processing in the primate cerebral cortex.
Cerebral Cortex (New York, NY: 1991). 1991;1(1):1-47.

Hayes BC, Doherty E, Grehan A, Madigan C, McGarvey C, Mulvany S, Matthews TG, King
MD. Neurodevelopmental outcome in survivors of hypoxic ischemic encephalopathy without
cerebral palsy. *European Journal of Pediatrics*. 2018 Jan 1;177(1):19-32.

Holcomb MJ, Davis AS. NEPSY-II. In *Encyclopedia of Child Behavior and Development 2011*
(pp. 1006-1008). Springer, Boston, MA.

Jacobs S, Hunt R, Tarnow-Mordi W, Inder T, Davis P. 181 cooling for newborns with hypoxic
ischaemic encephalopathy. *Pediatric Research*. 2005 Aug;58(2):385.

Jary S, Smit E, Liu X, Cowan FM, Thoresen M. Less severe cerebral palsy outcomes in infants
treated with therapeutic hypothermia. *Acta Paediatrica*. 2015 Dec;104(12):1241-7.

Korkman M, Kirk U, Kemp S. NEPSY-II: Clinical and interpretive manual. San Antonio, TX:
The Psychological Corporation. 2007.

1
2
3
4
5
6 Lee AC, Kozuki N, Blencowe H, Vos T, Bahalim A, Darmstadt GL, Niermeyer S, Ellis M,
7
8 Robertson NJ, Cousens S, Lawn JE. Intrapartum-related neonatal encephalopathy incidence
9
10 and impairment at regional and global levels for 2010 with trends from 1990. *Pediatric*
11
12 *Research*. 2013 Dec 20;74(S1):50.

13
14
15
16
17
18
19 Lindström K, Lagerroos P, Gillberg C, Fernell E. Teenage outcome after being born at term
20
21 with moderate neonatal encephalopathy. *Pediatric Neurology*. 2006 Oct 1;35(4):268-74.

22
23
24
25
26
27 Marlow N, Rose AS, Rands CE, Draper ES. Neuropsychological and educational problems at
28
29 school age associated with neonatal encephalopathy. *Archives of Disease in Childhood-Fetal*
30
31 *and Neonatal Edition*. 2005 Sep 1;90(5):380-7.

32
33
34
35
36
37 Mercuri E, Atkinson J, Braddick O, Anker S, Cowan F, Rutherford M, Pennock J, Dubowitz
38
39 L. Visual function in full-term infants with hypoxic-ischaemic encephalopathy.
40
41 *Neuropediatrics*. 1997 Jun;28(03):155-61.

42
43
44
45
46
47
48 Mercuri E, Haataja L, Guzzetta A, Anker S, Cowan F, Rutherford M, Andrew R, Braddick O,
49
50 Cioni G, Dubowitz L, Atkinson J. Visual function in term infants with hypoxic-ischaemic
51
52 insults: correlation with neurodevelopment at 2 years of age. *Archives of Disease in Childhood-*
53
54 *Fetal and Neonatal Edition*. 1999 Mar 1;80(2):F99-104.

1
2
3 Milner D, Goodale M. The visual brain in action. Oxford University Press; 2006 Oct 12.
4
5
6
7
8

9 Hazeldine B, Thyagarajan B, Grant M, Chakkarapani E. Survey of nutritional practices during
10 therapeutic hypothermia for hypoxic-ischaemic encephalopathy. *BMJ paediatrics open*.
11 2017;1(1).
12
13
14
15
16
17
18

19 Shankaran S, Pappas A, McDonald SA, Vohr BR, Hintz SR, Yolton K, Gustafson KE, Leach
20 TM, Green C, Bara R, Huitema CM. Childhood outcomes after hypothermia for neonatal
21 encephalopathy. *New England Journal of Medicine*. 2012 May 31;366(22):2085-92.
22
23
24
25
26
27
28
29

30 Spencer J, O'Brien J, Riggs K, Braddick O, Atkinson J, Wattam-Bell J. Motion processing in
31 autism: evidence for a dorsal stream deficiency. *Neuroreport*. 2000 Aug 21;11(12):2765-7.
32
33
34
35
36
37
38

39 Tonks J, Williams WH, Mounce L, Harris D, Frampton I, Yates P, Slater A. 'Trails B or not
40 Trails B?' Is attention-switching a useful outcome measure?. *Brain Injury*. 2011 Sep
41 1;25(10):958-64.
42
43
44
45
46
47
48

49 Tonks J, Williams WH, Slater A, Frampton I. Is damage to the pre-frontal cortex dormant until
50 adolescence, or difficult to detect? Looking for keys that unlock executive functions in children
51 in the wrong place. *Medical Hypotheses*. 2017 Oct 1;108:24-30.
52
53
54
55
56
57
58
59
60

1
2
3 Tonks J, Yates P, Slater A, Williams WH, Frampton I. Visual-spatial functioning as an early
4 indicator of socioemotional difficulties. *Developmental Neurorehabilitation*. 2009 Jan
5
6
7
8 1;12(5):313-9.
9

10
11
12
13 Tonks J, Yates PJ, Williams HW, Frampton I, Slater A. Measurement Issues:
14 Neuropsychological assessment with children and adolescents; unlocking the mysticism,
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

114 Tonks J, Yates PJ, Williams HW, Frampton I, Slater A. Measurement Issues:
115 Neuropsychological assessment with children and adolescents; unlocking the mysticism,
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160

114 May;19(2):151-8.

114 Young MP. Objective analysis of the topological organization of the primate cortical visual
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160

114 Young MP. Objective analysis of the topological organization of the primate cortical visual
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160

114 system. *Nature*. 1992 Jul;358(6382):152.