

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/207288>

Please be advised that this information was generated on 2019-12-31 and may be subject to change.



Demethylated hopanoids in '*Ca. Methylomirabilis oxyfera*' as biomarkers for environmental nitrite-dependent methane oxidation

Nadine T. Smit^{a,*}, Darci Rush^a, Diana X. Sahonero-Canavesi^a, Monique Verweij^a, Olivia Rasigraf^{b,1}, Simon Guerrero Cruz^{b,2}, Mike S.M. Jetten^b, Jaap S. Sinninghe Damsté^{a,c}, Stefan Schouten^{a,c}

^a NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry, and Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands

^b Department of Microbiology, Institute for Water and Wetland Research, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, the Netherlands

^c Department of Earth Sciences, Faculty of Geosciences, Utrecht University, P.O. Box 80.121, 3508 TA Utrecht, the Netherlands

ARTICLE INFO

Article history:

Received 11 April 2019

Received in revised form 2 July 2019

Accepted 16 July 2019

Available online 17 July 2019

Keywords:

Demethylated hopanoids

Candidatus Methylomirabilis oxyfera

Nitrite-dependent methane oxidation

Trisnorhopanoids

3-Methyl-22,29,30-trisnorhopan-21-one

Methylation

ABSTRACT

Hopanoids are lipids that are widespread in the bacterial domain and well established molecular biomarkers in modern and paleo environments. In particular, the occurrence of ¹³C-depleted 3-methylated hopanoids are characteristic of aerobic bacteria involved in methane oxidation. Previously the intra-aerobic methanotroph '*Candidatus Methylomirabilis oxyfera*' ('*Ca. M. oxyfera*'), which performs nitrite-dependent methane oxidation in anoxic environments, has been shown to synthesize bacteriohopanepolyols (BHPs) and their 3-methylated counterparts. However, since '*Ca. M. oxyfera*' does not utilize methane as a carbon source, its biomass and lipids do not show the characteristic ¹³C-depletion. Therefore, the detection of '*Ca. M. oxyfera*' in various environments is challenging, and still underexplored. Here, we re-investigated the hopanoid content of '*Ca. M. oxyfera*' bacteria using enrichment cultures. We found the GC-amenable hopanoids of '*Ca. M. oxyfera*' to be dominated by four demethylated hopanoids of which only one, 22,29,30-trisnorhopan-21-one, had been identified previously. The three novel hopanoids were tentatively identified as 22,29,30-trisnorhopan-21-ol, 3-methyl-22,29,30-trisnorhopan-21-one and 3-methyl-22,29,30-trisnorhopan-21-ol. These unique demethylated hopanoids are most likely biosynthesized directly by '*Ca. M. oxyfera*' bacteria. Bioinformatical analysis of the '*Ca. M. oxyfera*' genome revealed potential candidate genes responsible for the demethylation of hopanoids. For the sensitive detection of the four trisnorhopanoid biomarkers in environmental samples, a multiple reaction monitoring (MRM) method was developed and used to successfully detect the trisnorhopanoids in a peatland where the presence of '*Ca. M. oxyfera*' had been confirmed previously by DNA-based analyses. These new biomarkers may be a novel tool to trace nitrite-dependent methane oxidation in various (past) environments.

© 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Hopanoids are pentacyclic triterpenoids that modify the properties of membranes in microorganisms and can serve as molecular biomarkers in modern and paleo environments (Ourisson et al., 1979; Ourisson and Albrecht, 1992). They are biosynthesized by different bacterial phyla and have been used to trace the presence of these bacteria as far back as the Proterozoic (Brocks et al., 2005).

It was previously believed that hopanoids indicate oxic environmental conditions, however this changed with the discovery of hopanoid biosynthesis in anaerobic bacteria such as fermentative bacteria (Neunlist et al., 1985; Llopiz et al., 1992), anammox bacteria (Sinninghe Damsté et al., 2004), sulfate reducing bacteria (SRB) (*Desulfovibrio*) (Blumenberg et al., 2006, 2012) and bacteria of the *Geobacter* genus (Eickhoff et al., 2013). The molecular structures of hopanoids synthesized by bacteria are structurally very diverse and range from simple C₃₀ hopanols, such as diplopterol, to extended C₃₅ polyols known as bacteriohopanepolyols (BHPs). During diagenesis, these hopanoids can be abiotically or microbially altered (Rohmer and Ourisson, 1976; Albaiges and Albrecht, 1979) resulting in demethylation, aromatization, sulfuration, skeleton rearrangements and loss or transformation of

* Corresponding author.

E-mail address: nadine.smit@nioz.nl (N.T. Smit).

¹ Present address: German Research Centre for Geosciences (GFZ), Section 3.7 Geomicrobiology, Potsdam, Germany.

² Present address: Global Center of Expertise Microbiology & Adventitious Viral Agents Merck, Oss, the Netherlands.

functional groups. Both BHPs and derived diagenetic products are extensively used as lipid biomarkers for (specific) microbial communities and their metabolic role in modern and ancient ecosystems (e.g., Brocks and Pearson, 2005; Talbot et al., 2007).

One important example is the use of hopanoids to trace microbes involved in methane oxidation. BHPs with penta- and hexa-functionalized side chains containing an amine group at the C-35 position are characteristic of methane-oxidizing bacteria (MOB) from the alpha- and gammaproteobacterial classes (Talbot and Farrimond, 2007; van Winden et al., 2012). Another important structural feature in some hopanoids is the methylation at the C-3 position of the A-ring. This methylation was first shown in acetic acid bacteria (Rohmer and Ourisson, 1976) and later in MOB (Zundel and Rohmer, 1985). Subsequently, C-3 methylated hopanoids were used as biomarkers for the presence of aerobic methanotrophy in ancient sediments (Summons and Jahnke, 1992). However, Welander and Summons (2012) identified the *hpnR* gene responsible for 3-methylation in hopanoids and showed that it is widely distributed among various bacterial taxa, including those not involved in the methane cycle. Nevertheless, the origin of hopanoids from MOB can be inferred from their depletion in ^{13}C ($\delta^{13}\text{C}$ below ca. -40‰) as usually ^{13}C -depleted CH_4 is used as carbon source for their biomass (Summons et al., 1994; Jahnke et al., 1999). Consequently, C-3 methylated hopanoids can be used as an indicator for methanotrophy if accompanied by additional evidence of ^{13}C -depleted carbon isotope values.

An apparent exception to this ^{13}C depletion is the intra-aerobic methanotroph, '*Candidatus Methylopirabilis oxyfera*' ('*Ca. M. oxyfera*'), which belongs to the NC10 phylum and occurs in clusters with '*Ca. Methanoperedens nitroreducens*' archaea (ANME-2d) (Raghoebarsing et al., 2006; Ettwig et al., 2010; Vaksmaa et al., 2017). '*Ca. M. oxyfera*' lives under anoxic conditions and reduces nitrite to nitric oxide, and subsequently is believed to produce its own intracellular oxygen from nitric oxide for the intra-aerobic oxidation of methane (Ettwig et al., 2010). Thus, '*Ca. M. oxyfera*' directly connects the methane and nitrogen cycle (Ettwig et al., 2010). In contrast to most other MOB lipids, '*Ca. M. oxyfera*' lipids were shown not to be depleted in ^{13}C from methane (Kool et al., 2012, 2014), as they autotrophically fix carbon dioxide via the Calvin cycle (Rasigraf et al., 2014). These lipids include 3-methyl hopanoids, first recognized through genomic analysis (Welander and Summons, 2012), such as the rare BHP-hexol and a novel BHP identified as 3-methyl-BHP-hexol (Kool et al., 2014). Further studies of the lipid inventory of '*Ca. M. oxyfera*' showed that they produce rare methylated fatty acids such as $10\text{MeC}_{16:0}$ and $10\text{MeC}_{16:1\Delta 7}$ (Kool et al., 2012). However, the use of fatty acids as biomarkers for nitrite-dependent methane oxidation is limited to more recent geological time periods because these lipids are not resistant to alteration processes (Atlas and Bartha, 1973; Wenger et al., 2002).

Here, we re-investigated the hopanoid content of '*Ca. M. oxyfera*' biomass but now focused on those amenable to gas chromatography (GC) analysis. These biomarkers were then compared to GC-amenable hopanoids detected in a core from a peatland where the occurrence of '*Ca. M. oxyfera*' had been confirmed previously.

2. Material and methods

2.1. '*Ca. M. oxyfera*' enrichment cultures

An enrichment culture of '*Ca. M. oxyfera*' bacteria was obtained from a bioreactor operated under the conditions described previously by Ettwig et al. (2009). The reactor contained bright flocculent biomass which originated from active '*Ca. M. oxyfera*'

bacteria, and black material from the bottom of the bioreactor that was most likely dead '*Ca. M. oxyfera*' biomass. The bacteria in the bioreactor contained ca. 67% of '*Ca. M. oxyfera*' while the rest was composed of a mix of ANME-2d archaea and different minor bacteria phyla (e.g., *Anaerolinea thermophila*, *Caldithrix abyssi* or *Melioribacter roseus*) at abundances less than 7% for each phylum. The mineral medium containing $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (0.24 g/L), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.16 g/L), KHCO_3 (0.1 g/L), NaNO_2 (1.39 g/L), KH_2PO_4 (0.01 g/L), 1 mL of trace element solution and 3 mL of iron stock solution was sparged with Ar/CO_2 mixture (95:5, v/v) and supplied to the bioreactor at a rate of 1.5–2 L per day. The trace element solution contained $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (1.44 g/L), $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.6 g/L), CuSO_4 (4 g/L), $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (0.95 g/L), H_3BO_3 (0.07 g/L), $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (1 g/L), $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$ (0.1 g/L), $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (0.48 g/L), SeO_2 (0.13 g/L), CeCl_2 (0.12 g/L) and nitrilotriacetic acid (NTA, 30 g/L) as chelating agent. The iron stock solution contained $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (5 g/L) and NTA (10.31 g/L) as chelating agent. The steady state concentration of nitrite in the bioreactor was kept at 0–5 mg/L. Biomass from a second reactor enriched from Italian paddy field soils consisted of less '*Ca. M. oxyfera*' cell material (30%), more '*Ca. Methanoperedens nitroreducens*' (ANME-2d) archaea (30%) and other microbial communities (40%) as described previously (Vaksmaa et al., 2017).

2.2. Peatland core

A peatland core from the Brunsummerheide peatland, Netherlands (BRH) was sampled and divided into nine sections from 51 to 102 cm depth (i.e. 51–60 cm, from 60 to 95 cm, 95–102 cm) (Kool et al., 2012; Kool et al., 2014). Previous quantitative polymerase chain reaction (qPCR) analysis of '*Ca. M. oxyfera*' specific 16S rRNA genes revealed the peak of their abundance at 70–90 cm depth. In this zone both methane and nitrate showed depletion in pore water concentrations (Zhu et al., 2012).

2.3. Fluorescence in situ hybridization

Fluorescence in situ hybridization (FISH) of '*Ca. M. oxyfera*' enrichments from the two bioreactors was performed for estimation of bacterial and archaeal abundance. Liquid samples of 1.5 mL from an active enrichment culture were centrifuged for 5 min at 7000g (EppendorfTM 5424, Hamburg) and biomass pellets were washed 2 times with 1 mL phosphate-buffered saline (PBS: 130 mM NaCl and 10 mM phosphate buffer at pH 7.4). After washing, the samples were fixed with 4% paraformaldehyde at a 3:1 (v/v) ratio to the sample at 4 °C overnight. FISH was performed as previously described (Ettwig et al., 2008). Formamide concentration in the hybridization buffer was set to 35%. The following oligonucleotide probes were used: DAMOBACT-0193 (CGC TCG CCC CCT TTG GTC) specific for '*Ca. M. oxyfera*'-like bacteria; DAMOARCH-0641 (GGT CCC AAG CCT ACC AGT) specific for '*Ca. M. nitroreducens*'-like archaea; EUB 338 (S-D-Bact-0338-a-A-18) (Amann et al., 1990), EUB 338 II (S-D-Bact_0338-b-A-18) and EUB 338 III (S-D-Bact-0338-c-A-18) (Daims et al., 1999) for most bacteria; S-D-Arch-0915-a-A-20 for most archaea. Images were collected with a Zeiss Axioplan 2 epifluorescence microscope equipped with a CCD camera and processed with the Axiovision software package (Zeiss, Germany).

2.4. Genomic analysis

Identification of potential genes involved in hopanoid biosynthesis in '*Ca. M. oxyfera*' (NCBI: taxid671143) was carried out on the NCBI (National Center for Biotechnology Information) server with the PSI-BLAST algorithm (Schäffer et al., 2001) using characterized proteins from different organisms as queries (Supplementary Table S1). Operon analysis and visualization was performed

by using the Gene Context Tool NG (<http://biocomputo2.ibt.unam.mx/gctng/>) and protein analysis was conducted with Uniprot (UniProt Consortium, 2018).

2.5. Hopanoid analyses

2.5.1. Extraction of hopanoids

Freeze dried biomass of the '*Ca. M. oxyfera*' enrichments as well as freeze dried BRH peat core material (Kool et al., 2012) were extracted using a modified Bligh and Dyer technique (Schouten et al., 2008; Bale et al., 2013). The samples were ultrasonically extracted with a solvent mixture containing methanol (MeOH), dichloromethane (DCM) and phosphate buffer (2:1:0.8, v/v/v). After sonication (10 min) and centrifugation, the solvent layer was collected and the residue re-extracted twice. The combined solvent layers were separated from the aqueous layer by adding additional DCM and phosphate buffer to achieve a ratio of MeOH, DCM and phosphate buffer (1:1:0.9, v/v/v). The separated organic bottom layer was removed and collected while the aqueous layer was washed two more times with DCM. The combined DCM layers were dried under a continuous flow of N₂.

Aliquots of the total lipid extracts (TLEs) (ca. 0.5–3 mg) were base hydrolyzed (saponified), with 2 mL of 1 N KOH in MeOH solution, refluxed for 1 h at 130 °C, the pH adjusted to 5 with a 2 N HCL in MeOH solution, separated with 2 mL bidistilled water and 2 mL DCM and washed with DCM two more times. The combined DCM layers were dried over a Na₂SO₄ column. Afterwards the saponified fraction was methylated with diazomethane in diethyl ether, filtered over a small silica column with ethyl acetate and silylated with pyridine and N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA) at 60 °C for 20 min. These total lipid extracts (TLEs) were analyzed using GC and GC–mass spectrometry (MS).

2.5.2. Instrumental analysis

GC–MS analyses of the TLEs of different '*Ca. M. oxyfera*' biomass and BRH peat core material were performed on an Agilent Technologies GC–MS Triple Quad 7000C in full scan and multiple reaction monitoring (MRM) mode. The chromatography was carried out using a CP-Sil5 CB column (25 m × 0.32 mm with a film thickness of 0.12 μm; Agilent Technologies) with helium as carrier gas (constant flow 2 mL/min). For each sample, 1 μL was injected on-column at 70 °C, the temperature was increased by 20 °C/min to 130 °C, raised further by 4 °C/min to 320 °C, which was held for 10 min for full scan mode and for the MRM analysis. GC–MS full scan analysis was conducted over a mass range of *m/z* 50–850, the gain was set to 3 and with a scan rate of 700 ms. Compounds were identified by comparison with previously published data (Zundel and Rohmer, 1985; Sinninghe Damsté et al., 2004; Kool et al., 2012).

A new GC–MS–MS MRM method was developed to detect hopanoid ketones and alcohols in environmental samples using transitions and collision energies shown in Table 1. The MS1 and MS2 resolution was set on wide, the gain on 10 and the dwell time on 10 ms for all scans. As collision gases in the MRM scan mode, He was admitted at a flow of 1.5 mL/min and N₂ at a flow of 1 mL/min.

3. Results and discussion

3.1. Lipid analysis of '*Ca. M. oxyfera*' reveals novel demethylated hopanoids

The microbial community composition and microstructure of the '*Ca. M. oxyfera*' enrichment culture was assessed by FISH. Based on the average fluorescence signal, we estimated that approximately two-thirds of the biomass granules were composed

Table 1

Specifications of the MRM method for the four '*Ca. M. oxyfera*' trisnorhopanoids, showing the target transition (*m/z*), qualitative transition (*m/z*) and optimized collision energy (V).

	Target transition (<i>m/z</i>)	Qualitative transition (<i>m/z</i>)	Collision energy (V)
trisnorhopan-21-one	384.3 → 191.1	384.3 → 369.3	10
trisnorhopan-21-ol	458.3 → 182.1	458.3 → 237.2	15
3-methyl-trisnorhopan-21-one	398.4 → 205.2	398.4 → 383.3	10
3-methyl-trisnorhopan-21-ol	472.4 → 182.1	472.4 → 237.2	15

of bacteria related to '*Ca. M. oxyfera*' while the rest was composed of a mix of ANME-2d archaea and different minor bacteria phyla (e.g., *Anaerolinea thermophila*, *Caldithrix abyssi* or *Melioribacter roseus*) (Supplementary Fig. S1). The lipid composition of this biomass was dominated by a series of fatty acids with high relative abundances of C_{16:0} (22%), 10MeC_{16:1Δ7} (5%) and 10MeC_{16:0} (37%) fatty acids as described previously (Kool et al., 2012). We also detected a series of hopanoids using mass chromatograms for *m/z* 191 and *m/z* 205 revealing diploptene, diplopterol, C₃₂ 17β,21β-hopanoic acid and some of their 3-methylated homologues (Fig. 1). These hopanoids, especially diploptene and diplopterol, are found in a diverse range of aerobic and anaerobic bacteria (Rohmer et al., 1984; Summons et al., 1994; Sinninghe Damsté et al., 2004; Blumenberg et al., 2006). Furthermore, these hopanoids have been found in various environments like oxygen minimum zones (Wakeham et al., 2007), marine hydrocarbon seeps (Thiel et al., 2003; Niemann et al., 2006), lacustrine sediments (Innes et al., 1997), peatlands (van Winden et al., 2010) or soils (Crossman et al., 2005).

The GC–MS analysis also revealed the presence of four more minor peaks which we tentatively identified as hopanoids with relative abundances each about 5–36% compared to the most abundant hopanoid diplopterol, and 0.1% compared to the most abundant fatty acids, the 10MeC_{16:0} and C_{16:0} (Figs. 1 and 2). Of these four, 22,29,30-trisnorhopan-21-one (compound 1-TNHone in Fig. 1) was tentatively identified based on a published mass spectrum (Simoneit, 1977). This hopanoid was first identified in sediments from the Black Sea (Simoneit, 1977) and has since been found in various marine sediment and water column samples, where it is believed to be formed by aerobic diagenesis of BHPs (Conte et al., 1998; Botz et al., 2007).

Directly eluting after 22,29,30-trisnorhopan-21-one is a compound (labelled 2 in Fig. 1) with a diagnostic fragment of *m/z* 191 suggesting a non-A-ring methylated hopanoid (Fig. 2B). The difference of 74 Da between the molecular ions of 22,29,30-trisnorhopan-21-one and unknown compound 2, as well as ions at *m/z* 73 and 75, suggests the presence of a TMSi-derivatized alcohol. This is confirmed by the second diagnostic fragment of *m/z* 237, compared to the *m/z* 163 in the 22,29,30-trisnorhopan-21-one, suggesting that the alcohol is located at the D- or E-ring position. The fragment at *m/z* 367 indicates that the alcohol is likely at the E-ring position (Fig. 2B). This hopanoid was therefore tentatively identified as 22,29,30-trisnorhopan-21-ol (TNHol).

A similar doublet of peaks (compounds 1a and 2a in Fig. 1) eluted after the 22,29,30-trisnorhopan-21-one and 22,29,30-trisnorhopan-21-ol, respectively. Both mass spectra (Fig. 2C and D) show the diagnostic fragment of *m/z* 205, which indicates a methylation of the A-ring. Furthermore, both have molecular ions which are 14 Da heavier (*m/z* 398 and *m/z* 472) than that of 22,29,30-trisnorhopan-21-one and 22,29,30-trisnorhopan-21-ol, respectively (Fig. 2A and B, respectively). The first eluting




Fig. 1. Gas chromatogram of saponified and silylated total lipid extract from ‘*Ca. M. oxyfera*’ biomass. Enlarged inserts show the TIC (total ion chromatogram) of the hopanoids, the EIC (extracted ion chromatogram) of m/z 191 specific for hopanoids and m/z 205 specific for methylated-hopanoids indicating the main hopanoids in ‘*Ca. M. oxyfera*’ biomass. 3 = diploptene, 5 = diploptero, 6 = C_{32} 17 β ,21 β -hopanoic acid, 7 = C_{32} 17 β ,21 β -30-hydroxy hopanoic acid and one unknown hopanoid (4) as well as their methylated counterparts in the top EIC of m/z 205. New tentatively identified hopanoids for ‘*Ca. M. oxyfera*’ are 1 = 22,29,30-trisnorhopan-21-one (TNHone), 2 = 22,29,30-trisnorhopan-21-ol (TNHol), 1a = 3-methyl-22,29,30-trisnorhopan-21-one (3-Me TNHone) and 2a = 3-methyl-22,29,30-trisnorhopan-21-ol (3-Me TNHol).

compound 1a shows the diagnostic ion of the D/E-ring fragmentation of 22,29,30-trisnorhopan-21-one (m/z 163; Fig. 2C), whereas compound 2a shows that of 22,29,30-trisnorhopan-21-ol (m/z 237; Fig. 2D). Therefore, we tentatively identify these two compounds as the methylated counterparts of 22,29,30-trisnorhopan-21-one (3-Me TNHone) and 22,29,30-trisnorhopan-21-ol (3-Me TNHol), respectively. Previous studies have shown the presence of the HpnR gene responsible for the 3-methylation of hopanoids in the ‘*Ca. M. oxyfera*’ genome (Welander and Summons, 2012), as well as the presence of 3-methylated BHPs of BHP-hexol, BHP-pentol and BHP-tetrol in ‘*Ca. M. oxyfera*’ biomass (Kool et al., 2014), which would suggest that the methylated trisnorhopanoids identified here are methylated at the C-3 position. This is further supported by the elution order of these compounds: 2-methylated hopanoids elute directly after the non-methylated analogue, whereas the 3-methylated hopanoids elute ca. 1–2 min later (Summons and Jahnke, 1990; Farrimond et al., 2004). Here, the two methylated trisnorhopanoids elute 1.5 min later than their non-methylated homologues suggesting 3-methylation. Thus, we tentatively identi-

fied these two methylated compounds as 3-methyl-22,29,30-trisnorhopan-21-one (3-Me TNHone; Fig. 2C) and 3-methyl-22,29,30-trisnorhopan-21-ol (3-Me TNHol; Fig. 2D).

3.2. Origin of (3-methyl) 22,29,30-trisnorhopan-21-one and 22,29,30-trisnorhopan-21-ol

To the best of our knowledge, the three novel hopanoids (22,29,30-trisnorhopan-21-ol, 3-methyl-22,29,30-trisnorhopan-21-one and 3-methyl-22,29,30-trisnorhopan-21-ol) have not been identified before in any other bacterial cultures nor have they previously been detected in environmental samples. Since ‘*Ca. M. oxyfera*’ biomass derives from highly enriched, but not pure cultures, it cannot be fully excluded that other bacteria in the bioreactor produce these hopanoids. However, this possibility seems rather unlikely due to the low relative abundances of the other bacterial species present in the enrichment culture: only ten species had abundances >1% (and always less than 7%) in the enrichment culture. The most abundant bacteria, besides ‘*Ca. M. oxyfera*’, were *Anaerolinea thermophila* (6.5%), *Caldithrix abyssi* (5.5%), *Gemmatimonas aurantiaca* (3.2%) and *Melioribacter roseus* (1.7%). These species were previously investigated for their fatty acid inventory but not for hopanoids (Sekiguchi et al., 2003; Zhang et al., 2003; Podosokorskaya et al., 2013), although none of them has been shown to contain the gene responsible for hopanoid synthesis (e.g., Sinninghe Damsté et al., 2017) or hopanoid methylation at the C-3 position (Welander and Summons, 2012). Furthermore, we investigated another enrichment culture from a methanotrophic bioreactor fed with nitrite and methane which contained a lower proportion of ‘*Ca. M. oxyfera*’ (30%) and higher abundance of other species, mainly ANME-2D archaea (30%) (Vaksmas et al., 2017). This showed a much lower abundance of the trisnorhopanoids (Fig. 3), suggesting that ‘*Ca. M. oxyfera*’ is the likely source of these hopanoids.

Of the four trisnorhopanoids, only the 22,29,30-trisnorhopan-21-one has been reported previously, i.e. in anaerobic enrichment cultures of planctomycetes by Sinninghe Damsté et al., (2004) and in various environmental samples such as the anoxic brine-filled basins in the Mediterranean (ten Haven et al., 1987), anoxic sediments (Pancost and Sinninghe Damsté, 2003), in coals and shales from Indonesia (Hoffmann et al., 1984) and Green River oil shale kerogen (Barakat and Yen, 1990). These studies have suggested that 22,29,30-trisnorhopan-21-one is a diagenetic or microbial alteration product of BHPs such as bacteriohopanetetrol (BHT) and diploptero formed under oxic conditions (Santos et al., 1994; Conte et al., 2003). Moreover, 22,29,30-trisnorhopan-21-one is believed to be an indicator for high bio-productivity when found in the water column or sediments (Simoneit, 1977; Botz et al., 2007). However, since ‘*Ca. M. oxyfera*’ biomass was incubated under strictly anaerobic conditions, aerobic degradation processes can be excluded. We also analyzed black cell material from the bottom of the ‘*Ca. M. oxyfera*’ enrichment bioreactor, which most likely represents dead cell material from ‘*Ca. M. oxyfera*’. This material had no elevated relative abundances of the novel trisnorhopanoids compared to the more readily degradable fatty acids (Fig. 3), suggesting that the trisnorhopanoids are not formed after cell death. Together, these results suggest that the trisnorhopanoids in ‘*Ca. M. oxyfera*’ bacteria are more likely formed via a direct biosynthesis rather than by degradation processes in the bioreactor itself.

3.3. Potential biosynthesis of demethylated hopanoids in ‘*Ca. M. oxyfera*’

To further investigate the potential direct biosynthesis of the trisnorhopanoids, we explored biosynthetic genes that could be




Fig. 2. Mass spectra with fragmentation and molecular structures of trisnorhopanoids in '*Ca. M. oxyfera*' biomass. The mass spectra of 22,29,30-trisnorhopan-21-one was described previously by [Simoneit \(1977\)](#).

involved in the demethylation of the hopanoids (i.e., 22,29,30-trisnorhopan-21-one, 22,29,30-trisnorhopan-21-ol and their 3-methyl homologues) in '*Ca. M. oxyfera*'. This demethylation of the C₂₂, C₂₉ and C₃₀ carbon atoms can either take place before the cyclization of squalene to a hopane or thereafter. Squalene biosynthesis in bacteria involves a three-step reaction encoded by the HpnD, HpnC and HpnE genes and are usually found in one operon ([Pan et al., 2015](#)). We found the gene that potentially codes for HpnD (DAMO_1512) in the genome of '*Ca. M. oxyfera*', but we did not find any homologous sequence for HpnC nor HpnE using BLAST searches ([Supplementary Table S1](#)). Previous studies suggested that these steps can alternatively be performed by farnesyl diphosphate farnesyl transferase (FdfT) activities as found in for example acidobacteria, gammaproteobacteria or cyanobacteria ([Lee and Poulter, 2008](#); [Ohtake et al., 2014](#); [Sinninghe Damsté et al., 2017](#)). The gene DAMO_2922 is annotated as a farnesyl diphosphate farnesyl transferase and could be responsible for the squalene formation by this mechanism. Thus, a demethylated intermediate could potentially be formed by an alternative pathway to the three-step reaction by HpnD, HpnC and HpnE genes prior the cyclization reaction by the squalene synthetase DAMO_0045 gene and thus directly form the trisnorhopanoids.

Alternatively, and perhaps more likely, the demethylation of hopanoids could occur after the squalene cyclization in '*Ca. M. oxyfera*'. We speculate that such a process can be caused by oxidative activities similar to the removal of the C-4 methyl from lanosterol to generate 4-desmethyl lanosterol in bacteria catalyzed by SdmA and SdmB proteins ([Lee et al., 2018](#)). However, a BLAST search for SdmA and SdmB homologue proteins in the '*Ca. M. oxyfera*' genome did not indicate the presence of these two proteins, suggesting that the SdmA-SdmB proteins are only present in bacteria

(e.g., *Methylococcus capsulatus*, *Stigmatella aurantiaca* or *Methylosphaera hansonii*) that produce the C-4 demethylation in lanosterol (e.g., [Bird et al., 1971](#); [Schouten et al., 2000](#); [Bode et al., 2003](#)). In eukaryotes, the C-4 sterol demethylation is performed by other proteins i.e. ERG25, ERG26 and ERG27 ([Bard et al., 1996](#); [Gachotte et al., 1998](#); [Gachotte et al., 1999](#)). In this process, a C-4 sterol methyl oxidase (ERG25), a C-3 sterol dehydrogenase (C-4 decarboxylase) (ERG26) and a 3-keto sterol reductase (ERG27) are involved in an iterative mechanism for the demethylation of sterols. However, no homologue sequence for ERG25 and ERG27 were found in the '*Ca. M. oxyfera*' genome, although a BLAST algorithm search using ERG26 (P53199) from the yeast *Saccharomyces cerevisiae* retrieved DAMO_0933 gene, which potentially codes for a NAD-dependent epimerase/dehydratase ([Supplementary Table S1](#)). DAMO_0933 is part of the predicted operon *mox-DAMO_0930* which is composed of genes that codes for a protein of unknown function (DAMO_0930), a predicted glycosyltransferase (DAMO_0931), a potential methyltransferase (DAMO_0932), the NAD-dependent epimerase/dehydratase (DAMO_0933), a short-chain alcohol dehydrogenase (DAMO_0934) and a potential oxidoreductase (DAMO_0935). This combination of enzymes suggests that the trisnorhopanoids can be formed by a sequential oxidative process of diplopterol catalyzed by the genes in the DAMO_0930 operon ([Supplementary Fig. S2](#)). We propose that the oxidoreductase activity coded by the DAMO_0935 gene in combination with the dehydrogenase activity coded by the DAMO_0934 gene and the potential dehydratase coded by DAMO_0933 could be involved in a sequential demethylation process ([Supplementary Fig. S2](#)). Future work should involve the verification of the proposed enzymatic processes for the demethylation of hopanoids to better understand the mechanism




Fig. 3. Gas chromatograms of saponified and silylated total lipid extracts from (A) ‘*Ca. M. oxyfera*’ active biomass (bright flocculent cell material), (B) ‘*Ca. M. oxyfera*’ dead biomass (black cell material from bioreactor bottom) and (C) ‘*Ca. M. oxyfera*’ (30%) with enriched ANME-2D archaea (30%) from Italian paddy field soils. The four novel trisnorhopanoids 1 = 22,29,30-trisnorhopan-21-one (TNHone), 2 = 22,29,30-trisnorhopan-21-ol (TNHole), 1a = 3-methyl-22,29,30-trisnorhopan-21-one (3-Me TNHone) and 2a = 3-methyl-22,29,30-trisnorhopan-21-ol (3-Me TNHole) as well as the typical archaeal lipids archaeol and OH-archaeol (sn2-hydroxy-archaeol) are annotated in the chromatograms.

involved in the synthesis of trisnorhopanoids in ‘*Ca. M. oxyfera*’ and potentially other demethylated hopanoids in bacterial phyla.

3.4. Environmental occurrence of novel ‘*Ca. M. oxyfera*’ trisnorhopanoids

To investigate whether the novel hopanoids detected in ‘*Ca. M. oxyfera*’ are also present in the environment, we developed an MRM method to sensitively detect the trisnorhopanoids in environmental samples. For this, product ion scans of the molecular ions for 22,29,30-trisnorhopan-21-one, 22,29,30-trisnorhopan-21-ol and their 3-methylated counterparts were conducted using GC-MS-MS. The selection of the target ions was based on the most selective and abundant ions in these product scans (Fig. 2). The collision energies were then optimized to provide an optimal abundance for the different transitions within the MRM method (Table 1). The resulting MRM chromatogram shows substantially enhanced signal to noise ratio compared to the full scan chromatograms (Figs. 1 and 3) and extracted ion chromatograms of m/z 191 and m/z 205 (Fig. 1) and thus provides a substantially enhanced sensitivity (Fig. 4A).

Application of this MRM method to a peatland core from the Brunsummerheide, which previously revealed the presence of ‘*Ca. M. oxyfera*’ based on DNA and fatty acids (Kool et al., 2012, 2014), showed the presence of the novel hopanoids (Fig. 4B). However, the peat showed a different relative abundance of the four hopanoids than the ‘*Ca. M. oxyfera*’ biomass, i.e. with low abundance of the 3-methyl-22,29,30-trisnorhopan-21-one and a higher relative abundance of 22,29,30-trisnorhopan-21-one and 22,29,30-trisnorhopan-21-ol as well as a dominant 3-methyl-22,29,30-trisnorhopan-21-ol (Fig. 4B). Due to the lack of standards to quantify these hopanoids in the MRM transitions a hopanoid ratio was




Fig. 4. MRM (metastable reaction monitoring) chromatograms of the four distinct trisnorhopanoids (1 = 22,29,30-trisnorhopan-21-one; 2 = 22,29,30-trisnorhopan-21-ol; 1a = 3-methyl-22,29,30-trisnorhopan-21-one; and 2a = 3-methyl-22,29,30-trisnorhopan-21-ol) in (A) ‘*Ca. M. oxyfera*’ biomass and (B) Brunsummerheide (BRH) peat from 90 to 95 cm depth.

developed, i.e. the 3-Me trisnorhopanone/(3-Me trisnorhopanone + trisnorhopanone) ratio (Fig. 5D). This is based on the assumption that 22,29,30-trisnorhopan-21-one can also originate from bacteria other than ‘*Ca. M. oxyfera*’ as it has been reported in many environments (ten Haven et al., 1987; Botz et al., 2007), in contrast to its 3-methylated version. A peak of the 3-Me trisnorhopanone/(3-Me trisnorhopanone + trisnorhopanone) ratio occurs at 70–90 cm depth, and corresponds to the peak in ‘*Ca. M. oxyfera*’ cell abundance data based on qPCR 16S rRNA gene analysis as well as the specific 10MeC_{16:0} and 10MeC_{16:1} fatty acids (Fig. 5B and C; Kool et al., 2012). The 3-methyl-trisnorhopan-21-one is present in low abundance in the upper section of the core (50–70 cm depth), which might be related to the low abundance of NC10 cells in this section (Kool et al., 2012; Zhu et al., 2012). Furthermore, temporal variations of the water level in the peatland could have shifted the depth of the oxic–anoxic interface, and therefore the ‘*Ca. M. oxyfera*’ niche. This remnants of past NC10 abundance may have been picked up by our more sensitive MRM method for trisnorhopanoids compared to that of the specific fatty acids reported by Kool et al. (2012). Thus, this ratio of trisnorhopanoids may be a complimentary biomarker tool for the detection of ‘*Ca. M. oxyfera*’ in the environment. Moreover, these trisnorhopanoids may be better preserved over geological timescales than fatty acids and thus be a tool for the detection of past intra-aerobic methanotrophy in paleo-environments, in contrast to $\delta^{13}\text{C}$ analysis of hopanoids (Kool et al., 2014; Rasigraf et al., 2014).

3.5. Implications

Our results may have consequences for how we view the sources of demethylated hopanoids detected in present and past environments. Up to now, hopanoid demethylation processes were mainly attributed to diagenetic alteration such as microbial reworking of organic matter for bisnor- and trisnorhopanes or heavy biodegradation of petroleum for the presence of 25-norhopanes (Moldowan et al., 1984; Noble et al., 1985; Moldowan and McCaffrey, 1995). For example 18 α (H)-22,29,30-trisnorhopane (Ts) and the 17 α (H)-22,29,30-trisnorhopane (Tm),




Fig. 5. Depth profile of the Brunsummerheide (BRH) peatland core, showing (A) NO_3^- and CH_4 concentrations in pore water; (B) NC10 bacteria 16S rRNA gene copy number abundance; (C) specific fatty acids (FAs) of '*Ca. M. oxyfera*' (Kool et al., 2012); and (D) the novel hopanoid ratio 3-Me TNH/(3-Me TNH + TNH) of '*Ca. M. oxyfera*'. The maximum in RNA abundance of '*Ca. M. oxyfera*' is indicated by the grey area.

are thought to derive from diplopterol by a simple side-chain cleavage followed by an acid-catalyzed methyl shift (Seifert and Moldowan, 1978) and are often used to determine the source and the thermal maturity of petroleum (Seifert and Moldowan, 1978; Peters et al., 2005). It has been suggested that the 25-demethylation of 28,30-bisnorhopane to 25,28,30-trisnorhopane, common biomarker lipids in petroleum and organic-rich sediments (e.g., Noble et al., 1985; Peters et al., 2005), occurs during advanced stages of petroleum biodegradation in which the 28,30-bisnorhopane probably derives from a hopene precursor having a 17(18) double bond (Rullkötter et al., 1982; Volkman et al., 1983). A tentative identification of 28,30-bisnorhop-17(18)-ene as a precursor has been made previously in sediments from the Gulf of California (Rullkötter et al., 1982). However, it was also suggested that 28,30-bisnorhopane and 25,28,30-trisnorhopane were biosynthesized directly since they are not found as sulfur-bound hydrocarbons (Schoell et al., 1992; Schouten et al., 2001a). The ^{13}C -content of both compounds and their distribution strongly suggest that they are derived from anaerobic bacteria or those living under low oxygen conditions (Schoell et al., 1992; Schouten et al., 2001b). Our finding of demethylated hopanoids in '*Ca. M. oxyfera*' as well as those reported in the anaerobic planctomycetes (Sinninghe Damsté et al., 2004) now suggest that demethylated hopanoids may not only be formed by diagenetic processes, but also biosynthesized directly.

4. Conclusions

The intra-aerobic methanotroph '*Ca. Methyloirabilis oxyfera*' synthesizes a series of unique demethylated hopanoids identified as 22,29,30-trisnorhopan-21-one, 22,29,30-trisnorhopan-21-ol, 3-methyl-22,29,30-trisnorhopan-21-one and 3-methyl-22,29,30-

trisnorhopan-21-ol. These unique hopanoids suggest a possible demethylation process of hopanoids by '*Ca. M. oxyfera*'. This is further supported by the finding of potential candidate genes responsible for the demethylation of hopanoids in the '*Ca. M. oxyfera*' genome. For the sensitive detection of these four hopanoids, an MRM method was developed and successfully applied to an environmental setting where '*Ca. M. oxyfera*' was previously detected by specific fatty acids and genomic analysis. The novel trisnorhopanoids and the developed MRM method offer a new tool to investigate the presence of '*Ca. M. oxyfera*' and nitrite-dependent methane oxidation in modern and past environments. Moreover, the identification of demethylated hopanoids in living bacterial biomass gives a new perspective on the origin and sources of demethylated hopanoids found in the geological record.

Author contributions

NTS, DR and SS planned research. MJM, OR and SCG provided biomass and FISH analysis data. NTS performed lipid analysis and NTS and SS interpreted the data. MV, SS and NTS developed MRM method. DXSC analyzed genome data for potential biosynthesis pathways of demethylated hopanoids. NTS wrote the paper with input from all authors.

Declaration of Competing Interest

The authors declared that there is no conflict of interest.

Acknowledgements

We thank Irene Rijnstra for laboratory assistance and helpful comments in compound identification as well as Laura Villanueva

for discussions about hopanoid biosynthesis pathways. We also thank Annika Vaksmaa for providing methanotrophic ANME-2D/‘Ca. M. oxyfera’ biomass for comparison. We thank Martin Blumenberg and anonymous reviewer, as well as the Associate Editor for constructive comments. This study received funding from the Netherlands Earth System Science Center (NESSC) and Soehngen Institute for Anaerobic Microbiology (SIAM) through Gravitation grants (024.002.001 and 024.002.002) from the Dutch Ministry for Education, Culture and Science. DXSC received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement n° 694569 – MICROLIPIDS) and MSMJ by ERC AG ecomom 339880.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.orggeochem.2019.07.008>.

Associate Editor—John K. Volkman

References

- Albaiges, J., Albrecht, P., 1979. Fingerprinting marine pollutant hydrocarbons by computerized gas chromatography-mass spectrometry. *International Journal of Environmental Analytical Chemistry* 6, 171–190.
- Amann, R.L., Binder, B.J., Olson, R.J., Chisholm, S.W., Devereux, R., Stahl, D.A., 1990. Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Applied and Environmental Microbiology* 56, 1919–1925.
- Atlas, R., Bartha, R., 1973. Inhibition by fatty acids of the biodegradation of petroleum. *Antonie van Leeuwenhoek* 39, 257–271.
- Bale, N., Villanueva, L., Hopmans, E., Schouten, S., Sinninghe Damsté, J., 2013. Different seasonality of pelagic and benthic Thaumarchaeota in the North Sea. *Biogeosciences* 10, 7195–7206.
- Barakat, A., Yen, T., 1990. Distribution of pentacyclic triterpenoids in Green River oil shale kerogen. *Organic Geochemistry* 15, 299–311.
- Bard, M., Bruner, D., Pierson, C., Lees, N., Biermann, B., Frye, L., Koegel, C., Barbuch, R., 1996. Cloning and characterization of ERG25, the *Saccharomyces cerevisiae* gene encoding C-4 sterol methyl oxidase. *Proceedings of the National Academy of Sciences* 93, 186–190.
- Bird, C., Lynch, J., Pirt, F., Reid, W., Brooks, C., Middleditch, B., 1971. Steroids and squalene in *Methylococcus capsulatus* grown on methane. *Nature* 230, 473.
- Blumenberg, M., Hoppert, M., Krüger, M., Dreier, A., Thiel, V., 2012. Novel findings on hopanoid occurrences among sulfate reducing bacteria: Is there a direct link to nitrogen fixation? *Organic Geochemistry* 49, 1–5.
- Blumenberg, M., Krüger, M., Nauhaus, K., Talbot, H.M., Oppermann, B.I., Seifert, R., Pape, T., Michaelis, W., 2006. Biosynthesis of hopanoids by sulfate-reducing bacteria (genus *Desulfovibrio*). *Environmental Microbiology* 8, 1220–1227.
- Bode, H.B., Zeggel, B., Silakowski, B., Wenzel, S.C., Reichenbach, H., Müller, R., 2003. Steroid biosynthesis in prokaryotes: identification of myxobacterial steroids and cloning of the first bacterial 2,3(S)-oxidosqualene cyclase from the myxobacterium *Stigmatella aurantiaca*. *Molecular Microbiology* 47, 471–481.
- Botz, R., Schmidt, M., Wehner, H., Hufnagel, H., Stoffers, P., 2007. Organic-rich sediments in brine-filled Shaban- and Kebrut deeps, northern Red Sea. *Chemical Geology* 244, 520–553.
- Brocks, J.J., Love, G.D., Summons, R.E., Knoll, A.H., Logan, G.A., Bowden, S.A., 2005. Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea. *Nature* 437, 866.
- Brocks, J.J., Pearson, A., 2005. Building the biomarker tree of life. *Reviews in Mineralogy and Geochemistry* 59, 233–258.
- Conte, M., Dickey, T., Weber, J., Johnson, R., Knap, A., 2003. Transient physical forcing of pulsed export of bioactive material to the deep Sargasso Sea. *Deep Sea Research Part I: Oceanographic Research Papers* 50, 1157–1187.
- Conte, M.H., Weber, J., Ralph, N., 1998. Episodic particle flux in the deep Sargasso Sea: an organic geochemical assessment. *Deep Sea Research Part I: Oceanographic Research Papers* 45, 1819–1841.
- Crossman, Z., Ineson, P., Evershed, R., 2005. The use of ¹³C labelling of bacterial lipids in the characterisation of ambient methane-oxidising bacteria in soils. *Organic Geochemistry* 36, 769–778.
- Daims, H., Brühl, A., Amann, R., Schleifer, K.-H., Wagner, M., 1999. The domain-specific probe EUB338 is insufficient for the detection of all bacteria: Development and evaluation of a more comprehensive probe set. *Systematic and Applied Microbiology* 22, 434–444.
- Eickhoff, M., Birgel, D., Talbot, H.M., Peckmann, J., Kappler, A., 2013. Bacteriohopanoid inventory of *Geobacter sulfurreducens* and *Geobacter metallireducens*. *Organic Geochemistry* 58, 107–114.
- Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M.M., Schreiber, F., Dutilh, B.E., Zedelius, J., De Beer, D., 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464, 543–548.
- Ettwig, K.F., Shima, S., Van De Pas-Schoonen, K.T., Kahnt, J., Medema, M.H., Op Den Camp, H.J., Jetten, M.S., Strous, M., 2008. Denitrifying bacteria anaerobically oxidize methane in the absence of Archaea. *Environmental Microbiology* 10, 3164–3173.
- Ettwig, K.F., Van Alen, T., van de Pas-Schoonen, K.T., Jetten, M.S., Strous, M., 2009. Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum. *Applied and Environmental Microbiology* 75, 3656–3662.
- Farrimond, P., Talbot, H., Watson, D., Schulz, L., Wilhelms, A., 2004. Methylhopanoids: molecular indicators of ancient bacteria and a petroleum correlation tool. *Geochimica et Cosmochimica Acta* 68, 3873–3882.
- Gachotte, D., Barbuch, R., Gaylor, J., Nickel, E., Bard, M., 1998. Characterization of the *Saccharomyces cerevisiae* ERG26 gene encoding the C-3 sterol dehydrogenase (C-4 decarboxylase) involved in sterol biosynthesis. *Proceedings of the National Academy of Sciences* 95, 13794–13799.
- Gachotte, D., Sen, S., Eckstein, J., Barbuch, R., Krieger, M., Ray, B., Bard, M., 1999. Characterization of the *Saccharomyces cerevisiae* ERG27 gene encoding the 3-keto reductase involved in C-4 sterol demethylation. *Proceedings of the National Academy of Sciences* 96, 12655–12660.
- Hoffmann, C., Mackenzie, A., Lewis, C., Maxwell, J., Oudin, J., Durand, B., Vandenbroucke, M., 1984. A biological marker study of coals, shales and oils from the Mahakam Delta, Kalimantan, Indonesia. *Chemical Geology* 42, 1–23.
- Innes, H.E., Bishop, A.N., Head, I.M., Farrimond, P., 1997. Preservation and diagenesis of hopanoids in recent lacustrine sediments of Priest Pot, England. *Organic Geochemistry* 26, 565–576.
- Jahnke, L.L., Summons, R.E., Hope, J.M., Des Marais, D.J., 1999. Carbon isotopic fractionation in lipids from methanotrophic bacteria. II: The effects of physiology and environmental parameters on the biosynthesis and isotopic signatures of biomarkers. *Geochimica et Cosmochimica Acta* 63, 79–93.
- Kool, D.M., Talbot, H.M., Rush, D., Ettwig, K., Sinninghe Damsté, J.S., 2014. Rare bacteriohopanepolyps as markers for an autotrophic, intra-aerobic methanotroph. *Geochimica et Cosmochimica Acta* 136, 114–125.
- Kool, D.M., Zhu, B., Rijpstra, W.I.C., Jetten, M.S., Ettwig, K.F., Sinninghe Damsté, J.S., 2012. Rare branched fatty acids characterize the lipid composition of the intra-aerobic methane oxidizer “*Candidatus Methylopirabilis oxyfera*”. *Applied and Environmental Microbiology* 78, 8650–8656.
- Lee, A.K., Banta, A.B., Wei, J.H., Kiemle, D.J., Feng, J., Giner, J.-L., Welander, P.V., 2018. C-4 sterol demethylation enzymes distinguish bacterial and eukaryotic sterol synthesis. *Proceedings of the National Academy of Sciences* 115, 5884–5889.
- Lee, S., Poulter, C.D., 2008. Cloning, solubilization, and characterization of squalene synthase from *Thermosynechococcus elongatus* BP-1. *Journal of Bacteriology* 190, 3808–3816.
- Llopiz, P., Neunlist, S., Rohmer, M., 1992. Prokaryotic triterpenoids: O- α -D-glucuronopyranosyl bacteriohopanetetrol, a novel hopanoid from the bacterium *Rhodospirillum rubrum*. *Biochemical Journal* 287, 159–161.
- Moldowan, J.M., McCaffrey, M.A., 1995. A novel microbial hydrocarbon degradation pathway revealed by hopane demethylation in a petroleum reservoir. *Geochimica et Cosmochimica Acta* 59, 1891–1894.
- Moldowan, J.M., Seifert, W.K., Arnold, E., Clardy, J., 1984. Structure proof and significance of stereoisomeric 28,30-bisnorhopanes in petroleum and petroleum source rocks. *Geochimica et Cosmochimica Acta* 48, 1651–1661.
- Neunlist, S., Holst, O., Rohmer, M., 1985. Prokaryotic triterpenoids: The hopanoids of the purple non-sulphur bacterium *Rhodomicrobium vannielii*: an aminotriol and its aminoacyl derivatives, N-tryptophanyl and N-orithinyl aminotriol. *European Journal of Biochemistry* 147, 561–568.
- Niemann, H., Lösekann, T., De Beer, D., Elvert, M., Nadalig, T., Knittel, K., Amann, R., Sauter, E.J., Schlüter, M., Klages, M., 2006. Novel microbial communities of the Haakon Mosby mud volcano and their role as a methane sink. *Nature* 443, 854–858.
- Noble, R., Alexander, R., Kagi, R.I., 1985. The occurrence of bisnorhopane, trisnorhopane and 25-norhopanes as free hydrocarbons in some Australian shales. *Organic Geochemistry* 8, 171–176.
- Ohtake, K., Saito, N., Shibuya, S., Kobayashi, W., Amano, R., Hirai, T., Sasaki, S., Nakano, C., Hoshino, T., 2014. Biochemical characterization of the water-soluble squalene synthase from *Methylococcus capsulatus* and the functional analyses of its two DXXD (E) D motifs and the highly conserved aromatic amino acid residues. *The Federation of European Biochemical Societies Journal* 281, 5479–5497.
- Ouirsson, G., Albrecht, P., 1992. Hopanoids. 1. Geohopanoids: the most abundant natural products on Earth? *Accounts of Chemical Research* 25, 398–402.
- Ouirsson, G., Albrecht, P., Rohmer, M., 1979. The hopanoids: palaeochemistry and biochemistry of a group of natural products. *Pure and Applied Chemistry* 51, 709–729.
- Pan, J.-J., Solbiati, J.O., Ramamoorthy, G., Hillerich, B.S., Seidel, R.D., Cronan, J.E., Almo, S.C., Poulter, C.D., 2015. Biosynthesis of squalene from farnesyl diphosphate in bacteria: three steps catalyzed by three enzymes. *American Chemical Society Central Science* 1, 77–82.
- Pancost, R.D., Sinninghe Damsté, J.S., 2003. Carbon isotopic compositions of prokaryotic lipids as tracers of carbon cycling in diverse settings. *Chemical Geology* 195, 29–58.
- Peters, K.E., Peters, K.E., Walters, C.C., Moldowan, J., 2005. *The Biomarker Guide*. Cambridge University Press.
- Podoskorsky, O.A., Kadnikov, V.V., Gavrillov, S.N., Mardanov, A.V., Merkel, A.Y., Karnachuk, O.V., Ravin, N.V., Bonch-Osmolovskaya, E.A., Kublanov, I.V., 2013. Characterization of *Melioribacter roseus* gen. nov., sp. nov., a novel facultatively anaerobic thermophilic cellulolytic bacterium from the class *Ignavibacteria*, and

- a proposal of a novel bacterial phylum *Ignavibacteriae*. *Environmental Microbiology* 15, 1759–1771.
- Raghoebarsing, A.A., Pol, A., van de Pas-Schoonen, K.T., Smolders, A.J., Ettwig, K.F., Rijpstra, W.I.C., Schouten, S., Sinninghe Damsté, J.S., Op den Camp, H.J., Jetten, M.S., 2006. A microbial consortium couples anaerobic methane oxidation to denitrification. *Nature* 440, 918.
- Rasigraf, O., Kool, D.M., Jetten, M.S., Sinninghe Damsté, J.S., Ettwig, K.F., 2014. Autotrophic carbon dioxide fixation via the Calvin-Benson-Bassham cycle by the denitrifying methanotroph "*Candidatus Methyloirabilis oxyfera*". *Applied and Environmental Microbiology*. 04199-04113.
- Rohmer, M., Bouvier-Nave, P., Ourisson, G., 1984. Distribution of hopanoid triterpenes in prokaryotes. *Microbiology* 130, 1137–1150.
- Rohmer, M., Ourisson, G., 1976. Structure des bactériohopane-tétrols d'*Acetobacter xylinum*. *Tetrahedron Letters* 17, 3633–3636.
- Rullkötter, J., von der Dick, H., Welte, D., 1982. Organic petrography and extractable hydrocarbons of sediment from the Gulf of California. Deep-Sea Drilling Project Leg 64. Initial Reports of the Deep Sea Drilling Project 64, 837–853.
- Santos, V., Billett, D.S., Rice, A.L., Wolff, G.A., 1994. Organic matter in deep-sea sediments from the Porcupine Abyssal Plain in the north-east Atlantic Ocean. I – Lipids. *Deep Sea Research Part I: Oceanographic Research Papers* 41, 787–819.
- Schäffer, A.A., Aravind, L., Madden, T.L., Shavirin, S., Spouge, J.L., Wolf, Y.I., Koonin, E. V., Altschul, S.F., 2001. Improving the accuracy of PSI-BLAST protein database searches with composition-based statistics and other refinements. *Nucleic Acids Research* 29, 2994–3005.
- Schoell, M., McCaffrey, M., Fago, F., Moldowan, J., 1992. Carbon isotopic compositions of 28,30-bisnorhopanes and other biological markers in a Monterey crude oil. *Geochimica et Cosmochimica Acta* 56, 1391–1399.
- Schouten, S., Bowman, J.P., Rijpstra, W.I.C., Sinninghe Damsté, J.S., 2000. Sterols in a psychrophilic methanotroph, *Methylophosphaera hansonii*. *Federation of European Microbiological Societies Microbiology Letters* 186, 193–195.
- Schouten, S., De Loureiro, M.R., Sinninghe Damsté, J.S., de Leeuw, J.W., 2001a. Molecular biogeochemistry of Monterey sediments, Naples Beach, California: I. Distributions of hydrocarbons and organic sulfur compounds. In: Isaacs, C.M., Rullkötter, J. (Eds.), *The Monterey Formation: From Rocks to Molecules*. Columbia University Press, New York, pp. 150–174.
- Schouten, S., Schoell, M., Sinninghe Damsté, J.S., Summons, R.E., de Leeuw, J.W., 2001b. Molecular biogeochemistry of Monterey sediments, Naples Beach, California: II. Stable carbon isotopic compositions of free and sulphurbound carbon skeletons. In: Isaacs, C.M., Rullkötter, J. (Eds.), *The Monterey Formation: From Rocks to Molecules*. Columbia University Press, New York, pp. 175–188.
- Schouten, S., Hopmans, E.C., Baas, M., Boumann, H., Standfest, S., Könneke, M., Stahl, D.A., Sinninghe Damsté, J.S., 2008. Intact membrane lipids of "*Candidatus Nitrosopumilus maritimus*", a cultivated representative of the cosmopolitan mesophilic group I crenarchaeota. *Applied and Environmental Microbiology* 74, 2433–2440.
- Seifert, W.K., Moldowan, J.M., 1978. Applications of steranes, terpanes and monoaromatics to the maturation, migration and source of crude oils. *Geochimica et Cosmochimica Acta* 42, 77–95.
- Sekiguchi, Y., Yamada, T., Hanada, S., Ohashi, A., Harada, H., Kamagata, Y., 2003. *Anaerolinea thermophila* gen. nov., sp. nov. and *Caldilinea aerophila* gen. nov., sp. nov., novel filamentous thermophiles that represent a previously uncultured lineage of the domain Bacteria at the subphylum level. *International Journal of Systematic and Evolutionary Microbiology* 53, 1843–1851.
- Simoneit, B.R., 1977. The Black Sea, a sink for terrigenous lipids. *Deep Sea Research* 24, 813–830.
- Sinninghe Damsté, J.S., Rijpstra, W.I.C., Dedys, S.N., Foesel, B.U., Villanueva, L., 2017. Phenotype and genotyping of hopanoid production in Acidobacteria. *Frontiers in Microbiology* 8, 968.
- Sinninghe Damsté, J.S., Rijpstra, W.I.C., Schouten, S., Fuerst, J.A., Jetten, M.S., Strous, M., 2004. The occurrence of hopanoids in planctomycetes: implications for the sedimentary biomarker record. *Organic Geochemistry* 35, 561–566.
- Summons, R., Jahnke, L., 1992. Hopanes and hopanes methylated in ring-A: correlation of the hopanoids of extant methylotrophic bacteria with their fossil analogues. In: *Biomarkers in Sediments and Petroleum*. Prentice Hall, Englewood Cliffs, pp. 182–200.
- Summons, R.E., Jahnke, L.L., 1990. Identification of the methylhopanes in sediments and petroleum. *Geochimica et Cosmochimica Acta* 54, 247–251.
- Summons, R.E., Jahnke, L.L., Roksandic, Z., 1994. Carbon isotopic fractionation in lipids from methanotrophic bacteria: relevance for interpretation of the geochemical record of biomarkers. *Geochimica et Cosmochimica Acta* 58, 2853–2863.
- Talbot, H.M., Farrimond, P., 2007. Bacterial populations recorded in diverse sedimentary biohopanoid distributions. *Organic Geochemistry* 38, 1212–1225.
- Talbot, H.M., Rohmer, M., Farrimond, P., 2007. Structural characterisation of unsaturated bacterial hopanoids by atmospheric pressure chemical ionisation liquid chromatography/ion trap mass spectrometry. *Rapid Communications in Mass Spectrometry* 21, 1613–1622.
- ten Haven, H., Baas, M., de Leeuw, J., Maassen, J., Schenck, P., 1987. Organic geochemical characteristics of sediments from the anoxic brine-filled Tyro basin (eastern Mediterranean). *Organic Geochemistry* 11, 605–611.
- Thiel, V., Blumenberg, M., Pape, T., Seifert, R., Michaelis, W., 2003. Unexpected occurrence of hopanoids at gas seeps in the Black Sea. *Organic Geochemistry* 34, 81–87.
- UniProt Consortium, 2018. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Research* 47, D506–D515.
- Vaksmas, A., Guerrero-Cruz, S., van Alen, T.A., Cremers, G., Ettwig, K.F., Lüke, C., Jetten, M.S., 2017. Enrichment of anaerobic nitrate-dependent methanotrophic "*Candidatus Methanoperedens nitroreducens*" archaea from an Italian paddy field soil. *Applied Microbiology and Biotechnology* 101, 7075–7084.
- van Winden, J.F., Kip, N., Reichart, G.-J., Jetten, M.S., Op den Camp, H.J., Sinninghe Damsté, J.S., 2010. Lipids of symbiotic methane-oxidizing bacteria in peat moss studied using stable carbon isotopic labelling. *Organic Geochemistry* 41, 1040–1044.
- van Winden, J.F., Talbot, H.M., Kip, N., Reichart, G.-J., Pol, A., McNamara, N.P., Jetten, M.S., Op den Camp, H.J., Sinninghe Damsté, J.S., 2012. Bacteriohopanepolyl signatures as markers for methanotrophic bacteria in peat moss. *Geochimica et Cosmochimica Acta* 77, 52–61.
- Volkman, J.K., Alexander, R., Kagi, R.L., Rullkötter, J., 1983. GC–MS characterisation of C₂₇ and C₂₈ triterpanes in sediments and petroleum. *Geochimica et Cosmochimica Acta* 47, 1033–1040.
- Wakeham, S.G., Amann, R., Freeman, K.H., Hopmans, E.C., Jørgensen, B.B., Putnam, I. F., Schouten, S., Sinninghe Damsté, J.S., Talbot, H.M., Woeckel, D., 2007. Microbial ecology of the stratified water column of the Black Sea as revealed by a comprehensive biomarker study. *Organic Geochemistry* 38, 2070–2097.
- Welanders, P.V., Summons, R.E., 2012. Discovery, taxonomic distribution, and phenotypic characterization of a gene required for 3-methylhopanoid production. *Proceedings of the National Academy of Sciences* 109, 12905–12910.
- Wenger, L.M., Davis, C.L., Isaksen, G.H., 2002. Multiple controls on petroleum biodegradation and impact on oil quality. *Society of Petroleum Engineers Reservoir Evaluation & Engineering* 5, 375–383.
- Zhang, H., Sekiguchi, Y., Hanada, S., Hugenholtz, P., Kim, H., Kamagata, Y., Nakamura, K., 2003. *Gemmatimonas aurantiaca* gen. nov., sp. nov., a Gram-negative, aerobic, polyphosphate-accumulating micro-organism, the first cultured representative of the new bacterial phylum *Gemmatimonadetes* phyl. nov. *International Journal of Systematic and Evolutionary Microbiology* 53, 1155–1163.
- Zhu, B., van Dijk, G., Fritz, C., Smolders, A.J., Pol, A., Jetten, M.S., Ettwig, K.F., 2012. Anaerobic oxidation of methane in a minerotrophic peatland: enrichment of nitrite-dependent methane-oxidizing bacteria. *Applied and Environmental Microbiology* 78, 8657–8665.
- Zundel, M., Rohmer, M., 1985. Prokaryotic triterpenoids: 1. 3β-Methylhopanoids from *Acetobacter* species and *Methylococcus capsulatus*. *European Journal of Biochemistry* 150, 23–27.