

Open Research Online

The Open University's repository of research publications and other research outputs

Gait Rehabilitation for Neurological Conditions using Wearable Devices

Conference or Workshop Item

How to cite:

Islam, Riasat; Holland, Simon; Price, Blaine and Mulholland, Paul (2019). Gait Rehabilitation for Neurological Conditions using Wearable Devices. In: Workgroup on Interactive Systems in Health, CHI 2019: Conference on Human Factors in Computing Systems, 5 May 2019, Glasgow, United Kingdom.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: Accepted Manuscript

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Gait Rehabilitation for Neurological Conditions using Wearable Devices

Riasat Islam

The Open University
Milton Keynes, MK7 6AA, UK
riasat.islam@open.ac.uk

Simon Holland

The Open University
Milton Keynes, MK7 6AA, UK
simon.holland@open.ac.uk

Blaine Price

The Open University
Milton Keynes, MK7 6AA, UK
blaine.price@open.ac.uk

Paul Mulholland

The Open University
Milton Keynes, MK7 6AA, UK
paul.mulholland@open.ac.uk

ABSTRACT

Many people with neurological conditions such as stroke, brain injury or multiple sclerosis suffer from an impaired walking gait pattern. Gait improvement can lead to better fluidity in walking, improved health outcomes, greater independence, and enhanced quality of life. Existing lab-based studies with wearable haptic devices have shown that rhythmic haptic cueing can cause immediate improvements to gait features such as temporal symmetry, stride length and walking speed. However, such wearable haptic devices are unsuitable for self-managed use, and to move this approach from out of the lab into the wild, numerous design challenges need to be addressed. We are designing, developing, and testing a system to provide haptic rhythmic cues for sustainable self-managed long-term use outside the lab by survivors of stroke, and other neurological conditions, in their everyday lives.

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).

*WISH'19 Poster, CHI 2019 Conference on Human Factors in Computing Systems
May 4-9, 2019, Glasgow, Scotland, UK*

KEYWORDS

Wearables; gait; rehabilitation; stroke; neurology; physiotherapy; haptic; devices;

INTRODUCTION

Conditions such as Stroke, Multiple Sclerosis (MS), Traumatic Brain Injury (TBI), and Acquired Brain Injury (ABI) can all cause motor impairment, affecting basic tasks like walking and standing. To improve individual health and well-being for patients it would be helpful to be able to both assess and treat their gait outside the clinic or lab. This would have the potential to allow researchers, clinicians, caregivers, and physiotherapists to devise and deliver personalised rehabilitation plans [1]. Consequently, wearable sensing and feedback systems now offer considerable potential for the assessment and treatment of gait abnormalities outside of clinical settings [2]. Many survivors with motor impairment develop gait abnormalities, such as hemiparetic, hemiplegic, diplegic and neuropathic [3]. The exact nature and effects of such gait abnormalities are varied; however, common results include bone degeneration, joint pain, increase in higher vertical forces in the legs, loss of bone mineral density and increased risk of falls and fractures [4]. Overall these factors cause reduced mobility, reduced quality of life, lack of confidence and independence. Gait rehabilitation is a high priority for survivors of such conditions [5], though funding and availability for such therapy is often limited.

BIOLOGICAL ENTRAINMENT AND RHYTHMIC CUEING

Gait rehabilitation using rhythmic cueing can be performed via three sensory channels: auditory, visual and haptic [6]. These modalities have all shown improvements in gait features in the lab, especially auditory and haptic. The principle behind rhythmic auditory cueing and rhythmic haptic cueing is primarily biological entrainment [7], and its interaction with the Neural Central Pattern Generator [8]. In outline, the regular sequence of cues acts as a template for the brain to entrain with [9] typically leading to improvements in certain gait features, in particular increased temporal symmetry of steps, increased stride length and speed of walking [7].

Rhythmic cueing based on audio stimuli is a proven method for improving gait via entrainment [10], but in practice, its use is generally limited to the lab [7]. Haptic cueing is of particular interest because it offers an unobtrusive, invisible, sociable, safe alternative [11]. Other potential advantages of the haptic modality include the ability to deliver cues to the appropriate leg at the appropriate time, taking advantage of the proprioceptive and kinaesthetic senses [7,12]. However, existing systems that combine gait tracking and haptic cueing are unsuitable for long-term self-managed use outside the lab.

LONGITUDINAL PILOT STUDY

We carried out what is to the best of our knowledge the first two-week long pilot study on the self-managed use of wearable haptic devices for gait rehabilitation via entrainment in outdoor settings. The study involved a brain injury survivor who had reduced ability on both sides of his body and

therefore, significant gait impairments. On Day 1 of the study, we measured his baseline pre-intervention baseline measurements in the lab using x-OSC inertial measurement units (IMU) [14] on both ankles. The measurements were taken while walking 10 metres in a straight line for about 6-8 times. The IMU data was logged wirelessly to a laptop for later analysis. While he was walking in the lab, he was also video recorded to later show it to physiotherapists to evaluate his gait. The participant was asked to walk in synchrony to the haptic rhythm at a suitable outdoor setting for at least five minutes for about 10 days within two weeks.

To provide the haptic cue for previous in-lab studies, we have used a family of wearable haptic devices developed in-house, generally known as the Haptic Bracelets [15], however, for this longitudinal pilot study we have provided a commercial off-the-shelf wearable haptic device that can provide rhythmic haptic cue only. In addition, the participant was also provided with an iPhone running the Health Kit [16] and Moves [17] applications, recording their step count, walking distance, duration and route. After the two-week period, he returned to the lab and gait measurements were taken again using IMU sensors, similar to Day 1. He was also asked a number of interview questions related to usability and wearability of the devices, study compliance, challenges he faced and perceived changes in his gait. On comparing data between pre-intervention and post-intervention gait measurements, significant improvements in gait characteristics were observed. In addition, expert gait evaluation by a physiotherapist have also showed significant improvements on the following gait characteristics: stride length, temporal symmetry and walking speed.

CONCLUSION

This study has revealed numerous design challenges for users with characteristics such as reduced dexterity, hemiparesis, reduced sensitivity and limited proprioception. Future design aims include simplicity, wearability, unobtrusiveness, low maintenance, low cost, oversight, where appropriate, by clinicians, accuracy and completeness of the longitudinal data. For this pilot study, neither the participant nor the researchers had access to the data during the two-week period. As an extension to this study, we are planning to provide a customised wearable haptic device, that will provide the haptic rhythm and also be able to track activity and compliance over the length of the study. We believe that wearable devices are able to both assess and, where appropriate, treat gait outside the clinic or lab have the potential to make a substantial contribution to the gait rehabilitation of survivors of neurological conditions. The design of these systems will need to be adapted to be suitable for a combination of self-management and therapist-overseen use. At present, most stroke survivors, for example, do not receive any rehabilitative therapy at all beyond a few months after their stroke, as continued gait training is very expensive. Thus, such a system for community ambulators, where real time continuous gait data is collected over extended periods, in parallel

with the provision of haptic cueing, has potential to act as a highly desirable self-managed physiotherapy mechanism beyond care homes and indoor physiotherapy sessions. Gait rehabilitation can lead to many benefits, most importantly improved confidence and independence, and improved quality of life and well-being, additionally reduction of costs associated with care and rehabilitation.

REFERENCES

- [1] Dax Steins, Helen Dawes, Patrick Esser, and Johnny Collett. 2014. Wearable accelerometry-based technology capable of assessing functional activities in neurological populations in community settings: a systematic review. *Journal of NeuroEngineering and Rehabilitation* 11: 36.
- [2] Daniel K. Y. Chen, Markus Haller, and Thor F. Besier. 2017. Wearable lower limb haptic feedback device for retraining Foot Progression Angle and Step Width. *Gait & Posture* 55: 177–183.
- [3] Philip D Thompson. 2004. Gait disorders. *Neurology in Clinical Practice: Principles of diagnosis and management* 1: 323.
- [4] Matthew Shatzer and Howard Choi. 2017. *Physical Medicine and Rehabilitation Pocketpedia*. Springer Publishing Company.
- [5] Jeremiah J. Tate and Clare E. Milner. 2010. Real-Time Kinematic, Temporospatial, and Kinetic Biofeedback During Gait Retraining in Patients: A Systematic Review. *Physical Therapy* 90, 8: 1123–1134.
- [6] Pete B. Shull, Wisit Jirattigalachote, Michael A. Hunt, Mark R. Cutkosky, and Scott L. Delp. 2014. Quantified self and human movement: A review on the clinical impact of wearable sensing and feedback for gait analysis and intervention. *Gait & Posture* 40, 1: 11–19.
- [7] Theodoros Georgiou, Simon Holland, and Janet van der Linden. 2016. Wearable Haptic Devices for Post-stroke Gait Rehabilitation. *Proceedings of the 2016 ACM International Joint Conference on Pervasive and Ubiquitous Computing: Adjunct*, ACM, 1114–1119.
- [8] Michael H. Thaut, Gerald C. McIntosh, Spiros G. Prassas, and Ruth R. Rice. 1992. Effect of Rhythmic Auditory Cueing on Temporal Stride Parameters and EMG Patterns in Normal Gait. *Journal of Neurologic Rehabilitation* 6, 4: 185–190.
- [9] Michael H Thaut, Gerald C McIntosh, and Volker Hoemberg. 2015. Neurobiological foundations of neurologic music therapy: rhythmic entrainment and the motor system. *Frontiers in psychology* 5: 1185.
- [10] Michael H. Thaut and Mutsumi Abiru. 2010. Rhythmic Auditory Stimulation in Rehabilitation of Movement Disorders: A Review Of Current Research. *Music Perception* 27, 4: 263–269.
- [11] Riasat Islam, Simon Holland, Theodoros Georgiou, Blaine Price, and Paul Mulholland. 2018. A longitudinal rehabilitation case study for hemiparetic gait using outdoor rhythmic haptic cueing via a wearable device. In: *27th European Stroke Conference*, 11 - 13 April 2018, Athens, Greece.
- [12] Simon Holland, Rachel L. Wright, Alan Wing, Thomas Crevoisier, Oliver Hödl, and Maxime Canelli. 2014. A Gait Rehabilitation Pilot Study Using Tactile Cueing Following Hemiparetic Stroke. *Proceedings of the 8th International Conference on Pervasive Computing Technologies for Healthcare*, ICST (Institute for Computer Sciences, Social- Informatics and Telecommunications Engineering), 402–405.
- [13] Smart, Wearable & Vibrating Metronome | Soundbrenner Pulse. Retrieved March 7, 2018 from <https://www.soundbrenner.com/>.
- [14] x-OSC IMUs from X-IO Technologies, UK. Retrieved March 7, 2018 from <http://x-io.co.uk/x-osc/>.
- [15] Anders Bouwer, Simon Holland, and Mat Dalglish. 2013. The Haptic Bracelets: Learning Multi-Limb Rhythm Skills from Haptic Stimuli While Reading. In *Music and Human-Computer Interaction*. Springer, London, 101–122.
- [16] HealthKit - Apple Developer. Retrieved March 9, 2018 from <https://developer.apple.com/healthkit/>.
- [17] Moves - Activity Diary for iPhone and Android. Retrieved March 9, 2018 from <https://moves-app.com/>.