



This document is a postprint version of an article published in Science of the Total Environment © Elsevier after peer review. To access the final edited and published work see <https://doi.org/10.3168/jds.2018-14579>.

1 **This work has been published in JOURNAL OF DAIRY SCIENCE:**
2 **Mahmmod YS, Nonnemann B, Svennesen L, Pedersen K, Klaas IC. Typeability of MALDI-**
3 **TOF assay for identification of non-aureus staphylococci associated with bovine**
4 **intramammary infections and teat apex colonization. J Dairy Sci. 2018 Oct; 101 (10): 9430-**
5 **9438. doi: 10.3168/jds.2018-14579.**

8 **INTERPRETIVE SUMMARY**

9 **TYPEABILITY OF MALDI-TOF FOR NON-AUREUS STAPHYLOCOCCI**

10 **IDENTIFICATION**

11 **MAHMMOD**

12 MALDI-TOF has never been investigated for NAS from non-milk samples. We evaluated the
13 typeability of MALDI-TOF for identification of NAS associated with bovine IMI and teat apex
14 colonization. Proportion of NAS isolates correctly identified from milk (91%) was higher than
15 proportion of isolates correctly identified from teat skin (68%). In total, 93% of isolates were
16 successfully identified as NAS, while the remaining (7%) were shown to be other bacterial species
17 using MALDI-TOF. MALDI-TOF is efficient in NAS identification from milk but it may not always
18 be acceptable for routine identification of NAS from non-milk samples. Nucleic-acid based tools is
19 vital for accurate species identification of some NAS species.

21

22 **TYPEABILITY OF MALDI-TOF FOR NON-AUREUS STAPHYLOCOCCI**
23 **IDENTIFICATION**

24
25 **Typeability of MALDI-TOF Assay for Identification of Non-Aureus Staphylococci associated**
26 **with Bovine Intramammary Infections and Teat Apex Colonization**

27
28 Yasser S. Mahmmod^{*,†,‡¹}, Bettina Nonnemann[§], Line Svennesen^{*}, Karl Pedersen[§], and Ilka Christine
29 Klaas^{*,#}

30
31 ^{*} Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences,
32 University of Copenhagen, 1870 Frederiksberg C, Denmark

33 [†] Infectious Diseases, Department of Animal Medicine, Faculty of Veterinary Medicine, Zagazig
34 University, 44511-Zagazig, Sharkia Province, Egypt

35 [‡] IRTA, Centre de Recerca en Sanitat Animal (CReSA, IRTA-UAB), Campus de la Universitat
36 Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain (present)

37 [§] National Veterinary Institute, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

38 [#] DeLaval International AB, Tumba, Sweden (present)

39
40 ¹ Corresponding author

41 -----
42 Yasser S. Mahmmod

43 IRTA, Centre de Recerca en Sanitat Animal (CReSA, IRTA-UAB), Campus de la Universitat
44 Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain

45 Email: yasser@sund.ku.dk; yasser.mahmmod@irta.cat; yasserpcr@gmail.com

46 Mobile: 0034-612567320

47 **ABSTRACT**

48 Matrix-assisted laser desorption/ionization time of flight (**MALDI-TOF**), a culture-dependent assay,
49 has recently been implemented for routine identification of non-*aureus* Staphylococcus (**NAS**)
50 species from milk, but the assay has never been investigated for NAS from non-milk or
51 environmental samples. The objective of this study was to evaluate the typeability of MALDI-TOF
52 assay for the identification and differentiation of bovine-associated NAS species on aseptically
53 collected quarter-milk and teat skin in dairy herds. In eight herds, 14 to 20 cows with elevated
54 somatic cell count were randomly selected for teat skin swabs and foremilk samples from right hind
55 and left front quarters. Teat skin swabs and milk samples were collected aseptically for preliminary
56 identification using bacterial culture on chromogenic and calf blood agars. Colonies from milk and
57 teat skin samples with suspicion of having NAS were identified to species-level by MALDI-TOF
58 assay. Out of 511 isolates from 284 quarters (142 cows), 78% (n = 399) were identified by MALDI-
59 TOF. The percentage of correctly identified NAS from milk (91% 105/115) using MALDI-TOF was
60 higher than the percentage from teat skin (68%, 268/396). Out of the identified isolates, 93% (n =
61 373) were successfully identified as NAS, while the remaining 26 (7%) were shown to be other
62 bacterial species. Out of 26 non-NAS isolates, one originated from milk (*Corynebacterium stationis*),
63 while 25 originated from teat skin representing *Aerococcus viridans* (n = 7), *Bacillus pumilis* (n =
64 13), *Enterococcus saccharolyticus* (n = 1), *Clostridium septicum* (n = 1), *Corynebacterium stationis*
65 (n = 2), and *Corynebacterium casei* (n = 1). MALDI-TOF identified 85% (98/115) and 62%
66 (245/396) of the isolates in the first test. Isolates that were not identified to species-level at first test
67 were subjected to a second test, and here 47% (8/17) and 32% (48/151) from milk and teat skin,
68 respectively, were identified. After two rounds of MALDI-TOF, 22% (n = 112) of the isolates were
69 not identified representing 103 from teat skin and nine from milk. Eighteen isolates without
70 identification by MALDI-TOF were successfully identified to species-level using sequencing, where
71 16 were correctly identified as NAS, while the other two were *Corynebacterium stationis*.

72 In conclusion, MALDI-TOF is a reliable assay for identification and typeability of NAS species from
73 aseptically collected quarter milk samples. The assay may be used for identification of NAS species
74 from teat skin swabs. However, confirmation using nucleic-acid based tools is vital for accurate
75 species identification of some species and strains.

76

77 **KEYWORDS:** non-aureus staphylococci; bovine mastitis; teat skin colonization; phenotypic
78 identification

INTRODUCTION

79

80 Non-*aureus* Staphylococci (NAS) are a heterogeneous group of bacterial species (Schukken et al.,
81 2009), which regarded as a common cause of IMI in dairy herds (Zadoks and Watts, 2009).
82 Moreover, NAS are abundantly colonizing the teat skin, teat apex and teat canal and hence, many
83 studies have shown that teat colonization with NAS could have a significant role in initiation or
84 development of IMI with NAS in dairy cows (Leroy et al., 2015; De Visscher et al., 2016). Recent
85 studies documented that some species are more important than others in relation to udder health
86 (Supré et al., 2011; De Visscher et al., 2016). Furthermore, De Visscher et al. (2014) provided
87 evidence that the group of NAS species is comprised of environmental, opportunistic and host-
88 adapted species, which differ in ecology. Additionally, the authors concluded that some of the
89 extramammary niches, such as the teat apex, might act as infection sources for IMI-causing NAS.
90 Except for *Staphylococcus xylosus*, an association was observed between teat canal colonization and
91 IMI by all NAS species, in which the majority of IMI were preceded by teat canal colonization
92 (Quirk et al., 2012).

93

94 NAS are known to be a common teat apex colonizer (Braem et al., 2013; Falentin et al., 2016), and
95 are among the mastitis-causing bacteria most likely to enter the mammary gland through the teat
96 orifice (Fox and Norell, 1994) resulting in the establishment of IMI (Pyörälä and Taponen, 2009;
97 Piepers et al., 2010). Braem et al. (2012) identified staphylococci among the bacterial genera with
98 the highest percentage (31%) of colonized teat apices and they were detected with equal prevalence
99 from teat apices of non-infected, subclinically infected, and clinically infected quarters. Therefore, it
100 is crucial to identify and differentiate the NAS species colonizing teat skin or inhabiting milk causing
101 IMI to understand their epidemiology and to evaluate the clinical relevance and feasibility of species-
102 specific infection control measures (Zadoks and Watts, 2009). Furthermore, it is important for the
103 routine microbiological laboratories, because rapid and correct identification of mastitis causing

104 pathogens will influence the choice of antibiotic before the final determination of antibiotic
105 resistance of the isolate (Nagy et al., 2014). For many years, the species identification of NAS relied
106 on phenotypic characteristics, which is difficult, time consuming, laborious, and often inaccurate
107 (Watts et al., 1991; Vanderhaeghen et al., 2014). Although biochemical assays such as API systems
108 are widely used for identification of NAS, the accuracy and speed is not optimal (Taponen et al.,
109 2006; Capurro et al., 2009; Sampimon et al., 2009; Park et al., 2011).

110

111 Matrix-assisted laser desorption/ionization time of flight mass spectrometry (**MALDI-TOF**) is a
112 rapid method, which is able to identify a great variety of the isolated bacteria based on the
113 composition of conserved ribosomal proteins (Kliem and Sauer, 2012). This technique is based on
114 the acquisition of protein (ribosomal proteins) “fingerprints” directly from intact microorganisms,
115 since such profiles vary considerably among microorganisms (Singhal et al., 2015). The assay
116 provides a new diagnostic platform, which overcomes the limitations of traditional diagnostics for
117 NAS being time consuming and laborious, or the need of sugar fermentation or test kits (Watts et al.,
118 1991; Capurro et al., 2009; Vanderhaeghen et al., 2014; Taponen et al., 2016). The technique is
119 increasingly used in human medicine and, recently, it has been expanded as a routine diagnostic tool
120 in veterinary medicine (Randall et al., 2015; Pizauro et al., 2017). In the recent decade, research
121 studies showed that MALDI-TOF is a powerful and reliable diagnostic tool for identification and
122 discrimination of mastitis causing pathogens including NAS from bovine mastitis samples (Tomazi
123 et al., 2014; Gonçalves et al., 2014; Cameron et al., 2017a, b), and spiked milk samples (Barreiro et
124 al., 2017). MALDI-TOF assay was validated against other routine diagnostics for NAS such as Vitek
125 2 compact system (Elbehiry et al., 2016) and PCR (Pizauro et al., 2017) where it showed a better
126 performance in identification and discrimination of NAS species from bovine mastitis. To the best of
127 our knowledge there is no available literature describing the performance of MALDI-TOF assay for
128 identification of NAS from non-milk cow samples or environmental samples in dairy herds. The

129 objective of this study was to evaluate the typeability of MALDI-TOF assay for the identification
130 and differentiation of bovine-associated NAS species on quarter level from aseptically collected milk
131 (IMI) and teat skin (teat apex colonization) habitats in dairy herds.

132

133

MATERIAL AND METHODS

134 *Study population*

135 Eight dairy herds with Danish Holstein cows were selected for participating in a project on
136 *Streptococcus agalactiae* and *Staphylococcus aureus* IMI. To be eligible for inclusion in the present
137 study, herds had to have automatic milking systems (AMS) with ≥ 3 milking robots and bulk tank
138 milk PCR cycle threshold value ≤ 32 for *Streptococcus agalactiae*. About 30 to 40 lactating dairy
139 cows were selected randomly from each herd on the basis of the criteria of having no clinical
140 mastitis, SCC $\geq 200,000$ cells/mL at the preceding milk recording, and not subjected to antibiotic
141 therapy during the four weeks prior to sample collection. From each cow with an odd laboratory
142 running number, teat skin swab and aseptic milk samples were taken from right hind and left front
143 quarters (Mahmmod et al., 2018).

144

145 *Sampling Procedures*

146 Each herd was visited once to collect teat swab samples and aseptically collected quarter foremilk
147 samples for bacterial culture. The farmers were asked to separate the selected cows for sampling.
148 Cows were fixed in head lockers or tied. Teat swab samples were collected according to the modified
149 wet-dry method (Paduch et al., 2013). Briefly, the teat skin was sampled after cleaning with dry
150 tissue paper. The first rayon swab (DaklaPack, Glostrup, Denmark) was moistened with $\frac{1}{4}$ Ringer's
151 solution (Merck, Darmstadt, Germany) and rotated 360° around the teat about one cm from the teat
152 canal orifice. The same procedure was carried out with a dry swab (second). Immediately after

153 sampling, the tips of both swabs were transferred into one tube with 2 mL of sterile ¼ Ringer's
154 solution.

155 Quarter milk samples were collected directly after harvesting the teat swab samples according to
156 National Mastitis Council (1999) guidelines. Briefly, the teat end was thoroughly disinfected with
157 cotton swabs drenched with ethanol (70%). Individual quarter foremilk samples were then
158 aseptically collected in sterile screw-cap plastic tubes. New latex gloves were worn at each sampling
159 procedure and were changed after each cow. Tubes containing the teat swabs and aseptically
160 collected milk samples were stored at 5°C in ice boxes until the samples were delivered to the
161 microbiological laboratory within 24h. All study activities including farm visits, collection of
162 samples and laboratory examination were carried out during the period from February to May 2017.

163

164 *Laboratory Procedures*

165 *Bacteriological examination*

166 Bacterial culture for milk samples was conducted in accordance with National Mastitis Council
167 recommendations (1999). After vortexing, 0.01 mL of the milk sample from each quarter was
168 streaked using disposable, calibrated bacterial loops on a calf blood agar plate and simultaneously
169 another 0.01 mL of the same sample was streaked on chromogenic agar plates selective for
170 staphylococci species (*SASELECT*, Bio-Rad, Marnes-la-Coquette, France). Bacterial culture of teat
171 swab samples was performed according to the procedures of Paduch et al. (2013). Briefly, the teat
172 swab sample was vortexed before removing the swab tips from the tubes. The agar plates were
173 inoculated with 0.1 mL of a swab solution prepared with ¼ Ringer's solution. The inoculum was
174 spread with a sterile Drigalski spatula onto the agar surface of a calf blood agar plate and
175 *SASELECT* media simultaneously for each quarter.

176 All the inoculated plates were marked by the laboratory running number and sequence of the quarter
177 and were incubated aerobically at 37°C for 48h and examined for growth of NAS colonies after 24

178 and 48h. *Staphylococcus* species were presumptively identified on the basis of the phenotypic
179 characteristics of their colonies including shape (round, glossy) and color on the selective media. Up
180 to three different NAS species per sample, according to the colony color on the *SASELECT* media,
181 were considered for further identification at species-level. Cut-off \geq five CFU on the plate was
182 regarded as an acceptable cut-off point for definition the positivity of NAS according to Thorberg et
183 al. (2009) from milk and teat skin samples based on bacterial culture.

184

185 ***MALDI-TOF Assay and Species Identification***

186 Isolates of NAS species that were identified based on bacterial culture were subcultured individually
187 on new calf blood agar plates and incubated for 24 h at 37°C to be submitted freshly to MALDI-TOF
188 Bruker Biotyper software system (Microflex LT, Bruker Daltonics GmbH, Bremen, Germany) using
189 an Autoflex Speed for identification of NAS at species-level. The bacteria were prepared for mass
190 spectrometry analysis according to a standard extraction protocol using formic acid (Bizzini et al.,
191 2010), as recommended by the manufacturer. A sterile wooden applicator was used to pick the
192 material from a single bacterial colony followed by smearing a thin film of colony material onto a
193 MTP 384 ground steel target plate (Bruker Daltonics). Subsequently, each spot was overlaid with 1.0
194 μ L of a saturated α -cyano-4-hydroxycinnamic acid in 50% acetonitrile, and 2.5% trifluoroacetic acid
195 solution and air-dried at room temperature. Mass spectra were obtained on an Autoflex Speed
196 calibrated with the recommended Bruker *Escherichia coli* Bacterial Test Standard for Mass
197 Spectrometry.

198 Spectra of the analyzed bacteria were obtained in positive linear mode, within the mass range of 2 to
199 20 kDa. Spectra were acquired using the flexControl 3.4 program, and subsequently, analyzed using
200 MALDI Bruker Real-Time Classification 3.1 software equipped with the BDAL database (Bruker
201 Daltonics) comprised with 6903 reference spectra combined with verified local spectra from
202 National Veterinary Institute, Denmark. The software compares the 10 closest spectra in the database

203 to the entry and since expansion of the Bruker BDAL database is an ongoing process; local spectra
204 were generated from isolates obtained by the Diagnostic laboratory at DTU and identified by
205 biochemical methods and/or 16S rDNA sequencing to meet the requirements for both animal and
206 human reference spectra in the database. Improvement of the local database was conducted and
207 isolates submitted as reference strains for the extended database were implemented. These strains
208 were identified either by 16S rDNA sequencing or by biochemical methods as described by
209 Andresen et al. (2005). The 47 custom Main SPectra (MSPs) were created according to Nonnemann
210 et al. (2013) (Supplementary material 1).

211 The analysis for each isolate was run in triplicate. If the isolate was not identified to species-level in
212 the first run, it was subcultured and resubmitted for MALDI-TOF to exclude the reason of “no
213 identification” due to handling of the samples and to confirm that the non-identified isolates were
214 either due to new species or to limitation of the database. After two rounds of MALDI-TOF, the
215 unidentified isolates were considered as “no possible identification”. A cut-off score ≥ 1.7 was
216 regarded as a reliable threshold for the bacterial identification of NAS at species-level according to
217 Cameron et al. (2017a, b).

218

219 ***Identification by 16S rDNA sequencing***

220 A set of NAS isolates without possible identification by MALDI-TOF were subjected to 16S rDNA
221 sequencing for identification. Those NAS isolates were selected on the basis of having a cut-off
222 threshold < 1.7 in the triplicate runs of the two rounds of MALDI-TOF. One bacterial colony was
223 suspended in 1 mL PBS and centrifuged for 5 min at 10,000 g. The supernatant was removed and the
224 pellet resuspended in 100 μ L water. The sample was boiled for 10 min and immediately placed on
225 ice. The lysate was centrifuged for 2 min at 20,000g. The concentration of the DNA was quantified
226 on a NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) and
227 then was adjusted to 50 ng/ μ L.

228 The 16S rRNA gene was amplified by PCR in two separate products with following primers; 8f –
229 804r (5'- AGA GTT TGA TC (AC) TGG CTC AG -3' and 5'- GTA TTA CCG CGG CTG CTG G-
230 3') and (5'- CCA GCA GCC GCG GTA ATA C – 3' and 5'- GTT ACC TTG TTA CGA CTT CAC -
231 3') The amplification program ran as follows: denaturation at 94°C for 6 min, 33 cycles 94°C for 45
232 sec, 56°C for 45 sec, and 72°C for 90 sec, final extension 72°C for 10 min in a reaction volume of 50
233 µL. Amplified PCR products, 796 bp and 990 bp expected length, respectively, were verified by E-
234 Gel (Invitrogen). Fourty µL PCR product was purified with MinElute PCR purification kit (Qiagen)
235 and eluted in 20 µL EB buffer. The amplified 16s rRNA gene was sequenced on an ABI 3130
236 Genetic analyzer using BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystem)
237 according to manufacturer protocol. The resulting sequences were compared to the described
238 sequences by blasting with the GenBank database (<http://www.ncbi.nlm.nih.gov>) as previously
239 described (Strube et al. 2015).

240

241 *Statistical analysis*

242 A logistic mixed regression model with herd and cow treated as random intercept was performed to
243 investigate if teat apex colonization with a specific NAS species increased the odds of IMI with the
244 given species in the corresponding quarter. Different models were therefore performed for each of
245 the NAS IMI species recovered from the quarter milk samples. Statistical analysis was carried out in
246 R version 3.3.3 (The R Foundation for Statistical Computing). Results were considered significant if
247 $P \leq 0.05$.

248

248 **RESULTS**

249 Out of 150 eligible cows, 16 quarters were excluded for being dry or sampled twice. Out of 284
250 quarters of 142 cows, 228 quarters harbored at least one NAS species \geq five CFU/0.01 mL identified
251 by culture. Table 1 shows the number and distribution of NAS isolates from milk and teat skin
252 samples submitted to MALDI-TOF assay. Out of total 511 culture isolates, 399 (78%) were

253 identified to species-level by MALDI-TOF, while the remaining 112 (22%) were not identified after
254 two submissions to MALDI-TOF and were labeled as “no possible identification”. Out of the 399
255 isolates identified by MALDI-TOF, 373 (93%) were successfully identified as NAS, while the
256 remaining 26 isolates (7%) were diagnosed to be other bacterial species such as *Aerococcus viridans*,
257 *Bacillus pumilus*, *Enterococcus saccharolyticus*, *Clostridium septicum* and *Corynebacterium casei*,
258 Table 1. Out of the 26 isolates identified as other bacteria, one originated from milk and 25 from teat
259 skin. Out of the 373 isolates, 28% were isolated from milk (n = 105) and 72% from teat skin (n =
260 268). In the first round of MALDI-TOF, 85% (98/115) of the submitted isolates were identified from
261 milk, while the corresponding percentage was much lower for isolates from teat skin (62%,
262 245/396). In the second round, 47% (8/17) were identified from milk while it was 32% (48/151) for
263 teat skin isolates. The number of isolates without possible identification after two rounds of MALDI-
264 TOF from teat skin (n = 103) was about 10 times higher than that the number of isolates from milk
265 (n = 9).

266

267 In total, MALDI-TOF identified 16 different NAS species, 15 species from teat skin and 10 species
268 from milk, Table 2. In milk samples, *S. epidermidis* (49.5%, 52/105) was the most frequently
269 isolated NAS species, while *S. arlettae* and *S. warneri* (1/105) were the least frequently isolated
270 species. In teat skin swabs, *S. equorum* (43.3%, 116/268) was the most common species, while *S.*
271 *vitulinus* and *S. warneri* (1/268) were the least frequently isolated species. *S. equorum* was the most
272 frequently isolated NAS species from both left front (35.5%, 63/183) and right hind (30%, 57/190)
273 quarters. Teat apex colonization with *S. chromogenes*, *S. equorum*, *S. cohnii*, *S. epidermidis*, *S.*
274 *haemolyticus* and *S. xylosum* was not found to significantly increase the odds of IMI with these NAS
275 species, Table 2. Eighteen isolates without identification by MALDI-TOF were successfully
276 identified using sequencing analysis, where 89% (n = 16) were correctly identified as NAS, while the
277 other 11% (n = 2) were *Corynebacterium stationis*, Table 3.

278

279

DISCUSSION

280 To the best of our knowledge, this is the first study to evaluate the application of MALDI-TOF for
281 identification and differentiation of NAS species on quarter level from milk and teat skin in dairy
282 cows milked with AMS. Our findings showed that MALDI-TOF was able to identify 92% (106/115)
283 of the submitted culture isolates originating from milk samples indicating that it is a reliable assay
284 for the rapid and accurate identification of NAS species from milk samples. This finding is similar to
285 previous reports (Banach et al., 2016; Cameron et al., 2017a, b; Goetz et al., 2017; Savage et al.,
286 2017). For instance, Elbehiry et al. (2016) showed that MALDI-TOF correctly identified 100%
287 (44/44) of NAS species isolated from milk, while Tomazi et al. (2014) demonstrated that the
288 sensitivity of MALDI-TOF for the identification of NAS isolated from milk was 95.4% in
289 comparison to PCR-RFLP. Similarly, MALDI-TOF showed a better performance (93.2%) for
290 identification of 234 NAS representing 20 different species than Phoenix (75.6%) and Vitek-2
291 (75.2%) (Dupont et al., 2010). Loonen et al., (2012) reported a good performance of MALDI-TOF
292 for identification of NAS with a correct-identification rate of 99.3%. A lower identification
293 percentage (78%) was reported by Ayeni et al. (2017) based on 171 isolates (13 species) using
294 MALDI-TOF. However, the authors found that a drawback in identifying NAS with MALDI-TOF
295 was the inability to identify *S. gallinarum* in their study because it was absent from the MALDI-TOF
296 database at the time of study.

297

298 After two rounds of triplicate MALDI-TOF the overall number of isolates remaining without
299 possible identification was 22% (112/511) representing 103 from teat skin and nine from milk. This
300 is similar to Cameron et al. (2017b), who demonstrated that 40 isolates were not identified to
301 species-level after the first round of duplicate MALDI-TOF, but after being subjected to a second
302 round MALDI-TOF resulted in the identification of all except seven isolates. In line with that finding

303 Pizauro et al. (2017) reported that not all NAS found in buffalo milk could be identified by MALDI-
304 TOF. Additionally, Banach et al. (2016) reported that six isolates of NAS were not identified to
305 species-level by MALDI-TOF but were classified by means of routine bacteriological testing, and
306 comprised of *S. sciuri* (three strains), *S. xylosus* (two strains), and *S. equorum* (one strain). This
307 finding is in agreement with previous reports about the limitations of MALDI-TOF and could have
308 been caused by limited detection capacity of the bacteria due to a limited database of different
309 bacteria (Moussaoui et al., 2010; Barreiro et al., 2017; Cameron et al., 2017a). Furthermore, some
310 studies have shown that the conditions of bacterial growth, preparation of samples, number of
311 reference strains, and version of the software Biotyper may be a reason for the variability of NAS
312 identification (Benagli et al., 2011; Tomazi et al., 2014).

313

314 Our findings indicate that MALDI-TOF provides a valuable tool for identification and typeability of
315 NAS species from milk samples, while for isolates from non-milk samples, the assay showed a
316 limited performance. Furthermore, in both the first and second submissions, the performance of
317 MALDI-TOF for identification of isolates originating from milk (85% and 47%) was higher than the
318 assay performance in identification of isolates from teat skin (62% and 32%, respectively). One
319 possible explanation is that these unidentified bacteria from teat skin come from a natural teat skin
320 microbiota (commensal bacteria), which have not been previously included in the BDAL database.
321 For that reason, they may be out of interest of the microbiological diagnosticians and, therefore were
322 not considered in the database of MALDI-TOF. Moreover, our findings showed that the vast
323 majority of the unidentifiable NAS isolates were originated from teat skin. A reasonable explanation
324 could be (a) those isolates are new NAS species. Supré et al. (2010) have classified 10 non-motile,
325 Gram-stain-positive, coagulase-negative staphylococci isolated from bovine milk and teat apices as
326 *S. devriesei* sp. nov using 16S rRNA gene and four housekeeping genes (*rpoB*, *hsp60*, *tuf* and *dnaJ*)
327 in combination with tRNA-intergenic spacer length analysis. Another reason (b) these isolates could

328 be known NAS species but are not included in our BDAL database. It may be worth to mention that
329 the database of MALDI-TOF we used in this study, is not based only on the commercial version of
330 BDAL database. Our database was updated regularly because the assay is being used on research and
331 routine diagnostic service purposes for identification and differentiation of bacterial pathogens in
332 different samples types from both human and animals. We think that adding additional microbial
333 spectra, MSPs for some staphylococci species will improve the identification capacity of MALDI-
334 TOF assay both from milk and teat skin. In line with that statement, Cameron et al. (2017b) showed
335 that using a custom reference spectra expanded database, which included an additional 13 in-house
336 created reference spectra, isolates were identified by MALDI-TOF mass spectrometry with 99.2%
337 (854/861) typeability and 99.4% (849/854) accuracy.

338

339 Moreover, the BDAL database is mainly bacterial isolates originated from humans and then it
340 gradually extended to cover veterinary isolates (Tomazi et al., 2014). In support to that, Randall et al.
341 (2015) stated that the majority of reference spectra included in commercial MALDI-TOF databases
342 are derived from human isolates and, consequently, small differences between human and animal
343 isolates of the same bacterial species may influence the results obtained when testing some isolates
344 from animals. Remarkably, 89% of 18 isolates without possible identification by MALDI-TOF were
345 correctly identified as NAS using PCR indicating that confirmation using nucleic-acid tools is
346 essential for those suspected NAS isolates of environmental origin. By examination of the genetic
347 diversity among NAS species using molecular typing methods, Piessens et al. (2012) identified five
348 genotypes among *S. chromogenes* in six dairy farms. Cameron et al. (2017a) added that these
349 genotypes would have different spectra and would require their own entry within the database. We
350 updated the MALDI-TOF database on the basis of the findings in this study. Adding the spectra for
351 eight NAS species based on both prior knowledge of the species and results of the assessment of the
352 database, the authors were able to eliminate the unidentified risk from 8% unidentified to 0%

353 unidentified. Therefore, in future studies, we will expect better agreement in the performance of
354 MALDI-TOF between teat skin and milk samples.

355 Consequently, MALDI-TOF is a valuable routine diagnostic tool for identification and
356 differentiation of mastitis pathogens but it has some limitations with regard to different detection
357 capacity of the pathogens depending on the bacterial species, and limited database of pathogens
358 (Moussaoui et al., 2010; Barreiro et al., 2017; Cameron et al., 2017a). In line with that statement, our
359 findings showed that the isolates without possible identification by MALDI-TOF were successfully
360 identified by PCR, and sequencing analysis confirming the limited database of MALDI-TOF.
361 However, the database can continuously be expanded to accommodate new species and spectra.
362 Furthermore, the MALDI-TOF assay depends on initial microbiological culture, which is time-
363 consuming and laborious (i.e., plate preparation, sterilization of materials, time of incubation,
364 biochemistry tests) (Barreiro et al., 2010; Cameron et al., 2017a). Additionally, the reading is not
365 constant and differs by changing the cut-off (identification score) for species-level identification
366 (Barreiro et al., 2017; Cameron et al., 2017a) and extra rounds of identification may be necessary for
367 achieving accurate identification as shown in this study. Moreover, some previous studies have
368 shown that the conditions of bacterial growth, preparation of samples, number of reference strains or
369 spectra, database and version of the Biotyper may be a reason for the variability of NAS
370 identification (Benagli et al., 2011; Tomazi et al., 2014). Out of the initially unidentifiable isolates on
371 first round of MALDI-TOF, our findings showed that the assay identified 8/17 (47%; milk) and
372 48/151 (32%; teat skin) on the second round of identification. A plausible argument could be related
373 to the variation in the conditions of bacterial growth, extraction procedure and handling and
374 preparation of samples between the first and second rounds of MALDI-TOF. MALDI-TOF may be
375 sensitive to the sample preparations and handling due to individual variations. This may also explain
376 some of the deviations between different rounds, supporting previous reports (Benagli et al., 2011;
377 Tomazi et al., 2014).

378

379 Remarkably, we noticed that NAS isolates originating from teat skin may require extra rounds for
380 identification by MALDI-TOF in comparison to NAS isolates originating from milk. That
381 phenomenon is supported by the marked high number of isolates from teat skin remaining
382 “unidentifiable” after the first and second rounds of identification. A possible explanation could be
383 that bacterial isolates from the environment may have developed an extra layer or capsule of protein
384 material as a mean of protection against unfavorable environmental conditions.

385

386 MALDI-TOF as culture-dependent assay relies on direct analysis of proteins from bacterial cell
387 extracts of the reference strains included in the database (Liang et al., 1996). So, when a given NAS
388 strain is tested, the species of the reference strain with the closest match is retained for identification
389 of the tested strain (Ayeni et al., 2017). Therefore, an up-to-date database is essential for bacterial
390 identification and more spectra of appropriate reference strains of NAS should be added to the
391 database for accurate identification (van Veen et al., 2010; Cameron et al., 2017a). The same
392 conclusion was supported by Murugaiyan et al. (2014) and Randall et al. (2015), who reported that
393 the ongoing supplementation of the Bruker database library should further improve the utility of
394 MALDI-TOF in routine veterinary diagnostic laboratories. Murugaiyan et al. (2014) demonstrated
395 that 17 isolates initially diagnosed as *S. intermedius* with the current content of the BDAL database
396 were identified as *S. pseudintermedius* by applying the in-house reference spectra extended version
397 indicating that updating the reference spectra library allowed species identification of NAS.

398

399

CONCLUSION

400 MALDI-TOF provides a valuable and efficient platform for identification and typeability of NAS
401 species from aseptically collected quarter milk samples. MALDI-TOF may be used for identification

402 of NAS species from teat skin swabs. However, confirmation using nucleic-acid based tools is
403 essential for accurate identification of some NAS species and strains.

404

405

ACKNOWLEDGMENTS

406 The study was part of the research project ‘STOPMAST’ financed by the Danish Milk Levy
407 Foundation. We gratefully acknowledge the efforts of Nanna Skjølstrup and Louise Mathiasen for
408 their help with sampling and examination of samples by bacterial culture. Thanks to the Danish
409 farmers for their help and making their cows available for our study. Thanks to the milk quality
410 technicians at SEGES helping taking milk samples and to the laboratory technicians for their
411 technical and logistic support. Prof. Carsten Enevoldsen is acknowledged for his contribution in
412 proofreading of this work. Yasser Mahmmod was supported by the Islamic Development Bank Merit
413 Scholarship Program (IDB-MSP), Jeddah, Saudi Arabia.

414

415

REFERENCES

- 416 Andresen, L.O., P. Ahrens, L. Daugaard, V. Bille-Hansen. 2005. Exudative epidermitis in pigs
417 caused by toxigenic *Staphylococcus chromogenes*. *Vet. Microbiol.* 105: 291–300.
- 418 Ayeni, F.A., C. Andersen, and N. Nørskov-Lauritsen. 2017. Comparison of growth on mannitol salt
419 agar, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, VITEK® 2
420 with partial sequencing of 16S rRNA gene for identification of coagulase-negative
421 staphylococci. *Microb. Pathog.* 105: 255-259.
- 422 Banach, T., M. Bochniarz, P. Łyp, Ł. Adaszek, W. Wawron, B. Furmaga, M. Skrzypczak, J. Ziętek,
423 and S. Winiarczyk. 2016. Application of matrix-assisted laser desorption ionization time-of-
424 flight mass spectrometry for identification of coagulase-negative staphylococci isolated from
425 milk of cows with subclinical mastitis. *Pol. J. Vet. Sci.* 19:627-632.

426 Barreiro, J.R., C.R. Ferreira, G.B. Sanvido, M., Kostrzewa, T. Maier, B. Wegemann, V. Böttcher,
427 M.N. Eberlin, and M.V. dos Santos. 2010. Short communication: Identification of subclinical
428 cow mastitis pathogens in milk by matrix-assisted laser desorption/ionization time-of-flight
429 mass spectrometry. *J. Dairy Sci.* 93:5661-7.

430 Barreiro, J.R., J.L. Gonçalves, P.A. Braga, A.G. Dibbern, M.N. Eberlin, and M.V. dos Santos. 2017.
431 Non-culture-based identification of mastitis-causing bacteria by MALDI-TOF mass
432 spectrometry. *J. Dairy Sci.* 100:2928-2934.

433 Benagli, C., V. Rossi, M. Dolina, M. Tonolla, and O. Petrini. 2011. Matrix-assisted laser desorption
434 ionization-time of flight mass spectrometry for the identification of clinically relevant bacteria.
435 *PLoS ONE* 6:e16424.

436 Bizzini, A., C. Durussel, J. Bille, G. Greub, and G. Prod'hom. 2010. Performance of matrix-assisted
437 laser desorption/ ionization-time of flight mass spectrometry for identification of bacterial
438 strains routinely isolated in a clinical microbiology laboratory. *J. Clin. Microbiol.* 48:1549–
439 1554.

440 Braem, G., S. De Vliegher, B. Verbist, M. Heyndrickx, F. Leroy, and L. De Vuyst. 2012. Culture-
441 independent exploration of the teat apex microbiota of dairy cows reveals a wide bacterial
442 species diversity. *Vet. Microbiol.* 157:383-90.

443 Braem, G., S. De Vliegher, B. Verbist, V. Piessens, E. Van Coillie, L. De Vuyst, and F. Leroy. 2013.
444 Unraveling the microbiota of teat apices of clinically healthy lactating dairy cows, with special
445 emphasis on coagulase-negative staphylococci. *J. Dairy Sci.* 96:1499–1510.

446 Cameron, M., H.W. Barkema, J. De Buck, S. De Vliegher, M. Chaffer, J. Lewis, and G.P. Keefe.
447 2017a. Identification of bovine-associated coagulase-negative staphylococci by matrix-assisted
448 laser desorption/ionization time-of-flight mass spectrometry using a direct transfer protocol. *J.*
449 *Dairy Sci.* 100:2137-2147.

450 Cameron, M.J. Perry, J.R. Middleton, M. Chaffer, J. Lewis, and G.P. Keefe. 2017b. Evaluation of
451 MALDI-TOF mass spectrometry and a custom reference spectra expanded database for the
452 identification of bovine-associated coagulase-negative staphylococci. *J. Dairy Sci.* 101:1–6.

453 Capurro, A., K. Artursson, K.P. Waller, B. Bengtsson, H. Ericsson-Unnerstad, and A. Aspan. 2009.
454 Comparison of a commercialized phenotyping system, antimicrobial susceptibility testing, and
455 *tuf* gene sequence-based genotyping for species-level identification of coagulase-negative
456 staphylococci isolated from cases of bovine mastitis. *Vet. Microbiol.* 134:327–333.

457 De Visscher, A., K. Supré, F. Haesebrouck, R.N. Zadoks, V. Piessens, E. Van Coillie, S. Piepers, and
458 S. De Vliegher. 2014. Further evidence for the existence of environmental and host-associated
459 species of coagulase-negative staphylococci in dairy cattle. *Vet. Microbiol.* 172:466-74.

460 De Visscher, A., S. Piepers, F. Haesebrouck, and S. De Vliegher. 2016. Intramammary infection with
461 coagulase-negative staphylococci at parturition: Species-specific prevalence, risk factors, and
462 effect on udder health. *J. Dairy Sci.* 99:6457-69.

463 Dupont, C.1., V. Sivadon-Tardy, E. Bille, B. Dauphin, J.L. Beretti, A.S. Alvarez, N. Degand, A.
464 Ferroni, M. Rottman, J.L. Herrmann, X. Nassif, E. Ronco, and E. Carbonnelle. 2010.
465 Identification of clinical coagulase negative staphylococci isolated in microbiology laboratories
466 by MALDI-TOF mass spectrometry and two automates, *Clin. Microbiol. Infect.* 16:998e1004.

467 Elbehiry, A., M. Al-Dubaib, E. Marzouk, S. Osman, and H. Edrees. 2016. Performance of MALDI
468 biotyper compared with Vitek™ 2 compact system for fast identification and discrimination of
469 *Staphylococcus* species isolated from bovine mastitis. *Microbiology Open*, 5:1061–1070.

470 Falentin, H., L. Rault, A. Nicolas, D.S. Bouchard, J. Lassalas, P. Lamberton, J.M. Aubry, P.G.
471 Marnet, Y. Le Loir, and S. Even. 2016. Bovine teat microbiome analysis revealed reduced alpha
472 diversity and significant changes in taxonomic profiles in quarters with a history of mastitis.
473 *Front. Microbiol.* 7:480.

474 Fox, L.K., and R.J. Norell. 1994. *Staphylococcus aureus* colonization of teat skin as affected by
475 postmilking teat treatment when exposed to cold and windy conditions. *J. Dairy Sci.* 77:2281–
476 2288.

477 Goetz, C., Y.D.N. Tremblay, D. Lamarche, A. Blondeau, A.M. Gaudreau, J. Labrie, F. Malouin, and
478 M. Jacques. 2017. Coagulase-negative staphylococci species affect biofilm formation of other
479 coagulase-negative and coagulase-positive staphylococci. *J. Dairy Sci.* 100:6454-6464.

480 Gonçalves, J.L., T. Tomazi, J.R. Barreiro, P.A. Braga, C.R. Ferreira, J.P. Araújo Junior, M.N.
481 Eberlin, and MV. dos Santos. 2014. Identification of *Corynebacterium* spp. isolated from bovine
482 intramammary infections by matrix-assisted laser desorption ionization-time of flight mass
483 spectrometry. *Vet. Microbiol.* 173:147-51.

484 Kliem, M., and S. Sauer. 2012. The essence on mass spectrometry based microbial diagnostics. *Curr.*
485 *Opin. Microbiol.* 15:397–402

486 Leroy, F., E. Van Coillie, G. Braem, V. Piessens, B. Verbist, L. De Vuyst, and S. De Vliegher. 2015.
487 Short communication: Subtyping of *Staphylococcus haemolyticus* isolates from milk and
488 corresponding teat apices to verify the potential teat-skin origin of intramammary infections in
489 dairy cows. *J. Dairy Sci.* 98:7893-8.

490 Liang, X., K. Zheng, M.G. Qian, and D.M. Lubman. 1996. Determination of bacterial protein
491 profiles by matrix-assisted laser desorption/ionization mass spectrometry with high-performance
492 liquid chromatography, *Rapid Commun. Mass Spectrom.* 10:1219e1226.

493 Loonen, A.J.M., A.R. Jansz, J.N.B. Bergland, M. Valkenburg, P.F.G. Wolffs, and A.J.C. van den
494 Brule. 2012. Comparative study using phenotypic, genotypic, and proteomics methods for
495 identification of coagulase-negative staphylococci, *J. Clin. Microbiol.* 50:1437e1439.

496 Mahmmud, Y., I. Klaas, L. Svennesen, K. Pedersen, and H. Ingmer 2018. Coagulase negative
497 staphylococci distribution in dairy herds with automatic milking system and their crosstalk with

498 Staphylococcus aureus from intramammary infections and teat apex. National Mastitis Council
499 57th Annual Meeting, 31st January to 2nd February 2018, Tucson, Arizona, US.

500 Moussaoui, W., B. Jaulhac, A.M. Hoffmann, B. Ludes, M. Kostrzewa, P. Riegel, and G. Prévost
501 2010. Matrix-assisted laser desorption ionization time-of-flight mass spectrometry identifies 90
502 % of bacteria directly from blood culture vials. Clin. Microbiol. Infect. 16:1631–1638.

503 Murugaiyan, J., B. Walther, I. Stamm, Y. Abou-Elnaga, S. Brueggemann-Schwarze, S. Vincze, L.H.
504 Wieler, A. Lubke-Becker, T. Semmler, and U. Roesler. 2014. Species differentiation within the
505 Staphylococcus intermedius group using a refined MALDI-TOF MS database. Clin. Microbiol.
506 Infect. 20:1007–1015.

507 Nagy, E., M. Abrók, N. Bartha, L. Bereczki, E. Juhász, G. Kardos, K. Kristóf, C. Miszti, and E.
508 Urbán. 2014. Special application of matrix-assisted laser desorption ionization time-of-flight
509 mass spectrometry in clinical microbiological diagnostics. Orv. Hetil, 155:1495-503.

510 National Mastitis Council, 1999. Laboratory Handbook on Bovine Mastitis. National Mastitis
511 Council, Madison, WI.

512 Nonnemann, B., M. Tvede, T. Bjarnsholt. 2013. Identification of pathogenic microorganisms directly
513 from positive blood vials by matrix-assisted laser desorption/ionization time of flight mass
514 spectrometry. APMIS 123:871-877.

515 Paduch, J.H., E. Mohr and V. Kromker. 2013. The association between bedding material and the
516 bacterial counts of *Staphylococcus aureus*, *Streptococcus uberis* and coliform bacteria on teat
517 skin and in teat canals in lactating dairy cattle. J. Dairy Res. 80:159-164.

518 Park, J.Y., L.K. Fox, K.S. Seo, M.A. McGuire, Y.H. Park, F.R. Rurangirwa, W.M. Sicho, and G.A.
519 Bohach. 2011. Comparison of phenotypic and genotypic methods for the species identification
520 of coagulase-negative staphylococcal isolates from bovine intramammary infections. Vet.
521 Microbiol. 147:142-8.

522 Piepers, S., G. Opsomer, H.W. Barkema, A. de Kruif, and S. De Vliegher. 2010. Heifers infected
523 with coagulase-negative staphylococci in early lactation have less cases of clinical mastitis and a
524 higher milk production in their first lactation than non-infected heifers. *J. Dairy Sci.* 93:2014–
525 2024.

526 Piessens, V., S. De Vliegher, B. Verbist, G. Braem, A. Van Nuffel, L. De Vuyst, M. Heyndrickx, and
527 E. Van Coillie. 2012. Intra-species diversity and epidemiology varies among coagulase-negative
528 *Staphylococcus* species causing bovine intramammary infections. *Vet. Microbiol.* 155:62-71.

529 Pizauro, L.J.L., C.C. de Almeida, G.A. Soltes, D. Slavic, O.D. Rossi-Junior, F.A. de Ávila, L.F.
530 Zafalon, and J.I. MacInnes. 2017. Species level identification of coagulase negative
531 *Staphylococcus* spp. from buffalo using matrix-assisted laser desorption ionization-time of flight
532 mass spectrometry and cydB real-time quantitative PCR. *Vet. Microbiol.* 204:8-14.

533 Pyörälä, S., and S. Taponen. 2009. Coagulase-negative staphylococci - emerging mastitis pathogens.
534 *Vet. Microbiol.* 134:3-8.

535 Quirk, T., L.K. Fox, D.D. Hancock, J. Capper, J. Wenz, and J. Park. 2012. Intramammary infections
536 and teat canal colonization with coagulase-negative staphylococci after postmilking teat
537 disinfection: species-specific responses. *J. Dairy Sci.* 95:1906-12.

538 Randall, L.P., F. Lemma, M. Koylass, J. Rogers, R.D. Ayling, D. Worth, M. Klita, A. Steventon, K.
539 Line, P. Wragg, J. Muchowski, M. Kostrzewa. and A.M. Whatmore. 2015. Evaluation of
540 MALDI-ToF as a method for the identification of bacteria in the veterinary diagnostic
541 laboratory. *Res. Vet. Sci.* 101:42-49.

542 Sampimon, O.C., R.N. Zadoks, S. De Vliegher, K. Supré, F. Haesebrouck, H.W. Barkema, J. Sol,
543 and T.J.G.M. Lam. 2009b. Performance of API Staph ID 32 and Staph-Zym for identification of
544 coagulase-negative staphylococci isolated from bovine milk samples. *Vet. Microbio.* 136:300–
545 305.

546 Savage, E., S. Chothe, V. Lintner, T. Pierre, T. Matthews, S. Kariyawasam, D. Miller, D. Tewari,
547 and B. Jayarao. 2017. Evaluation of Three Bacterial Identification Systems for Species
548 Identification of Bacteria Isolated from Bovine Mastitis and Bulk Tank Milk Samples.
549 Foodborne Pathog. Dis. 14:177-187.

550 Schukken, Y.H., R.N. González, L.L. Tikofsky, H.F. Schulte, C.G. Santisteban, F.L. Welcome, G.J.
551 Bennett, M.J. Zurakowski, and R.N. Zadoks. 2009. CNS mastitis: Nothing to worry about? Vet.
552 Microbiol. 134:9-14.

553 Singhal, N., M. Kumar, P.K. Kanaujia, and J.S. Viridi. 2015. MALDI-TOF mass spectrometry: an
554 emerging technology for microbial identification and diagnosis. Front. Microbiol. 6:791.

555 Strube, M.L., H.C. Ravn, H.C. Ingerslev, A.S. Meyer, and M. Boye. 2015. *In situ* prebiotics for
556 weaning piglets: *in vitro* production and fermentation of potato galacto-rhamnogalacturonan.
557 Appl. Environ. Microbiol. 81:1668–1678.

558 Supré, K., F. Haesebrouck, R.N. Zadoks, M. Vaneechoutte, S. Piepers, and S. De Vliegher. 2011.
559 Some coagulase-negative *Staphylococcus* species affect udder health more than others. J. Dairy
560 Sci. 94:2329–2340.

561 Supré, K., S. De Vliegher, I. Cleenwerck, K. Engelbeen, S. Van Trappen, S. Piepers, O.C.
562 Sampimon, R.N. Zadoks, P. De Vos, and F. Haesebrouck. 2010. *Staphylococcus devriesei* sp.
563 nov., isolated from teat apices and milk of dairy cows. Int. J. Syst. Evol. Microbiol. 60:2739-44.

564 Taponen, S., H. Simojoki, M. Haveri, H.D. Larsen, and S. Pyörälä. 2006. Clinical characteristics and
565 persistence of bovine mastitis caused by different species of coagulase-negative staphylococci
566 identified with API or AFLP. Vet. Microbiol. 115:199-207.

567 Taponen, S., S. Nykäsenoja, T. Pohjanvirta, A. Pitkälä, and S. Pyörälä. 2016. Species distribution
568 and *in vitro* antimicrobial susceptibility of coagulase-negative staphylococci isolated from
569 bovine mastitic milk. Acta Vet. Scand. 58:12.

570 Thorberg, B.M., M.L. Danielsson-Tham, U. Emanuelson, K. Persson Waller. 2009. Bovine
571 subclinical mastitis caused by different types of coagulase-negative staphylococci. *J. Dairy Sci.*
572 92:4962-4970.

573 Tomazi, T., J.L. Gonçalves, J.R. Barreiro, P.A. de Campos Braga, L.F. Prada e Silva, M.N. Eberlin,
574 and M.V. dos Santos. 2014. Identification of coagulase-negative staphylococci from bovine
575 intramammary infection by Matrix-Assisted Laser Desorption Ionization-Time of Flight mass
576 spectrometry. *J. Clin. Microbiol.* 52:1658–1663.

577 van Veen, S.Q., E.C. Claas, and E.J. Kuijper. 2010. High-throughput identification of bacteria and
578 yeast by matrix-assisted laser desorption ionization time of flight mass spectrometry in
579 conventional medical microbiology laboratories, *J. Clin. Microbiol.* 48:900e907.

580 Vanderhaeghen, W., S. Piepers, F. Leroy, E. Van Coillie, F. Haesebrouck, and S. De Vlieghe. 2014.
581 Invited review: Effect, persistence, and virulence of coagulase-negative *Staphylococcus* species
582 associated with ruminant udder health. *J. Dairy Sci.* 97:5275–5293.

583 Watts, J.L., C.H. Ray, and P.J. Washburn. 1991. A convenient method for differentiation of
584 coagulase-negative staphylococci isolated from bovine mammary glands. *J. Dairy Sci.* 74:426–
585 428.

586 Zadoks, R.N., and J.L. Watts. 2009. Species identification of coagulase-negative staphylococci:
587 Genotyping is superior to phenotyping. *Vet. Microbiol.* 134:20–28.

588

589 **Table 1.** Number and distribution of NAS isolated from milk and teat skin of dairy cows in eight
 590 dairy herds submitted to MALDI-TOF assay for species identification of NAS.

Sample type	MALDI-TOF identification				No possible identification n	Bacterial species (n) wrongly identified as NAS on culture
	First round		Second round			
	Submitted	Identified	Submitted	Identified		
Milk	115	98	17	8	9	<i>Corynebacterium stationis</i> (1)
Teat skin	396	245	151	48	103	<i>Aerococcus viridans</i> (7); <i>Corynebacterium casei</i> (1); <i>Bacillus pumilus</i> (13); <i>Clostridium septicum</i> (1); <i>Enterococcus saccharolyticus</i> (1); <i>Corynebacterium stationis</i> (2)
Overall	511	343	168	56	112	26

591

592 **Table 2.** Description of 16 species of NAS analyzed by MALDI-TOF and their association from
 593 aseptic quarter milk and teat skin habitats collected from 142 cows (284 quarters) in eight dairy herds
 594 with automatic milking systems.

NAS species ^a	Number (%)	Sample type (%)		OR ^c (95% CI)	P-value [*]
		Milk	Teat skin		
<i>S. b. arlettae</i>	12 (3.2)	1 (0.9)	11 (4.1)	---	---
<i>S. capitis</i>	3 (0.8)	---	3 (1.1)	---	---
<i>S. cohnii</i>	43 (11.5)	5 (4.8)	38 (14.2)	2.23 (0.11 - 15.6)	0.48
<i>S. epidermidis</i>	60 (16.1)	52 (49.5)	8 (3.0)	0.88 (0.05 - 5.07)	0.90
<i>S. haemolyticus</i>	58 (15.6)	16 (15.2)	42 (15.7)	1.13 (0.17 - 4.24)	0.55
<i>S. hominis</i>	17 (4.6)	3 (2.9)	14 (5.2)	---	---
<i>S. piscifermentans</i>	2 (0.5)	---	2 (0.8)	---	---
<i>S. saprophyticus</i>	5 (1.3)	---	5 (1.9)	---	---
<i>S. sciuri</i>	9 (2.4)	---	9 (3.4)	---	---
<i>S. simulans</i>	2 (0.5)	2 (1.9)	---	---	---
<i>S. succinus</i>	2 (0.5)	---	2 (0.8)	---	---
<i>S. vitulinus</i>	1 (0.3)	---	1 (0.4)	---	---
<i>S. warneri</i>	2 (0.5)	1 (0.9)	1 (0.4)	---	---
<i>S. chromogenes</i>	16 (4.3)	11 (10.5)	5 (1.9)	NA ^d	NA
<i>S. equorum</i>	122 (32.7)	6 (5.7)	116 (43.3)	NA	NA
<i>S. xylosus</i>	19 (5.1)	8 (7.6)	11 (4.1)	NA	NA
<i>Other NAS (S. chromogenes, S. equorum, S. xylosus)</i> ^e	157 (42.1)	25 (23.8)	132 (49.3)	0.40 (0.09 - 1.69)	0.21
Total	373 (100)	105 (100)	268 (100)	---	---

595 ^a *Staphylococcus arlettae*, *S. warneri*, and *S. hominis* were not considered in the statistical analysis
 596 because of the few number of observations (< 5), while *S. capitis*, *S. piscifermentans*, *S.*
 597 *saprophyticus*, *S. sciuri*, *S. simulans*, *S. succinus*, and *S. vitulinus* were not isolated from milk and/or
 598 teat skin.

599 ^b S= *Staphylococcus*, ^c OR= Odds ratio; ^d NA= not applicable; ^{*} significance at P < 0.05

600 ^e *Other NAS (S. chromogenes, S. equorum, S. xylosus)*: they are grouped together for a valid
 601 statistical analysis because the mixed model does not work for each species separately.

602 **Table 3.** Collection of bacterial isolates (n=18) from bovine teat skin suspected to be NAS on culture
603 and with no possible identification by MALDI-TOF after two rounds of submissions and were
604 correctly identified by 16S sequencing.

NAS species identified by sequencing	Number (%)
<i>Jeotgalicoccus psychrophilus</i>	1 (5.6)
<i>Corynebacterium stationis</i>	2 (11.1)
<i>Staphylococcus cohnii</i>	7 (38.9)
<i>Staphylococcus devriesei</i>	1 (5.6)
<i>Staphylococcus equorum</i>	3 (16.7)
<i>Staphylococcus arlettae</i>	2 (11.1)
<i>Staphylococcus vitulinus</i>	2 (11.1)
Total	18 (100)

605