



**This document is a postprint version of an article published in Food Research International © Elsevier after peer review. To access the final edited and published work see <https://doi.org/10.1016/j.foodres.2017.12.052>**

Manuscript Number:

Title: MLVA subtyping of *Listeria monocytogenes* isolates from meat products and meat processing plants

Article Type: Research Paper

Keywords: *Listeria monocytogenes*, typing, MLVA, MLST, meat industries, meat products

Corresponding Author: Dr. Teresa Aymerich, Dr

Corresponding Author's Institution: IRTA

First Author: belen Martin, Dr

Order of Authors: belen Martin, Dr; Sara Bover-Cid; Teresa Aymerich, Dr

Abstract: *Listeria monocytogenes* is widely distributed in meat products and the meat-processing industry thus posing a risk to consumers. The aim of this study was to evaluate the suitability of the multilocus variable-number tandem-repeat analysis (MLVA) for use as a *L. monocytogenes* subtyping technique for surveillance and routine control in meat products and meat processing plants. A collection of 113 isolates (including control strains and isolates from meat products and meat processing plants) were subject to MLVA analysis using two different platforms for fragment sizing: 1.) ABI 3730xl DNA analyzer (Life Technologies) as the reference method and 2.) the QIAxcel Advanced System (Qiagen). Although discrepancies in fragment sizing were observed it was possible to standardize the results in order to assign the same allele for a given fragment independently of the platform used for fragment sizing. MLVA and multilocus sequence typing (MLST) results were compared and yielded Simpson's diversity indices of 0.907 and 0.872, respectively. The congruence between both typing methods was measured with the adjusted Wallace coefficient (AW). Using MLVA as the primary method, AW= 0.946 suggested that MLVA can predict the sequence type with high accuracy. Given its discriminatory power and high throughput, MLVA could be considered a rapid, reliable, and high-throughput alternative to existing subtyping methods for surveillance and control of *L. monocytogenes* in the meat-processing industry.

## Highlights

QIAxcel sizing platform is a trustworthy alternative to automated sequencers

Twenty-four MLVA unique profiles were determined among the meat industry isolates

MLVA had more discriminatory power than MLST, especially for serotype 1/2c isolates

MLVA can predict the sequence type with high accuracy

MLVA is a reliable method to type and trace *L.monocytogenes* in the meat industry

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**MLVA subtyping of *Listeria monocytogenes* isolates from meat products and meat processing plants**

Belén Martín, Sara Bover-Cid and Teresa Aymerich\*

IRTA, Food Safety Program. Finca Camps i Armet, 17121 Monells, Spain.

\*Corresponding author at: Food Safety Program, IRTA. Finca Camps i Armet, 17121 Monells, Spain.

E-mail address: [teresa.aymerich@irta.cat](mailto:teresa.aymerich@irta.cat) (T. Aymerich).

## ABSTRACT

1  
2  
3        *Listeria monocytogenes* is widely distributed in meat products and the meat-  
4  
5 processing industry thus posing a risk to consumers. The aim of this study was to  
6  
7 evaluate the suitability of the multilocus variable-number tandem-repeat analysis  
8  
9 (MLVA) for use as a *L. monocytogenes* subtyping technique for surveillance and  
10  
11 routine control in meat products and meat processing plants. A collection of 113  
12  
13 isolates (including control strains and isolates from meat products and meat processing  
14  
15 plants) were subject to MLVA analysis using two different platforms for fragment sizing:  
16  
17 1.) ABI 3730xl DNA analyzer (Life Technologies) as the reference method and 2.) the  
18  
19 QIAxcel Advanced System (Qiagen). Although discrepancies in fragment sizing were  
20  
21 observed it was possible to standardize the results in order to assign the same allele  
22  
23 for a given fragment independently of the platform used for fragment sizing. MLVA and  
24  
25 multilocus sequence typing (MLST) results were compared and yielded Simpson's  
26  
27 diversity indices of 0.907 and 0.872, respectively. The congruence between both typing  
28  
29 methods was measured with the adjusted Wallace coefficient (AW). Using MLVA as  
30  
31 the primary method, AW= 0.946 suggested that MLVA can predict the sequence type  
32  
33 with high accuracy. Given its discriminatory power and high throughput, MLVA could be  
34  
35 considered a rapid, reliable, and high-throughput alternative to existing subtyping  
36  
37 methods for surveillance and control of *L. monocytogenes* in the meat-processing  
38  
39 industry.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

*Listeria monocytogenes* is a ubiquitous bacterium that is known as the causative agent of human listeriosis, an important foodborne disease with a high fatality rate particularly in high-risk population such as the elderly, immunocompromised patients, pregnant woman and newborn infants. In The European Union, there has been an increasing trend of listeriosis during the period 2008-2015. A total of 1,524 confirmed human listeriosis cases were reported in 2015 with a fatality rate of 17.7% (EFSA-ECDC, 2016).

*L. monocytogenes* is widely distributed in food-processing environments (Ferreira, Wiedmann, Teixeira, & Stasiewicz, 2014; Martin et al., 2014; Paoli, Bhunia, & Bayles, 2005; Tompkin, 2002) thus posing a risk of contamination of food products.

Contaminated foods are consider the main vehicle for listeriosis (Scallan et al., 2011) particularly ready-to-eat (RTE) foods, which are intended to be consumed without further processing. Outbreaks and sporadic cases of listeriosis are generally associated to the consumption of those RTE foods such as soft cheese, smoked fish, vegetables and meat and meat products (Buchanan, Gorris, Hayman, Jackson, & Whiting, 2017; EFSA-ECDC, 2016).

Characterization of *L. monocytogenes* strains is needed in order to determine its virulence potential, for surveillance purposes and epidemiological tracking (Kathariou, 2002; Swaminathan & Gerner-Smidt, 2007). Among the 13 described serotypes of *L. monocytogenes*, serotypes 1/2a, 1/2b and 4b are implicated in most cases of human listeriosis and outbreaks (Clark et al., 2010; Doumith, Buchrieser, Glaser, Jacquet, & Martin, 2004; EFSA-ECDC, 2015, 2016; Pagotto, Ng, Clark, & Farber, 2006). On the other hand, serotype 1/2c are commonly described in food-processing environments and food products (Gelbicova & Karpiskova, 2009; Martin et al., 2014; Thévenot et al., 2006) but it has rarely been implicated in human listeriosis cases (Orsi, den Bakker, & Wiedmann, 2011).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Molecular typing methods are crucial for the identification and monitoring clonal groups of *L. monocytogenes* along the food chain. There are many molecular methods currently available for typing *L. monocytogenes* isolates differing in discriminatory power and epidemiological concordance (EFSA BIOHAZ Panel, 2013; van Belkum et al., 2007). Pulsed-field gel electrophoresis (PFGE) has been long considered the “gold standard” method for *L. monocytogenes* subtyping because of its high discriminatory power, reproducibility and repeatability (Gerner-Smidt et al., 2006; Lukinmaa, Aarnisalo, Suihko, & Siitonen, 2004). However, PFGE is considered a laborious and time-consuming technique with limited data portability (EFSA BIOHAZ Panel, 2013; Heir, Lindstedt, Vardund, Wasteson, & Kapperud, 2000). Nowadays, sequence-based typing methods provide unambiguous and portable data that can be useful not only for typing purposes, but also for the study of the population structure and evolution of this pathogen (Nightingale, 2010). Multi-locus sequence typing (MLST) is a well established sequenced-based typing method for studying population genetics of *L. monocytogenes* (Salcedo, Arreaza, Alcalá, de la Fuente, & Vazquez, 2003) providing an easy and unambiguous inter-laboratory exchange of data through public databases (Nightingale, 2010). Multi-virulence-locus sequence typing (MVLST) has also been used as a sequence-based approach for *L. monocytogenes* genotyping showing an excellent epidemiological concordance (Y. Chen, Zhang, & Knabel, 2007; Zhang, Jayarao, & Knabel, 2004). Nevertheless these techniques generally show a limited discriminatory power (den Bakker, Didelot, Fortes, Nightingale, & Wiedmann, 2008; EFSA BIOHAZ Panel, 2013; Ragon et al., 2008) for their use in the surveillance of *L. monocytogenes*. The advent of next generation sequencing technologies has dramatically reduced the cost of DNA sequencing making whole-genome sequencing (WGS) a convenient tool for molecular epidemiology and foodborne outbreak investigations (Datta, Laksanalamai, & Solomotis, 2013); thus, it is rapidly becoming the method of choice for *L. monocytogenes* genotyping in national reference laboratories. On the other hand, WGS is still prohibitive for most routine laboratories and generates massive amount of

1 data requiring intensive bioinformatic analysis, especially for testing a large number of  
2 isolates. Recently, multiple-locus variable number of tandem repeat analysis (MLVA)  
3 has been emerged as a powerful method to subtype food-borne pathogens such as  
4 *Salmonella enterica* serotypes Typhimurium and Enteritidis (Heck, 2009; Lindstedt,  
5 Heir, Gjernes, & Kapperud, 2003), *Escherichia coli* O157:H7 (Cooley et al., 2007) and  
6 *L. monocytogenes* (Lindstedt et al., 2008; Miya et al., 2008; Murphy et al., 2007;  
7 Sperry, Kathariou, Edwards, & Wolf, 2008). The approach is based on the detection of  
8 the number of tandem repeats at multiple variable-number tandem repeat loci  
9 distributed along the genome. Typically, multiplex PCR is used to amplify the tandem  
10 repeats and flanking regions and the amplification products are sized using capillary  
11 electrophoresis. MLVA is considered an easy and low-cost method which provides  
12 rapid and portable results with a high discriminatory power (Lindstedt et al., 2008;  
13 Sperry et al., 2008).

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In this study, we have applied MLVA to subtype a collection of *L. monocytogenes* strains isolated from meat-processing plants using both an automatic sequencer and/or capillary electrophoresis for amplicon sizing. MLVA results were compared to those previously obtained with MLST to evaluate the discriminatory power of the technique for its implementation in systematic environmental and product monitoring in meat processing plants.

## 2. Material and methods

### 2.1. *L. monocytogenes* isolates

A total of 106 isolates of *L. monocytogenes* were obtained from meat products and meat processing plants. Ninety-six isolates were collected from 18 meat processing plants, including 53 isolates from RTE meat products (fermented sausages, dry ham, blood sausages and other cured pork products), 10 isolates from raw meat products (beef and pork) and 33 isolates from food contact surfaces. In addition, 10 isolates



1 from the IRTA collection (recovered from meat products) were also analyzed. Seven  
2 control strains of *L. monocytogenes*, EGDe, ScottA, ATCC 35152 (equivalent to  
3 CIP104794), ATCC 19112 (equivalent to SLCC2372), ATCC 19114 (equivalent to  
4 SLCC2374), ATCC 19117 and CECT4032 (equivalent to F646/86) were used to  
5 calibrate the MLVA method.  
6  
7  
8  
9

10  
11 The serotype and sequence type (ST) of the 106 isolates were previously  
12 determined by us (Martin et al., 2014). For control strains, ST and serotype was  
13 obtained from the Listeria MLST Database hosted at the web site of the Institut Pasteur  
14 (<http://bigsd.b.pasteur.fr/listeria/>) and from Chenal-Francisque et al. (2011) and  
15 Cantinelli et al. (2013).  
16  
17  
18  
19  
20  
21

## 22 2.2. DNA extraction from *L. monocytogenes* isolates.

23  
24 Genomic DNA was extracted using DNeasy® blood and Tissue kit (Qiagen GmbH,  
25 Hilden, Germany) and the QIAcube® automated sample preparation system (Qiagen).  
26 Cultures were grown overnight at 37°C in Tryptic Soy Broth (BD, Sparks, MD) and 1 ml  
27 was centrifuged at 9,000 x g for 5 min; the pellets obtained were resuspended in 180 µl  
28 of lysis buffer (35 mg/ml lysozyme in 20 mM Tris-HCl pH 8.0, 2 mM EDTA, 1.2% Triton)  
29 and the enzymatic lysis step was programmed at 37°C for 1 h. DNA was eluted in 150  
30 µl of AE buffer (10 mM Tris-HCl, 0.5 mM EDTA pH 9.0), quantified using Quant-It™  
31 high sensitivity DNA assay kit (Invitrogen, Merelbeke, Belgium) and adjusted to 20  
32 ng/µl. The DNA was split into aliquots and stored at -20°C for further use in PCR.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

## 47 2.3. MLVA

48  
49 The MLVA procedure was conducted using the eight primer pairs proposed by  
50 Sperry et al. (Sperry et al., 2008; Sperry, Kathariou, Edwards, & Wolf, 2009) and two  
51 different multiplex-PCR protocols were performed. The first protocol consisted in two  
52 4-plex PCR reactions as in the reference protocol but using different dyes to label  
53 forward primers (Table 1), the Type-it Microsatellite PCR Kit (Qiagen), 0.2 µM of each  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 primer (all purchased at Life Technologies) and 2 µl of DNA. After amplification,  
2 fluorescent PCR products were resolved by automated capillary electrophoresis on an  
3 ABI 3730xl DNA analyzer (Life Technologies) with GeneScan-500 LIZ size standard  
4 (Life Technologies) using commercial GeneScan service (Macrogen Inc). Fragment  
5 sizes obtained were then analyzed using the Peakscanner software version 1.0  
6 (ThermoFisher Scientific). The second protocol consisted in 3 different multiplex-PCR  
7 reactions of 25 µl final volume using also the Type-it Microsatellite PCR Kit and the  
8 same primers (0.2 µM each) but not labelled. PCR reaction one (PCR1) contained  
9 primers for amplification of locus Lm-2, Lm-23 and Lm-32, the second PCR (PCR2)  
10 contained primers for locus Lm-3 and Lm-5 and the third PCR (PCR3) contained  
11 primers for locus Lm-8, Lm-10 and Lm-11. Thermal cycling conditions consisted of a  
12 denaturation step at 95°C for 5 min and 35 cycles of 30 s at 94°C, 30 s at 50°C and  
13 30 s at 72°C and a final extension step at 72°C for 5 min. All amplification reactions  
14 were performed in a thermal cycler GeneAmp PCR System 2700. The high-resolution  
15 capillary electrophoresis device QIAxcel Advanced System (Qiagen) was used to  
16 determine the size of PCR products. After PCR amplification, PCR plates were directly  
17 loaded onto the system. The QIAxcel DNA High Resolution kit was used applying the  
18 method OM800 (separation time of 920 s at 5 kV) with the Alignment Marker 15-600  
19 bp (Qiagen). Each QIAxcel run included a QX DNA size Marker 25-600 bp (Qiagen).  
20 Fragment sizing was performed using software QIAxcel Screengel v 1.4.0 (Qiagen).  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

#### 45 *2.4. Data analysis*

46  
47  
48 Estimated fragment size, dye (when present), peak height, and area data for each  
49 isolate were exported into Bionumerics Software v. 7.6 (Applied Maths, Sint-Martens-  
50 Latem, Belgium). With the first protocol (fragments size determined with ABI 3730xl  
51 DNA analyzer), seven control strains (EGDe, ScottA, ATCC 35152, ATCC 19112,  
52 ATCC 19114, ATCC 19117 and CECT4032) were used to calibrate the system and  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

adjust the offsets (Table 1) to obtain the same copy number for each VNTR locus as the reference study (Sperry et al., 2008). Null alleles were coded as negative (-2).

Results obtained with the second protocol (fragments size determined with QIAxcel advanced System) were calibrated according the results obtained with the automated sequencer. A look-up table was constructed for allele assignment containing the different alleles obtained for each locus and the corresponding observed fragment size (Table 2). This table was used as a mapping list in Bionumerics MLVA plugin for determining VNTR copy numbers.

The relationship among *L. monocytogenes* isolates was studied with the minimum spanning tree (MST) using categorical coefficient and the unweighted-pair group method with arithmetic mean (UPGMA). Both analysis were performed with Bionumerics software. Simpson's diversity index (SID) as described by Hunter and Gaston (1988) was used to measure genetic diversity among isolates. An index greater than 0.90 is considered desirable if the typing results are to be interpreted with confidence (Hunter & Gaston, 1988). The degree of congruence among MLVA and MLST was calculated using the adjusted Wallace coefficient (AW), which indicates the probability that two strains classified as the same type by one typing method will also be identified as the same type by the other method (Carrico et al., 2006). Both index were calculated using the Comparing Partitions tool available online (<http://www.comparingpartitions.info/>).

### 3. Results

#### 3.1. Optimization of *L. monocytogenes* MLVA

A total of 106 isolates previously recovered from meat processing plants and meat products (RTE and raw products) were submitted to MLVA typing. In addition, 7 control strains were also analyzed to first establish the correct offset to obtain the same copy number for each locus. *L. monocytogenes* ATCC 19117 was not analyzed in the study

1 carried out by Sperry et al. (2008), but as the complete genome sequence of this strain  
2 (Sumrall, Klumpp, Shen, & Loessner, 2016) together with that of strains EGDe (Glaser  
3 et al., 2001) and ATCC 19112 (Kuenne et al., 2013) have been determined, the real  
4 fragment size of the amplified fragments (determined *in silico*) were also taken into  
5 account for the establishment of the offsets. The MLVA typing system using ABI 3730xl  
6 for fragment sizing could be adjusted to obtain the same MLVA profiles previously  
7 published except for strain EGDe, which showed a different copy number for locus Lm-  
8 23 (20 instead of 19). In our hands, fragment analysis of locus Lm-23 in EGDe using  
9 ABI 3730xl automated sequencer was 229.70 bp, instead of 221.99 bp as previously  
10 described by Sperry et al. (2008) using a different sizing platform. Nevertheless, *in*  
11 *silico* MLVA analysis of EGDe showed a fragment of 231 bp for locus Lm-23, closer to  
12 our results and for that reason we decided to maintain the difference in that locus.  
13  
14

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Fragment sizing was also carried out using QIAxcel Advanced System. Overall, discrepancies between QIAxcel results and the reference values obtained with ABI 3730xl were observed (Table 2), being QIAxcel values consistently lower than the reference values. Differences augmented along with the increase of the PCR fragment length. Highest differences were observed among PCR fragments longer than 380 bp (mainly locus Lm-2), showing a difference from expected values of 4 to 14 bp. To overcome discrepancies among sizing platforms, a look-up table containing fragment size ranges for all alleles was constructed taking into account the results obtained with the reference method (using ABI 3730xl automated sequencer). This look-up table was used in Bionumerics software to assign the number of repeats at each locus. Using this approach, the concordance between both sizing methods was 100%.

### 3.2. *L. monocytogenes* distribution in meat products and meat processing plants by MLVA.

A total of 27 different MLVA profiles (Fig. 1) were obtained considering all the isolates from meat products, food processing plants and control strains (N=113).

1 Without control strains (N=106), the number of MLVA unique profiles was 24. Most  
2 common MLVA profiles were 19-3-5-4-6-2-20-13 assigned to MLVA Type (MT) 1  
3 (22.3% of isolates) and 15-3-5-4-1-4-37-14 (MT2, 15.2% of isolates). Isolates from the  
4 IRTA collection (recovered from RTE meat products in different years) displayed 7  
5 different MTs, but common to the MTs displayed by others isolates from meat products  
6 and /or food-contact surfaces (MT1, MT2, MT7, MT8 and MT16), except for MT15  
7 (which included two IRTA collection isolates) and MT19 (one single isolate).  
8  
9

10 The suitability of MLVA to establish the contamination patterns of *L. monocytogenes*  
11 in meat processing plants was also evaluated. Fig. 2 shows the dendrogram generated  
12 with the isolates (N=96) recovered from the 18 meat processing plants evaluated on  
13 the basis of MLVA data, the source information, in addition to serotype and ST. The  
14 isolates showed 21 different MTs distributed throughout the meat processing plants.  
15 MT1 was the most frequent genotype, represented by 23 isolates and detected in 12 of  
16 the plants studied. MT2 (corresponding to ST121) was represented by 16 isolates  
17 recovered from several plants (8 out of 18). In contrast, 11 different MTs were only  
18 detected in one single plant. Although, MLVA identified more types than MLST, similar  
19 results were obtained regarding to the source tracking of isolates, and 25 of the  
20 isolates recovered from meat products showed the same genotype as isolates  
21 collected from food contact surfaces of their respective production plants.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 3.3. Evaluation of the MLVA subtyping method in comparison with MLST

44 The discriminatory power of the MLVA subtyping method was evaluated comparing  
45 the results with the previously obtained by MLST (Martin et al., 2014). The number of  
46 MLVA unique profiles obtained was 27 corresponding to the 22 different allelic profiles  
47 previously identified by MLST. Most STs could be clearly differentiated or even further  
48 subtyped into different MTs; ST9 could be split in four different MTs and ST8, ST3 and  
49 ST1 in two unique MTs each. In contrast, ST122 (corresponding to control strain  
50 ATCC 19112) could not be distinguished from ST9 isolates.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Clustering analysis of MLVA data was highly consistent with MLST analysis and  
2 interestingly, isolates belonging to the same ST clustered together. MLVA identified  
3 some clonal complexes (CC) in accordance with MLST data, CC9 (that included ST9,  
4 ST625 and ST35 isolates), CC8 (including ST8 and ST16 isolates), CC3 (composed  
5 by two MTs of ST3 isolates) CC1 (composed by three MTs) and CC87 (composed by  
6 three MTs). In contrast, two different ST isolates (ST12 and ST204) clustered together,  
7 while by MLST they belong to different CCs. Table 3 shows the different MLVA profiles  
8 found in relation with the serotypes and STs previously described. Simpson's diversity  
9 index (SID) was calculated based on different genotypes obtained to assess  
10 discrimination ability of both MLST and MLVA. The diversity of the complete panel of  
11 isolates (N=113) was SID=0.907 by MLVA and slightly lower by MLST (SID=0.872).  
12 Differences in diversity were observed among serotypes (Table 3). Serotype 1/2a  
13 (N=41) demonstrated the highest diversity, regardless the subtyping method used  
14 (SID=0.796 by MLVA and SID=0.778 by MLST). Serotype 1/2b (N=19) also showed a  
15 high diversity index (SID=0.754) when using MLVA, but clearly lower by MLST  
16 (SID=0.556). Serotype 4b (N=14) showed also a higher diversity index by MLVA than  
17 by MLST, SID=0.659 and 0.513 respectively. However, the most relevant difference  
18 was in serotype 1/2c (N=37), where diversity was almost double by MLVA (SID=0.493)  
19 compared to MLST (SID=0.246). The adjusted Wallace coefficients (AW) obtained  
20 were  $AW_{MLVA \rightarrow MLST} = 0.946$  (CI=0.857-1.000) and  $AW_{MLST \rightarrow MLVA} = 0.660$  (CI=0.518-0.802).  
21 These values indicated that ST was well predicted by MLVA but MT was less well  
22 predicted by MLST.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

#### 49 **4. Discussion**

50  
51  
52 Molecular typing of *L. monocytogenes* isolates has an important role in meat-  
53 processing plants in order to identify contamination and spread routes of this pathogen.  
54  
55 Currently typing methods used including PFGE, MLST and recently WGS, present  
56  
57 several drawbacks for their use in food control laboratories, such as the intensively  
58  
59  
60  
61  
62  
63  
64  
65

1 work required and the high cost. MLVA is an alternative molecular typing method with  
2 practical advantages including its rapidity, ease of use and low cost (EFSA BIOHAZ  
3 Panel, 2013; van Belkum et al., 2007). In this study, we have optimized a MLVA  
4 protocol to discriminate among *L. monocytogenes* isolates recovered from meat  
5 products and meat processing plants using two different sizing technologies and  
6 comparing the results with those of MLST.  
7  
8  
9  
10  
11  
12

13 MLVA is based on the accurate sizing of PCR amplified DNA fragments by  
14 capillary electrophoresis. Nevertheless, important differences between measured and  
15 real fragment lengths are commonly observed (Larsson, Torpdahl, Nielsen, & Group,  
16 2013) mainly attributed to the different capillary electrophoresis platforms and reagents  
17 used for fragment sizing (Hyytia-Trees, Lafon, Vauterin, & Ribot, 2010). In this sense,  
18 the use of a standardized MLVA method to normalized fragment size data is a must.  
19 Efforts for standardization have been carried out for MLVA analysis of some food-borne  
20 pathogens (Hyytia-Trees et al., 2010; Larsson et al., 2013) and although several MLVA  
21 schemes for typing *L. monocytogenes* have been published (S. Chen, Li, Saleh-Lakha,  
22 Allen, & Odumeru, 2011; Chenal-Francisque et al., 2013; Lindstedt, 2005; Murphy et  
23 al., 2007; Saleh-Lakha et al., 2013) none of them has been established as the standard  
24 method. Among the MLVA methods available, we chose the MLVA scheme developed  
25 by Sperry et al. (2008) intended to be implemented in Pulsenet Laboratories and  
26 adapted it to be used with ABI 3730xl. Offsets had to be adjusted in order to maintain  
27 the copy numbers of VNTR loci previously published to obtain comparable results.  
28 Nevertheless, it was not possible to calculate an offset for locus Lm-23 to maintain the  
29 copy number of this locus for all control strains (EGDe, showed a higher copy number,  
30 20 instead of 19). Some authors recommends the use of an established set of  
31 calibration strains to convert the discrepancies between real (sequenced) and  
32 measured fragment length to obtain comparable results between laboratories  
33 independently of the capillary electrophoresis platform used (Larsson et al., 2013). This  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 implies that each new allele must be verified by sequencing but it would avoid errors in  
2 copy number assignation due to discrepancies on measured fragment lengths.  
3

4  
5       Apart from the results obtained using ABI 3730xl, MLVA was also carried out using  
6  
7 QIAxcel Advanced System (developed by Qiagen), an automated capillary  
8  
9 electrophoresis platform that constitutes an alternative to the methods based on  
10  
11 separation of fluorescent-labelled PCR fragments using automated sequencers. This  
12  
13 platform has been successfully used for MLVA genotyping of several bacterial species  
14  
15 other than *L. monocytogenes* (De Santis et al., 2013; Gauthier et al., 2015;  
16  
17 Nikolayevskyy et al., 2016; Takahashi et al., 2014) leading to shorter turnaround times,  
18  
19 technical simplicity and cost effectiveness what can be of great interest for small food  
20  
21 control laboratories. In this study, discrepancies on size fragment were observed  
22  
23 between both platforms, mainly in fragments higher than 380 bp. The results obtained  
24  
25 with ABI 3730xl were taken as reference values and a look-up table containing  
26  
27 fragment size ranges was constructed for allele assignment as previously suggested to  
28  
29 overcome discrepancies on fragment size data among platforms (Hyytia-Trees et al.,  
30  
31 2010). In this way, concordance between both sizing platforms was 100%, concluding  
32  
33 that QIAxcel is an effective and affordable alternative to automated sequencers that  
34  
35 exhibit adequate performance after accurate validation.  
36  
37  
38  
39  
40

41 MLVA was compared with the widely used MLST scheme of Institute Pasteur that  
42  
43 establishes a common nomenclature for *L. monocytogenes* genotypes. High level of  
44  
45 congruence was obtained between both molecular typing methods suggesting that  
46  
47 MLVA can be routinely used as an alternative method to MLST. Similar results of  
48  
49 agreement between MLVA and MLST was also obtained by other authors (Chenal-  
50  
51 Francisque et al., 2013) although a different MLVA scheme was applied for genotyping  
52  
53 *L. monocytogenes* isolates recovered from different sources (mainly human). Most STs  
54  
55 could be differentiated by MLVA and all MLST clonal complexes could be also  
56  
57 identified. This level of congruence could be explained by a similar level of variation in  
58  
59  
60  
61  
62  
63  
64  
65



1 both MLVA and MLST markers (Chenal-Francisque et al., 2013) and by the low level of  
2 genetic recombination in *L. monocytogenes* (Cantinelli et al., 2013; den Bakker et al.,  
3 2010; Ragon et al., 2008). Recombination is more prevalent in lineage II than in lineage  
4 I, which is more affected by positive selection (Cantinelli et al., 2013; den Bakker et al.,  
5 2008). Regarding most prevalent STs found, MLVA confirmed the clonal origin of  
6 ST121 isolates whereas ST9 was more diversified by MLVA than by MLST.  
7

8 Interestingly, MLVA could also distinguished one single ST9 isolate that showed a  
9 different virulence type (VT95) from the others as reported in a previous study (Martin  
10 et al., 2014) . From our results, it is possible to deduce ST from the MT obtained  
11 although this encompass the MLST analysis of every newly described MT.  
12

13 Discriminatory power of MLVA varies depending on the MLVA scheme used and the  
14 collection of isolates evaluated. In our study, MLVA showed a higher discriminatory  
15 power than MLST and differences were observed regarding the serotype and ST of the  
16 isolates. Several studies showed a discriminatory power of MLVA similar or even  
17 higher than PFGE (Li et al., 2013; Saleh-Lakha et al., 2013) whereas others showed  
18 lower discrimination of MLVA (Chenal-Francisque et al., 2013; Sperry et al., 2008).  
19 Chenal- Francisque et al. (2013) showed that discrimination power of MLVA is  
20 dependent on the clonal complex, being highly efficient in the discrimination of CC9  
21 and CC4 isolates. In accordance with this results, we also showed a high diversity  
22 among ST9 isolates, the most prevalent among the isolates studied, what could  
23 underline the utility of MLVA for subtyping *L. monocytogenes* isolates in meat-  
24 processing environments to identify the sources of contamination and improve the  
25 safety of meat products.  
26

## 27 **5. Conclusions**

28 Our results show that MLVA could be used as a rapid and reliable subtyping  
29 method for *L. monocytogenes* isolates in meat-processing plants. It permits to track the  
30 spread of clones and the persistence of particular subtypes and can be an efficient and  
31

1 high- throughput tool to improve food safety control and *L. monocytogenes*  
2 surveillance.  
3  
4  
5  
6

## 7 **Acknowledgements**

8  
9  
10 This work was supported by the Spanish Science Ministry project CDS 2007-0016  
11 and by the CERCA Programme/Generalitat de Catalunya. We would like to thank  
12 Javier Yangüela for providing strains recovered from meat processing plants. We also  
13 thank Montse Badia and Carmen Raya for their technical assistance.  
14  
15  
16  
17  
18  
19

## 20 **References**

- 21  
22  
23 Buchanan, R. L., Gorris, L. G. M., Hayman, M. M., Jackson, T. C., & Whiting, R. C.  
24 (2017). A review of *Listeria monocytogenes*: An update on outbreaks, virulence,  
25 dose-response, ecology, and risk assessments. *Food Control*, 75, 1-13.  
26  
27  
28  
29  
30 Cantinelli, T., Chenal-Francisque, V., Diancourt, L., Frezal, L., Leclercq, A., Wirth, T.,  
31 Brisse, S. (2013). "Epidemic Clones" of *Listeria monocytogenes* are widespread  
32 and ancient clonal groups. *Journal of Clinical Microbiology*, 51(11), 3770-3779.  
33  
34  
35  
36 Carrico, J. A., Silva-Costa, C., Melo-Cristino, J., Pinto, F. R., de Lencastre, H.,  
37 Almeida, J. S., & Ramirez, M. (2006). Illustration of a common framework for  
38 relating multiple typing methods by application to macrolide-resistant  
39 *Streptococcus pyogenes*. *Journal of Clinical Microbiology*, 44(7), 2524-2532.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Chen, S., Li, J., Saleh-Lakha, S., Allen, V., & Odumeru, J. (2011). Multiple-locus  
variable number of tandem repeat analysis (MLVA) of *Listeria monocytogenes*  
directly in food samples. *International Journal of Food Microbiology*, 148(1), 8-  
14.
- Chen, Y., Zhang, W., & Knabel, S. J. (2007). Multi-virulence-locus sequence typing  
identifies single nucleotide polymorphisms which differentiate epidemic clones

1 and outbreak strains of *Listeria monocytogenes*. *Journal of Clinical*  
2 *Microbiology*, 45(3), 835-846.  
3

4 Chenal-Francisque, V., Diancourt, L., Cantinelli, T., Passet, V., Tran-Hykes, C., Bracq-  
5  
6 Dieye, H., Brisse, S. (2013). Optimized Multilocus Variable-Number Tandem-  
7 Repeat Analysis Assay and Its Complementarity with Pulsed-Field Gel  
8 Electrophoresis and Multilocus Sequence Typing for *Listeria monocytogenes*  
9 Clone Identification and Surveillance. *Journal of Clinical Microbiology*, 51(6),  
10 1868-1880.  
11

12 Clark, C. G., Farber, J., Pagotto, F., Ciampa, N., Dore, K., Nadon, C., Ng, L. K. (2010).  
13 Surveillance for *Listeria monocytogenes* and listeriosis, 1995-2004.  
14  
15 *Epidemiology and Infection*, 138(4), 559-572.  
16

17 Cooley, M., Carychao, D., Crawford-Miksza, L., Jay, M. T., Myers, C., Rose, C.,  
18  
19 Mandrell, R. E. (2007). Incidence and tracking of *Escherichia coli* O157:H7 in a  
20 major produce production region in California. *Plos One*, 2(11), e11159.  
21

22 Datta, A. R., Laksanalamai, P., & Solomotis, M. (2013). Recent developments in  
23  
24 molecular sub-typing of *Listeria monocytogenes*. *Food Additives and*  
25  
26 *Contaminants Part a-Chemistry Analysis Control Exposure & Risk Assessment*,  
27  
28 30(8), 1437-1445.  
29

30 De Santis, R., Ancora, M., De Massis, F., Ciammaruconi, A., Zilli, K., Di Giannatale, E.,  
31  
32 Lista, F. (2013). Molecular strain typing of *Brucella abortus* isolates from Italy by  
33  
34 two VNTR allele sizing technologies. *Molecular Biotechnology*, 55(2), 101-110.  
35

36 den Bakker, H. C., Cummings, C. A., Ferreira, V., Vatta, P., Orsi, R. H., Degoricija, L.,  
37  
38 Wiedmann, M. (2010). Comparative genomics of the bacterial genus *Listeria*:  
39  
40 Genome evolution is characterized by limited gene acquisition and limited gene  
41  
42 loss. *Bmc Genomics*, 11, 688-708.  
43

44 den Bakker, H. C., Didelot, X., Fortes, E. D., Nightingale, K. K., & Wiedmann, M.  
45  
46 (2008). Lineage specific recombination rates and microevolution in *Listeria*  
47  
48 *monocytogenes*. *Bmc Evolutionary Biology*, 8, 277-290.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Doumith, M., Buchrieser, C., Glaser, P., Jacquet, C., & Martin, P. (2004). Differentiation of the major *Listeria monocytogenes* serovars by Multiplex PCR. *J. Clin. Microbiol.*, 42(8), 3819-3822.
- EFSA-ECDC. (2015). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2013. *EFSA Journal*, 13(1), 3991-4153.
- EFSA-ECDC. (2016). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2015. *EFSA Journal*, 13(12), 4329.
- EFSA BIOHAZ Panel. (2013). Scientific Opinion on the evaluation of molecular typing methods for major food-borne microbiological hazards and their use for attribution modelling, outbreak investigation and scanning surveillance: Part 1 (evaluation of methods and applications). *EFSA Journal*, 11(12), 3508-3584.
- Ferreira, V., Wiedmann, M., Teixeira, P., & Stasiewicz, M. J. (2014). *Listeria monocytogenes* persistence in food-associated environments: epidemiology, strain characteristics, and implications for public health. *Journal of Food Protection*, 77(1), 150-170.
- Gauthier, M., Bidault, F., Mosnier, A., Bablishvili, N., Tukvadze, N., Somphavong, S., Berland, J.-L. (2015). High-throughput mycobacterial interspersed repetitive-unit-variable-number tandem-repeat genotyping for *Mycobacterium tuberculosis* epidemiological studies. *Journal of Clinical Microbiology*, 53(2), 498-503.
- Gelbicova, T., & Karpiskova, R. (2009). Occurrence and characteristics of *Listeria monocytogenes* in ready-to-eat food from retail market in the Czech Republic. *Czech Journal of Food Sciences*, 27, 3-7.
- Gerner-Smidt, P., Hise, K., Kincaid, J., Hunter, S., Rolando, S., Hyytia-Trees, E., PulseNet, T. (2006). PulseNet USA: A five-year update. *Foodborne Pathogens and Disease*, 3(1), 9-19.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Glaser, P., Frangeul, L., Buchrieser, C., Rusniok, C., Amend, A., Baquero, F., Cossart, P. (2001). Comparative genomics of *Listeria* species. *Science*, 294(5543), 849-852.
- Heck, M. (2009). Multilocus variable number of tandem repeats analysis (MLVA) - a reliable tool for rapid investigation of *Salmonella typhimurium* outbreaks. *Eurosurveillance*, 14(15), 2-2.
- Heir, E., Lindstedt, B. A., Vardund, T., Wasteson, Y., & Kapperud, G. (2000). Genomic fingerprinting of shigatoxin-producing *Escherichia coli* (STEC) strains: comparison of pulsed-field gel electrophoresis (PFGE) and fluorescent amplified-fragment-length polymorphism (FAFLP). *Epidemiology and Infection*, 125(3), 537-548.
- Hunter, P. R., & Gaston, M. A. (1988). Numerical Index of the Discriminatory Ability of Typing Systems: an Application of Simpson's Index of Diversity. *Journal of Clinical Microbiology*, 26(11), 2465-2466.
- Hyytia-Trees, E., Lafon, P., Vauterin, P., & Ribot, E. M. (2010). Multilaboratory validation study of standardized multiple-locus variable-number tandem repeat analysis protocol for shiga toxin-producing *Escherichia coli* O157: a novel approach to normalize fragment size data between capillary electrophoresis platforms. *Foodborne Pathogens and Disease*, 7(2), 129-136.
- Kathariou, S. (2002). *Listeria monocytogenes* virulence and patogenicity, a food safety perspective. *Journal of Food Protection*, 65, 1811-1829.
- Kuenne, C., Billion, A., Abu Mraheil, M., Strittmatter, A., Daniel, R., Goesmann, A., Chakraborty, T. (2013). Reassessment of the *Listeria monocytogenes* pan-genome reveals dynamic integration hotspots and mobile genetic elements as major components of the accessory genome. *BMC Genomics*, 14.
- Larsson, J. T., Torpdahl, M., Nielsen, E. M., & Group, M. W. (2013). Proof-of-concept study for successful inter-laboratory comparison of MLVA results. *Eurosurveillance*, 18(35), 20-27.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Li, X. J., Huang, B. X., Eglezos, S., Graham, T., Blair, B., & Bates, J. (2013). Identification of an optimized panel of variable number tandem-repeat (VNTR) loci for *Listeria monocytogenes* typing. *Diagnostic Microbiology and Infectious Disease*, 75(2), 203-206.
- Lindstedt, B. A. (2005). Multiple-locus variable number tandem repeats analysis for genetic fingerprinting of pathogenic bacteria. *Electrophoresis*, 26(13), 2567-2582.
- Lindstedt, B. A., Heir, E., Gjernes, E., & Kapperud, G. (2003). DNA fingerprinting of *Salmonella enterica* subsp *enterica* serovar Typhimurium with emphasis on phage type DT104 based on variable number of tandem repeat loci. *Journal of Clinical Microbiology*, 41(4), 1469-1479.
- Lindstedt, B. A., Tham, W., Danielsson-Tham, M.-L., Vardund, T., Helmersson, S., & Kapperud, G. (2008). Multiple-locus variable-number tandem-repeats analysis of *Listeria monocytogenes* using multicolour capillary electrophoresis and comparison with pulsed-field gel electrophoresis typing. *Journal of Microbiological Methods*, 72(2), 141-148.
- Lukinmaa, S., Aarnisalo, K., Suihko, M. L., & Siitonen, A. (2004). Diversity of *Listeria monocytogenes* isolates of human and food origin studied by serotyping, automated ribotyping and pulsed-field gel electrophoresis. *Clinical Microbiology and Infection*, 10(6), 562-568.
- Martin, B., Perich, A., Gomez, D., Yangüela, J., Rodriguez, A., Garriga, M., & Aymerich, T. (2014). Diversity and distribution of *Listeria monocytogenes* in meat processing plants. *Food Microbiology*, 44, 119-127.
- Miya, S., Kimura, B., Sato, M., Takahashi, H., Ishikawa, T., Suda, T., Wiedmann, M. (2008). Development of a multilocus variable-number of tandem repeat typing method for *Listeria monocytogenes* serotype 4b strains. *International Journal of Food Microbiology*, 124(3), 239-249.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Murphy, M., Corcoran, D., Buckley, J. F., O'Mahony, M., Whyte, P., & Fanning, S. (2007). Development and application of Multiple-Locus Variable Number of tandem repeat Analysis (MLVA) to subtype a collection of *Listeria monocytogenes*. *International Journal of Food Microbiology*, 115(2), 187-194.
- Nightingale, K. (2010). *Listeria monocytogenes*: knowledge gained through DNA sequence-based subtyping, implications, and future considerations. *Journal of Aoac International*, 93(4), 1275-1286.
- Nikolayevskyy, V., Trovato, A., Broda, A., Borroni, E., Cirillo, D., & Drobniewski, F. (2016). MIRU-VNTR genotyping of *Mycobacterium tuberculosis* strains using Qiaxcel technology: a multicentre evaluation study. *Plos One*, 11(3).
- Orsi, R. H., den Bakker, H. C., & Wiedmann, M. (2011). *Listeria monocytogenes* lineages: Genomics, evolution, ecology, and phenotypic characteristics. *International Journal of Medical Microbiology*, 301(2), 79-96.
- Pagotto, F., Ng, L. K., Clark, C., & Farber, J. (2006). Canadian listeriosis reference service. *Foodborne Pathogens and Disease*, 3(1), 132-137.
- Paoli, G. C., Bhunia, A. K., & Bayles, D. O. (2005). *Listeria monocytogenes*. In P. M. Fratamico, A. K. Bhunia & J. L. Smith (Eds.), *Foodborne pathogens. Microbiology and molecular biology*. Norfolk, UK: Caister Academic Press.
- Ragon, M., Wirth, T., Hollandt, F., Lavenir, R., Lecuit, M., Le Monnier, A., & Brisse, S. (2008). A new perspective on *Listeria monocytogenes* evolution. *Plos Pathogens*, 4(9), e1000146.
- Salcedo, C., Arreaza, L., Alcalá, B., de la Fuente, L., & Vazquez, J. A. (2003). Development of a multilocus sequence typing method for analysis of *Listeria monocytogenes* clones. [Article]. *Journal of Clinical Microbiology*, 41(2), 757-762.
- Saleh-Lakha, S., Allen, V. G., Li, J., Pagotto, F., Odumeru, J., Taboada, E., Chen, S. (2013). Subtyping of a large collection of historical *Listeria monocytogenes* strains from Ontario, Canada, by an improved multilocus variable-number

1 tandem-repeat analysis (MLVA). *Applied and Environmental Microbiology*,  
2 79(20), 6472-6480.  
3

4 Scallan, E., Hoekstra, R. M., Angulo, F. J., Tauxe, R. V., Widdowson, M. A., Roy, S. L.,  
5  
6 Griffin, P. M. (2011). Foodborne illness acquired in the United States-Major  
7 pathogens. *Emerging Infectious Diseases*, 17(1), 7-15.  
8

9  
10 Sperry, K. E. V., Kathariou, S., Edwards, J. S., & Wolf, L. A. (2008). Multiple-locus  
11 variable-number tandem-repeat analysis as a tool for subtyping *Listeria*  
12 *monocytogenes* strains. *Journal of Clinical Microbiology*, 46(4), 1435-1450.  
13  
14

15  
16 Sperry, K. E. V., Kathariou, S., Edwards, J. S., & Wolf, L. A. (2009). Multiple-locus  
17 variable-number tandem-repeat analysis as a tool for subtyping *Listeria*  
18 *monocytogenes* strains (vol 46, pg 1435, 2008). *Journal of Clinical*  
19 *Microbiology*, 47(6), 1984-1984.  
20  
21

22  
23 Sumrall, E., Klumpp, J., Shen, Y., & Loessner, M. J. (2016). Genome sequences of five  
24 nonvirulent *Listeria monocytogenes* serovar 4 strains. *Genome*  
25 *announcements*, 4(2), e00179-00116  
26  
27

28  
29 Swaminathan, B., & Gerner-Smidt, P. (2007). The epidemiology of human listeriosis.  
30 *Microbes and Infection* 9, 1236-1243.  
31  
32

33  
34 Takahashi, H., Ohshima, C., Nakagawa, M., Thanatsang, K., Phraephaisarn, C.,  
35 Chaturongkasumrit, Y., Kimura, B. (2014). Development of new multilocus  
36 variable number of tandem repeat analysis (MLVA) for *Listeria innocua* and its  
37 application in a food processing plant. *Plos One*, 9(9).  
38  
39

40  
41 Thévenot, D., Delignette-Muller, M. L., Christieans, S., Leroy, S., Kodjo, A., & Vernozzy-  
42 Rozand, C. (2006). Serological and molecular ecology of *Listeria*  
43 *monocytogenes* isolates collected from 13 French pork meat salting-curing  
44 plants and their products. *International Journal of Food Microbiology*, 112(2),  
45 153-161.  
46  
47

48  
49 Tompkin, R. B. (2002). Control of *Listeria monocytogenes* in the food-processing  
50 environment. *Journal of Food Protection*, 65(4), 709-725.  
51  
52  
53  
54  
55  
56  
57



1 van Belkum, A., Tassios, P. T., Dijkshoorn, L., Haeggman, S., Cookson, B., Fry, N. K.,  
2 Esgem. (2007). Guidelines for the validation and application of typing methods  
3 for use in bacterial epidemiology. *Clinical Microbiology and Infection*, 13, 1-46.  
4  
5 Zhang, W., Jayarao, B. M., & Knabel, S. J. (2004). Multi-virulence-locus sequence  
6 typing of *Listeria monocytogenes*. *Applied and Environmental Microbiology*,  
7 70(2), 913-920.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure Captions

1  
2  
3  
4  
5 **Fig.1.** Minimum spanning tree analysis of the 113 isolates of *L. monocytogenes*  
6  
7 based on MLVA data. Each circle corresponds to one MT and the size of the circle  
8  
9 reflects the number of strains within that type. Numbers inside each circle represent the  
10  
11 ST of the isolates. The connection lines between circles differ based on the number of  
12  
13 different alleles between the corresponding MTs; bold lines correspond to one different  
14  
15 allele, plain lines to two alleles, dashed lines to three alleles and dotted lines to 4 or  
16  
17 more alleles. Grey zone surrounding some circles indicate that these types belong to  
18  
19 the same clonal complex. Control strains were also included in the analysis and they  
20  
21 are indicated in the figure.  
22  
23  
24  
25  
26  
27


28 **Fig.2.** UPGMA dendrogram of the 96 Isolates collected from meat-processing  
29  
30 plants and either from the food-contact surfaces sampled or the meat products  
31  
32 elaborated in each plant. Characteristics of each isolates including serotype, MT (MT),  
33  
34 sequence type (ST), source and meat-processing plant of isolation are indicated. Some  
35  
36 isolates presenting the same characteristics are grouped together and number of  
37  
38 isolates (N) are indicated.  
39  
40  
41

42 <sup>a</sup> MT, MLVA Type.

43  
44  
45 <sup>b</sup> Source: FCS, Food-contact surface. When indicated (clean), food-contact  
46  
47 surfaces were sampled prior processing but after cleaning and disinfection; RMP, Raw  
48  
49 meat product; RTE, Ready-to-eat meat product.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure

[Click here to download high resolution image](#)



Figure




Table 1. Parameters for the calculation of VNTR loci copy numbers with the ABI 3730xl platform.

VNTR locus	Offset	Repeat size (bp)	Min value	Max Value	Tolerance	Reaction group /fluorescent dye
Lm-2	291	6	11	20	3	R1 / VIC
Lm-8	185	15	3	4	7	R1/ NED
Lm-10	316	6	3	9	3	R1/ 6-FAM
Lm-11	100	12	1	6	6	R1/ 6-FAM
Lm-3	200	9	1	9	4	R2/ VIC
Lm-15	318	12	1	7	6	R2/ 6-FAM
Lm-23	169	3	15	42	1.5	R2/ NED
Lm-32	83	6	10	21	3	R2/ 6-FAM



	10	198.8	198.8	198.8	-	201	201	200.9	-	1
	18	222.7	222.7	222.7	-	224	224	223.9	-	1
	20	228.3	229.9	228.6	0.49	229	232	230.2	0.63	40
	23	236.7	236.9	236.8	0.08	235	237	236.5	0.99	4
	29	252.7	255.9	254.2	0.92	253	255	253.6	0.72	19
	33	265.9	267.2	266.4	0.54	265	267	265.8	0.85	10
	35	271.1	272.3	271.4	0.52	271	271	270.6	0.15	7
	37	280.0	281.0	280.3	0.45	282	283	282.3	0.49	17
	41	290.1	291.2	290.7	0.45	293	294	293.7	0.38	9
	42	295.1	295.1	295.1	.	297	297	297.0	-	1
Lm-32	-2	-	-	-	-	-	-	-	-	1
	13	160.7	161.0	160.8	0.08	163	165	163.7	0.64	57
	14	166.2	166.4	166.3	0.06	169	171	169.7	0.58	21
	17	183.7	183.9	183.8	0.06	186	188	187.2	0.75	24
	18	189.7	189.8	189.7	0.03	192	195	193.5	0.88	10

<sup>a</sup> Standard deviation

Table 3. MLVA subtyping of *L. monocytogenes* isolates in relation with serotypes and sequence types (ST).

Serotype	Simpson's index (SID)		ST	MLVA Type	MLVA profile <sup>a</sup>	N <sup>b</sup>
	MLST	MLVA				
1/2a	0.778	0.796	431	23	17-3-5-4-1-6-18-14	1
			11	12	16-3-5-1-1-2-NA-14	3
			121	2	15-3-5-4-1-4-37-14	17
			155	6	16-3-5-4-1-2-29-13	6
			204	22	17-3-5-NA-1-2-20-13	1
			26	19	18-3-5-4-2-3-23-13	1
			31	24	16-3-5-NA-3-4-10-13	1
			12	27	15-3-5-NA-1-2-20-13	1
			16	21	17-3-5-NA-4-2-41-13	1
			8	11	16-3-5-NA-5-2-41-13	3
				7	17-3-5-NA-5-2-41-13	5
			35	18	18-3-5-4-4-2-20-13	1
			1/2b	0.556	0.754	224
3	8	16-3-5-5-8-5-29-17				4
	4	16-3-5-5-9-5-29-17				7
87	26	15-3-7-5-3-4-35-17				1
	5	16-3-7-5-3-4-35-17				6
1/2c	0.246	0.493	122	1	19-3-5-4-6-2-20-13	1
			625	10	19-3-5-4-5-2-20-13	4
			9	1	19-3-5-4-6-2-20-13	25
				9	19-3-5-4-9-2-20-13	4
				15	18-3-5-4-6-2-20-13	2
				17	19-3-5-4-8-2-20-13	1
4b	0.513	0.659	1 <sup>c</sup>	16	15-4-5-5-1-3-33-18	2
				3	15-4-5-6-1-3-33-18	8
			2	13	15-3-3-5-1-5-23-17	2
			194	14	18-4-5-5-3-4-29-17	2
4a			136	42	11-3-5-NA-NA-2-42-NA	1
4d			2	5	15-3-3-5-1-5-23-17	1



<sup>a</sup> MLVA allele profile was presented in an allelic string as follows: Lm-2/Lm-8/Lm-10/Lm-11/Lm-3/Lm-15/Lm-23/Lm-32. NA denotes a null allele at this locus

<sup>b</sup> Number of isolates that share the same MLVA type.



<sup>c</sup> The ST of one isolate showing MT16 (CECT4032) was not determined, therefore it has not been considered to calculate SID value for MLST in serotype 4b.


**\*Graphical Abstract**




*L. monocytogenes* isolates



MLVA Typing using two platforms



**1) Reference method: ABI 3730xl DNA analyzer (Life Technologies)**



**2) QIAXcel Advanced System (Qiagen)**

