



This is the peer reviewed version of the following article: Giménez Papiol, G., Roque, A., & Estévez, A. (2017). Variations in Bacterial Community of Rearing Water and Gut of Common Dentex, *Dentex dentex* (Linnaeus 1758), Larvae Using Three Microalgae Management Approaches. *Journal Of The World Aquaculture Society*, 49(5), 953-963. doi:10.1111/jwas.12437, which has been published in final form at <https://doi.org/10.1111/jwas.12437>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions

1 Variations In Bacterial Community Of Rearing Water And Gut Of Common
2 Dentex, Dentex dentex Linnaeus 1758, Larvae Using Three Microalgae
3 Management Approaches

4

5 Gemma Giménez Papiol^{1, 2*}, Ana Roque², Alicia Estévez²

6 ¹Laboratory of Toxicology and Environmental Health, Environmental Analysis
7 and Management Group, Chemical Engineering Department, Universitat Rovira
8 i Virgili, Països Catalans 26, 43007 Tarragona, Spain.

9 ²Institut de Recerca i Tecnologia Agroalimentàries (IRTA), Ctra. Poble Nou Km.
10 5.5, 43540 Sant Carles de la Ràpita, Tarragona, Spain

11

12 Phone: +34 977 55 85 46

13 *corresponding author: gemma.gimenez@fundacio.urv.cat

14

15 Running title: Bacterial community in common dentex larvae rearing

16

17

Abstract

18 **Microorganisms present in the rearing water colonize the gut of first**
19 **feeding larvae and represent the first barrier against opportunistic**
20 **pathogens. The aim of the experiments presented herein was to standardize**
21 **a protocol for the management of rearing water and microalgae suitable for**
22 **the larval rearing of common dentex. In Experiment 1, bacteria-algae**
23 **interactions were tested using a monospecific microalgal community,**
24 **“Tetraselmis chuii”, suitable for nutritional experiments and with known**
25 **antibacterial activity. In Experiment 2, the evolution of the bacterial**
26 **community and larval performance (growth and survival) were monitored**
27 **daily, in three conditions: 1) “Mature water”; T. chuii was added 5 days**
28 **before the rearing of common dentex larvae, 2) “Green water”; T. chuii was**
29 **added 1 day before, and 3) “Clear water”; no T. chuii addition. The results**
30 **show the influence of the presence of T. chuii on the evolution of the**
31 **bacterial communities, both in terms of bacterial density and morphology,**
32 **and indicate “Green water” is the most suitable water treatment for**
33 **management of larval rearing for common dentex.**

34

35 Keywords

36 common dentex larvae, intensive rearing, Microalgae-bacteria interactions,
37 rearing water

38

- 39 Abbreviations
- 40 CFU: Colony Forming Units
- 41 DAPI: 4',6-diamidino-2-phenylindole
- 42 dph: days post-hatching
- 43 DW: dry weight
- 44 ind: individuals
- 45 MA: Marine Agar
- 46 TCBS: Thiosulfate Citrate Bile Salts Sucrose
- 47

48 Common dentex larvae, as with other marine fish larvae
49 (Chantanachookhin et al. 1991), develop immune organs during metamorphosis,
50 or onwards (Santamaría 2001). Nevertheless, the specific immune system in fish
51 larvae is usually not active until some time after the formation of the thalamus,
52 the thymus and the spleen (Manning and Tattner 1985; Magnadóttir et al. 2005).
53 Microorganisms present in the rearing water colonize the gut of larvae at first
54 feeding and represent the first barrier against opportunistic pathogens. Therefore,
55 the composition of the microbial community of the rearing water affects the
56 composition of the gut microbiota, although some differences are detected
57 depending on fish species (Cahill 1990). An adequate microbial control during the
58 egg stage and early development can improve larval survival, growth, and fish
59 quality (Douillet and Pickering 1999).

60 Interactions between bacteria and microalgae are complex, multiple, and
61 have a significant impact on aquaculture production. Microalgae are able to
62 produce growth promoters and inhibitors, besides several chemical clues, that
63 affect the population density and composition of their microscopic counterpart. It
64 is hypothesized that an adequate selection of a microalgae-bacteria consortium
65 would improve aquaculture productivity, efficiency and sustainability (Natrah et al
66 2014).

67 Interactions between larvae and bacteria are complex, species-specific,
68 and not fully understood (Olafsen 2001). Negative interactions, i.e. pathologies,
69 have been the subject of most studies. In the last decades, studies on positive
70 interactions, i. e. probiotics, have become more common (Gómez-Gil et al. 2000;
71 Gatesoupe 2002). Vadstein et al. (1993) and Skjermo and Vadstein (1999)

72 suggest a bulk of strategies to control the microbial community and its interactions
73 with larvae based mainly on three principles: 1) non-selective reduction of
74 bacteria, 2) selective enhancement of bacteria, and 3) improvement of larval
75 resistance. Enhancement of bacteria can also be made non-selectively by means
76 of the “mature water” technique, which relies on phytoplankton-bacteria
77 interactions (Skjermo et al. 1997).

78 Strict pathogens have not been identified and/or associated to high larval
79 mortality episodes in common dentex (Company et al. 1999; Crespo et al. 2001),
80 but well known opportunistic pathogens, mainly from the “Vibrio” genera, are
81 present naturally in marine water, and their growth can be favoured by larval
82 culturing conditions (Bergh 1996). Growth of these opportunistic bacteria
83 coincides with the development of the fish immune system (Santamaría 2001),
84 making this a period of growth for common dentex larvae where they are more
85 vulnerable. In addition, they are very sensitive to the inherent stress of artificial
86 rearing conditions (Rigos et al. 1998).

87 The aim of the following experiments was to standardize a protocol for the
88 management of rearing water and microalgae suitable for common dentex larvae.
89 The mature water technique used in these studies is not as described by Skjermo
90 et al. (1997); instead, UV-filtered seawater with microalgae was maintained in
91 tanks some days before larval stocking in order to obtain microbially mature
92 rearing water with bacteria associated to microalgae culture stocks. “Tetraselmis
93 chuii” was chosen mainly due to its suitability for nutritional experimental
94 purposes and its known antibacterial activity against some species of Vibrio and
95 “Listonella” (Riquelme and Avendaño-Herrera 2003).

96

97

Materials and Methods

98

Experiment 1

99 The data gathered in Experiment 1 (Fig. 1) was used for the design of
100 Experiment 2. Six 500 L cylindroconical, black-bottomed tanks connected to the
101 same IRTAmar recirculation unit (Carbó et al. 2001) were used to check the
102 dynamics of bacterial and microalgae communities in the water during 7 days.

103 Four 35 L cylindrical PVC containers (“baskets”) were immersed in each
104 holding tank, in order to mimic the experimental conditions of common dentex,
105 “Dentex dentex”, larvae rearing. All the baskets had the bottom and three 10x10
106 cm lateral windows covered with a 150 µm diameter mesh, and were provided
107 with an air-lift system and aeration supply. All the tanks were filled with UV-filtered
108 seawater, and no water exchange was performed during the entire test.
109 Tetraselmis chuii (57×10^3 cells/mL) was added to three of the tanks, the
110 remaining three were used as control tanks. Temperature, salinity, oxygen, and
111 pH were checked daily; nitrites and ammonia were checked on day 4. Irradiance
112 was evaluated at the beginning of the experiment, measured at the water surface
113 in the middle of the tank with a luxometer (Lutron LX-101 lux meter, Lutron
114 Electronics Enterprise Co. Ltd., Taiwan).

115 Fifty mL of water were taken periodically from the middle of the rearing
116 tanks, at 18 cm under the water surface. Water samples for T. chuii density were
117 taken once per day. Water samples for microbial analysis were taken twice per
118 day.

119 Tetraselmis chuii density was determined with a coulter/particle counter
120 (Multisizer 3, Beckman, COULTER, Miami, Florida, US). Samples for microbial

121 analysis were fixed with 25% glutaraldehyde (Sigma Aldrich, Madrid, Spain)
122 achieving 10% of the sample volume, and stored at 4°C until analysis. Five
123 hundred µL of 4',6-diamidino-2-phenylindole (DAPI; Sigma Aldrich, Madrid,
124 Spain) were added to a subsample of 5 mL. After 10 min in darkness, the sample
125 was filtered through a 0.2 µm black polycarbonate filter (Nucleopore Track-Etch,
126 Whatman, Maidstone, England); the filter was put on a slide glass and kept at -
127 20°C in darkness until analysis. Bacteria were counted using an epifluorescent
128 microscope (Leica Microsystems, GmbH, Wetzlar, Germany) where filters were
129 exposed to 365 nm light in order to induce DAPI's fluorescence (Kepner and
130 Pratt, 1994).

131 *Experiment 2*

132 Experiment 2 lasted for 16 days (Fig. 1). The effect of three types of rearing
133 water on common dentex larval survival and growth were tested using six 500 L
134 cylindroconical, black-bottomed tanks connected to the same recirculation unit
135 (Carbó et al. 2001); two tanks per type of rearing water. Each tank contained 4
136 baskets as described in Experiment 1. The types of water were: 1) "Mature
137 water"; tanks were filled with UV-filtered seawater and T. chuii was added 5 days
138 before the rearing of 0 days post-hatching (dph) common dentex larvae, 2)
139 "Green water"; tanks were filled with UV-filtered seawater and T. chuii was added
140 1 day before the rearing of 0 dph common dentex larvae, and 3) "Clear water";
141 two tanks were filled with UV-filtered seawater the same day of the rearing of 0
142 dph common dentex larvae, and no microalga was added. Once common dentex
143 larvae were stocked, routine management was followed, including rotifer addition
144 to the rearing tanks when larvae were 3 dph (experimental day 8), partial

145 recirculation of water through the recirculation units when larvae were 3 dph
146 (experimental day 8, 30% tank volume renewed daily), and full recirculation of
147 water when larvae were 7 dph (experimental day 14, 100% tank volume renewed
148 daily).

149 Temperature, salinity, oxygen, and pH were checked daily. Nitrites and
150 ammonia were checked on experimental days 8 and 14. Irradiance was evaluated
151 at the beginning of the experiment, measured at the water surface in the middle
152 of the tank, with a luxometer (Lutron LX-101 lux meter, Lutron Electronics
153 Enterprise Co. Ltd., Taiwan). Every day, 50 mL of water were taken from the
154 middle of the rearing tanks (n = 6) at 18 cm under the water surface for microbial
155 analysis.

156 A single batch of floating eggs was incubated in a basket immersed in a
157 500 L black-bottomed tank, at 19 ± 1 °C, until larvae hatched 48 h later. A sub-
158 sample of eggs was incubated in 96 well EIA plates at 19 ± 1 °C in darkness to
159 determine the batch quality, based on hatching rate and daily mortality (Giménez
160 et al. 2006a). Some freshly spawned eggs were directly plated onto Marine Agar
161 (MA) or saline Thiosulfate Citrate Bile Salts Sucrose (TCBS; Scharlau Chemie,
162 Barcelona, Spain) medium with a sterilised Pasteur pipette (Naess and Bergh
163 1994). The results were qualitative, of presence or absence of total heterotrophic
164 bacteria (growth in MA) or presumptive Vibrio spp. (growth in TCBS) on the
165 surface of the eggs.

166 Larvae at 0 dph were stocked at 40 larvae/L in each basket; larvae were
167 fed twice per day 10 individuals (ind)/mL of enriched rotifer, "Brachionus

168 rotundiformis” (2h, 20°C seawater, 250 ind/mL, 0.1 gr/L of Easy Selco, INVE,
169 Belgium) from 3 dph until the end of the experiment (larvae of 11 dph).

170 At the end of the experiment, larvae were counted in order to determine
171 larval survival. At 0, 7 and 11 dph, 20 larvae per basket were sampled to obtain
172 dry weight (DW) data, and 20 additional larvae per tank (five per basket) were
173 sampled for quantification of gut bacteria. The protocol was a modification from
174 Muroga et al. (1987) and Bergh (1996): larvae’s skin surface was sterilized with
175 0.05% iodine (Sigma Aldrich, Madrid, Spain) for 3 minutes and rinsed three times
176 with autoclaved seawater before pooling and homogenization in 10 mL of
177 autoclaved seawater, and follow with the plate count protocol.

178 Colony forming units (CFU) of total heterotrophs and presumptive Vibrio
179 spp. were determined by the plate count method: samples of water or
180 homogenates of larvae were serially diluted in autoclaved seawater and 100 µL
181 of each serial dilution were plated onto MA (for total heterotrophic bacteria), or
182 TCBS medium (for presumptive Vibrio spp.). Three plates per sample and dilution
183 were incubated for 24h at 25°C, and the number of bacterial colonies growing in
184 the plates (CFU) were counted. Counts between 30 and 300 CFU per plate were
185 used for final calculation of bacterial load in seawater (CFU/mL) or per larva
186 (CFU/ind) using the following formulae:

187
$$\text{CFU/mL} = \text{Average of plate counts} \times \text{DF}$$

188
$$\text{CFU/ind} = ((\text{Average of plate counts} \times \text{DF}) \times V) / N$$

189 Where DF is the dilution factor, V is the volume where D. dentex larvae
190 were homogenized (10 mL) and N is the number of D. dentex larvae (20).

191 Results of larvae DW, bacterial load in seawater and bacterial load in
192 larvae gut were analysed by One-way analysis of variance ($P < 0.05$) and a post
193 hoc pair-wise multiple comparison of the mean using Tukey's test ($P < 0.05$,
194 StatgraphicsPlus 4.1, Microsoft Inc).
195

196

Results and Discussion

197 The bacteria-algae interactions are as complex, species-specific, and
198 partially understood, as the larvae-bacteria interactions (Olafsen 2001, Natrah et
199 al. 2014). In addition, all variations of these algae-bacteria-larvae interactions
200 occur at the same time and are interconnected.

201 Bacteria can be introduced to the rearing tanks through four main routes: 1)
202 water, 2) microalgae, 3) eggs / larvae surfaces and 4) live prey.

203 The bacterial community in the water has been monitored during the
204 present experiments. The flux of population composition of the microbiological
205 community was observed to be dynamic. In both experiments, physico-chemical
206 conditions did not differ between experimental groups (Table 1). In Experiment 1,
207 bacteria-algae interactions were tested using a monospecific microalgal
208 community; the only difference between experimental groups was the addition of
209 microalgae at the beginning of the experiment, but this difference dramatically
210 affected the evolution of the bacterial community (Figs. 2 and 3). Tetraselmis chuii
211 densities were between 57×10^3 and 59×10^3 cells/mL during the experiment,
212 while bacterial densities in tanks with T. chuii were significantly higher than those
213 in tanks without microalgae, mainly from day 5 onwards (Fig. 2). Tetraselmis chuii
214 was added at densities used for routine rearing of common dentex larvae in IRTA
215 facilities; no fertilizer or nutrients were added to the rearing tanks, consequently,
216 T. chuii growth was not promoted. Bacteria in tanks with T. chuii can grow on the
217 organic matter produced by the microalgae, such as extracellular polymeric
218 substances (EPS) known to be excreted by microalgae (Joyce and Utting 2015;
219 Natrah et al. 2014), but bacteria in tanks without T. chuii do not have this

220 additional source of nutrients. Results from DAPI staining showed two different
221 morphologies: bacteria associated to microalgae were larger in size and usually
222 found linked in chains, while bacteria in seawater without microalgae addition
223 were smaller and appeared isolated (Fig. 3). These results suggest 5 days as the
224 minimum time required for detecting changes in the bacterial community in the
225 tanks with T. chuii, consequently, this time lapse was selected for the design of
226 “mature” water in Experiment 2.

227 Curves plotted using data from total heterotrophic CFU/mL were similar
228 between water types used in Experiment 2, although more similar between the
229 water types with T. chuii (“mature” and “green” water, Figs. 4 and 5) than between
230 those and clear water (Fig. 6): bacterial load increased shortly after larvae and
231 rotifers were added to the rearing tanks and it decreased to similar levels after
232 100% water renewal started 9 days after the larval addition (on experimental day
233 14). Presumptive Vibrio spp. CFU/mL were not detectable until one day after the
234 addition of common dentex larvae to the tanks in any of the water types,
235 regardless of the confinement time of this water; afterwards, they followed a trend
236 similar to that obtained from total heterotrophic bacteria in each water type (Figs.
237 4, 5 and 6).

238 Total heterotrophic bacteria reached the maximum in “clear water” (2.5×10^5
239 CFU/mL) when common dentex larvae were 3 dph and rotifers were added to the
240 tanks; in “mature” and “green” water, the maximum appeared when common
241 dentex larvae were 5 dph, and reached 10 times lower densities (5.5×10^4 and
242 6.9×10^5 CFU/mL, respectively). Presumptive Vibrio spp. also reached their
243 maximum when common dentex larvae were 5 dph in “mature” and “green”

244 waters (experimental day 10; 2.5×10^4 and 1.7×10^4 CFU/mL, respectively), while
245 in “clear” water there were two peaks, one when common dentex larvae were 4
246 dph (experimental day 9; 1.2×10^4 CFU/mL) and the second when they were 7
247 dph (experimental day 14; 6×10^3 CFU/mL), with lower presumptive Vibrio spp.
248 CFU/mL than in “green” and “mature” water.

249 The increase of CFU/mL of total heterotrophic bacteria and presumptive
250 Vibrio spp. shortly after rotifer addition to the tanks were expected based on
251 previous results of bacterial load in live prey (Giménez et al. 2006b). The present
252 data suggest a buffer effect of T. chuii, and/or the bacterial community present in
253 their culture, on the proliferation of total heterotrophic bacteria and presumptive
254 Vibrio spp. Common aquatic opportunistic pathogens can be detected, such as
255 presumptive Vibrio spp., and are the target of disinfection protocols for finfish
256 larvae (Giménez et al. 2009) and live prey (Giménez et al. 2006b). A
257 management based on the use of microalgae, combined with water exchange,
258 can be more effective than disinfecting the tanks when larvae are already
259 stocked, mainly in the case of common dentex larvae, which are very sensitive to
260 disinfectants (Giménez et al. 2009) and stress (Rigos et al. 1998).

261 There was no bacterial growth on any of the eggs directly plated onto MA
262 and TCBS; therefore, the surface of the eggs was considered virtually free of
263 bacteria and their contribution to the increase of bacterial load in the rearing water
264 is considered very low. Can et al. (2012) report the presence of total heterotrophic
265 bacteria and presumptive Vibrio spp. in common dentex eggs kept in tanks for
266 72h. It is possible that, despite the lack of detection of bacterial growth in the
267 present study, once the larvae have hatched, the high concentration of nutrients

268 from the eggs promote the increase of fast growing bacteria such as presumptive
269 Vibrio spp. It could explain the increase in bacterial load in the rearing water when
270 larvae were added, especially presumptive Vibrio spp. Common dentex eggs do
271 not survive washing several times on a mesh, nor the standard iodine disinfection
272 (unpublished data) described for "Sparus aurata", "Dicentrarchus labrax" (Moretti
273 et al. 1999) or "Hippoglossus hippoglossus" eggs (Bergh and Jelmert 1996), and
274 are more sensitive to ozone treatment than Sparus aurata, Dicentrarchus labrax
275 and "Pagrus pagrus" (Can et al. 2012), probably due to their thinner chorion
276 (Iconomidou et al. 2000). Consequently, disinfection of common dentex eggs
277 does not seem necessary, based on the results of bacterial load and their
278 sensitivity to disinfection procedures.

279 The study of larvae-bacteria interactions was affected by the low quality of
280 the batch of eggs used in the experiment, which possibly affected negatively the
281 results in larval survival (Table 2). Mortality of 5 dph starved larvae kept in EIA
282 plates was high (89.1%), compared to the published average mortality rate of
283 29.2% and minimum mortality rate of 4.2% obtained during a spawning season
284 from the same broodstock (Giménez et al. 2006a). There were no significant
285 differences in larval survival, i.e. larvae were at similar stocking densities. They
286 also shared the same genetic background, and were fed the same live prey.
287 Larvae reared in "green" water were significantly bigger ($38.8 \pm 11 \mu\text{g}$, $P < 0.05$)
288 than larvae reared in "mature" and "clear" water (Table 2). These data suggest
289 "green" water as the most suitable technique for experimental larval rearing of
290 common dentex, although the experiment ended when larvae were 11 dph,
291 before the formation of the immunological organs was completed (Santamaría

292 2001). The beneficial effects of controlling the microbial community in the rearing
293 water during early larval rearing will probably be more evident later in
294 development, during or after metamorphosis.

295 There can be a relationship between the trends of total heterotrophic CFU
296 per larvae and in the rearing water because the larvae gut is colonized by the
297 bacteria present in the rearing water during their early development (Bergh 1996).
298 There was no bacterial colonization in the gut of 0 dph common dentex larvae
299 (Table 2)utilizing the protocol for determining gut bacteria, adapted from Muroga
300 et al. (1987) and Bergh (1996) . These results are in agreement with common
301 dentex larval development, because at 0 dph their gut is closed, neither the mouth
302 nor the anus are opened until 3 dph (Santamaría 2001). Between 7 and 11 dph,
303 a reduction of total heterotrophic CFU/ind was observed in larvae reared in
304 “green” water. Opposite results were observed in gut microbiota of larvae reared
305 in “clear” and “mature” water. When larvae were 7 dph (experimental day 12)
306 there was a higher density of total heterotrophic CFU/mL in “green” water than in
307 “mature” and “clear” water (Figs. 4, 5 and 6), when larvae were 11 dph
308 (experimental day 16) the concentrations of total heterotrophic CFU/mL were
309 similar in the three types of rearing water, lower than at experimental day 12 for
310 “green” water and higher for “mature” and “clear” water. Presumptive Vibrio spp.
311 CFU/ind were similar at 11 dph among all rearing water types, although at 7 dph
312 larvae reared in “mature” water had more presumptive Vibrio spp. CFU/ind than
313 larvae reared in “green” or “clear” water. The hypothesis of gut colonization,
314 suitable for results on total heterotrophic bacteria, is not suitable for presumptive

315 Vibrio spp. because there is no relationship with the results of presumptive Vibrio
316 spp. in the rearing tanks (Figs. 4, 5 and 6)

317 Total heterotrophic CFU/ind and presumptive Vibrio spp. CFU/ind,
318 obtained in the present experiment, are within the range found by Muroga et al.
319 (1987) for Pagrus major and "Acanthopagrus schlegeli" at similar degree days
320 but much lower than those found in "Scophthalmus maximus" (Nicolas et al. 1989;
321 Salvesen et al. 1999; Makridis et al. 2000) and in "Paralichthys dentatus" (Eddy,
322 and Jones 2002). Comparisons with published results (Table 3) must take into
323 account that studies on the effects of microbiological environment on larval
324 performance have been mainly carried out with cold water species. These
325 species are cultured at a lower water temperature and undergo a longer larval
326 development than common dentex larvae.

327

328

Conclusions

329 Addition of microalgae to the rearing tanks, and the duration of their
330 confinement before the onset of larvae rearing, affect the composition and
331 dynamics of the microbial communities in the rearing water. The addition of larvae
332 and rotifers resulted in an increase in the microbial load of the rearing water, both
333 total heterotrophs and presumptive Vibrio spp., but the presence of T. chuii
334 seems to buffer the bacteria proliferation.

335 Disinfection of eggs and live prey before their addition to the rearing tanks
336 might be good practices for the reduction of bacterial load in the rearing water,
337 but egg disinfection is not suitable for common dentex given their sensitivity to
338 disinfectants, and not relevant taking into account the low bacterial load on their
339 egg surface .

340 Results on larval growth suggest “green” water as the most suitable
341 technique for the experimental design and for rearing of common dentex.

342

Acknowledgments

343 Funding provided by the Spanish Instituto Nacional de Investigación Agraria y
344 Alimentaria (INIA) for GGP fellowship, and by a project from Junta Nacional
345 Asesora de Cultivos Marinos (JACUMAR). We wish to thank M. Monllaó and J.
346 Canoura (Centre d'Aqüicultura IRTA) for their help and advise in larval rearing,
347 and K. B. Andree for final review of the manuscript.

348

349

Literature Cited

- 350
- 351 **Bergh, O. and A. Jelmert.** 1996. Iodophor disinfection of eggs of Atlantic Halibut.
352 Journal of Aquatic Animal Health 8: 135 – 145.
- 353 **Bergh, O. 1996.** Ecological relations between bacteria and early life stages of
354 fish, with special emphasis on the Atlantic halibut, Hippoglossus
355 hippoglossus in aquaculture. PhD Thesis. Department of Microbiology,
356 University of Norway, Bergen (Norway). 100 pages.
- 357 **Cahill, M. M. 1990.** Bacterial flora of fishes: a review. Microbial Ecology 19: 21 –
358 41.
- 359 **Can, E., Karacalar, U., Saka, S., Firat, K.** 2012. Ozone disinfection of eggs from
360 gilthead seabream Sparus aurata, Sea bass Dicentrarchus labrax, red
361 porgy, and common dentex Dentex dentex. Journal of Aquatic Animal
362 Health 24 (2): 129 – 133. DOI: 10.1080/08997659.2012.675925
- 363 **Carbó, R., Estévez, A., Furones, M. D.** 2001. Intelligent and multifunctional
364 recirculation system. Its application in research at CA – IRTA. EAS Special
365 Publication, 32: 171 – 172.
- 366 **Chantanachookhin, C., Seikai, T., Tanaka, M.** 1991. Comparative study of the
367 ontogeny of the lymphoid organs in three species of marine fish.
368 Aquaculture 99: 143 – 155.
- 369 **Company, R., Sitjà-Bobadilla, A., Pujalte, M. J., Garay, E., Alvarez-Pellitero,**
370 **P., Pérez-Sánchez, J.** 1999. Bacterial and parasitic pathogens in cultured
371 common dentex, Dentex dentex L. Journal of Fish Diseases 22: 299 – 309.

- 372 **Crespo, S., Martín de Mateo, M., Santamaría, C.A., Sala, R., Grau, A., Pastor,**
373 **E.** 2001. Histopathological observations during larval rearing of common
374 dentex Dentex dentex L. (Sparidae). Aquaculture, 192: 121 – 132.
- 375 **Douillet, P. A. and P. L. Pickering.** 1999. Seawater treatment for larval culture
376 of the fish Sciaenops ocellatus Linnaeus (red drum). Aquaculture 170: 113
377 – 126.
- 378 **Eddy, S. D. and S. H. Jones.** 2002. Microbiology of summer flounder
379 Paralichthys dentatus fingerling production at a marine fish hatchery.
380 Aquaculture 211: 9 – 28.
- 381 **Gatesoupe, F. J.** 2002. Probiotic and formaldehyde treatments of Artemia nauplii
382 as food for larval pollack, Pollachius pollachius. Aquaculture 212: 347 –
383 360.
- 384 **Giménez, G., Estévez, A., Lahnsteiner, F., Zecevic, B., Bell, J. G.,**
385 **Henderson, R. j., Piñera, J. A., Sánchez-Prado, J. A.** 2006a. Egg quality
386 criteria in common dentex (Dentex dentex). Aquaculture 260: 232 – 243.
- 387 **Giménez, G., Padrós, F., Roque, A., Estévez, A., Furones, D.** 2006b. Bacterial
388 load reduction of live prey for fish larval feeding using Ox-Aquaculture.
389 Aquaculture Research 37: 1130 – 1139.
- 390 **Giménez-Papiol, G., Padrós, F., Roque, A., Estévez, A., Furones, D.** 2009.
391 Effects of a peroxide-based comercial product on bacterial load of larval
392 rearing water and on larval survival of two species of Sparidae under
393 intensive culture: preliminary study. Aquaculture Research 40: 504 – 508.

- 394 **Gómez-Gil, B., Roque, A., Turnbull, J. F.** 2000. The use and selection of
395 probiotic bacteria for use in the culture of larval aquatic organisms.
396 *Aquaculture* 191: 259 – 270.
- 397 **Iconomidou, V. A., Chryssikos, D. G., Gionis, V. Pavlidis, M. A., Paipetis, A,**
398 **Hamodrakas, S. J.** 2000. Secondary structure of chorion proteins of the
399 teleostean fish Dentex dentex by ATR FT-IR and FT-Raman spectroscopy.
400 *Journal of Structural Biology*, 132: 112 – 122.
- 401 **Joyce, A. and S. Utting.** 2015. The role of exopolymers in hatcheries: an
402 overlooked factor in hatchery hygiene and feed quality. *Aquaculture* 446:
403 122 – 131.
- 404 **Kepner, R. L and J. R. Pratt.** 1994. Use of fluorochromes for direct enumeration
405 of total bacteria in environmental samples: past and present.
406 *Microbiological Reviews* 58 (4): 603 – 615.
- 407 **Magnadottir, B., Lange, S., Gudmundsdottir, S., Bøgwald, J, Dalmo, R. A.**
408 2005. Ontogeny of humoral immune parameters in fish. *Fish and Shellfish*
409 *Immunology* 19: 429 – 439.
- 410 **Makridis, P., Fjellheim, A. J., Skjermo, J., Vadstein, O.** 2000. Colonization of
411 the gut in first feeding turbot by bacterial strains added to the water or
412 bioencapsulated in rotifers. *Aquaculture International* 8: 367 – 380.
- 413 **Manning, M. J. and M. F. Tattner.** 1985. *Fish Immunology*. Academic Press Inc.,
414 London, 374 pages.

- 415 **Moretti, A., Pedini, M., Cittolin, G. Guidastrì, R.** 1999. Manual on Hatchery
416 Production of Seabass and Gilthead Seabream. Food and Agriculture
417 Organization of the United Nations, Vol. 1.
- 418 **Muroga, K., Higashi, M., Keitoku, H.** 1987. The isolation of intestinal microflora
419 of farmed red seabream (Pagrus major) and black seabream
420 (Acanthopagrus schlegeli) at larval and juvenile stages. Aquaculture 65:
421 79 – 88.
- 422 **Naess, T. and O. Bergh.** 1994. Calanoid copepod resting eggs can be surface
423 disinfected. Aquacultural Engineering 13: 1 – 9.
- 424 **Natrah, F. M., Bossier, P.; Sorgeloos, P., Yusoff, F. Md., Defoirdt, T.** 2014.
425 Significance of microalgal-bacterial interactions for aquaculture. Reviews
426 in Aquaculture 6: 48 – 61.
- 427 **Nicolas, J. L., Robic, E., Ansquer, D.** 1989. Bacterial flora associated with a
428 trophic chain consisting of microalgae, rotifers and turbot larvae: influence
429 of bacteria on larval survival. Aquaculture 83: 237 – 248.
- 430 **Olafsen, J. A.** 2001. Interactions between fish larvae and bacteria in marine
431 aquaculture. Aquaculture 200: 223 – 247.
- 432 **Rigos, G., Grigorakis, K., Nengas, I. Christophilogiannis, M., Yiadnisi, M.,**
433 **Kout-Somidou, A., Andriopoulou, A, Alexis, M.** 1998. Stress-related
434 pathology seems a significant obstacle for the intensive farming of
435 common dentex, Dentex dentex (Linnaeus, 1758). Bulletin of the
436 European Association of Fish Pathology 18: 15 – 18.

- 437 **Riquelme, C. E., Avendaño-Herrera, R. E.** 2003. Interacción bacteria-microalga
438 en el ambiente marino y uso potencial en acuicultura. Revista Chilena de
439 Historia Natural 76: 725 – 736.
- 440 **Salvesen, I., Skjermo, J., Vadstein, O.** 1999. Growth of turbot (Scophthalmus
441 maximus L.) during first feeding in relation to the proportion of r/K-
442 strategists in the bacterial community of the rearing water. Aquaculture
443 175: 337 – 350.
- 444 **Santamaría, C. A.** 2001. Desarrollo de la larva de dentón, Dentex dentex
445 (Linnaeus, 1758): Estudio cuantitativo del crecimiento, aspectos
446 histológicos y organogénesis. PhD thesis. Departament de Biologia
447 Animal, de Biologia Vegetal i d'Ecologia. Universitat Autònoma de
448 Barcelona. Bellaterra, Barcelona, 261 pages.
- 449 **Skjermo, J., Salvesen, I., Oie, G. Olsen, Y., Vadstein, O.** 1997. Microbially
450 matured water: a technique for selection of a non-opportunistic bacterial
451 flora in water that may improve performance of marine larvae. Aquaculture
452 International 5: 13 – 28.
- 453 **Skjermo, J., Vadstein, O.** 1999. Techniques for microbial control in the intensive
454 rearing of marine larvae. Aquaculture 177: 333 – 343.
- 455 **Vadstein, O., Oie, G., Olsen, Y. Salvesen, I., Skjermo, J. Skjak-Braek, G.**
456 1993. A strategy to obtain microbial control during larval development of
457 marine fish. In: Reinertsen, Dahle, Jørgensen, Tvinnereim (Eds.). Fish
458 Farming Technology. Balkema, Rotterdam: 69 – 75.

459 **Verner-Jeffreys, D. W., Shields, R. J., Bricknell, I. R., Birkbeck, T. H. 2003.**
460 Changes in the gut-associated microflora during the development of
461 Atlantic halibut (Hippoglossus hippoglossus L.) larvae in three British
462 hatcheries. *Aquaculture* 219: 21 – 42.
463

464

Table legends

465 Table 1. Rearing conditions maintained in Experiment 1 and Experiment 2.

466 Table 2. Egg quality data, survival, DW, and bacterial load results of larvae reared
467 in the different rearing water types of Experiment 2. a, b and ab superscripts
468 denote significant differences ($P < 0.05$). *counts lower than 30 CFU per plate.

469 Table 3. Summary of bibliographical data of gut-associated bacterial flora hosted
470 by fish larvae of different species. U = unspecified bacteria; H = total
471 heterotrophic bacteria; V = presumptive Vibrio spp.

472

473

Figure legends

474 Figure 1. Set up of Experiments 1 and 2. Black and white bars are the time scale
475 of experiment, in days. Colour of the cells indicate the status of tanks: white =
476 empty tank; dark grey = tank with water, no water exchange; grey = tank with
477 water, 30% water exchange per day; light grey = tank with water, 100% water
478 exchange per day. Samplings: * = microbiological sample; T = T. chuii sample; X
479 = checking of temperature, salinity, oxygen, and pH; L = D. dentex larvae sample;
480 N = checking of nitrites and ammonia.

481 Figure 2. Bacteria density (cells/mL) obtained in Experiment 1, counted at the
482 microscope using DAPI staining. a and b superscripts denote significant
483 differences ($P < 0.05$) between types of rearing water at the same sampling time.

484 Figure 3. Images showing bacterial communities in Experiment 1 at the beginning
485 and the end of the experiment in each type of rearing water. Bacteria are stained
486 using DAPI (cocci and rods). Images A and B: bacteria communities in the tanks
487 with (A) and without (B) T. chuii at the beginning of the experiment, time 0h.
488 Images C and D: bacteria communities in the tanks with (C) and without (D) T.
489 chuii at the end of the experiment, time 146h.

490 Figure 4. Evolution of Heterotrophic bacteria and presumptive Vibrio spp.
491 densities (CFU/mL) in “mature water” from Experiment 2.

492 Figure 5. Evolution of Heterotrophic bacteria and presumptive Vibrio spp.
493 densities (CFU/mL) in “green water” from Experiment 2.

494 Figure 6. Evolution of Heterotrophic bacteria and presumptive Vibrio spp.
495 densities (CFU/mL) in “clear water” from Experiment 2.

496 TABLE 1.

497

	Experiment 1		Experiment 2		
	Control	With <u>T. chuii</u>	Clear water	Green water	Mature water
Initial <u>T. chuii</u> density (cells/mL)	0	57,425 ± 454	0	55,508 ± 378	56,251 ± 851
Temperature (°C, mean ± SD)	17.7 ± 0.5	17.8 ± 0.6	18.8 ± 0.5	18.5 ± 0.0	18.3 ± 0.1
Salinity (g/L, mean ± SD)	34.3 ± 0.1	34.2 ± 0.1	34.8 ± 0.2	34.9 ± 0.1	35.0 ± 0.1
Oxygen (mg/L, mean ± SD)	8.7 ± 0.1	8.8 ± 0.2	9.1 ± 0.4	9.0 ± 0.3	9.5 ± 0.4
pH (mean ± SD)	8.3 ± 0.1	8.3 ± 0.1	7.9 ± 0.3	8.2 ± 0.0	8.2 ± 0.0
Nitrites (mg/L, maximum value)	0.02	0.04	0.01	0.02	0.04
Ammonia (mg/L, maximum value)	0	0.06	0	0.02	0.06
Photoperiod (hours light:hours darkness)				16:8	
Irradiance (μmol/m ² s)				3.7	

498

499 TABLE 2.

	Clear water	Green water	Mature water
Hatching rate (EIA plates results, %)		94.8	
Mortality of starved 5 dph larvae (EIA plate results, %)		89.1	
Survival at 11 dph (% , mean \pm SD)	2.8 \pm 1.7	2.9 \pm 1.5	1.7 \pm 0.6
Initial DW (μ g, mean \pm SD)		29.2 \pm 1.5	
7 dph DW (μ g, mean \pm SD)	19.3 \pm 1.6 ^b	22.2 \pm 5 ^a	21.8 \pm 4.1 ^a
11 dph DW (μ g, mean \pm SD)	31.1 \pm 10.5 ^b	38.8 \pm 11 ^a	33.1 \pm 7.3 ^{ab}
Total heterotrophic CFU/ind (mean \pm SD) at 0 dph	0	0	0
Total heterotrophic CFU/ind (mean \pm SD) at 7 dph	77 \pm 8	301 \pm 134	250 \pm 143
Total heterotrophic CFU/ind (mean \pm SD) at 11 dph	563 \pm 105	132 \pm 10	634 \pm 133
Presumptive <i>Vibrio</i> spp. CFU/ind (mean \pm SD) at 0 dph	0	0	0
Presumptive <i>Vibrio</i> spp. CFU/ind (mean \pm SD) at 6 dph	*	*	167 \pm 60
Presumptive <i>Vibrio</i> spp. CFU/ind (mean \pm SD) at 11 dph	42 \pm 22	30 \pm 15	42 \pm 21

500

501

502 TABLE 3.

Species	Larval age (dph)	Water temperature (°C)	Day degrees	Bacterial density (CFU/ind)	Reference
<u>Pagrus major</u>	4	18.4	92	55 (U)	Muroga et al. 1987
	18	20.6	380.4	16000 (U)	
<u>Acanthopagrus schlegeli</u>	4	19.2	96	1.3 (U)	Muroga et al. 1987
	18	20.0	456	4600 (U)	
<u>Scophthalmus maximus</u>	4	not available	not available	12000 (H)	Nicolas et al. 1989
				820 (V)	
				18000 (H)	
	6			1400 (V)	
<u>Scophthalmus maximus</u>	1	15	29	2500 – 21000 (U)	Salvesen et al. 1999
	5	18	98	1900 – 31000 (U)	
	12		224	53200 - 868000 (U)	
<u>Scophthalmus maximus</u>	1	16	30	74000 – 34000 (U)	Makridis et al. 2000
	2	18	48	*Bioencapsulation	
	3		66		
	6		120		
	9		174		
	12		228		
<u>Paralichthys dentatus</u>	1 – 90	not available	-	10 ³ - 10 ⁴ (U)	Eddy and Jones 2002
<u>Hippoglossus hippoglossus</u>	5	not available	-	<5 (U)	Verner-Jeffreys et al. 2003
<u>Dentex dentex</u>	0	19	19	0 (H)	Present experiments
				0 (V)	

6

133

77 – 301 (H)
167 (V)

11


228

132 – 634 (H)
30 – 42 (V)


503

504 FIGURE 1.

EXPERIMENT 1



EXPERIMENT 2




505

506

507

508 FIGURE 2.




509

510 FIGURE 3.


511

512

513



514 FIGURE 4.




515

516

517 FIGURE 5.


●— Heterotrophic bacteria
○— Presumptive *Vibrio* spp.



518

519

520 FIGURE 6.



521