

CORRECTION

Correction to: Trend of Changes in Serum Albumin and Its Relation with Sex, Age, and BMI Following Laparoscopic Mini-gastric Bypass Surgery in Morbid Obese Cases

Mehrdad Karimi^{1,2} · Ali Kabir² · Masoumeh Nejatifar³ · Abdolreza Pazouki^{2,4,5}

Published online: 13 November 2017

© Springer Science+Business Media, LLC, part of Springer Nature 2017

Correction to: OBES SURG<https://doi.org/10.1007/s11695-017-2912-2>

Due to a production error the bottom portion of Fig. 3 was missing in the version of this article originally published online. The corrected version of the article has now been published. The corrected version of the figure follows:

The online version of the original article can be found at <https://doi.org/10.1007/s11695-017-2912-2>

✉ Ali Kabir
aikabir@yahoo.com

Mehrdad Karimi
karimi.mehrdad89@yahoo.com

Masoumeh Nejatifar
mejatifar@gmail.com

Abdolreza Pazouki
apazouki@yahoo.com


¹ Department of Surgery, Shahrekord University of Medical Sciences, Sharekord, Iran

² Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Room 255, Central Building, Hemmat Expway, Tehran, Iran

³ Faculty of Health, ShahidBeheshti University of Medical Sciences, Tehran, Iran

⁴ Center of Excellence for Minimally Invasive Surgery Training, Iran University of Medical Sciences, Tehran, Iran

⁵ Center of Excellence of European Branch of International Federation for Surgery of Obesity, Tehran, Iran



Age (year)	18-39			≥ 40		
Baseline serum albumin (g/dl)	3.50-3.90	4.00-4.90	≥ 5.00	3.50-3.90	4.00-4.90	≥ 5.00

Groups of Age-Baseline serum albumin level