

ORIGINAL ARTICLE

Prevalence of high-level gentamicin-resistant *Enterococcus faecalis* and *Enterococcus faecium* in an Iranian hospital

M. EMANEINI^{1,2}, B. KHORAMIAN³, F. JABALAMELI², R. BEIGVERDI², K. ASADOLLAHI⁴, M. TAHERIKALANI⁵, A.R. LARI¹
¹ Antimicrobial Resistance Research Center, Iran University of Medical Sciences, Tehran, Iran; ² Department of Microbiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran; ³ Department of Clinical Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran; ⁴ Department of Social Medicine, Faculty of Medicine, Ilam University of Medical Sciences, Ilam, Iran; ⁵ Razi Herbal Medicines Researches Center, Department of Microbiology, School of Medicine, Lorestan University of Medical Sciences, Khoramabad, Iran

Keywords

Enterococcus • HLGR • RAPD-PCR

Summary

This study was designed to determine the molecular characteristics and antimicrobial resistance of enterococcal strains isolated from patients admitted to an Iranian Hospital. Enterococcal strains were isolated from the burn patients. All strains were screened for genes encoding resistance to aminoglycoside [*aac(6')-Ie-aph(2'')*-Ia, *aph(3')*, *ant(4')*], resistance to vancomycin (*vanA*, *vanB*), resistance to tetracycline (*tetK*, *tetL*, *tetM*, *tetO*), and resistance to erythromycin (*ermA*, *ermB*, *ermC*) by PCR and multiplex PCR-based methods. Genetic diversity was evaluated via Random Amplified Polymorphic DNA (RAPD)-PCR. All enterococcal isolates showed complete sensitivity to vancomycin with MIC $\leq 0.5 \mu\text{g/ml}$. Resistance to gentamicin, tetracycline, erythromycin, ciprofloxacin or quinopristin-dalfopristin was detected, whilst more than 96.2% of isolates were high-level gentamicin-

resistant (HLGR) and multiple drug resistant. The most prevalent aminoglycoside resistance gene was *aac(6')-Ie-aph(2'')*-Ia, that was found in 96.2% (26/27) of the isolates. The most prevalent tetracycline resistance genes were *tetM*, found in 85.1% (23/27) followed by *tetL* and *tetO* found in 7.4% (2/27) of the isolates. The *ermA* and *ermB* genes were detected in 33.3% (9/27) and 44.4% (12/27) of the isolates respectively. RAPD-PCR analysis yielded 17 distinct profiles among 27 investigated isolates. One cluster of isolates shared the same RAPD pattern, while 16 isolates had unique RAPD pattern. Our study showed that during the examination time period one RAPD genotype was the common type and was disseminated among patients in the burn unit. Interestingly, most of these strains had an identical or very similar antibiotic and gene resistance pattern.

Introduction

Enterococci are Gram-positive, facultative anaerobic bacteria that are members of the normal flora in the human gastrointestinal tract but have been recognized as important pathogens worldwide [1, 2]. *Enterococcus faecalis* and *Enterococcus faecium* are the most prevalent species isolated from the urinary tract and wound infections, which affect mainly patients admitted to the medical care centers [3-6]. The disruption of the normal skin barrier, the immunocompromised state, prolonged hospitalization and antibiotic therapy put burns patients at a high risk of acquiring nosocomial infections including enterococcal infections [7]. Enterococcal infections can be difficult to treat because they have a remarkable ability of adaptation when are exposed to antibiotics; they have intrinsic resistance to several antimicrobial agents and have a tremendous capacity to acquire high levels of resistance to antibiotics [8, 9]. Vancomycin-resistant enterococci (VRE) and high-level gentamicin-resistant (HLGR) isolates have emerged as important pathogen in Iran as well as in other countries,

which create serious challenges for treating the infected patients [9-11].

There are several reports on the endemicity of VRE and HLRG in Iran but there is a lack of information on enterococcal stains isolated from burn centers [3, 12]. This study was designed to determine the molecular characteristics and antimicrobial resistance of enterococcal strains isolated from patients in a burn center in Iran.

Materials and methods

BACTERIAL ISOLATES

Twenty-seven enterococcal isolates were collected from wound specimens of patients with burn injury, during April to September 2012, in a burn hospital in Tehran. Only one isolate per patient was included in the study. Identification of enterococci was performed based on a series of conventional microbiological tests including Gram reaction, catalase reaction, presence of pyrrolidonyl arylamidase (PYR), growth on bile-aesculin agar and 6.5% NaCl media, motility test, arginine decarboxylation in Moeller decarboxylase

media, pyruvate utilization, and fermentation of arabinose, raffinose, mannitol and ribose [3]. To confirm the identity of the isolates as *E. faecalis* or *E. faecium* the *ddl_E* gene was amplified by a polymerase chain reaction (PCR) method as described by Dutka-Malen et al. [13].

ANTIMICROBIAL SUSCEPTIBILITY TESTING

Antibiotic susceptibility of the strains was performed by the disk diffusion method. The antibiotics (Mast Group Ltd., Merseyside, UK) tested were ciprofloxacin (5 µg), erythromycin (15 µg), tetracycline (30 µg), and quinupristin-dalfopristin (15 µg). High-level resistance to gentamicin was also determined by disk diffusion method on Mueller-Hinton agar (Conda S.A., Madrid, Spain) using 120 µg gentamicin disk. The minimal inhibitory concentration (MIC) of vancomycin was determined by the standard agar dilution test on *Brain Heart Infusion* agar (Conda S.A., Madrid, Spain). All Antibiotic susceptibilities were performed and interpreted according to the criteria of the Clinical and Laboratory Standards Institute (CLSI) guidelines [14].

DNA EXTRACTION AND GENE DETECTION

DNA extraction was performed as described previously and this DNA was used as a template for PCR analysis [11]. All strains were screened for genes encoding resistance to aminoglycoside [*aac(6')-Ie-aph(2'')-Ia*, *aph(3')*, *ant(4')*], vancomycin (*vanA*, *vanB*), tetracycline (*tetK*, *tetL*, *tetM*, *tetO*), and resistance to erythromycin (*ermA*, *ermB*, *ermC*) by PCR and multiplex PCR-based methods, using specific primers [3, 11,12, 15, 16].

RANDOM AMPLIFIED POLYMORPHIC DNA (RAPD)-PCR

The RAPD-PCR assay was carried out in 25 µl reaction volumes containing 0.5 µM of each primer (5'-AGCGGGCCAA-3' and 5'-ACGGCCGACC-3'), 12.5 µl of PCR master (Sinaclone Inc, Iran) and 5 µl of DNA template. Cycling conditions were as follow: initial denaturation at 95°C for 5 min, followed by 30 cycles of 95°C for 1 min, 38.5°C for 1 min, and 72°C for 2 min, and final extension at 72°C for 10 min. PCR products were then separated by electrophoresis in 1.4% agarose gel with 0.5X TBE buffer. DNA bands were observed by staining with KBC power load dye (Kawsar Biotech Co. Iran) and photographed under UV illumination. RAPD patterns were analyzed by visual inspection. Only the obvious, prominent and reproducible bands from repeated experiments (at least twice) were considered and any pattern differing by one or more bands was classified as a distinct RAPD type.

Results

Distribution of species according to conventional microbiological tests and PCR method included 19 (70.3%) *E. faecalis* and 8 (29.7%) *E. faecium*. The prevalence of antimicrobial resistance pattern and RAPD types have been summarized in Table I. In the present study, no VRE was

recovered and all enterococcal isolates showed complete sensitivity to vancomycin with MIC ≤ 0.5 µg/ml.

All isolates at least exhibited resistance to one antibiotic and 6 different profiles were observed on the basis of their antibiotic resistance patterns. High-level resistance to gentamicin and multiple drug resistance were observed in 96.2% of isolates. The most frequent profile was resistance to erythromycin, gentamicin, tetracycline, quinupristin-dalfopristin and ciprofloxacin that was observed in 74% of isolates.

The most prevalent aminoglycoside resistance gene was *aac(6')-Ie-aph(2'')-Ia*, found in 96.2% (26/27) of the isolates. These genes were detected in all HLGR isolates. The *ant(4')* gene was detected in 62.9% (17/27) of isolates. The most prevalent tetracycline resistance gene was *tetM*, found in 85.1% (23/27) of the isolates followed by *tetL* and *tetO* found in 7.4% (2/27) of the isolates. The *ermA* and *ermB* genes were detected in 33.3% (9/27) and 44.4% (12/27) of the isolates respectively. The *aph(3')*, *tetK*, *ermC*, *vanA*, and *vanB* genes were not detected in any of the enterococcal isolates in this study.

To determine the degree of clonality among enterococcal isolates, the RAPD typing was used. RAPD analysis yielded 17 distinct profiles among 27 investigated isolates (Fig. 1). One clusters of isolates shared the same RAPD patterns, while 16 isolates had unique RAPD patterns. A dominant RAPD type designated as A, that consisted 11 isolates of *E. faecalis* and were HLGR.

Discussion

Among Enterococci isolates, *E. faecalis* and *E. faecium* are the most common species caused enterococcal nosocomial infections [1]. In our study, the most prevalent species was *E. faecalis* (70.3%) followed by *E. faecium* (29.7%) isolated from burn wound infections. Similar results were reported for clinical isolates by other studies [3, 11, 16, 17].

In the treatment of enterococcal infections, the combination of gentamicin with a beta lactam antibiotic or a glycopeptide is used to obtain a synergistic bactericidal effect. However, strains that are highly resistant to gentamicin are no longer susceptible to the combination therapy [18]. Several studies have demonstrated that VRE have caused both outbreaks and endemic infections in burn units [5, 6]. However, in this study, there were no VRE isolates and more than 96% of isolates were HLGR and 74% of them were simultaneously resistant to erythromycin, tetracycline, quinupristin-dalfopristin and ciprofloxacin. Our study is the first investigation on enterococcal resistance among burn patients in Iran and the findings indicated that vancomycin keeps its therapeutic effects against enterococcal infections.

Resistance against aminoglycosides among enterococci is commonly due to enzymatic modification. Genes encoding aminoglycoside-modifying enzymes (AMEs) are often carried on transposable elements. One of the most prevalent AMEs genes among enterococci is *aac(6')-Ie-aph(2'')-Ia* encoding a bifunctional enzyme *Aac(6')-Ie-Aph(2'')*-Ia which confers resistance to virtually all of the clinically available aminoglycosides including gentamicin [11, 19].

Tab. I. The distribution of antimicrobial resistance genes, virulence genes and biofilm production among enterococcal species.

Isolate	Date of Isolation	Species	Resistance pattern	Resistance genes	RAPD type
1	20/4/2012	<i>E. faecalis</i>	E*,GM, TET,SYN,CIP	<i>tetM,ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
2	22/4/2012	<i>E. faecium</i>	GM, TET, SYN	<i>tetM, aac(6')-le-aph(2'')-la, ant(4')</i>	B
3	24/4/2012	<i>E. faecium</i>	E,GM, TET,SYN,CIP	<i>tetM, aac(6')-le-aph(2'')-la</i>	H
4	26/4/2012	<i>E. faecium</i>	E,GM, TET	<i>tetL, tetM, ermB, aac(6')-le-aph(2'')-la, ant(4')</i>	G
5	26/4/2012	<i>E. faecalis</i>	E,GM, TET	<i>tetM,ermB, aac(6')-le-aph(2'')-la</i>	P
6	1/5/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM, ermB, aac(6')-le-aph(2'')-la</i>	A
7	4/5/2012	<i>E. faecium</i>	E,GM, TET	<i>tetL, tetM, ermB, aac(6')-le-aph(2'')- la, ant(4')</i>	J
8	8/5/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>ermB, aac(6')-le-aph(2'')-la</i>	A
9	15/5/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM, aac(6')-le-aph(2'')-la</i>	N
10	2/6/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermB, aac(6')-le-aph(2'')-la</i>	A
11	12/7/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
12	14/7/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
13	14/7/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM, ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
14	21/7/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM, ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
15	23/7/2012	<i>E. faecium</i>	TET	<i>tetM</i>	M
16	25/7/2012	<i>E. faecium</i>	E,GM, TET,SYN,CIP	<i>tetM,ermB, aac(6')-le-aph(2'')-la</i>	F
17	7/8/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM, ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
18	7/8/2012	<i>E. faecalis</i>	GM,SYN,CIP	<i>tetM,ermB, aac(6')-le-aph(2'')-la</i>	E
19	14/8/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	D
20	14/8/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermB, aac(6')-le-aph(2'')-la, ant(4')</i>	I
21	21/8/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermA, aac(6')-le-aph(2'')-la, ant(4')</i>	A
22	25/8/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,ermA aac(6')-le-aph(2'')-la, ant(4')</i>	A
23	11/9/2012	<i>E. faecalis</i>	E,GM, TET,SYN,CIP	<i>tetM,tetO, ermB, aac(6')-le-aph(2'')-la, ant(4')</i>	K
24	11/9/2012	<i>E. faecium</i>	E, GM	<i>aac(6')-le-aph(2'')-la, ant(4')</i>	Q
25	16/9/2012	<i>E. faecalis</i>	E, GM,TET,SYN,CIP	<i>tetM, ermB, aac(6')-le-aph(2'')-la</i>	C
26	16/9/2012	<i>E. faecalis</i>	E, GM,TET,SYN,CIP	<i>tetM aac(6')-le-aph(2'')-la, ant(4')</i>	L
27	16/9/2012	<i>E. faecium</i>	E, GM,TET,SYN,CIP	<i>tetM,tetO, ermB, aac(6')-le-aph(2'')-la, ant(4')</i>	O

* E, Erythromycin; G, Gentamicin; SYN, Quinupristin-dalfopristin; TET, Tetracycline; CIP, Ciprofloxacin

In the current study, as in many other reports, the *aac(6')-le-aph(2'')-la* gene was the most prevalent AME gene, encountered in 96.2% of isolates followed by the *ant(4')* gene and was detected in 62.9% of isolates [16, 20]. Acquired resistance to tetracyclines and macrolides in enterococci is often by mobile genetic elements [20]. In the present study, as in many previous reports, the most common


tetracycline and erythromycin resistance mechanism was mediated by the *tetM* and *ermB* genes, respectively [21-25]. To take preventive measures and apply infection control, identification of the source of infection and nature of outbreaks are required. RAPD-PCR has become a reliable tool for the differentiation and identification of enterococci from clinical origin [26-28]. In this study, RAPD-PCR profiles showed that eleven *E. faecalis* isolates shared identical banding patterns. In addition, the results of RAPD typing corresponded with antibiogram findings and the resistance genes patterns for eight of these eleven isolates. This could be explained by the predominance or an outbreak of a particular clonal group of *E. faecalis* with high-level gentamicin resistance in the assessed burn unit.

In conclusion, we found that over the examination time period one RAPD genotype was the common type and was disseminated among patients in the burn unit. Interestingly, most of these strains had an identical or very similar antibiotic and gene resistance pattern.

Acknowledgments

This research has been supported by Iran University of Medical Sciences & Health Services grant number: 17989/91-03-13 and Tehran University of Medical Scienc-

Fig. 1. RAPD-PCR profiles of *Enterococcus faecalis* and *E. faecium* isolates. Lanes 1 and 2; DNA size marker (100bp), Lanes A-Q; Enterococcal isolates with different RAPD patterns.



es & Health Services grant 95-02-30-32393. Authors deny any existing or potential conflicts of interest of a financial, personal or any other nature that could affect or bias their research.

Authors' contributions

ME: study design, project and protocol development, data analysis and manuscript writing. B.K., F.J. and R.B.: collected data and performed microbiological and molecular experiments. K.A.: was involved in the editing of the manuscript. M.T.: was performed the data quality control and manuscript writing. A.R.L.: conceived, designed and coordinated the research. All authors read and approved the final draft of the manuscript.

References

- [1] Cetinkaya Y, Falk P, Mayhall CG. *Vancomycin-resistant enterococci*. Clin Microbiol Rev 2000;13:686-707.
- [2] Klare I, Konstabel C, Mueller-Bertling S, Werner G, Strommenger B, Kettlitz C, Borgmann S, Schulte B, Jonas D, Serr A, Fahr AM, Eigner U, Witte W. *Spread of ampicillin/vancomycin-resistant Enterococcus faecium of the epidemic-virulent clonal complex-17 carrying the genes esp and hyl in German hospitals*. Eur J Clin Microbiol Infect Dis 2005;24:815-25.
- [3] Fatholahzadeh B, Hashemi FB, Emaneini M, Aligholi M, Nakhjavani FA, Kazemi B. *Detection of Vancomycin Resistant Enterococci (VRE) isolated from urinary tract infections (UTI) in Tehran, Iran*. Daru 2006;14:141-5.
- [4] Titze-de-Almeida R, Van Belkum A, Felipe MS, Zanella RC, Top J, Willems RJ. *Multilocus sequence typing of hospital-associated Enterococcus faecium from Brazil reveals their unique evolutionary history*. Microb Drug Resist 2006;12:121.
- [5] Wibbenmeyer L, Williams I, Ward M, Xiao X, Light T, Latenser B, Lewis R, Kealey GP, Herwaldt L. *Risk factors for acquiring vancomycin-resistant Enterococcus and methicillin-resistant Staphylococcus aureus on a burn surgery step-down unit*. J Burn Care Res 2010;31:269-79.
- [6] Altoparlak U, Koca O, Ozkurt Z, Akcay MN. *Incidence and risk factors of vancomycin-resistant enterococcus colonization in burn unit patients*. Burns 2011;37:49-53.
- [7] Branski LK, Al-Mousawi A, Rivero H, Jeschke MG, Sanford AP, Herndon DN. *Emerging infections in burns*. Surg Infect (Larchmt) 2009;10:389-97.
- [8] Kaye KS, Kaye D. *Multidrug-resistant pathogens: mechanisms of resistance and epidemiology*. Curr Infect Dis Rep 2000;2:391-8.
- [9] Nishimoto Y, Kobayashi N, Alam MM, Ishino M, Uehara N, Watanabe N. *Analysis of the prevalence of tetracycline resistance genes in clinical isolates of Enterococcus faecalis and Enterococcus faecium in a Japanese hospital*. Microb Drug Resist 2005;11:146-53.
- [10] Hryniewicz W, Zareba T, Kawalec M. *Susceptibility patterns of Enterococcus spp. isolated in Poland during 1996*. Int J Antimicrob Agents 1998;10:303-7.
- [11] Emaneini M, Aligholi M, Aminshahi M. *Characterization of glycopeptides, aminoglycosides and macrolide resistance among Enterococcus faecalis and Enterococcus faecium isolates from hospitals in Tehran*. Pol J Microbiol 2008;57:173-8.
- [12] Emaneini M, Hashemi FB, Aligholi M, Fatholahzadeh B, Kazemi B, Sadeghi F. *Detection of vanB genotype enterococci in Iran*. Int J Antimicrob Agents 2005;26:98-9.
- [13] Dutka-Malen S, Evers S, Courvalin P. *Detection of glycopeptide resistance genotypes and identification to the species level of clinically relevant enterococci by PCR*. J Clin Microbiol 1995;33:1434.
- [14] Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing; twenty-third informational supplement*. M100-S23, 2013.
- [15] Emaneini M, Bigverdi R, Kalantar D, Soroush S, Jabalameli F, Noorazar Khoshgnab B, Asadollahi P, Taherikalani M. *Distribution of genes encoding tetracycline resistance and aminoglycoside modifying enzymes in Staphylococcus aureus strains isolated from a burn center*. Ann Burns Fire Disasters 2013;30:26:76-80.
- [16] Udo EE, Al-Sweih N, Phillips OA, Chugh TD. *Species prevalence and antibacterial resistance of enterococci isolated in Kuwait hospitals*. J Med Microbiol 2003;52:163-8.
- [17] Titze-de-Almeida R, Rollo Filho M, Nogueira CA, Rodrigues IP, Eudes Filho J, Nascimento RS, Ferreira RF 2nd, Moraes LM, Boelens H, Van Belkum A, Felipe MS. *Molecular epidemiology and antimicrobial susceptibility of Enterococci recovered from Brazilian intensive care units*. Braz J Infect Dis 2004;8:197-205.
- [18] Kaçmaz B, Aksoy A. *Antimicrobial resistance of enterococci in Turkey*. Int J Antimicrob Agents 2005;25:535-8.
- [19] Chow JW. *Aminoglycoside resistance in enterococci*. Clin Infect Dis 2000;31:586-9.
- [20] Hegstad K, Mikalsen T, Coque TM, Werner G, Sundsfjord A. *Mobile genetic elements and their contribution to the emergence of antimicrobial resistant Enterococcus faecalis and Enterococcus faecium*. Clin Microbiol Infect 2010;16:541-54.
- [21] Vignaroli C, Zandri G, Aquilanti L, Pasquaroli S, Biavasco F. *Multidrug-resistant enterococci in animal meat and faeces and co-transfer of resistance from an Enterococcus durans to a human Enterococcus faecium*. Curr Microbiol 2011;62:1438-47.
- [22] Aarestrup FM, Agero Y, Gerner-Smidt P, Madsen M, Jensen LB. *Comparison of antimicrobial resistance phenotypes and resistance genes in Enterococcus faecalis and Enterococcus faecium from humans in the community, broilers, and pigs in Denmark*. Diagn Microbiol Infect Dis 2000;37:127-37.
- [23] De Leener E, Martel A, Decostere A, Haesebrouck F. *Distribution of the erm (B) gene, tetracycline resistance genes, and Tn1545-like transposons in macrolide- and lincosamide-resistant enterococci from pigs and humans*. Microb Drug Resist 2004;10:341-5.
- [24] Nishimoto Y, Kobayashi N, Alam MM, Ishino M, Uehara N, Watanabe N. *Analysis of the prevalence of tetracycline resistance genes in clinical isolates of Enterococcus faecalis and Enterococcus faecium in a Japanese hospital*. Microb Drug Resist 2005;11:146-53.
- [25] Reyes J, Hidalgo M, Díaz L, Rincón S, Moreno J, Vanegas N, Castañeda E, Arias CA. *Characterization of macrolide resistance in Gram-positive cocci from Colombian hospitals: a countrywide surveillance*. Int J Infect Dis 2007;11:329-36.
- [26] van den Braak N, Power E, Anthony R, Endtz HP, Verbrugh HA, van Belkum A. *Random amplification of polymorphic DNA versus pulsed field gel electrophoresis of SmaI DNA macrorestriction fragments for typing strains of vancomycin-resistant enterococci*. FEMS Microbiol Lett 2000;192:45-52.
- [27] Monstein HJ, Quednau M, Samuelsson A, Ahmè S, Isaksson B, Jonasson J. *Division of the genus Enterococcus into species groups using PCR-based molecular typing methods*. Microbiology 1998;144:1171-9.
- [28] Gzyl A, Augustynowicz E. *Technical aspects of random amplified polymorphic DNA (RAPD) technique in genotyping of bacterial strains*. Acta Microbiol Pol 1999;48:243-59.

■ Correspondence: Mohammad Emaneini, Department of Microbiology, School of Medicine, Tehran University of Medical Sciences 100 Poursina St., Keshavarz Blvd., Tehran, Iran - Tel- Fax: 098 021 8895 5810 - E-mail: emaneini@tums.ac.ir

■ Received on August 2, 2015. Accepted on September 19, 2016.