

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/289243917>

Genetic and Morphological Diversity of the Genus *Penicillium* From Mazandaran and Tehran Provinces, Iran

Article in *Jundishapur Journal of Microbiology* · January 2016

DOI: 10.5812/jjm.28280

CITATIONS

2

READS

113

11 authors, including:



Mahdi Abastabar

84 PUBLICATIONS 377 CITATIONS

[SEE PROFILE](#)



Hossein Mirhendi

Isfahan University of Medical Sciences

230 PUBLICATIONS 2,793 CITATIONS

[SEE PROFILE](#)



Mohammad T Hedayati

Mazandaran University of Medical Sciences

138 PUBLICATIONS 2,351 CITATIONS

[SEE PROFILE](#)



Tahereh Shokohi

Mazandaran University of Medical Sciences

182 PUBLICATIONS 843 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Genotyping and antifungal susceptibility pattern of clinical and environmental isolates of *Aspergillus nidulans* [View project](#)



Diagnosis and identification of mucormycosis in patients suspected to invasive fungal infection using conventional, semi-nested PCR and sequencing [View project](#)

Genetic and Morphological Diversity of the Genus *Penicillium* From Mazandaran and Tehran Provinces, Iran

Mahdi Abastabar,^{1,*} Hossein Mirhendi,² Mohammad Taghi Hedayati,¹ Tahereh Shokohi,¹ Ali Rezaei-Matehkolaei,³ Rasoul Mohammadi,⁴ Hamid Badali,¹ Maryam Moazeni,¹ Iman Haghani,¹ Aynaz Ghojoghi,⁵ and Javad Akhtari^{6,7}

¹Invasive Fungi Research Center (IFRC), Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, IR Iran

²Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, IR Iran

³Department of Medical Mycology, School of Medicine, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

⁴Department of Medical Parasitology and Mycology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, IR Iran

⁵Iran University of Medical Sciences and Health Services, Tehran, Iran

⁶Immunogenetic Research Center, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, IR Iran

⁷Department of Nanobiomedicine, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, IR Iran

*Corresponding author: Mahdi Abastabar, Invasive Fungi Research Center (IFRC), Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, IR Iran. Tel: +98-911211347, Fax: +98-1133543248, E-mail: Mabastabar@gmail.com

Received 2015 March 2; Revised 2015 July 20; Accepted 2015 July 26.

Abstract

Background: The genus *Penicillium* contains a large number of ubiquitous environmental taxa, of which some species are clinically important. Identification of *Penicillium* down to the species level is currently based on polyphasic criteria, including phenotypic features and genetic markers. Biodiversity of the genus *Penicillium* from Mazandaran and Tehran provinces has not been described.

Objectives: The current paper focused on the environmental biodiversity of *Penicillium* isolates within some areas of Mazandaran and Tehran provinces, based on morphological traits and the molecular data from partial sequence of the β -tubulin (BT2) gene.

Materials and Methods: A total of 400 strains were isolated from the environment and investigated using morphological tests and sequencing of BT2, in order to characterize the spectrum of the *Penicillium* species.

Results: Sequence analysis of BT2 and morphological criteria of 20 strains representative of 10 species showed that *Penicillium chrysogenum* was the most prevalent species (n = 6), followed by *P. polonicum* (n = 3), *P. glabrum* (n = 2), *P. palitans* (n = 2), *P. melanoconidium* (n = 2), and other species, including *P. expansum*, *P. canescense*, *P. griseofulvum*, *P. italicum*, and *P. raistrickii* with one case each.

Conclusions: It was shown that partial β -tubulin sequence, as a reliable genetic target, supported specific morphological criteria for identification of the *Penicillium* species. Like other assessments throughout the world, *P. chrysogenum* remains the most frequent environmental *Penicillium* species in Mazandaran and Tehran Provinces.

Keywords: Beta-Tubulin, PCR, DNA Sequencing, *Penicillium*

1. Background

The genus *Penicillium* comprises a large number of ubiquitous filamentous fungi, of which some are involved in human infections, ranging from mild to severe infections, especially in patients infected with human immunodeficiency virus (HIV), patients with underlying diseases, and intravenous drug abusers (1). Moreover, they lead to many complications, such as mycotoxicosis, allergies, and fungal sinusitis, and are frequently isolated from soil, dead plant materials, rotten wood, decaying vegetables, and foods (2). Based on phenotypic data, the genus of *Penicillium* is regarded to be related to ascomycota, because of its sexual reproduction by means of ascospores (3, 4). Members of the genus are of commercial and industrial importance due to their use in production of antibiotics, anti-tumoral, anti-fungal, anti-insect, and anti-viral compounds, as well as extracellular enzymes (2).

To date, it has been revealed that various sections of *Penicillium* encompass more than 250 species (5). The most common species of *Penicillium*, *P. chrysogenum*, is found to be an agent of onychomycosis, keratomycosis, allergic bronchopulmonary mycosis, and asthma, which may be associated with indoor environments, deserts, dried foods, cheese, and is morphologically characterized by biverticillate, terverticillate, or quarter-verticillate conidiophores, floccose to velutinous in colony texture, and yellow exudate droplets (2, 6-8). However, these conventional methods were noted to be tedious, time consuming and unreliable for species delineation (2, 9, 10). Nevertheless, molecular-based methods, such as DNA sequencing, have provided powerful tools for precise identification of the *Penicillium* species.

Currently, species identification in this genus is based

on polyphasic criteria consisting of morphological and biochemical traits and molecular data from internally transcribed spacer (ITS) regions of ribosomal DNA (rDNA) and the partial β -tubulin (BT2) sequences (2, 9, 11). While the ITS-rDNA regions have some limitations, such as less resolution in differentiation of closely related species, due to a better species-specific resolution provided by the partial BT2 sequencing, application of this locus has substantially increased and is now an excellent marker for differentiation of *Penicillium* spp (2, 9, 12).

2. Objectives

Given that there was no comprehensive evaluation about the biodiversity of this genus in Iran, this study was focused on determining the distribution profile of *Penicillium* spp. in two provinces of Iran (Mazandaran and Tehran).

3. Materials and Methods

3.1. Fungal Strains

In total, 400 isolates used in this study were obtained from soil, air, cereals, and decaying vegetables from different regions of Mazandaran and Tehran provinces (Table 1). Stock cultures were maintained on slants of 2% malt extract agar (Difco, Detroit, MI, USA) and incubated at 25°C for one week.

3.2. Morphological Identification and Cultural Characterization

Strains were cultured on 2% malt extract agar (Difco, Detroit, MI, USA), Czapek yeast extract agar (Difco, Detroit, MI, USA), and yeast extract agar (Himedia, India) and incubated at 24°C in the dark for one week. Identification was performed primarily based on macroscopic and microscopic morphology. Microscopic observations were based on slide culture techniques using potato dextrose agar (PDA) (5). Mounts of four-day-old slide cultures were made in lactic acid or lactophenol cotton blue, and light micrographs were taken using a Nikon Eclipse 80i microscope (Nikon, Tokyo, Japan) equipped with a Nikon digital sight DS-Fi1 camera. Moreover, strains were evaluated based on phenotypic characters, i.e. color, texture, growth rate, pigmentation, conidiophores, conidia, metulae, and phialide morphology (2).

3.3. Molecular Characterization

The fungal mycelia were grown on 2% MEA plates for four days at 24°C. Subsequently, a sterile blade was used to scrape off the mycelium from the surface of the plate. Genomic DNA was extracted using an ultra clean microbial DNA Isolation Kit (MoBio Inc. Solana Beach, CA, USA), according to the manufacturer's instructions. DNA extracts were stored at -20°C prior to use.

The BT2 gene was partially amplified and sequenced using the universal fungal primers as follows: Bt2a (5'-GGTAACCAAATCGGTGCTGCTTC-3') and Bt2b (5'-ACCCT-CAGTGTAGTGACCCTTGGC-3') (13, 14). PCR reactions were performed on a Corbett research thermal cycler, model CG1-96 (Sydney, Australia) in 50 μ L volume containing 25 ng of template DNA, 5 μ L reaction buffer (0.1 M Tris-HCl, pH 8.0, 0.5 M KCl, 15 mM MgCl₂, 0.1% gelatine, 1% Triton X-100), 0.2 mM of each dNTP, and 2.0 units Taq DNA polymerase (ITK diagnostics, Leiden, The Netherlands). Amplification of BT2 was performed with one cycle of five minutes at 95°C for primary denaturation, followed by 35 cycles of 95°C for 45 seconds, 60°C for 120 seconds, and 72°C for 60 seconds, with a final extension of 72°C for seven minutes. Amplicons were purified using GFX PCR DNA and a gel band purification kit (GE healthcare, Buckinghamshire, UK). PCR-sequencing was performed as follows: 95°C for one minute, followed by 30 cycles consisting of 95°C for 10 seconds, 50°C for five seconds and 60°C for two minutes. Reactions were purified with Sephadex G-50 fine columns (GE healthcare bio-sciences AB, Uppsala, Sweden), and PCR products were sent to Bioneer company (Bioneer, Daejeon, South Korea) for bidirectional sequencing with the primers mentioned previously.

3.4. Alignment and Phylogenetic Reconstruction

Sequence data obtained in this study were imported to MEGA software version 5 (<http://www.megasoftware.net>) and adjusted using ClustalW. Ambiguous regions were excluded from the alignment, and then each trimmed sequence was exported to a BLAST search of the NCBI databases for species recognition. They were compared with *Penicillium* sequences, and query sequences with high similarity ($\geq 99\%$) were considered as the same species.

The program, RAXML-VI-HPC v. 7.0.0, as implemented on the Cipres portal v. 1.10, was used for the tree search and the bootstrap analysis (GTRMIX model of molecular evolution and 500 bootstrap replicates). Bootstrap values equal to or greater than 70% were considered significant. Phylogenetic analyses were carried out based on the BT2 sequence in order to assess the phylogenetic placement of diverse *Penicillium* species, along with their relative sequences downloaded from GenBank, based on the Barreto et al. design (9). The phylogenetic tree was edited using Tree View v. 1.6.6. *Paecilomyces variotii* was used as the out-group.

4. Results

4.1. Morphological Identification and Cultural Characterization

All 400 *Penicillium* isolates were screened for phenotypic characters, and 20 strains representative of 10 morphological species, including *P. chrysogenum*, *P. polonicum*, *P. canescens*, *P. griseofulvum*, *P. italicum*, *P. raistrickii*, *P. expansum*,

P. melanoconidium, *P. palitans*, and *P. glabrum*, were identified and selected for molecular analysis. In Figure 1A - J, the colony characteristics of 20 representative isolates on CYA and YES after seven days at 25°C were exemplified.

Some species, such as *P. chrysogenum* and *P. italicum*, due to production of typical yellow and red exudates on CYA and YES, respectively, could easily be identified, while for others it was essential to apply three complementary specific media (MEA, YES, and CYA) for exact identification (Figure 1A - G). Except for *P. chrysogenum* and *P. expansum* with floccose colony texture, others were velutinous (Table 1 and Figure 1). Upon microscopic examination, nearly all identified species were one- or two-staged branched (biverticillate, terverticillate), except for *P. glabrum*, which typically produces no branching in conidiophore (monoverticillate) (Table 2).

The species isolated in this study had globose, smooth-wall conidia, cylindrical phialide, and metula in various sizes. Interestingly, the only observed finely roughened conidium was related to *P. canescens*, which had been isolated from soil.

4.2. Results of BT2 Amplification, Sequencing, and Phylogeny

Partial amplification of BT2 was successful in all isolates and yielded a single PCR band of approximately 450 base

pair (bp) in gel electrophoresis (Figure 2). Sequencing showed that the length of the isolates ranged between 445 to 460 bp.

Final identification of isolates was performed by comparing the obtained sequences with those of reference strains held in the open access validated CBS database for *Penicillium* (<http://www.cbs.knaw.nl/penicillium/>).

Sequences of all isolates were deposited in GenBank and the accession numbers of the sequences are given in Table 1. In Table 1 the results of sequence and phenotypic-based identification were compared. Figure 1 shows that phylogenetic reconstruction was successfully carried out for the BT2 region and represented the maximum parsimony phylogenetic tree reconstructed on then β -tubulin gene. The phylogenetic assessment showed that there were three major sections, consisting of *Viridicata*, *Chrysogena*, *Penicillium*, and one series, *Glabra* including *P. glabrum*.

4.3. Viridicata Section Dataset

The analysis showed that the environmental species including *P. melanoconidium* (represented by two strains), *P. polonicum* (represented by three strains), *P. palitans* (represented by two strains), and the related standard strains (represented by three strains) are monophyletic and well supported (96.4% bootstrap, Figure 3).

Table 1. Comparison of Cultural Characteristics and Sequencing for Species Identification in 20 Representative Isolates

Species Identification by Phenotypic Criteria	Number of Isolates	Source	Cultural Characteristics	Species Identification by Sequencing of B-Tubulin	Accession Number
<i>P. chrysogenum</i>	6	Soil, fruit, air	Floccose	<i>P. chrysogenum</i>	KP851943, KT285850, KT285859, KT285861, KT285865, KT285866
<i>P. polonicum</i>	3	Wheat	Velutinous	<i>P. polonicum</i>	KT285848, KT285856, KT285864
<i>P. glabrum</i>	2	Plant	Velutinous	<i>P. glabrum</i>	KT285852, KT285854
<i>P. palitans</i>	2	Bread	Velutinous	<i>P. palitans</i>	KT285849, KT285851
<i>P. melanoconidium</i>	2	Wheat	Velutinous	<i>P. melanoconidium</i>	KT285858, KT285860
<i>P. expansum</i>	1	Fruit	Floccose	<i>P. expansum</i>	KT285855
<i>P. raistrickii</i>	1	Soil	Velutinous	<i>P. raistrickii</i>	KT285857
<i>P. italicum</i>	1	Fruit	Velutinous	<i>P. italicum</i>	KT285853
<i>P. griseofulvum</i>	1	Soil	Velutinous	<i>P. griseofulvum</i>	KT285863
<i>P. canescens</i>	1	Soil	Velutinous	<i>P. canescens</i>	KT285862
Total	20				




Figure 1. A, *P. chrysogenum*. 7-day old colonies at CYA, MEA, YES. B, *P. canescens*, C, *P. griseofulvum*, D, *P. polonicum*, E, *P. glabrum*, F, *P. expansum*, G, *P. italicum*, H, *P. raistrickii*, I, *P. melanoconidium*, J, *P. palitans*.

Table 2. The Criteria Used in Phenotypic Based Identification^a

Species Identification by Phenotypic Criteria	Conidia	Phialide	Conidiophore Branching Pattern ^a	Metulae
<i>P. polonicum</i>	Smooth, globose, 3 μm	Cylindrical, 10 μm	Terverticillate, Biverticillate	Cylindrical, 10 μm
<i>P. glabrum</i>	Smooth, subglobose to elliptical, 3 μm	Cylindrical, 10 μm	Monoverticillate	not present
<i>P. expansum</i>	Smooth, ellipsoidal, 3 μm	Cylindrical, 8 μm	Terverticillate	Cylindrical, 11 μm
<i>P. raistrickii</i>	Smooth, globose, 2.5 μm	Cylindrical, 7-9 μm	Biverticillate	Cylindrical, 10-12 μm
<i>P. italicum</i>	Smooth, ellipsoidal, 3.5 μm	Cylindrical, 8-10 μm	Terverticillate	Cylindrical, 14 μm
<i>P. melanoconidium</i>	Smooth, globose, 3.5 μm	Cylindrical, 10-14 μm	Terverticillate	Cylindrical, 12 μm
<i>P. griseofulvum</i>	Smooth, ellipsoidal, 2 μm	Cylindrical, 4-5 μm	Terverticillate	Cylindrical, 9 μm
<i>P. chrysogenum</i>	Smooth, globose, 3 μm	Cylindrical, 7 μm	Bi, ter and quarterverticillate	Cylindrical, 10 μm
<i>P. canescens</i>	finely roughened, globose, 2 μm	Cylindrical, 8 μm	Biverticillate, Monoverticillate	Cylindrical, 10-16 μm
<i>P. palitans</i>	Smooth, globose, 3 - 4 μm.	Cylindrical, 10 μm	Terverticillate	Cylindrical, 10-14 μm

^aDefinitions: Monoverticillate, conidiophore without branching; Biverticillate, conidiophore one-stage branching; Terverticillate, conidiophore two-stage branch.


4.4. *Penicillium* Section Dataset

The phylogenetic tree (Figure 3) shows that the *Penicillium* section is composed of five species, *P. canescens*, *P. italicum*, *P. expansum*, *P. griseofulvum* and *P. raistrickii*, which are phylogenetically distinct from other species and sister to each other and standard strains in this section. The species of the *Penicillium* section were strongly (93.1%) well supported by bootstrap.

4.5. *Chrysogena* Section and *Glabra* Series Dataset


Among accepted members of the *Chrysogena* section and the *glabra* series, only *P. chrysogenum* and *P. glabrum*, respectively, were isolated from all the examined isolates. The phylogenetic tree (Figure 3) shows that six *P. chrysogenum* and one standard strain (UBOCC-A 101393) formed a well-supported monophyletic group nested within this section of *Penicillium*.

Figure 2. Gel Electrophoresis of PCR Products of Some *Penicillium* Species



Line 1, *P. chrysogenum*; Line 2, *P. canescens*; Line 3, *P. griseofulvum*; Line 4, *P. polonicum*; Line 5, *P. glabrum*; Line 6, *P. expansum*; Line 7, *P. italicum*; Line 8, *P. raistrickii*; N, negative control; M, DNA ladder.

Figure 3. Maximum Parsimony Tree Based on β -tubulin Sequences



Branch numbers are bootstrap values $\geq 70\%$, estimated based on 500 replicates.

5. Discussion

To evaluate biodiversity and phylogeny of the *Penicillium* species in Mazandaran and Tehran Provinces, the current study focused on morphology-based identification and sequencing of the β -tubulin gene. Except for the studies by Sabokbar et al. and Javadi et al. there has been no extensive attempt at this type of study, including reliable morphological and high-resolution molecular approaches for *Penicillium* identification, until now in Iran (15, 16). Over the past decades, focusing on macro- and microscopic characteristics of the colonies has been the main strategy for classification and identification of the *Penicillium* species (2, 17). In our study, which was morphologically based on three specific medium (CYA, MEA, and YES) (2), 10 species of *Penicillium* were identified, compared to Javadi et al. who did not focus on phenotypic characteristics and whose sampling was restricted, taken only from soil (16). Also, the study performed by Sabokbar et al. showed intra-species variation of *Penicillium* species isolated from air in Iran using RAPD-PCR, but did not consider identification of the *Penicillium* species (15).

In the present study, in descending arrangement of frequency, the isolates that were identified were: *P. chrysogenum*, *P. polonicum*, *P. palitans*, *P. glabrum*, *P. melanoconidium*, *P. expansum*, *P. raistrickii*, *P. italicum*, *P. griseofulvum*, and *P. canescens*. Contrary to ITS regions, the β -tubulin gene is known to have more variation in *Penicillia* and currently is considered a reliable alternative marker for differentiation of the *Penicillium* species (5). Since identification of all *Penicillium* species by morphology was stated to be very problematic and often impossible, relying on sequencing of the high resolution genetic marker, β -tubulin, allowed for accurate and rapid recognition of the isolates in our study (2, 14). Similar to Barreto et al. parsimony analysis was used in our study to determine the phylogeny of the isolates (9). The *Viridicata* section is substantially divided into five series including *Viridicata*, *Camemberti*, *Corymbifera*, *Verrucosa*, and *Solita* (9). In the current survey, only species related to two former series, such as *P. melanoconidium*, *P. polonicum*, and *P. palitans*, were isolated, of which *P. melanoconidium* and *P. polonicum* are regarded to be prone to cereals such as barley, wheat, rye, oats, rice, and cause mycotoxicosis symptoms in humans (2).

Also, *P. palitans* may produce some significant toxins such as cyclopiazonic acid and fumigaclavine on foods, and the former is proven to be an agent of organ damage in mammals (2). In the *Penicillium* section, *P. griseofulvum*, as the main producer of griseofulvin, which is placed in ser. *Urticolae* alongside *P. raistrickii* and *P. canescens*, had completely identical sequences with the standard strain (100%). *Penicillium expansum* and *P. italicum*, two members of the *Expansa* and *Italica* series, respectively, had 99.9% identity to the standard sequences (PSN118 and AB688988.1). *Penicillium chrysogenum*, the predominant isolate in our study, is assigned to section *Chrysogena*, ser. *Chrysogena* (9). The species belonging to this section vary from one geographic site to another. Whereas *P. chrysogenum*, *P. spinulosum*, and *P. ox-*

alicum ranked as the first infrequency in Kansas, *P. citrinum* was the main agent isolate from the Taipei area (18). *Penicillium chrysogenum* was the predominant agent among all assessed isolates in this study, and this finding was concordant with other reports from around the world (19). Extra evaluation, with emphasis on polyphasic procedures, not only phenotypic methods but also molecular approaches focusing on a combination of partial β -tubulin, calmodulin, and ITS sequence, are required to clarify the complete spectrum of the *Penicillium* species in Iran.

5.1. Conclusions

The current investigation found a high level of genetic variability in the β -tubulin gene in various *Penicillium* species isolated from soil, fruits, food, and grains of Mazandaran and Tehran Provinces. This study reconfirmed the view that BT2 PCR-sequencing is a reliable and applicable diagnostic tool for differentiating and the molecular taxonomy of closely related *Penicillium* species.

Acknowledgments

This work was financially supported by the invasive fungi research center (IFRC), faculty of medicine, Mazandaran University of Medical Sciences, Sari, Iran and Tehran University of Medical Sciences, Tehran, Iran.

Footnotes

Author Contribution: Hossein Mirhendi developed the study concept and design; Mahdi Abastabar and Iman Haghani contributed to acquisition of data; Ali Rezaei-Matehkolaei, Rasoul Mohammadi, Maryam Moazeni and Aynaz Ghoghghi analyzed the data; Mahdi Abastabar and Hamid Badali wrote the manuscript; Mohammad Taghi Hedayati and Tahereh Shokohi contributed to critical revision of the manuscript; Javad Akhtari contributed to statistical analysis; Hossein Mirhendi contributed to administrative, technical, and material support and study supervision.

Funding/Support: This research was financially supported by a grant of Mazandaran University of Medical Sciences and Tehran University of Medical Sciences.

References

1. Le T, Wolbers M, Chi NH, Quang VM, Chinh NT, Lan NP, et al. Epidemiology, seasonality, and predictors of outcome of AIDS-associated *Penicillium marneffei* infection in Ho Chi Minh City, Viet Nam. *Clin Infect Dis*. 2011;52(7):945–52. doi:10.1093/cid/cir028. [PubMed: 21427403]
2. Frisvad JC, Samson RA. Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A guide to identification of food and airborne terverticillate *Penicillia* and their mycotoxins. *Stud in mycol*. 2004;49(1):C174.
3. Cheeseman K, Ropars J, Renault P, Dupont J, Gouzy J, Branca A, et al. Multiple recent horizontal transfers of a large genomic region in cheese making fungi. *Nat Commun*. 2014;5:2876. doi:10.1038/ncomms3876. [PubMed: 24407037]
4. Haga DI, Burrows SM, Iannone R, Wheeler MJ, Mason R, Chen J, et al. Ice nucleation and its effect on the atmospheric transport of fungal spores from the classes Agaricomycetes, Ustilaginomycetes, and Eurotiomycetes. *Atmos Chem Phys Discuss*. 2014;14:5013–59.

5. Houbraken J, Samson RA. Phylogeny of Penicillium and the segregation of Trichocomaceae into three families. *Stud Mycol.* 2011;**70**(1):1-51. doi:10.3114/sim.2011.70.01. [PubMed: 22308045]
6. Arora R, Gupta S, Raina UK, Mehta DK, Taneja M. Penicillium keratitis in vernal Keratoconjunctivitis. *Indian J Ophthalmol.* 2002;**50**(3):215-6. [PubMed: 12355698]
7. Hajoui FZ, Zeroual Z, Ghfir B, Moustachi A, Lyagoubi M, Aoufi S. [The mould onychomycosis in Morocco: about 150 isolated cases in 20 years]. *J Mycol Med.* 2012;**22**(3):221-4. doi:10.1016/j.mycmed.2012.05.001. [PubMed: 23518078]
8. Chowdhary A, Agarwal K, Kathuria S, Gaur SN, Randhawa HS, Meis JF. Allergic bronchopulmonary mycosis due to fungi other than Aspergillus: a global overview. *Crit Rev Microbiol.* 2014;**40**(1):30-48. doi:10.3109/1040841X.2012.754401. [PubMed: 23383677]
9. Barreto MC, Houbraken J, Samson RA, Frisvad JC, San-Romão MV. Taxonomic studies of the Penicillium glabrum complex and the description of a new species P. subericola. *Fungal Divers.* 2011;**49**(1):23-33. doi:10.1007/s13225-011-0090-4.
10. Seifert KA. Progress towards DNA barcoding of fungi. *Mol Ecol Resour.* 2009;**9** Suppl s1:83-9. doi:10.1111/j.1755-0998.2009.02635.x. [PubMed: 21564968]
11. Houbraken JAMP, Frisvad JC, Samson RA. Taxonomy of Penicillium citrinum and related species. *Fungal Divers.* 2010;**44**(1):117-33.
12. Houbraken J, Frisvad JC, Samson RA. Taxonomy of Penicillium section Citrina. *Stud Mycol.* 2011;**70**(1):53-138. doi:10.3114/sim.2011.70.02. [PubMed: 22308046]
13. Glass NL, Donaldson GC. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol.* 1995;**61**(4):1323-30. [PubMed: 7747954]
14. Abastabar M, Mirhendi H, Rezaei-Matehkolaei A, Shidfar MR, Kordbacheh P, Makimura K. Restriction analysis of beta-tubulin gene for differentiation of the common pathogenic dermatophytes. *J Clin Lab Anal.* 2014;**28**(2):91-6. doi:10.1002/jcla.21649. [PubMed: 24395510]
15. Sabokbar A, Bakhtiari A, Khosravi A, Zanjani LS. Intraspecies Molecular Segregation of Penicillium Species Isolated from Air in Iran Using Rapid Polymerase Chain Reaction Method. *Glob Vet.* 2012;**8**:119-23.
16. Javadi MA, TajickGhanbary MA, Tazick Z. Isolation and Molecular Identification of Soil Inhabitant Penicillia. *Annals Biol Res.* 2012;**3**:5758-61.
17. Visagie CM, Houbraken J, Frisvad JC, Hong SB, Klaassen CH, Perrone G, et al. Identification and nomenclature of the genus Penicillium. *Stud Mycol.* 2014;**78**:343-71. doi:10.1016/j.simyco.2014.09.001. [PubMed: 25505353]
18. Shen HD, Lin WL, Tam MF, Wang SR, Tzean SS, Huang MH, et al. Characterization of allergens from Penicillium oxalicum and P. notatum by immunoblotting and N-terminal amino acid sequence analysis. *Clin Exp Allergy.* 1999;**29**(5):642-51. [PubMed: 10231324]
19. Scott J, Untereiner WA, Wong B, Straus NA, Malloch D. Genotypic variation in Penicillium chrysogenum from indoor environments. *Mycologia.* 2004;**96**(5):1095-105. [PubMed: 21148929]