

Trauma Mon. 2015 November; 20(4): e29033.

doi:10.5812/traumamon.29033

Published online 2015 November 23.

Research Article

Emergency Nurses' Requirements for Disaster Preparedness

Hesam Seyedin,¹ Zahra Abbasi Dolatabadi,^{1,*} and Fatemeh Rajabifard²¹School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, IR Iran²Research Center of Future Studies, Kerman University of Medical Sciences, Kerman, IR Iran

*Corresponding author: Zahra Abbasi Dolatabadi, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, IR Iran. Tel: +98-2161054401, Fax: +98-218883334, E-mail: zahra_abbasi2000@yahoo.com

Received 2015 April 10; Revised 2015 June 2; Accepted 2015 July 1.

Abstract

Background: Natural and man-made disasters affect people, communities, and health systems. Nurses play a key role in the health system and must be prepared for appropriate response in disasters.

Objectives: This study was done to assess the current knowledge of nurses in emergency departments for disaster preparedness.

Patients and Methods: This cross-sectional study was conducted on 110 emergency nurses working in teaching hospitals affiliated to Iran University of Medical Sciences, Tehran, Iran. A questionnaire was used to collect data and the data were then analyzed using descriptive and inferential statistics with SPSS software version 16.0.

Results: Based on the findings, most of the participants (64.5%) were aged 30 - 39 years old. 84% of the cases were female and 97.3% had a bachelor's degree. The average perceived knowledge of nurses was 2.43 ± 1.01 . The nurses had the highest familiarity with triage and lowest knowledge in epidemiology and decision-making.

Conclusions: Lack knowledge of nurses regarding response to disaster situations indicates inefficiencies in the current system. Therefore, it is recommended to organize more workshops, annual training courses, and maneuvers based on staff needs and formulate continuous education courses for nurses.

Keywords: Needs Assessment, Education, Emergency Department, Knowledge, Disasters, Emergency Preparedness, Nurse Staff

1. Background

Advancements in technology and industrialization increase the rate of natural and man-made disasters (1, 2). In 2014 alone, 202 disasters were recorded in Asia affecting 87760054 and injuring 10,107 people (3). Iran is vulnerable to a range of natural and man-made disasters (4, 5) and the capital Tehran is not an exception. It is located on active faults; earthquakes with high severity are expected (6). Moreover, man-made hazards such as oil and gas reservoir tank explosions (7), rising immigration to Tehran, presence of factories, and high population density can result in significant losses and casualties from disasters in this megacity (6).

The health care system plays a very important role in helping communities following disasters (8). After disasters, destruction and heavy damages to the healthcare infrastructures, buildings and management system are expected, which may result in death and injuries (9-12). The population in affected areas expect an efficient and timely response from the health care system. Nurses play a key role in the system (13) and must be prepared for appropriate response to disasters and crises (14-17).

Previous experiences and trainings of nurses affect their preparedness, increase their awareness, self-confi-

dence and skills in disaster response and decrease their vulnerability to unpredictable events (18-23). Studies show that most nurses are not ready to face mass casualties, and their preparedness, education and skills are not adequate for an appropriate response (15). Accordingly, the first step is to determine priorities and training needs of Emergency Department (ED) nurses as the front line responders (24, 25). This study sought to assess this issue.

2. Objectives

This study was performed to assess the current knowledge of nurses in EDs for disaster preparedness.

3. Patients and Methods

This cross-sectional study was conducted in teaching hospitals affiliated to Iran University of Medical Sciences in 2014. The study population ($n = 460$) included nurses in the EDs of 8 hospitals. With a confidence coefficient of 95% and a testing power of 90%, 110 participants were recruited using convenient sampling. The inclusion criteria were: employment in ED for at least six months, and having a bachelor or higher degree in nursing. Question-

naire completion was voluntary and anonymous. Those who volunteered, filled and signed consent forms.

A two-part questionnaire was used to collect the data. The first part covered demographic information of the nurses including age, sex, married status and work history. The second part consisted of 42 questions. In this questionnaire, knowledge and training needs of nurses in disaster response were examined in 8 domains namely: triage, communication, biological agent, decontamination, quarantine, isolation, reporting and accessing critical resources, psychological issues and vulnerable population, epidemiological and clinical decision and incident command system (ICS) (26, 27). A five-point Likert scale (1 = not familiar/very poor, 2 = poor/familiar, 3 = fair/familiar, 4 = good/familiar, 5 = very familiar/very good) was used to measure knowledge. The questionnaire was adapted from the studies of Wisniewski et al. (26). Translation and back-translation were used to develop a Persian version of the questionnaire. To test the validity, the instrument was reviewed by the research team and 15 experts. A pilot study was conducted with a sample (n = 30). The Cronbach's Alpha test indicated that items were internally consistent ($\alpha = 0.92$). The nurses entered the study with their own consent and they filled out and signed the consent form. The final instrument had 42 questions. The questionnaire was completed in the wards and the completion of questionnaire took around 30 minutes.

Questionnaires were distributed among the nurses by one of the researchers. Data were analyzed using SPSS version 16.0. Results are expressed as frequency, mean and standard deviation and inferential (Pearson correlation coefficient and independent t-test) statistics.

4. Results

Most of the participants were aged 30 - 39 years-old. Eighty-four percent of the nurses were females and 97.3% had a bachelor's degree. Only three participants had a postgraduate degree. Also, 59 nurses had less than five years experience in EDs (Table 1).

The average perceived knowledge of nurses was 2.43 ± 1.01 . The nurses had the highest familiarity with triage (2.77 ± 0.86). They had the highest knowledge in basic first aid in large-scale disasters, administering oxygen and artificial respiration with a mean of 2.97 ± 1.12 and the lowest familiarity in assessing the effectiveness of their activities in responding to the large-scale disasters (2.55 ± 1.02). Epidemiology and decision-making obtained the lowest score (2.47 ± 0.82). In this domain, general issues related to mass casualty management in large-scale disasters (ethical, cultural, legal and safety issues) and ability to identify the underlying disease resulting from exposure to chemical, biological and radiological agents obtained 2.52 ± 1.13 and 2.39 ± 1.05 , respectively (Table 2).

Table 1. Distribution of the Participants' Demographic Characteristics^{a,b}

Variables	Values	Mean \pm SD
Age, y		33.15 \pm 6.07
< 30	25 (22.7)	
30 - 39	71 (64.5)	
> 40	14 (12.7)	
Gender		-
Female	93 (84.5)	
Male	17 (15.5)	
Academic degree		-
BS	107 (97.3)	
MS	2 (1.8)	
PhD	1 (0.9)	
Experience in nursing, y		8.59 \pm 5.59
< 5	24 (21.8)	
6 - 10	45 (4.09)	
11 - 15	27 (24.5)	
16 <	14 (12.7)	
Experience in ED, y		5.20 \pm 3.68
< 5	59 (53.6)	
6 - 8	32 (29.1)	
9 <	19 (17.3)	

^aAbbreviations: BS, Bachelor of Science; ED, Emergency Department; MS, Master of Science degree; PhD, Doctor of Philosophy degree.

^bValues are presented as No. (%).

Table 2. Familiarity and Knowledge of Nurses in Different Fields of Disaster Preparedness^{a,b}

Domain	Mean \pm SD
ICS	2.51 \pm 0.84
Triage	2.77 \pm 0.86
Communication	2.58 \pm 0.90
Psychological domain and vulnerable groups	2.50 \pm 0.92
Decontamination, quarantine and isolation	2.58 \pm 0.86
Epidemiology and clinical decision	2.47 \pm 0.82
Reporting and accessing critical resources	2.59 \pm 0.89
Biological factors	2.49 \pm 0.85
The average total perceived knowledge	2.43 \pm 1.01

^aAbbreviation: ICS, incident command system.

^bn = 110.

5. Discussion

Researchers believe that formulating a comprehensive plan for disaster response must assess, training needs of individuals be (25, 27). This study surveyed nurse's educational needs in response to disasters and found a moderate response in disaster preparedness knowledge.

There was no relationship between nurses' demographic data (age, gender and their experience in ED) and their level of knowledge. Emergency nurses obtained the highest score in triage and lowest score in epidemiology and clinical decision making. As triage is performed in the daily activities of nurses in EDs, they have appropriate information regarding it.

Despite the establishment of disaster committees in hospitals, nurses are unfamiliar with ICS, reporting methods, and access to data resources. Wisniewski et al. (26) found that knowledge of nurses in disaster preparedness was lower than expected. Although nurses knew triage they had the lowest familiarity with communication. Similar results were obtained in the study of Garbutt et al. (27).

In a study conducted by Worrall (28), nurses obtained the highest scores in biological agent and lowest scores in ICS. McKibbin et al. (29) also showed that familiarity of nurses in disaster preparedness were in low. The nurses had little knowledge regarding epidemiological and biological agents. The studies of Al Khalaileh et al. (16) in Jordanian and O'Sullivan et al. (18) showed difficulty of nurses in dealing with mass casualties.

Lack of cooperation of two teaching hospitals and nurses' refusal to participate in the study (due to work) were limitations. These factors can affect the generalizability of the results.

As the health care system in Middle East countries are nearly the same (2, 30), the findings of this study are similar in these countries and the must opt to appropriate policies to deal with crises. Organizing annual training courses and maneuvers based on staff needs are appropriate strategies for preparedness of medical centers. Continuing education should be held for the nurses. Bioterrorism agents, use of personal protective equipment, epidemiology, psychological first aid, disaster management processes, mass casualties management and command systems are the required fields for education. Scenario-based drills and exercises are also suggested. Moreover, establishing a formal curriculum in mass casualty and disaster management to improve knowledge and skills in all nurses are essential.

Acknowledgments

This study was supported by vice chancellor for research and technologies of Iran University of medical sciences. The approval code is 21074-136-01-92. The authors express their appreciation to all those who helped in this study.

Footnotes

Authors' Contribution: Zahra Abbasi Dolatabadi: Data gathering, data analysis, and article adjustment. Hesam

Seyedin: Drafting of the manuscript. Fatemeh Rajabifard: Data analysis and article adjustment.

Funding/Support: This study was supported by the vice chancellor for research and technology of Iran University of Medical Sciences.

References

1. Coppola DP. *Introduction to International Disaster Management*. 2 ed. United Kingdom: Butterworth-Heinemann; 2011. p. 684.
2. Seyedin H, Zaboli R, Ravaghi H. Major incident experience and preparedness in a developing country. *Disaster Med Public Health Prep*. 2013;7(3):313-8. doi: 10.1001/dmp.2012.34. [PubMed: 22851618]
3. EM-DAT database advanced search [cited 2015 March 14];2015 Available from: http://www.emdat.be/advanced_search/index.html.
4. Seyedin SH, Jamali HR. Health information and communication system for emergency management in a developing country, Iran. *J Med Syst*. 2011;35(4):591-7. doi: 10.1007/s10916-009-9396-0. [PubMed: 20703530]
5. Forouzan AS, Baradarn Eftekhari M, Falahat K, Dejman M, Heidari N, Habibi E. Psychosocial needs assessment among Earthquake survivors in Lorestan province with an emphasis on the vulnerable groups. *Glob J Health Sci*. 2013;5(4):79-84. doi: 10.5539/gjhs.v5n4p79. [PubMed: 2377724]
6. Seyedin H, Abasi Dolat Abadi Z, Sorani M, Naghdi S, Rajabfard Mazraeno F. Vulnerability assessment of general hospitals of Tehran University of Medical Sciences. *J Health Promot Manage*. 2014;3(2):65-71.
7. Tehran municipality. The neighborhoods of Tehran. Tehrann: 2015. Available from: <http://www.mytehran.ir/Default.aspx>.
8. Jahangiri K, Izadkhan YO, Lari A. Hospital safety index (HSI) analysis in confronting disasters: A case study from Iran. *Int J Health Sys Dis Manage*. 2014;2(1):44-9.
9. Kenner CA, Pressler JL. Developing a culture of emergency preparedness education in nursing. *Nurse Educ*. 2014;39(1):1-3. doi: 10.1097/NNE.000000000000003. [PubMed: 24300248]
10. Goodwin veenema T. *Disaster nursing and emergency preparedness for chemical, biological and radiological terrorism and other hazards*. 2 ed. New York: Springer Publishing Company; 2007. p. 760.
11. Sadat Sajadi H, Zaboli R. Assessing hospital disaster preparedness in Tehran: Lessons learned on disaster and mass casualty management system. *Int J Health Sys Dis Manag*. 2014;2(4):220-4. doi: 10.4103/2347-9019.144405.
12. Khani-Jazani R, Molavi-Taleghani Y, Seyedin H, Vafae-Najar A, Ebrahimipour H, Pourtaieb A. Risk Assessment of Drug Management Process in Women Surgery Department of Qaem Educational Hospital (QEH) Using HFMEA Method (2013). *Iran J Pharm Res*. 2015;14(2):495-504. [PubMed: 25901157]
13. Karimi Z, Ashktorab T, Mohammadi E, Abedi HA. Using the hidden curriculum to teach professionalism in nursing students. *Iran Red Crescent Med J*. 2014;16(3):e15532. doi: 10.5812/ircmj.15532. [PubMed: 24829784]
14. Yin H, He H, Arbon P, Zhu J. A survey of the practice of nurses' skills in Wenchuan earthquake disaster sites: implications for disaster training. *J Adv Nurs*. 2011;67(10):2231-8. doi: 10.1111/j.1365-2648.2011.05699.x. [PubMed: 21615462]
15. Schultz CH, Koenig KL, Whiteside M, Murray R, National Standardized All-Hazard Disaster Core Competencies Task F. Development of national standardized all-hazard disaster core competencies for acute care physicians, nurses, and EMS professionals. *Ann Emerg Med*. 2012;59(3):196-208 e1. doi: 10.1016/j.annemergmed.2011.09.003. [PubMed: 21982151]
16. Al Khalaileh MA, Bond E, Alasad JA. Jordanian nurses' perceptions of their preparedness for disaster management. *Int Emerg Nurs*. 2012;20(1):14-23. doi: 10.1016/j.ienj.2011.01.001. [PubMed: 22243713]
17. Pourvakhshoori N, Norouzi K, Ahmadi F, Hosseini M, Khankeh H. Nursing in disasters: a review of existing models. *Health Emerg Dis*. 2014;161-4.
18. O'Sullivan TL, Dow D, Turner MC, Lemyre L, Corneil W, Krewski D, et al. Disaster and emergency management: Canadian nurses'

- perceptions of preparedness on hospital front lines. *Prehosp Disaster Med.* 2008;**23**(3):s11-8. [PubMed:18702283]
19. Hammad KS, Arbon P, Gebbie KM. Emergency nurses and disaster response: An exploration of South Australian emergency nurses' knowledge and perceptions of their roles in disaster response. *Australasian Emerg Nurs J.* 2011;**14**(2):87-94. doi: 10.1016/j.aenj.2010.10.002.
 20. Hsu EB, Thomas TL, Bass EB, Whyne D, Kelen GD, Green GB. Healthcare worker competencies for disaster training. *BMC Med Educ.* 2006;**6**:19. doi:10.1186/1472-6920-6-19. [PubMed:16549004]
 21. Duong K. Disaster education and training of emergency nurses in South Australia. *Australas Emerg Nurs J.* 2009;**12**:86-92. doi: 10.1016/j.aenj.2010.10.002.
 22. Mokhtari J, Khademol Hosseini SM, Zarchi AKarimi A, Naimabadi T, Saghafinia M. Survey of Nurses' Acquire and Retain CPR Cognitive Knowledge and Psychomotor Skills Following CPR Training. *Trauma Month.* 2007;**12**(3):263-71.
 23. Farahani B, Abbasi Dolatabadi Z. The effect of education related to brain death and organ donation on the attitude and knowledge of Iranian nursing students. *Organ tissue and cells.* 2013;**16**:53-7.
 24. Miller P. *An Assessment of Emergency Department Staff Knowledge of Emergency Preparedness.* Kentucky: Northern Kentucky University; 2011.
 25. Landry LG, Stockton A. Evaluation of a collaborative project in disaster preparedness. *Nurse Educ.* 2008;**33**(6):254-8. doi: 10.1097/01.NNE.0000334786.15052.d9. [PubMed:18981905]
 26. Wisniewski R, Dennik-Champion G, Peltier JW. Emergency preparedness competencies: assessing nurses' educational needs. *J Nurs Adm.* 2004;**34**(10):475-80. [PubMed:15577671]
 27. Garbutt SJ, Peltier JW, Fitzpatrick JJ. Evaluation of an instrument to measure nurses' familiarity with emergency preparedness. *Mil Med.* 2008;**173**(11):1073-7. [PubMed:19055181]
 28. Worrall J. Are emergency care staff prepared for disaster? *Emerg Nurse.* 2012;**19**(9):31-7. doi: 10.7748/en2012.02.19.9.31.c8943. [PubMed:22489365]
 29. McKibbin AE, Sekula K, Colbert AM, Peltier JW. Assessing the learning needs of South Carolina nurses by exploring their perceived knowledge of emergency preparedness: evaluation of a tool. *J Contin Educ Nurs.* 2011;**42**(12):547-58. doi: 10.3928/00220124-20111003-04. [PubMed:21985073]
 30. Seyedin H, Ryan J, Sedghi S. Lessons learnt from the past and preparedness for the future: how a developing country copes with major incidents. *Emerg Med J.* 2011;**28**(10):887-91. doi:10.1136/emj.2009.090555. [PubMed:20943839]