

Mutagenic Effects of Nanosilver Consumer Products: a new Approach to Physicochemical Properties

Masomeh Heshmati^a, Sepideh ArbabiBidgoli^{b*}, Samideh Khoei^c,
Seyed Mahdi Rezayat^{b,d} and Kazem Parivar^a

^aDepartment of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran. ^bDepartment of Toxicology and Pharmacology, Faculty of Pharmacy, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran. ^cMedical Physics Department, School of Medicine, Iran University of Medical Sciences, Tehran, Iran. ^dDepartment of Pharmacology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

Abstract

Serious concerns have been expressed about potential health risks of Nano silver containing consumer products (AgNPs) therefore regulatory health risk assessment on such nanoparticles has become mandatory for the safe use of AgNPs in biomedical products with special concerns to the mutagenic potentials. In this study, we examined the inhibitory and mutagenicity effects of AgNPs in three different sizes of three colloidal AgNPs by Minimal Inhibitory concentration (MIC), Minimal Bactericidal Concentration (MBC) and Bacterial Reverse Mutation Assay (Ames test). All samples were characterized by transmission electron microscopy (TEM), X-Ray Diffraction (XRD) and Dynamic Light Scattering (DLS). DLS analysis showed lack of large agglomeration of the AgNPs and TEM results showed the spherical AgNPs with the average sizes of 15, 19.6, 21.8 nms. Furthermore the XRD analysis showed the crystalline samples with a face centered cubic structure of pure silver. Ames test results on Colloidal silver nanoparticles showed lack of any mutation in TA100, TA98, YG1029S. typhimurium strains. In addition colloidal silver nanoparticles reduced the mutation rates in all three strains in a concentration dependent manner. This finding creates a new issue in the possible antimutagenic effects of colloidal AgNPs as a new pharmaceutical product which should be considered in future studies by focusing on the physicochemical properties of AgNPs.

Keywords: Mutagenicity; Nanosilver; AgNPs; Physicochemical Property; biomedical products.

Introduction

For more than a century, Silver nanoparticles (AgNPs) were used as effective antiseptic agents, but in recent decade they have found increasing global concerns because of their strong antibacterial activities in comparison

to silver and other antiseptic compounds (1). AgNPs containing products comprising silver nanoparticles, are attracting interest for a range of biomedical and pharmaceutical applications owing to their potent antibacterial activities (2). Moreover it has recently been demonstrated that AgNP has useful anti-inflammatory effects which may accelerate wound healing (1). The key to its broad-acting and potent antibacterial activity is the multifaceted mechanism by which AgNP acts

* Corresponding author:

E-mail: Sepideharbabi@yahoo.com




Figure 1. characterization of AgNPs by Transmission Electron Microscopy; A) Representative image of A-AgNP; B) Representative image of B-AgNP; C) Representative image of AgNPs water dispersion.

on microbes. This is utilized in antibacterial coatings on medical devices to reduce the rate of nosocomial infections. Many new synthesis methods have emerged and are being evaluated for AgNP production for medical applications (2).

This property has led the global and local markets to many new healthcare and cosmetic products, wound dressings, textiles, contraceptive devices, surgical instruments, bone prostheses, water purification systems, indoor air quality management and other products from AgNPs (3). Increased utilization of nanoparticles in recent years due to significant advances in nanotechnology have an impact on industrial technology with increased risk of human exposure to nanoparticles through occupational and environmental routes (4).

The antibacterial properties of AgNPs are mainly attributed to their high surface area to volume ratio but this potential may cause their higher reactivity to macromolecules especially DNA reactions, their possible genotoxic effects, mutagen induced health risks properties and carcinogenic properties as well (5). According to the possible toxic potentials of AgNPs (6), many studies have revealed the association between genotoxicity of AgNPs and their sizes. Although different size dependent genotoxic effects from AgNP are reported by different methods in a dose-dependent manner (7), other physicochemical properties including zeta potential, shapes (rods, triangles, spherical particle) and aggregation capacities have been

considered for their genotoxic potentials (8)

Out of different genotoxicity assessment methods, the Ames test is known as the most accurate and commonly used procedure to detect genotoxic carcinogens which cause two classes of gene mutation, base pair substitution and small frameshifts (9). Although this test is essential within the current battery of assays required for genotoxicity evaluation and one of key assays recommended by the United Kingdom expert advisory Committee on Mutagenicity (7), its efficiency in genotoxicity assessment of various kinds of nanoparticles has remained inconclusive. This study aimed to compare the mutagenic effects of a wide range of doses of three different commercially available colloidal AgNPs in similar sizes on the basis of their physicochemical properties especially by Ames Mutagenicity assay.

Experimental

Materials

AgNP commercial products

Three types of prevalent commercial products of colloidal silver nanoparticles were used for this study. The first commercial product of AgNP was a yellowish-brown product, purchased from a local manufacturer in Tehran and coded as A-AgNP. According to the information provided by manufacturer, it was a water-based colloid which contained 4000 mg/ml spherical uncoated AgNP. The size and zeta

potential of this sample was provided by TEM and DLS method respectively (Figure 1).

The second commercial AgNPs was donated by another local manufacturer in Isfahan and coded as B-AgNP. According to the information provided by manufacturer, it was water-based plant coated colloid contained 187.5 mg/ml of spherical silver nanoparticles. The size and zeta potential of this sample was provided by TEM and DLS method respectively (Figure 1).

The third colloidal silver nanoparticle was purchased from US Research nanomaterials co, Ltd (TX, USA). This product was a water based colloidal AgNP containing 1000 mg/ml spherical silver nanoparticles with 99.9% purity according to the information provided by the supplier. The size and zeta potential of this sample was provided by TEM and DLS method respectively (Figure 1).

Characterization of nanoparticles

The concentration of samples was determined by Atomic Absorption Spectroscopy (AAS). AgNPs were homogeneously dispersed in sterile distilled water by sonication for 30 minutes and filtered. Transmission electron microscopy (TEM) was used to estimate the size, shape and composition of the AgNPs. The distribution of the AgNPs, hydrodynamic size and zeta potential was evaluated using Dynamic light Scattering (model ZEN3600; Molvern instrument Ltd, Tokyo, Japan). The X-ray diffraction pattern was performed in order to characterize nanoparticle structure. Experimental methods were used to optimize the concentrations of each serial diluted samples and to assure the quality of provided data. All experiments were repeated at least three times to confirm the accuracy and reproducibility of this method. Standard gold nanoparticles with pre-determined size were used to validate the instrument. Both above parameters (size and zeta potential) were measured at least three times for each sample. The data were calculated as the average size or zeta potential of AgNPs.

Salmonella strains

The Salmonella Typhymurim strains used in this study were TA100, TA98 and YG1029 a bacterial O-acetyltransferase-overproducing strain of TA100(10). The YG1029 was cloned and its activity was first described and established by

T Nohmi and colleagues(11) who provided all strains as gift, from Biological Safety Research Center co, Ltd in Tokyo-Japan (2012).

MIC and MBC levels

The Minimal Inhibitory Concentration (MIC) value is the lowest concentration of AgNPs which prevents bacterial growths. Appropriate concentrations of each AgNPs samples were prepared according to their working manual, using muellerhintonbroth in the range of 31.2-0.015 $\mu\text{g/ml}$ with serial two fold dilutions of 1, 1/2, 1.4, 1/8, 1/16, 1/32, 1/64, 1/32, 1/64, 1/128, 1/256, 1/512, 1/1024 inoculate 1×10^8 colony forming units (CFU)/ml of tester strains. After 24h from incubation at 37°C, MIC values for each strain were determined by choosing the lowest concentrations in which no growth occurs. Additionally, the minimal bactericidal concentration (MBC) values were determined by sub-culturing the content of each tube without any growth, on the Mueller Hinton agar medium and looking for any bacterial growth(12). The MIC value was considered as the concentration with 99% bacterial growth inhibition and the MBC value was considered as the concentration with 100% of the inhibitory properties in comparison to the negative control.

Bacterial Reverse Mutation Assay

The pre-incubation Ames assay was performed according to the method of OECD 471(13). Before starting the main experiments, the mentioned strains were checked for their genetic integrity by histidine/biotin dependence, histidine dependence, biotin dependence, rfa marker (crystal violet) and the presence of plasmid pKM101 (Ampicillin resistance) tests (14; 15; 16). During all these preliminary experiments, the strains were grown overnight in nutrient broth for 16-18h in incubator at 37°C with a density of $1 - 2 \times 10^8$ (CFU) ml in presence of 25 $\mu\text{g/ml}$ Ampicillin for TA98, TA100 and 10 $\mu\text{g/ml}$ Tetracycline for YG1029). The top agar was supplemented with histidine/biotin and prepared by dissolving 0.6g of agar-agar and 0.6g NaCl in 100 ml distilled water. A sterilized aqueous solution of L-histidine and D-biotin (0.5 mM/L) was added to top agar medium immediately before applications(14; 16). In the

Table 1. Physicochemical characteristics of AgNPs.

Samples	Concentration ¹ (µg/ml)	Shape ²	TEM size(nm ²)	Hydrodynamic size(nm ³)	Zeta potential mV(pH) ²	Coting Style	Specific Gravity(cP)	pH	PDI
A-AgNP	4000	different	19.6	83.7	-12	uncoated	1.14	2.5	0.608
B-AgNP	187.5	spherical	21.8	52.56	-24.6	Plant coated	0.8872	4	0.516
C-AgNP	1000	spherical	15	62.4	-11.29	uncoated	ND ⁴	7	ND

¹Determined by Atomic Absorption Spectroscopy (AAS)

²Determined by Transmission Electron Microscopy (TEM)

³Determined by Dynamic Light Scattering (DLS)

⁴Not determined

next step, 100 µl of overnight cultured bacteria with concentration of $1 - 2 \times 10^8$ CFU/ml were incubated at 37°C for 45 min in a sterile glass tube containing 500 µlit sodium phosphate buffer (0.1 M, pH7.4) with the different concentrations of the colloidal AgNPs. The concentration of AgNPs used in this study was 3.12-0.0015 µg/plate. After incubation, 2 ml of Top agar supplemented with histidine/biotin (kept in 45°C water bath) and added to the mixture and mixed for 3 s using a vortex mixer, then poured on a plate of minimal glucose agar media(16). Plates were incubated for 48-72 h at 37°C and the revertant colonies were counted. Three equal plates were used for each concentration and each experiment was repeated three times to get maximum accuracy and reproducibility for this sensitive method(17).

Data analysis

For the Ames mutagenicity assay, positive responses required a dose-related increase in the number of revertant colonies/plate for each of strains and sodium azide at 5 µg/ml concentration (Sigma Aldrich, cat number: 438456) was used as positive control. Negative response was defined as no concentration related increase in the number of revertant colonies and distilled water was used as negative control in this study. Also, a positive control was defined as an agent with defined concentration which can double the number of revertant colonies/plate in comparison to the vehicle control(17). The Mutagenic Index (M.I) was described for each assay and calculated as the ratio between number of histidine revertants induced per plate of the test sample and spontaneous revertants of the negative control(18).

Statistical analysis was performed using IBM SPSS statistics software version 20 and one-way

ANOVA (LSD) was used to compare the colony counts of each plate in different concentrations and groups. Multifactorial analysis was carried out by considering the concentration as the main factor and p-values 0.05 was considered as statistically significant changes.

Results and Discussion

AgNPs characterization

As described above, three samples of commercial AgNPs were characterized by TEM, DLS and XRD methods (Table 1). The average particle size of aqueous colloidal silver nanoparticles A-AgNP, B-AgNP and C-AgNP was 19.6, 21.8 and 15 nm respectively. Shapes of AgNPs with 21.8 and 15 nm were spherical and different shapes were observed in AgNPs with 19.6 nm (Figure 1). All nanoparticle suspensions were stable and monodispersed with normal pH. The XRD showed crystalline pattern in all three samples. All samples had a face centered cubic structure of pure silver (Table 1)

MIC and MBC determination by broth serial macrodilution method

The inhibitory effects of different concentrations of AgNPs on three Salmonella strains were examined and described in Table 2-4. Most of our experiments showed clearly that MIC and MBC levels were equal in all three samples. MIC and MBC of A-AgNP (19.6 nm) in TA100, TA98 and YG1029 were 3.9, 3.9, 7.81 µg/ml respectively. MIC and MBC of B-AgNP (21.8 nm) in TA100, TA98 and YG1029 were also 3.9, 7.81, 15.6 µg/ml respectively. MIC and MBC of C-AgNP (15 nm) was equal to 3.9 µg/ml in TA98 but in TA100 and YG1029 were 3.9, 1.9 and 7.81, 3.9 µg/ml respectively.

Table 2. Mutagenicity of A-AgNPs in *S. Typhimurium* tester strains (TA98, TA100, YG1029)¹.

Mutagenic Index (SEM ± Mean)			Number of revertant colonies/Plate (SEM ± Mean)			Dose (µg/plate)
MI _{YG1029}	MI _{TA100}	MI _{TA98}	YG1029	TA100	TA98	
1.00 ± 0.00*	1.00 ± 0.00	1.00 ± 0.00	55.3 ± 2.6	197 ± 2.08	95.6 ± 2.9	0.00
0.68 ± 0.02*	0.86 ± 0.01*	0.79 ± 0.03*	37.3 ± 1.2	170 ± 1.2	±1.769.6	0.0015
0.84 ± 0.03*	0.77 ± 0.006*	0.75 ± 0.02*	46.3 ± 1.4	153 ± 1.73	2.03 73.3±	0.003
0.87 ± 0.05*	0.79 ± 0.008*	0.58 ± 0.02* ³	48.3 ± 0.88	156.3 ± 1.73	2.1 54.3±	0.006
0.9 ± 0.05*	0.82 ± 0.004*	1.00 ± 0.03	49.6 ± 0.66	162.3 ± 1.85	96.3 ± 2.02	0.012
0.64 ± 0.02*	0.83 ± 0.002*	0.85 ± 0.01*	35.6 ± 0.88	164.3 ± 1.3	79 ± 2.6	0.024
0.73 ± 0.01*	0.96 ± 0.003 ⁴	0.8 ± 0.01*	40.6 ± 1.2	190.3 ± 1.4	74.3 ± 2.02	0.048
0.76 ± 0.05*	0.87 ± 0.003	0.84 ± 0.08*	42.3 ± 0.88	172.6 ± 3.2	72.3 ± 1.6	0.097
0.84 ± 0.04*	0.89 ± 0.01	0.98 ± 0.03	46.6 ± 1.2	176.3 ± 1.6	91 ± 2.08	0.195
0.67 ± 0.04*	ND	ND	37 ± 1.1	T	T	0.39
ND			T			0.78
						1.56
						3.12
13.1 ± 0.35	4.8 ± 0.009	9.75 ± 0.16	726 ± 13.1	948 ± 10.7	926.6 ± 2.08	Positive control

¹Note: Above data represents the mean number of revertant colonies ± SEM from three independent experiments, each repeated three times.

²Note: T denotes the toxic concentration of A-AgNP. Doses higher than T could not be evaluated against; the positive control (sodium azide with 5 µg/ml) and control negative (distilled water) and described as Not Determined (ND).

³Highest inhibitory level was 0.58 ± 0.02.

*p ≤ 0.001, p ≤ 0.05 vs control.

Mutagenic potentials of AgNPs by Ames test

As described above, bacterial mutagenicity was assessed in *S. typhimurium* tester strain TA98 for detection of frameshift mutation, TA100 for measurement of base pair substitution and YG1029, that was derived from TA100, with the enzymatic activity of O-acetyltransferase for its point mutation capacities via metabolic activation. Details are described below:

Mutagenic Potentials of A-AgNPs

Significant increase in the number of revertant colonies was detected in positive control plates and the number compared with negative control (≤ 0.001). The Mutagenicity Index (MI) was recorded as 9.75 ± 0.16 for TA98, 4.8 ± 0.009 for TA 100 and 13.1 ± 0.35 for YG1029. This observation indicated that the test system is sensitive and specific to the mutagenic potentials of sodium azide in all three tester strains (Table 2).

The number of colonies and the MIs were calculated for 12 dilutions of A-AgNP. Concentrations upper than 0.39 µg/plate were considered toxic for TA98 and concentrations

upper than 0.78 µg/plate were considered toxic for TA100 and YG1029 (Table 2).

Out of 12 evaluated concentrations of A-AgNP, 8 concentrations (0.0015-0.0195 µg/plate) showed lack of mutagenic potentials when compared with positive and negative controls. Significant lower colony numbers and MIs in above dilutions in comparison to the colony numbers in TA98 (95.6 ± 2.9), TA100 (197 ± 2.08) and YG1029 (55.3 ± 2.6) showed inhibitory effects of Nanosilver with determined physicochemical properties in these strains. Moreover YG1029 showed more sensitivity to higher concentrations of A-AgNP in this study (Table 2).

Mutagenic Potentials of B-AgNPs

Significant increase in the number of revertant colonies was detected in positive control plates when the number compared with negative control (≤ 0.001). The Mutagenicity Index (MI) was recorded as 9.2 ± 0.13 for TA98, 4.5 ± 0.14 for TA 100 and 7.5 ± 0.19 for YG1029. This observation indicated that

Table 3. Mutagenicity of B-AgNP in *S. Typhimurium* tester strains (TA98, TA100, YG1029)¹.

Mutagenicity Index (SEM±Mean)			Number of revertant colonies/ Plate (SEM±Mean)			Dose (µg/plate)
MI _{YG1029}	MI _{TA100}	MI _{TA98}	YG1029	TA100	TA98	
1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	72 ± 1.4	196.6 ± 8.81	100 ± 0.57	0.00
0.02 ± 0.76*	0.78 ± 0.03*	0.78 ± 0.03*	55.3 ± 1.2	153 ± 1.7	71.3 ± 1.45	0.0015
0.72 ± 0.01*	0.88 ± 0.04*	0.88 ± 0.04*	52.3 ± 1.2	173.3 ± 2.02	61.3 ± 1.45	0.003
0.68 ± 0.01*	0.73 ± 0.04*	0.73 ± 0.04*	50 ± 1.5	143.3 ± 2.02	64.3 ± 2.1	0.006
0.79 ± 0.06*	0.98 ± 0.03*	0.98 ± 0.03*	57 ± 1.2	192.3 ± 1.8	73 ± 1.15	0.012
0.86 ± 0.01*	0.82 ± 0.03*	0.82 ± 0.03*	62 ± 2.1	161.3 ± 0.8	82 ± 1.52	0.024
0.79 ± 0.02*	0.89 ± 0.03*	0.89 ± 0.03*	58 ± 2.6	175.3 ± 1.4	70.6 ± 1.2	0.048
0.65 ± 0.02*	0.97 ± 0.03*	0.97 ± 0.03*	48 ± 2.5	192 ± 1.5	84 ± 1.73	0.097
0.82 ± 0.02*	0.92 ± 0.04*	0.92 ± 0.04*	59 ± 2.9	180 ± 1.4	94.3 ± 2.4	0.195
0.84 ± 0.03*	ND	ND	61 ± 3.8	T	T	0.39
0.85 ± 0.01*			62 ± 1.5			0.78
ND			T			1.56
						3.12
7.5 ± 0.19	4.5 ± 0.14	9.2 ± 0.13	580 ± 20.8	893 ± 12	926.6 ± 14.5	Positive control

¹Note: Above data represents the mean number of revertant colonies ± SEM from three independent experiments, each repeated three times.

²Note: T denotes the toxic concentration of A-AgNP. Doses higher than T could not be evaluated against; the positive control (sodium azide with 5 µg/ml) and control negative (distilled water) and described as Not Determined (ND).

*p ≤ 0.001, p ≤ 0.05 vs control.

the test system is sensitive and specific to the mutagenic potentials of sodium azide in all three tester strains (Table 3). The number of colonies and the MIs were calculated for 12 dilutions of A-AgNP. Concentrations upper than 0.39 µg/plate were considered toxic for TA98 and TA100 and concentrations upper than 1.56 µg/plate was considered toxic for YG1029 (Table 3).

Out of 12 evaluated concentrations of A-AgNP, 8 concentrations (0.0015-0.195 µg/plate) showed lack of mutagenic potentials when compared with positive and negative controls. Significant lower colony numbers and MIs in above dilutions in comparison to the colony numbers in TA98 (100 ± 0.57), TA196.6 (197 ± 8.81) and YG1029 (72 ± 1.4) showed inhibitory effects of Nanosilver with determined physicochemical properties in these strains. Moreover YG1029 showed more sensitivity to higher concentrations (0.39 and 0.78 µg/plate) of A-AgNP in this study (Table 3).

Mutagenic Potentials of C-AgNPs

Significant increase in the number of revertant colonies was detected in positive control plates when the number compared with negative control (≤ 0.001). The Mutagenicity Index (MI) was recorded as 9.8 ± 0.28 for TA98, 4.8 ± 0.05 for TA 100 and 8.3 ± 0.55 for YG1029.

This observation indicated that the test system is sensitive and specific to the mutagenic potentials of sodium azide in all three tester strains (Table 4). The number of colonies and the MIs were calculated for 12 dilutions of C-AgNP. Concentrations upper than 0.195 µg/plate were considered toxic for TA98 and TA100 and concentrations upper than 0.39 µg/plate was considered toxic for YG1029 (Table 4).

Out of 12 evaluated concentrations of A-AgNP, 8 concentrations (0.0015-0.195 µg/plate) showed lack of mutagenic potentials when compared with positive and negative controls. Significant lower colony numbers and MIs in

Table 4. Mutagenicity of C-AgNPs in *S. Typhimurium* tester strains (TA98, TA100, YG1029)¹.

Mutagenicity Index (SEM ± Mean)			Number of revertant colonies/Plate (SEM ± Mean)			
MI _{YG1029}	MI _{TA100}	MI _{TA98}	YG1029	TA100	TA98	
1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	69.9 ± 2.4	190.3 ± 2.6	197 ± 3.2	0.00
0.64 ± 0.05*	0.9 ± 0.01*	0.52 ± 0.01*	45 ± 2.8	174.3 ± 2.3	95.6 ± 2.3	0.0015
0.75 ± 0.01*	0.8 ± 0.01*	0.66 ± 0.004*	52.6 ± 1.4	160 ± 0.5	50.6 ± 1.7	0.003
0.97 ± 0.02	0.7 ± 0.01*	0.85 ± 0.02*	67.6 ± 1.2	143.3 ± 0.3	63.6 ± 0.8	0.006
0.87 ± 0.03	0.7 ± 0.001*	0.78 ± 0.01*	61.3 ± 4.09	139.6 ± 2.02	81.3 ± 1.00	0.012
0.71 ± 0.06*	0.7 ± 0.007*	0.94 ± 0.01	49.3 ± 2.72	151.6 ± 0.6	75 ± 1.2	0.024
0.93 ± 0.017	0.8 ± 0.01*	0.87 ± 0.01*	65.3 ± 3.17	170 ± 0.57	90.6 ± 1.4	0.048
0.99 ± 0.06	0.7 ± 0.01*	0.97 ± 0.02	69.3 ± 2.02	135 ± 2.6	83.3 ± 1.00	0.097
0.85 ± 0.02	0.7 ± 0.01*	0.92 ± 0.04	59.3 ± 0.88	143 ± 1.7	93 ± 0.8	0.195
0.70 ± 0.05*	ND	ND	49 ± 2.3	T	T	0.39
ND			T			0.78
						1.56
						3.12
8.3 ± 0.55	4.8 ± 0.05	9.8 ± 0.28	580 ± 25.1	926.6 ± 20.8	943.3 ± 17.6	Positive control

¹Note: Above data represents the mean number of revertant colonies ± SEM from three independent experiments, each repeated three times.

²Note: T denotes the toxic concentration of A-AgNP. Doses higher than T could not be evaluated against; the positive control (sodium azide with 5 µg/ml) and control negative (distilled water) and described as Not Determined (ND).


*p ≤ 0.001, p ≤ 0.05 vs control.

above dilutions in comparison to the colony numbers in TA98 (197 ± 3.2), TA100 (190.3 ± 2.6) and YG1029 (69.9 ± 2.4) showed inhibitory effects of Nanosilver colloidal solution with determined physicochemical properties in these strains. Moreover YG1029 showed more sensitivity to a higher concentrations 0.39 µg/plate of C-AgNP in this study (Table 4).


Our research focused on genotoxic effects of silver nanoparticles (AgNPs), which can be utilized in biomedical research, pharmaceutical products and environmental cleaning applications (19). Despite the widespread use of AgNPs in a wide range of biomedical products as a new group of health products, there are numerous knowledge gaps regarding their possible toxic potentials especially their mutagenic effects in human (8), (7). We described previously systemic toxic effects of AgNPs through dermal application in animal models (8), (1), (3) but limited studies on genotoxic effects of AgNPs with dermal exposure has motivated us to continue our toxicity evaluations in this new area and this new bacterial settings for nanomaterials.

We defined three distinctive statistical groups


as “A”, “B”, and “C” with 15-22 nm size and compared each one with negative control (Table 2-4) and positive control (Figure 2:2a, 2b, 2c). The rationale of this size selection was based on the results of one recent study which had introduced 20 nm AgNPs much more toxic than larger sizes due to their higher dissolution rate and larger surface area to volume ratio (20). Revertant colonies didn't induced by the samples in a wide range of nontoxic concentrations but low AgNP concentrations showed significant inhibitory effects on TA98, TA100 and YG1024. In accordance to mentioned results, larger size AgNPs with 40-50 nm diameter (7) and lower sizes of AgNP with 5 nm diameter (17) could stop the growth of TA100 and TA98, in a dose-dependent pattern but as a novel issue lower levels of revertant colonies even lower than the negative control responded properly. Although this preliminary finding suggests again the same result as before in a new range size, the clear and significant microbial inhibitory effects of all three samples (p < 0.001) suggest their possible antimutagenic effects of AgNPs with certain physicochemical properties which should be



2a: Mutagenicity of 19.6 nmAgNPs (A-AgNP)



2b: Mutagenicity of 21.8 nmAgNPs (B-AgNP).



2c: Mutagenicity of 15 nmAgNPs (B-AgNP).

Figure 2. Comparison of mutagenic effects of AgNPs with positive control in *S. typhimurium* tester strains (TA98, TA100, YG1029).

considered for future experiments on certain mutagenic agents.

Other than the impact of size in mutagenic capacity, we considered the direct role of concentration and the indirect roles of shape, coating, zeta potential, PDI, pH and viscosity in this classic bioassay (16). The role of nanoparticle shapes e.g. rods, triangles and spherical particles (21) on bacterial toxicity has been described before in concentrations lower than MIC and MBC values (22). The shape of nanoparticles may modify the toxicity of AgNPs by changing the extents of Ag⁺ ions (22) but other than the role of shape, impact of coating was considered for gram negative bacterium *Pseudomonas putida* and results showed no simple clear-cut relation between the toxicity of the different particles and their shapes and coatings. Actually the ion release kinetics could be considered as one of the major factors for genotoxic effects of AgNPs, an issue which was not determined by us (23), (24).

Our result showed weak effects of physicochemical properties on mutagenicity potentials of AgNPs as well as no simple clear-cut relation between coating and zeta potential of materials. There are several explanations for these negative Ames results. Some studies showed that the negative results may be due to the inability of AgNPs to penetrate through the bacteria cell wall and lack of bacterial cells to perform endocytosis and incorporate them (7; 17; 15) but the underlying factor was the agglomeration of AgNPs which makes the particles too large to transport through the pores in bacterial cell wall. This lack of uptake could potentially lead to false negative results.

Colloidal silver nanoparticles did not induce mutations in TA100, TA98, YG1029 *S. typhimurium* strains, so far they reduced the rate of mutations in all three strains in a concentration dependent manner, a new issue which should be analyzed in next studies to assess their antimutagenic potentials. Moreover these data suggest the technical weakness of Ames test to evaluate the mutagenicity of the AgNPs. It seems that more genotoxic methods e.g. micronucleus assay, comet assay, chromosome aberration assay are necessary to assess the nanoparticle genotoxicity and their distinctive characteristics

including size, shape, zeta potential, solubility, surface area.

Acknowledgement

This project was financially supported by Iran Nano Initiative Council (INIC) and was conducted in Islamic Azad University, Pharmaceutical Sciences Branch, Toxicogenomics lab. We wish to thank Prof. Takehiko Nohmi, Ph.D and Visiting Scientist at Scientist Emeritus, Biological Safety Research Center, National Institute of Health Sciences in Tokyo-Japan and his team for providing the strains in this study as gift. Authors are also thankful from Hossein Jamalifar, Microbiology Lab, Faculty of Pharmacy, Tehran University of Medical Sciences for his technical assistance and support.

References

- (1) Arbabi Bidgoli S, Mahdavi M, Rezayat M, Korani K, Amani A and Ziarati P. Toxicity assessment of nanosilver wound dressing in wistar rat. *Acta Med. Iran.* (2013) 51: 203-208.
- (2) Arora S, Jain J, Rajwade JM and Paknikar KM. Interactions of silver nanoparticles with primary mouse fibroblasts and liver cells. *Toxicol. Appl. Pharmacol.* (2009) 236: 310-318.
- (3) Korani M, Rezayat M and Arbabi Bidgoli S. Sub-chronic dermal toxicity of silver nanoparticles in guinea pig: special emphasis to heart, bone and kidney toxicities. *Iran. J. Pharm. Res.* (2013) 12: 511-519.
- (4) Song M, Shan Li Y, Kasai H and Kawai K. Metal nanoparticle-induced micronuclei and oxidative DNA damage in mice. *J. Clin. Biochem. Nutr.* (2012) 50: 211-216.
- (5) Wijnhoven S, Peijnenburg W, Herberts A, Hagens W, Oomen A, Heugens E, Roszek B, Bisschops J, Gosens I, Demeent D, Dekkers S, Dejong D, Van zijverden M and Egeertsma R. Nano-silver - a review of available data and knowledge gaps in human. *Nanotoxicol.* (2009) 3: 109-138.
- (6) Faedmaleki F, H Shirazi F, Salarian AA, Ahmadi Ashtiani H and Rastegar H. Toxicity Effect of Silver Nanoparticles on Mice Liver Primary Cell Culture and HepG2 Cell Line. *Iran J Pharm Res.* 2014 Winter; 13: 235-42.
- (7) Kim H, Park Y, Shin D, Min Oh S and Hyuck Chung K. Appropriate *in-vitro* methods for genotoxicity testing of silver nanoparticles. *Environ. Health Toxicol.* (2013) 28: 1-8.
- (8) Ames BN, Lee FD and Durston WE. An improved bacterial test system for the detection and classification

- of mutagens and carcinogens. *Proc. Natl. Acad. Sci. USA* (1973) 70: 782-786.
- (9) Sato G, Asakura S, Hakura A, Tsutsui-Hiyoshi Y, Kobayashi N and Tsukidate K. Assessment of potential mutagenic activities of a novel benzothiazole MAO-A inhibitor E2011 using *Salmonella typhimurium* YG1029. *Mutat. Res.* (2000) 472: 163-9.
 - (10) Watanabe M, Igarashi T, Kaminuma T, Sofuni T and Nohmi T. *N*-hydroxyarylamine *O*-acetyltransferase of *Salmonella typhimurium*: proposal for a common catalytic mechanism of arylamine acetyltransferase enzymes. *Environ. Health Perspect.* (1994) 102: 83-89.
 - (11) Wendline N and Gasser R.A. macrodilution well method for minimum inhibitory concentration and minimum bactericidal concentration determination of antibacterial against *Borrelia burgdorferi in-vitro*. *Spirochetes and Tick-borne Dis.* (1997) 4:7-10.
 - (12) Ames, B.N., McCann, J. and Yamasaki, E. Methods for Detecting Carcinogens and Mutagens with the Salmonella/Mammalian-Microsome Mutagenicity Test. *Mutation Res.* (1975), 31,347-364.
 - (13) Kumar A, Pandey AK, Singh SS, Shanker R and Dhawan A. Cellular uptake and mutagenic potential of metal oxide nanoparticles in bacterial cells. *Chemosphere* (2011) 83: 1124-32.
 - (14) Doak SH, Manshian B, Jenkins GJS and Singh N. *In-vitro* genotoxicity testing strategy for nanomaterials and the adaptation of current OECD guidelines. *Mutat. Res.* (2012) 745: 104–111.
 - (15) Mortelmans K and Zeiger E. The Ames *Salmonella*/microsome mutagenicity assay. *Mutat. Res.* (2000) 455: 29-60.
 - (16) Li Y, Chen DH, Yan J, Chen Y, Mittelstaedt RA, Zhang Y, Biris AS, Heflich RH and Chen T. Genotoxicity of silver nanoparticles evaluated using the Ames test and *in-vitro* micronucleus assay. *Mutat. Res.* (2012), 745(1-2): 4-10.
 - (17) Lupi S, Marconi S, Paiaro E, Fochesato A and Gregorio P. Mutagenicity evaluation with Ames test of hydro-alcoholic solution of terpenes. *J. Prev. Med. Hyg.* (2009) 50: 170-174.
 - (18) Korani M, Rezayat SM, Minaee MB, Arbabi Bidgoli S, Gilani K and Adeli S. Acute and subchronic dermal toxicity of nanosilver in guinea pig. *Int. J. Nanomed.* (2011) 6: 61-8.
 - (19) Okafor F, Janen A, Kukhtareva T, Edwards V and Curley M. Green synthesis of silver nanoparticles, their characterization, application and antibacterial activity. *Int J Environ Res Public Health.* 2013; 10(10): 5221-38.
 - (20) Radniecki TS, Stankus DP, Naigh A, Nason JA and Semprini L. Influence of liberated silver from silver nanoparticles on nitrification inhibition of *Nitrosomonas europaea*. *Chemosphere* (2011) 85: 43-9.
 - (21) Matzke M, Jurkschat K and Backhaus T. Toxicity of differently sized and coated silver nanoparticles to the bacterium *Pseudomonas putida*: risks for the aquatic environment? *Ecotoxicology* (2014) 23:818-529.
 - (22) Visnapuu M, Joost U, Juganson K, Künnis-Beres K, Kahru A, Kisand V and Ivask A. Dissolution of silver nanowires and nanospheres dictates their toxicity to *Escherichia coli*. *Bio. Med. Res. Int.* (2013) 10: 1-9.
 - (23) Pan X, Redding JE, Wiley PA, Wen L, McConnell JS and Zhang B. Mutagenicity evaluation of metal oxide nanoparticles by the bacterial reverse mutation assay. *Chemosphere* (2010) 79: 113-116.
 - (24) Ahamed M, Alsalhi MS and Siddiqui MK. Silver nanoparticle applications and human health. *Clin. Chim. Acta* (2010) 411: 1841–1848.
 - (25) Sintubin L, De Gussem B, Van der Meeren P, Pycke BF, Verstraete W, Boon N. The antibacterial activity of biogenic silver and its mode of action. *Appl. Microbiol. Biotechnol.* (2011) 91: 152-162.
 - (26) Nakagawa Y, Wakuri S, Sakamoto K and Tanaka N. The photogenotoxicity of titanium dioxide particles. *Mutat. Res.* (1997) 394: 125-32.
 - (27) Mori T, Takada H, Ito S, Matsubayashi K, Miwa N and Sawaguchi T. Preclinical studies on safety of fullerene upon acute oral administration and evaluation for no mutagenesis. *Toxicol.* (2006) 225: 48-54.
 - (28) Sera N, Tokiwa H and Miyata N. Mutagenicity of the fullerene C60-generated singlet oxygen dependent formation of lipid peroxides. *Carcinogenesis* (1996) 17: 163-9.
 - (29) Flower NA, Brabu B, Revathy M, Gopalakrishnan C, Raja SV, Murugan SS and Kumaravel TS. Characterization of synthesized silver nanoparticles and assessment of its genotoxicity potentials using the alkaline comet assay. *Mutat. Res.* (2012) 742: 61-65.
 - (30) Asare N, Instanes C, Sandberg WJ, Refsnes M, Schwarze P, Kruszewski M, Brunborg G. Cytotoxic and genotoxic effects of silver nanoparticles in testicular cells. *Toxicol.* (2012) 291: 65-72.
 - (31) Dufour EK, Kumaravel T, Nohynek GJ, Kirkland D and Toutain H. Clastogenicity, photo-clastogenicity or pseudo-photo-clastogenicity: Genotoxic effects of zinc oxide in the dark, in pre-irradiated or simultaneously irradiated Chinese hamster ovary cells. *Mutat. Res.* (2006) 607: 215-24.