



## Bell's palsy following Growth hormone therapy in a patient with Prader-Willi syndrome: the first report

Setila Dalili<sup>1</sup>, Elham Bidabadi<sup>2</sup>, Babak Behnam<sup>3,4,5</sup>

<sup>1</sup>Department of Pediatric Endocrinology, Pediatric Growth Disorders Research Center, 17th Shahrivar Hospital, School of Medicine, Rasht University of Medical Sciences, Rasht, Iran

<sup>2</sup>Department of Pediatric Neurology, Pediatric Growth Disorders Research Center, 17th Shahrivar Hospital, School of Medicine, Rasht University of Medical Sciences, Rasht, Iran

<sup>3</sup>Medical Genetics Branch, National Human Genome Research Institute (NHGRI), National Institute of Health (NIH), Bethesda, Maryland, USA

<sup>4</sup>Department of Medical Genetics and Molecular Biology, Faculty of Medicine, Iran University of Medical Sciences (IUMS), Tehran, Iran

<sup>5</sup>Present address: NIH Undiagnosed Diseases Program, Common Fund, Office of the Clinical Director, NHGRI, National Institutes of Health, Bethesda, Maryland, USA

Corresponding Author: Babak Behnam

E-mail: Babak.Behnam@nih.gov; b\_behnam@yahoo.com

Genet. Mol. Res. 17 (1): gmr16039877

Received January 08, 2018

Accepted January 28, 2018

Published February 20, 2018

DOI <http://dx.doi.org/10.4238/gmr16039877>

Copyright © 2017 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution ShareAlike (CC BY-SA) 4.0 License.

**ABSTRACT.** Prader–Willi syndrome (PWS) is a complex genetic disorder with different manifestations in infancy and childhood including obesity, type 2 diabetes mellitus, mild to moderate intellectual impairment and learning disabilities. In this syndrome, growth hormone therapy improves outcomes. For the first time, here we report a 11-year-old boy with PWS who presented with three episodes of unilateral facial palsy after starting growth hormone therapy.

**KEY WORDS:** Prader-Willi syndrome, Bell's palsy, Growth Hormone

## INTRODUCTION

Prader–Willi syndrome (PWS) clinical manifestation may present in infancy and childhood with hypotonia, feeding problem, poor growth and development. Affected individuals in childhood develop as more appetite, hyperphagia, obesity, type 2 diabetes mellitus, mild to moderate intellectual impairment, learning disabilities, and some other manifestations.

Genetically, PWS occurred due to genomic imprinting resulting in dysfunctional genes in a specific region of chromosome 15. Briefly, some genes are actively expressed if only a copy is inherited from father, and a paternal chromosome 15 segment is deleted in 70% cases of PWS. Maternal uniparental disomy mainly occurred in the rest of cases (25%), besides some other rare causes for PWS including chromosomal rearrangement (translocation). Although growth hormone therapy has been recommended for PWS to improve the disease outcomes, it has some complications too (1,2).

Bell's palsy is a lower motor neuron unilateral facial paralysis or the evidence of an aural, a neurological, or a local cause. Bell's palsy has been reported upon IGF therapy in growth hormone insensitivity syndrome (GHIS), GH gene deletion, and diabetes mellitus (3,4). Also, an association and coincidence has been very recently reported between Bell's palsy and obesity (5), which the latter is a symptom of Prader-Willi syndrome too. Here, we report a 11-year-old boy with Prader-Willi syndrome. It is more likely an unfavorable effect of growth hormone therapy in children.

## CASE REPORT

A 11-year-old boy presented with unilateral facial palsy. The patient was first examined at the age of 4 years because of obesity, weakness, and poor development. At that time, and mainly due to a generalized and prolonged hypotonia, paternal genes on chromosome 15 were evaluated using an MLPA kit (MLPA-ME028-B1; MRC Holland, Amsterdam, The Netherlands) to detect copy number changes and aberrant methylation patterns in the 15q11 locus using an ABI Prism 310 genetic analyser. The evaluation was also carried out in all available family members and an aberrant methylation pattern was found in the region 15q11-q15 in the patient. Therefore, he was diagnosed with Prader Willi syndrome (PWS) either via uniparental disomy or imprinting defects, so a further molecular analysis was recommended to differentiate them using STR analysis and imprinting center sequencing.

Early diagnosis and treatment may improve quality of life in these patients, so we begin human growth hormone (HGH) administration based on Federal Drug Administration (FDA) approval at 9 years old. Linear growth, muscle tone and decreases body fat were improved by HGH replacement but a facial asymmetry was noted with a unilateral peripheral facial nerve deficit without history of trauma or seizure by 6 months after HGH replacement therapy. All other cranial nerves and his blood pressure were normal, and there was no remarkable finding in physical exam.

The laboratory tests including fasting blood glucose and insulin, lipid profiles, complete blood count, serum electrolytes, liver function tests, coagulation tests and the cerebrospinal fluid were in normal ranges. Any intracranial abnormality or evidence of mastoiditis was not obtained in cerebral axial computed tomographic (CT) scan and axial T1- and T2-weighted magnetic resonance imaging (MRI). Growth hormone was discontinued, and the patient was treated with oral hydrocortisone. However, Bell's palsy was recurred three times more despite GH discontinuation. The results of biochemical and CBC tests are summarized in the table 1. Genetic testing was approved by the local ethics committees (**Rasht** University of Medical sciences). With informed consent from the parents and in accordance with the Declaration of Helsinki, genomic DNA was extracted from peripheral blood lymphocytes, and a further genetic assay was performed in the Molecular Genetics Laboratory.

Table 1. Patient's Test Results

| Test                              | Result | NL Range       |
|-----------------------------------|--------|----------------|
| Fasting Blood Sugar (FBS)         | 90     | 80-126 mg/dL   |
| Fasting insulin level             | 12     | 1.8-24.6 mU/L  |
| <i>Lipid profiles</i>             |        |                |
| HDL                               | 50     | 36-84 mg/dL    |
| LDL                               | 110    | 64-136 mg/dL   |
| Total Cholesterol                 | 200    | 125-244 mg/dL  |
| <i>Complete blood count (CBC)</i> |        |                |
| RBC                               | 4.5    | 4.0-4.9        |
| WBC                               | 8      | 4.5-10.5       |
| Hemoglobin                        | 12     | 11.0-13.3      |
| <i>Serum electrolytes</i>         |        |                |
| Sodium                            | 137    | 135-148 mmol/L |
| Chloride                          | 110    | 102-112 mmol/L |
| Potassium                         | 4      | 3.5-5.8 mmol/L |
| Phosphorus                        | 4.5    | 3.2-6.3 mg/dL  |
| <i>Liver function tests</i>       |        |                |
| Alkaline Phosphatase              | 200    | 115-336        |
| SGOT                              | 35     | 5-40 U/L       |
| SGPT                              | 36     | 7 to 56 U/L    |
| <i>Coagulation tests</i>          |        |                |
| Prothrombin Time (PT)             | 13     | 11-15 sec      |
| Partial Thromboplastin Time (PTT) | 48     | 42-54 sec      |

## DISCUSSION

The benefits of HGH treatment in patient with Prader-Willi syndrome are well established including increased linear growth, improved body composition, decreased body fat, increase motor function and psychomotor development (1). Side effects of GH was reported in children and adolescents not including Bell's palsy.

It is the first report of Bell's palsy in PWS which was improved by the hormone discontinuous. Despite the safety profile of growth hormone treatment, Bell's palsy association with or secondary to GH therapy has been previously reported in a case as a GH unfavorable side effect or a coincident (3). It may be a co-incidence of Bell's palsy and PWS, but also it may be a consequence of GH therapy as a side effect too. A predisposition or susceptibility to Bell's palsy could also be considered in some genetic diseases such as PWS. To confirm this possible tendency, and to better understanding of Bell's palsy in Prader-Willi syndrome, further studies are recommended.

## ACKNOWLEDGEMENT

This work was partly supported by the Intramural Research Program of the National Human Genome Research Institute (NHGRI), NIH.

## CONFLICT OF INTEREST

Authors have no conflict of interest

## ESTABLISHED FACTS AND NOVEL INSIGHTS

- It is the first, here we report a 11-year-old boy with PWS and facial (Bell's) palsy after starting growth hormone therapy.
- A predisposition or susceptibility to Bell's palsy could also be considered in some genetic diseases such as PWS

## REFERENCES

- Bell J, Parker K, Swinford R, Hoffman A, et al. (2010). Long-term safety of recombinant human growth hormone in children. *The Journal of Clinical Endocrinology & Metabolism*. 95(1): 167-177. <https://doi.org/10.1007/s12325-011-0063-8>
- Carrel AL, Myers SE, Whitman BY, Eickhoff J, et al. (2010). Long-term growth hormone therapy changes the natural history of body composition and motor function in children with Prader-Willi syndrome. *The Journal of Clinical Endocrinology & Metabolism*. 95(3): 1131-1136. <https://doi.org/10.1210/jc.2009-1389>
- Toledo Del Castillo B, Kyriakos G, Roldán Martín MB (2014). Growth hormone treatment and Bell's palsy: coincidence or consequence? *An Pediatr (Barc)*. 2014 Jan;80(1):61-62.
- Clemmons DR, Moses AC, McKay MJ, Sommer A, et al. (2000). The combination of insulin-like growth factor I and insulin-like growth factor-binding protein-3 reduces insulin requirements in insulin-dependent type 1 diabetes: evidence for in vivo biological activity. *J Clin Endocrinol Metab*. 85(4):1518-1524. <https://doi.org/10.1210/jc.85.4.1518>
- Koosmanee S, Tabrizi M, Naderi N, Rad AH (2016). Parental Anthropometric Indices and Obesity in Children. *Acta Medica Iranica*. 54(4):270-275.