



The Open Medicinal Chemistry Journal

Content list available at: <https://openmedicinalchemistryjournal.com>



RESEARCH ARTICLE

Design and Synthesis of WM5 Analogues as HIV-1 TAR RNA Binders

Jenny Desantis^{1,2}, Serena Massari^{1,*}, Alice Sosic³, Giuseppe Manfroni¹, Rolando Cannalire¹, Tommaso Felicetti¹, Christophe Pannecouque⁴, Barbara Gatto³ and Oriana Tabarrini¹

¹Department of Pharmaceutical Sciences, University of Perugia, 06123 Perugia, Italy

²Department of Chemistry, Biology and Biotechnology, University of Perugia, 06123 Perugia, Italy

³Department of Pharmaceutical and Pharmacological Sciences, University of Padova, 35131, Padova, Italy

⁴Rega Institute for Medical Research, Laboratory of Virology and Chemotherapy, K.U. Leuven, K.U. Leuven, B-3000 Leuven, Belgium

Abstract:

Background:

The 6-aminoquinolone **WM5**, previously identified by us, is among the most selective small molecules known as TAR RNA binders to show anti-HIV activity.

Methods:

Starting from **WM5**, a series of analogues modified at N-1, C-6 or C-7 position was prepared by inserting guanidine or amidine groups as well as other protonable moieties intended to electrostatically bind the phosphate backbone of TAR. All the compounds were tested for their ability to inhibit HIV-1 replication in MT-4 cells and in parallel for their cytotoxicity. The active compounds were also evaluated for their ability to interfere with the formation of the Tat-TAR complex using a Fluorescence Quenching Assay (FQA).

Results:

Some of the synthesized compounds showed an anti-HIV-1 activity in the sub-micromolar range with the naphthyridone derivatives being the most potent. Three of the synthesized derivatives were able to interact with the Tat-TAR complex formation presenting K_i values improved as compared to the values obtained with **WM5**.

Conclusion:

The addition of a pyridine-based protonable side chain at the N-1 position of the quinolone/naphthyridone core imparted to the compounds the ability to interfere with Tat-TAR complex formation and HIV-1 replication.

Keywords: Anti-HIV quinolones, Tat-mediated transcription inhibitors, Antiviral agents, Medicinal chemistry, WM5, Fluorescence quenching assay.

Article History

Received: November 5, 2018

Revised: January 26, 2019

Accepted: February 5, 2019

1. INTRODUCTION

In the past decades, considerable work has been done in exploiting small non-coding hairpin RNA fragments as drug targets. However, compared with the compounds targeting proteins, the identification of molecules that selectively bind to structured RNA motifs remains an important challenge [1 - 3]. One of the most studied RNA structures is the Transactivation response element (TAR) of HIV-1 genome [4 - 9], a short stem-bulge-loop structure located in the long terminal repeat

region at the 5' end of all nascent HIV-1 transcripts. Its specific interaction with the HIV-1 regulatory protein Tat is essential for viral gene expression, replication and pathogenesis [10, 11].

TAR RNA is extremely conserved among viral isolates making it a very attractive target. During the last two decades, many small molecules, aminoglycosides-conjugates, Tat-based peptidomimetics, and peptides have been reported as TAR binders [12 - 19].

Among the small molecules known as TAR RNA binder, the 6-aminoquinolone **WM5** Fig. (1), which was developed in our lab, inhibits the Tat-mediated transactivation by interfering selectively with the bulge region of TAR RNA [20].

* Address correspondence to this author at the Department of Pharmaceutical Sciences, University of Perugia, Via del Liceo 1, 06123, Perugia, Italy; Tel: +390755855134; E-mail: serena.massari@unipg.it

Numerous derivatives have been designed and synthesized starting from **WM5** with the aim to gain Structure-Activity Relationship (SAR) insights, reduce the cytotoxicity and further investigate the molecular mechanism of action [21]. As a consequence, many compounds endowed with good HIV-1 and HIV-2 inhibitory activity in both chronically infected and acutely infected cells have already been identified. However, a direct correlation between the anti-HIV activity and their ability to interfere with the Tat-TAR RNA complex was not always observed [22]. On the other hand, when the quinolone nucleus lacking both C-7 arylpiperazine and C-3 carboxylic functions was properly functionalized with protonable groups at the C-2 position, a series of 2-phenylquinolones was obtained showing an improved ability to displace Tat-TAR complex but at the expense of antiviral activity [23, 24].

Here, we came back to **WM5** preparing a series of analogues designed to enhance its ability to recognize the viral nucleic acid TAR RNA. Considering that TAR plays its essential role in the Tat-mediated transcription by specifically binding Tat and that in the Tat-TAR complex the arginine rich basic portion of Tat is responsible for the TAR binding, we decided to modify **WM5** by inserting guanidine or amidine groups as well as other protonable moieties intended to bind electrostatically the phosphate backbone of TAR. In the present paper, the synthesis of a series of **WM5** analogues modified at N-1, C-6 or C-7 position Fig. (1), and their anti-HIV-1 activity along with their cytotoxicity in MT-4 cells, are reported. The active compounds were also evaluated in a Fluorescence Quenching Assay (FQA) to determine their ability to interfere with the formation of the complex between TAR RNA and a truncated Tat peptide.

1.1. Design of WM5 Analogues

Starting from **WM5**, the C-6 amino group was replaced by a more basic guanidine in compound **1**, while at C-7 position an amidine moiety replaced the 2-pyridine ring in compound **2**, and an imidazoline ring was present at the C-4 position of the pyridine in compound **3** Fig. (1). Pyridine and (pyridinyl)piperazine moieties were inserted at the N-1 position through methyl, ethyl, or propyl chains, as in compounds **4**, **5** [25] and **6**, respectively Fig. (1). Moreover, by maintaining the propyl chain, the piperazine and the pyridine rings were alternatively deleted as in compounds **7** and **8**, respectively Fig. (1). We have previously reported that the 1,8-naphthyridone nucleus, when decorated with a 1,3-benzothiazolpiperazine moiety at the C-7 position, grants potent and selective anti-HIV activity [26, 27]. Thus, the same flexible protonable side chains were also inserted at N-1 position of the 1,8-naphthyridone scaffold, obtaining compounds **9** [25], **10** [25], **11**, and **12** Fig. (1).

2. MATERIALS AND METHODS

2.1. Chemistry

Starting materials, reagents, and solvents that were commercially available were used as supplied. The reactions

were monitored by TLC on silica gel 60F254 (Merck) and the compounds were visualized by UV and/or iodine. Flash chromatography columns were performed on Merck silica gel 60 (mesh 230-400). After extraction, organic solutions were dried using anhydrous Na₂SO₄, filtered, and evaporated to dryness at reduced pressure using a Büchi rotary evaporator. Yields are of pure products and were not optimized. Melting points were determined in capillary tubes (Büchi Electrothermal Mod. 9100) and are uncorrected. Elemental analyses were performed on a Fisons elemental analyzer, Model EA1108CHN, and the data for C, H and N are within ± 0.4% of the theoretical values. ¹H NMR and ¹³C NMR spectra were recorded at 200 MHz (Bruker Avance DPX-200) and 400 MHz (Bruker Avance DRX-400) using residual solvents such as dimethylsulfoxide (δ = 2.48) or chloroform (δ = 7.26) as an internal standard. Chemical shifts (δ) are reported in parts per million (ppm), and peak multiplicity are reported as s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), hept (heptet), m (multiplet), or brs (broad singlet). Target compounds **5** [25], **9** [26], **10** [27] have been synthesized as previously reported.

2.1.1. 6-(E)-(2,3-Di-Tert-Butoxyguanidino)-1-Methyl-4-Oxo-7-[4-(Pyridin-2-yl)Piperazin-1-yl]-1,4-Dihydroquinoline-3-Carboxylic Acid (13)

To a solution of **WM5** [20] (0.13 g, 3.29 mmol) in dry DMF (10 mL), Et₃N (0.07 g, 7.24 mmol, 0.1 mL), *N,N'*-di-(tert-butoxycarbonyl)thiourea (0.10 g, 3.45 mmol), and HgCl₂ (cat. amount) were added. The mixture was reacted at room temperature for 15 min. The solvent was subsequently concentrated under vacuum and the residue was treated with EtOH yielding a precipitate that was eliminated by filtration. The filtrate was evaporated to dryness obtaining a solid that was washed with hot EtOH (3 times), filtered off, and dried to give compound **13** (0.10 g, 10%): mp > 300 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.50 (s, 18H, CH₃), 3.05-3.25 (m, 4H, piperazine CH₂), 3.70-3.90 (m, 7H, CH₃ and piperazine CH₂), 6.60-6.75 (m, 2H, pyridine CH), 7.05 (s, 1H, H-8), 7.40 (bs, 1H, NH), 7.50 (t, *J* = 6.0 Hz, 1H, pyridine CH), 8.10 (d, *J* = 6.0 Hz, 1H, pyridine CH), 8.70 (s, 1H, H-5), 9.40 (s, 1H, H-2), 10.80 (bs, 1H, NH), 12.40 (s, 1H, COOH).

2.1.2. 6-Guanidino-1-Methyl-4-Oxo-7-(4-(Pyridin-2-yl) Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylic Acid (1)

HCl was bubbled through a solution of compound **13** (0.10 g, 1.74 mmol) in a mixture of absolute EtOH (10 mL) and CHCl₃ (10 mL), at room temperature. After 15 h, the obtained precipitate was collected by filtration, washed with a mixture of EtOH/CHCl₃ (1:1), and dried, to give **1** (0.018 g, 26%): mp > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 3.10-3.25 and 4.10-4.25 (m, each 4H, piperazine CH₂), 3.90 (s, 3H, CH₃), 7.00 (t, *J* = 6.4 Hz, 1H, pyridine CH), 7.30 (s, 1H, H-8), 7.50 (d, *J* = 9.5 Hz, 1H, pyridine CH), 8.05-8.10 (m, 2H, pyridine CH), 8.25 (bs, 1H, NH), 8.75 (s, 1H, H-5), 8.90 (s, 1H, H-2), 9.50 (bs, 1H, COOH). Anal. calcd. for C₂₁H₂₂N₆O₃: C 59.85, H 5.50, N 23.26, found: C 60.00, H 5.75, N 23.00.




Fig. (1). WM5 analogues characterized by protonable groups.

2.1.3. Ethyl 7-(4-Carbamidoylpiperazin-1-Yl)-1-Methyl-6-Nitro-4-Oxo-1,4-Dihydroquinoline-3-Carboxylate (15)

To a solution of synthon **14** [28] (0.70 g, 1.92 mmol) in dry DMSO (50 mL), methyl imidothiocarbamate sulfate (0.26 g, 0.959 mmol) was added. The reaction mixture was stirred at 120 °C for 6 h. After cooling, the precipitate obtained was filtered and dried, to give **15** (0.77 g, 99%): mp 179-181 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 3.20-3.30 and 3.60-3.70 (m, each 4H, piperazine CH₂), 3.90 (s, 3H, CH₃), 4.25 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 7.15 (s, 1H, H-8), 8.10 (bs, 3H, NH and NH₂), 8.60 (s, 1H, H-5), 8.70 (s, 1H, H-2).

2.1.4. 6-Amino-7-(4-Carbamidoylpiperazin-1-Yl)-1-Methyl-4-Oxo-1,4-Dihydroquinoline-3-Carboxylic Acid (2)

To a solution of **15** (0.15 g, 3.69 mmol) in H₂O (200 mL) and NH₄OH conc (5 mL), a solution of FeSO₄ (0.71 g, 2.58 mmol) in the minimum amount of H₂O was added and the mixture was stirred at room temperature. After 1 h, the reaction mixture was filtered over Celite® and the filtrate was

evaporated to dryness to give a residue that was crystallized from H₂O yielding ethyl 6-amino-7-(4-carbamimidoylpiperazin-1-yl)-1-methyl-4-oxo-1,4-dihydroquinoline-3-carboxylate (0.08 g, 58%), which was used in the next step without further purification. Thus, ethyl 6-amino-7-(4-carbamimidoylpiperazin-1-yl)-1-methyl-4-oxo-1,4-dihydroquinoline-3-carboxylate (0.08 g, 0.21 mmol) was dissolved in EtOH (3 mL) and 6N HCl (3 mL) was added. The reaction mixture was stirred at reflux for 7 h. After cooling, the precipitate obtained was filtrated, washed with EtOH, and purified by crystallization from MeOH/DMF, to give **2** (0.013 g, 17%): mp > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.90-3.20 and 3.60-3.90 (m, each 4H, piperazine CH₂), 3.90 (s, 3H, CH₃), 7.20 (s, 1H, H-8), 7.50 (s, 1H, H-5), 7.90 (bs, 3H, NH e NH₂), 8.80 (s, 1H, H-2). Anal. calcd. for C₁₆H₂₀N₆O₃: C 55.80, H 5.85, N 24.40, found: C 55.63, H 5.98, N 24.35.

2.1.5. 1-[4-(4,5-Dihydro-1H-Imidazol-2-Yl)Pyridin-2-Yl]Piperazine (18)

A mixture of **16** [29] (0.50 g, 26.1 mmol), ethylenediamine (0.17 g, 2.87 mmol), and *p*-toluenesulfonic acid (0.54 g, 2.87

mmol) was reacted under reflux for 3 h. After cooling, the reaction mixture was diluted with H₂O and then extracted with CHCl₃. The organic layers were dried and evaporated to dryness to give **18** (0.40 g, 66%) as an oil; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.70-2.80 and 3.20-3.30 (m, each 4H, piperazine CH₂), 3.65 (bs, 4H, imidazoline CH₂), 3.80 (bs, 1H, NH), 7.00 (d, *J* = 5.0 Hz, 1H, pyridine CH), 7.20 (s, 1H, pyridine CH), 8.20 (s, 1H, pyridine CH).

2.1.6. General Procedure for the C-7 Nucleophilic Substitution Reaction (Method A)

A mixture of the selected synthon (1.0 equiv), the appropriate base (3.0 equiv), and K₂CO₃ (3.0 equiv) in dry DMF was stirred at 40-80°C until no starting material was detected by TLC (2-72 h). After cooling, the reaction mixture was poured into ice/water yielding a precipitate which was washed with water and then with Et₂O, and further purified as reported in the description of the compounds.

2.1.7. Ethyl 7-(4-(4-(4,5-Dihydro-1h-Imidazol-2-yl)Pyridin-2-yl)Piperazin-1-yl)-1-Methyl-6-Nitro-4-Oxo-1,4-Dihydroquinoline-3-Carboxylate (19)

The title compound was obtained starting from **17** [20] and **18** using Method A (80 °C, 6 h) after crystallization from DMF in 88% yield: mp 255-256 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 3.30-3.40 and 3.70-3.75 (m, each 4H, piperazine CH₂), 3.80 (bs, 4H, imidazoline CH₂), 3.95 (s, 3H, CH₃), 4.25 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 7.05 (d, *J* = 5.0 Hz, 1H, pyridine CH), 7.10-7.15 (m, 1H, pyridine CH), 7.20 (s, 1H, H-8), 8.20 (d, *J* = 5.0 Hz, 1H, pyridine CH), 8.55 (s, 1H, H-2), 8.70 (s, 1H, H-5).

2.1.8. General Procedure for Reduction of C-6 Nitro Group (Method B)

A stirred solution of the selected 6-nitroderivative in DMF was hydrogenated over a catalytic amount of Raney nickel under atmospheric pressure at room temperature until no starting material was detected by TLC (15 min-3 h). The mixture was then filtered over Celite, and the filtrate was evaporated to dryness to afford a residue that was treated with EtOH/Et₂O yielding a solid that was filtered and dried.

2.1.9. Ethyl 6-Amino-7-(4-(4-(4,5-Dihydro-1h-Imidazol-2-yl)Pyridin-2-yl)Piperazin-1-yl)-1-Methyl-4-Oxo-1,4-Dihydroquinoline-3-Carboxylate (20)

The title compound was obtained starting from **19** using Method B (3 h) after purification by flash chromatography eluting with CH₂Cl₂/MeOH (9:1) in 58% yield: mp 300 °C (dec); ¹H NMR (200 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 3.05-3.10 and 3.50-3.55 (m, each 4H, piperazine CH₂), 3.80 (bs, 4H, imidazoline CH₂), 3.90 (s, 3H, CH₃), 4.20 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 5.25 (bs, 2H, NH₂), 7.00 (s, 1H, H-8), 7.15 (d, *J* = 5.0 Hz, 1H, pyridine CH), 7.45 (s, 1H, H-5), 7.60-7.65 (m, 1H, pyridine CH), 8.30 (d, *J* = 5.0 Hz, 1H, pyridine CH), 8.40 (s, 1H, H-2).

2.1.10. General Procedure for Basic Hydrolysis (Method C)

A suspension of selected ethyl ester (0.3 mequiv) in 4% NaOH (5 mL) was refluxed until no starting material could be detected by TLC (4-20 h). After cooling the reaction mixture was acidified to pH = 4 with 2N HCl, obtaining a precipitate which was filtered, washed with water, and purified as described below.

2.1.11. 6-Amino-7-{4-[4-(4,5-Dihydro-1h-Imidazol-2-yl)Pyridin-2-yl]Piperazin-1-yl}-1-Methyl-4-Oxo-1,4-Dihydroquinoline-3-Carboxylic Acid (3)

The title compound was obtained starting from **20** using Method C (15 h) after crystallization from DMF in 32% yield: mp > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.80-2.90 and 3.00-3.25 (m, each 4H, piperazine CH₂), 3.50 (bs, 4H, imidazoline CH₂), 3.90 (s, 3H, CH₃), 5.00 (bs, 1H, NH), 5.50 (bs, 2H, NH₂), 6.90 (s, 1H, H-8), 7.10-7.25 (m, 2H, pyridine CH), 7.50 (s, 1H, H-5), 8.25 (d, *J* = 5.0 Hz, 1H, pyridine CH), 8.75 (s, 1H, H-2), 11.20 (COOH). Anal. calcd. for C₂₃H₂₅N₇O₅: C 61.73, H 5.63, N 21.91, found: C 61.98, H 5.82, N 21.73.

2.1.12. General Procedure for the Cycloaracylation Reaction, Step 1 (Method D)

A mixture of acrylate (1.0 equiv) and appropriate amine (1.2 equiv) in a mixture of Et₂O/EtOH (3:1), was stirred at room temperature until no starting material was detected by TLC (30 min-2 h). The reaction mixture was then concentrated *in vacuo*, yielding a residue which was used in the consecutive step without further purification.

2.1.13. Ethyl 2-(2,4-Dichloro-5-Nitrobenzoyl)-3-(Pyridin-2-Ylmethylamino)Acrylate (22)

The title compound was obtained starting from **21** [30] and (pyridin-2-ylmethyl)amine using Method D (2 h) in 78% yield: mp 98-99 °C; ¹H-NMR (200 MHz, CDCl₃) δ 1.00 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 4.00 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 4.75 (d, *J* = 5.9 Hz, 2H, CH₂), 7.15-7.25 (m, 2H, pyridine CH), 7.50 (s, 1H, H-3'), 7.70-7.80 (m, 2H, pyridine CH and H-6'), 8.30 (d, *J* = 14.0 Hz, 1H, vinyl CH), 8.65-8.75 (m, 1H, pyridine CH), 11.40-11.50 (m, 1H, NH).

2.1.14. Ethyl 3-[(3-Bromopropyl)Amino]-2-(2,4-Dichloro-5-Nitrobenzoyl)Acrylate (23)

The title compound was obtained starting from **21** [30] and (3-bromopropyl)amine using Method D (30 min) as keto-enolic mixture in 66% yield: mp 66-67 °C; ¹H-NMR (200, MHz, DMSO-*d*₆) δ 0.75 and 0.90 (t, *J* = 7.0 Hz, 0.75 and 2.25H, CH₂CH₃), 2.10-2.25 (m, 2H, CH₂), 3.50-3.75 (m, 4H, CH₂), 3.90 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.20 (bs, 1H, NH), 8.10 (s, 2H, H-3' and H-6'), 8.20-8.25 (m, 1H, vinyl CH), 9.75 (bs, 0.75H, OH), 10.90 (bs, 0.25H, NH).

2.1.15. Ethyl 2-(2,4-Dichloro-5-Nitrobenzoyl)-3-[(3-(Pyridin-2-Ylamino)Propyl)Amino]Prop-2-Enoate (24)

The title compound was obtained as an oil, starting from **21** [30] and *N*-(pyridine-2-yl)propane-1,3-diamine [31] using Method D (30 min) in 29% yield; ¹H-NMR (200 MHz, CDCl₃)

δ 1.05 (t, $J = 7.1$ Hz, 3H, CH_2CH_3), 1.90-2.00 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 3.50-3.60 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 3.95-4.05 (q, $J = 7.1$ Hz, 2H, CH_2CH_3), 4.50 (bs, 1H, NH), 6.35-6.45 (m, 1H, pyridine CH), 6.60-6.70 (m, 1H, pyridine CH), 7.30-7.40 (m, 1H, pyridine CH), 7.55 (s, 1H, H-3'), 8.75 (s, 1H, H-6'), 8.20-8.30 (m, 2H, vinyl CH and pyridine CH), 11.25 (bs, 1H, NH).

2.1.16. General Procedure For The Cycloaracylation Reaction, Step 2 (Method E)

A mixture of acrylate intermediate (1.0 equiv) and K_2CO_3 (3.0 equiv) in dry DMF was heated at 80°C for 1 h. After cooling, the reaction mixture was poured into ice/water, giving a precipitate that was filtered, washed with water and then with Et_2O .

2.1.17. Ethyl 7-Chloro -6-Nitro-4 -Oxo-1- (Pyridin -2-Ylmethyl) -1,4- Dihydroquinoline- 3-Carboxylate (25)

The title compound was obtained starting from **22** using Method E in 86% yield: mp $228-230^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.30 (t, $J = 7.1$ Hz, 3H, CH_2CH_3), 4.45 (q, $J = 7.1$ Hz, 2H, CH_2CH_3), 5.40 (s, 2H, CH_2), 7.20-7.30 (m, 2H, pyridine CH), 7.55 (s, 1H, H-8), 7.65-7.75 (m, 1H, pyridine CH), 8.50-8.60 (m, 2H, pyridine CH and H-2), 8.95 (s, 1H, H-5).

2.1.18. Ethyl 1-(3-Bromopropyl)-7-Chloro-6-Nitro-4-Oxo-1,4-Dihydroquinoline-3-Carboxylate (26)

The title compound was obtained starting from **23** using Method E in 88% yield: mp $180-181^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 1.30 (t, $J = 7.0$ Hz, 3H, CH_2CH_3), 2.15-2.30 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 3.70 (t, $J = 7.5$ Hz, 2H, NCH_2), 4.25 (q, $J = 7.0$ Hz, 2H, CH_2CH_3), 4.50-4.60 (m, 2H, CH_2Br), 8.25 (s, 1H, H-8), 8.75 (s, 1H, H-5), 8.80 (s, 1H, H-2).

2.1.19. Ethyl 7- Chloro- 6-Nitro-4-Oxo-1- [3-(Pyridin-2-Ylamino) Propyl]-1,4- Dihydroquinoline-3- Carboxylate (27)

The title compound was obtained starting from **24** using Method E in 77% yield: mp $148-149^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.40 (t, $J = 7.1$ Hz, 3H, CH_2CH_3), 2.20-2.30 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 3.50-3.60 (m, 2H, NCH_2), 4.30-4.45 (m, 4H, CH_2NH and CH_2CH_3), 4.60-4.70 (m, 1H, NH), 6.40-6.50 and 6.55-6.65 and 7.40-7.50 (m, each 1H, pyridine CH), 7.75 (s, 1H, H-8), 8.00-8.10 (m, 1H, pyridine CH), 8.60 (s, 1H, H-2), 9.00 (s, 1H, H-5).

2.1.20. Ethyl 6- Nitro -4-Oxo-1- (Pyridin-2-Ylmethyl) -7-(4-(Pyridin2-yl) Piperazin-1-yl) -1,4- Dihydroquinoline -3- Carboxylate (28)

The title compound was obtained starting from **25** and 1-(2-pyridinyl)piperazine using Method A (80°C , 2 h) in 88% yield: mp $181-183^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.35 (t, $J = 7.0$ Hz, 3H, CH_2CH_3), 3.05-3.15 and 3.55-3.65 (m, each 4H, piperazine CH_2), 4.45 (q, $J = 7.0$ Hz, 2H, CH_2CH_3), 5.35 (s, 2H, CH_2), 6.60-6.70 (m, 2H, pyridine CH), 6.90 (s, 1H, H-8), 7.15-7.30 (m, 2H, pyridine CH), 7.40-7.50, 7.60-7.70, and 8.15-8.25 (m, each 1H, pyridine CH), 8.60-8.65 (m, 2H, H-2 and pyridine CH), 8.80 (s, 1H, H-5).

2.1.21. Ethyl 6-Nitro-4-Oxo-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1-[3-(4-(Pyridin2-yl)Piperazin-1-yl)Propyl]-1,4-Dihydroquinoline-3-Carboxylate (29) And Ethyl 1-(3-Bromopropyl)-6-Nitro-4-Oxo-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylate (30)

The title compounds were obtained starting from **26** and 1-(2-pyridinyl)piperazine using Method A (40°C , 72 h) after purification by flash chromatography eluting with $\text{CHCl}_3/\text{MeOH}$ (from 100:0 to 80:20) in 53% yield (compound **29**) and 23% yield (compound **30**); compound **29**: mp $89-91^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.40 (t, $J = 7.0$ Hz, 3H, CH_2CH_3), 2.05-2.10 and 2.40-2.45 (m, each 2H, CH_2), 2.50-2.60, 3.25-3.35, 3.50-3.60 and 3.75-3.85 (m, each 4H, piperazine CH_2), 4.25-4.45 (m, 4H, CH_2CH_3 and CH_2), 6.60-6.75 (m, 4H, pyridine CH), 6.90 (s, 1H, H-8), 7.45-7.55 and 8.15-8.25 (m, each 2H, pyridine CH), 8.60 (s, 1H, H-5), 8.80 (s, 1H, H-2); compound **30**: mp $209-210^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.40 (t, $J = 7.0$ Hz, 3H, CH_2CH_3), 2.30-2.50 (m, 2H, CH_2), 3.25-3.40 (m, 4H, piperazine CH_2), 3.60-3.80 (m, 6H, piperazine CH_2 and CH_2), 4.30-4.50 (m, 4H, CH_2CH_3 and CH_2), 6.60-6.75 (m, 2H, pyridine CH), 7.00 (s, 1H, H-8), 7.50 (t, $J = 6.4$ Hz, 1H, pyridine CH), 8.15-8.25 (m, 1H, pyridine CH), 8.40 (s, 1H, H-5), 8.80 (s, 1H, H-2).

2.1.22. Ethyl 6- Nitro -4-Oxo-1- [3-(Pyridin-2-Ylamino) Propyl]-7- (4-(Pyridin2-yl) Piperazin-1-yl) -1,4- Dihydroquinoline -3- Carboxylate (31)

The title compound was obtained starting from **27** and 1-(2-pyridinyl)piperazine using Method A (80°C , 3 h), in 100% yield: mp $124-125^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.40 (t, $J = 7.1$ Hz, 3H, CH_2CH_3), 2.20-2.30 (m, 2H, $\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}$), 3.10-3.20 (m, 4H, piperazine CH_2), 3.50-3.60 (m, 2H, $\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}$), 3.65-3.75 (m, 4H, piperazine CH_2), 4.25-4.40 (m, 4H, $\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}$ and CH_2CH_3), 4.75-4.85 (m, 1H, NH), 6.35-6.45 (m, 1H, pyridine CH), 6.55-6.70 (m, 3H, pyridine CH), 6.80 (s, 1H, H-8), 7.30-7.40, 7.45-7.55, 8.05-8.15, and 8.15-8.25 (m, each 1H, pyridine CH), 8.45 (s, 1H, H-5), 8.85 (s, 1H, H-2).

2.1.23. Ethyl 6-Nitro-4-Oxo-1-(3-(Piperazin-1-Yl)Propyl)-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylate (32)

Piperazine (0.08 g, 9.39 mmol) and DIPEA (0.12 g, 9.38 mmol) were added to a solution of **30** (0.13 g, 2.34 mmol) in dry DMF (5 mL), and the mixture was stirred at 80°C for 48 h. After cooling, the reaction mixture was evaporated to dryness and the residue was purified by flash chromatography eluting with $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ (from 90:10:0 to 79:20:1), to give **32** (0.08 g, 63%): mp $186-189^\circ\text{C}$; $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 1.15 (t, $J = 7.0$ Hz, 3H, CH_2CH_3), 1.90-2.00 and 2.10-2.15 (m, each 2H, CH_2), 2.20-2.30, 2.90-3.0, 3.10-3.20 and 3.70-3.80 (m, each 4H, piperazine CH_2), 4.10 (q, $J = 7.0$ Hz, 2H, CH_2CH_3), 4.20-4.30 (m, 2H, CH_2), 6.70 (t, $J = 6.5$ Hz, 1H, pyridine CH), 6.90 (d, $J = 9.3$ Hz, 1H, pyridine CH), 7.10 (s, 1H, H-8), 7.60 (t, $J = 6.5$ Hz, 1H, pyridine CH), 8.10-8.20 (m, 1H, pyridine CH), 8.60 (s, 1H, H-5), 8.70 (s, 1H, H-2).

2.1.24. Ethyl 6-Amino-4-Oxo-1-(Pyridin-2-Ylmethyl)-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylate (33)

The title compound was obtained starting from **28** using Method B (45 min) in 34% yield: mp 268-270 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.30 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.75-2.85 and 3.60-3.70 (m, each 4H, piperazine CH₂), 4.25 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 5.20 (bs, 2H, NH₂), 5.70 (s, 2H, CH₂), 6.70 (t, *J* = 5.0 Hz, 1H, pyridine CH), 6.85 (d, *J* = 8.5 Hz, 1H, pyridine CH), 7.00 (s, 1H, H-8), 7.30 (t, *J* = 5.2 Hz, 1H, pyridine CH), 7.35-7.45 (m, 2H, H-5 and pyridine CH), 7.60 and 7.80 (t, *J* = 7.0 Hz, each 1H, pyridine CH), 8.20 and 8.45 (d, *J* = 3.7 Hz, each 1H, pyridine CH), 8.75 (s, 1H, H-2).

2.1.25. Ethyl 6-Amino-4-Oxo-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1-[3-(4-(Pyridin2-yl)Piperazin-1-yl)Propyl]-1,4-Dihydroquinoline-3-Carboxylate (34)

The title compound was obtained starting from **29** using Method B (1 h) in 88% yield: mp 233-234 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.90-2.00 (m, 2H, CH₂), 2.20-2.40 (m, 6H, CH₂ and piperazine CH₂), 3.10-3.20 and 3.40-3.50 (m, each 4H, piperazine CH₂), 3.70-3.75 (m, 4H, piperazine CH₂), 4.20 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 4.45-4.50 (m, 2H, CH₂), 5.25 (bs, 2H, NH₂), 6.65-6.70 (m, 2H, pyridine CH), 6.80 and 6.90 (d, *J* = 5.0 Hz, each 1H, pyridine CH), 7.15 (s, 1H, H-8), 7.45-7.55 (m, 3H, H-5 and pyridine CH), 8.10-8.20 (m, 2H, pyridine CH), 8.50 (s, 1H, H-2).

2.1.26. Ethyl 6-Amino-4-Oxo-1-[3-(Pyridin-2-Ylamino)Propyl]-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylate (35)

The title compound was obtained starting from **31** using Method B (15 min) in 46% yield: mp 244-245 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.95-2.05 (m, 2H, NCH₂CH₂CH₂NH), 2.80-2.95 (m, 4H, piperazine CH₂), 3.20-3.30 (m, 2H, NCH₂CH₂CH₂NH), 4.60-4.75 (m, 4H, piperazine CH₂), 4.20 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 4.25-4.35 (m, 2H, NCH₂CH₂CH₂NH), 5.25 (s, 2H, NH₂), 6.40-6.50 and 6.65-6.75 (m, each 2H, pyridine CH), 7.00 (s, 1H, H-8), 7.25-7.35 (m, 1H, pyridine CH), 7.50-7.60 (m, 2H, pyridine CH and H-5), 7.85-7.95 and 8.10-8.20 (m, each 1H, pyridine CH), 8.50 (s, 1H, H-2).

2.1.27. Ethyl 6-Amino-4-Oxo-1-(3-(Piperazin-1-Yl)Propyl)-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylate (36)

The title compound was obtained starting from **32** using Method B (40 min) in 88% yield: mp 256-257 °C; ¹H NMR (200 MHz, methanol-*d*₄) δ 1.35 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.05-2.10 (m, 2H, CH₂), 2.40-2.50 (m, 6H, CH₂ and piperazine CH₂), 2.90-3.00, 3.20-3.30, and 3.70-3.80 (m, each 4H, piperazine CH₂), 4.35 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.60-4.65 (m, 2H, CH₂), 6.70 (t, *J* = 6.5 Hz, 1H, pyridine CH), 6.90 (d, *J* = 9.3 Hz, 1H, pyridine CH), 7.20 (s, 1H, H-8), 7.60 (t, *J* = 6.5 Hz, 1H, pyridine CH), 7.65 (s, 1H, H-5), 8.10-8.20 (m, 1H, pyridine CH), 8.70 (s, 1H, H-2).

2.1.28. 6-Amino-4-Oxo-1-(Pyridin-2-Ylmethyl)-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylic Acid (4)

The title compound was obtained starting from **33** using Method C (20 h) after purification by treatment with Et₂O in 64% yield: mp 287-288 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.85-2.95 and 3.65-3.75 (m, each 4H, piperazine CH₂), 5.45 (bs, 2H, NH₂), 5.90 (s, 2H, CH₂), 6.65 (t, *J* = 5.7 Hz, 1H, pyridine CH), 6.95 (d, *J* = 8.6 Hz, 1H, pyridine CH), 7.20 (s, 1H, H-8), 7.25-7.35 (m, 1H, pyridine CH), 7.50-7.60 (m, 3H, H-5 and pyridine CH), 7.80 (t, *J* = 7.1 Hz, 1H, pyridine CH), 8.20 and 8.50 (d, *J* = 3.3 Hz, each 1H, pyridine CH), 9.00 (s, 1H, H-2), 15.90 (s, 1H, COOH); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 45.08 (2C), 49.90 (2C), 58.54, 106.46, 106.88, 107.66, 108.38, 113.67, 122.78, 122.85, 123.78, 132.67, 137.85, 138.10, 142.33, 145.67, 147.55, 148.08, 150.01, 155.17, 159.43, 167.47, 176.88. Anal. calcd. for C₂₅H₂₄N₆O₃: C 65.78, H 5.30, N 18.41, found: C 70.03, H 5.32, N 18.48.

2.1.29. 6-Amino-4-Oxo-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1-[3-(4-(Pyridin2-yl)Piperazin-1-yl)Propyl]-1,4-Dihydroquinoline-3-Carboxylic Acid (6)

The title compound was obtained starting from **34** using Method C (6 h) after purification by crystallization from DMF in 20% yield: mp > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.15-2.25 and 3.0-3.15 (m, each 2H, CH₂), 3.25-3.70 (m, 10H, piperazine CH₂), 3.75-4.00 (m, 4H, piperazine CH₂), 4.25-4.45 and 4.65-4.80 (m, each 2H, piperazine CH₂ and CH₂), 5.50 (bs, 2H, NH₂), 6.75 and 6.95 (t, *J* = 6.4 Hz, each 1H, pyridine CH), 7.10 (d, *J* = 9.0 Hz, 1H, pyridine CH), 7.30 (s, 1H, H-8), 7.40 (d, *J* = 9.0 Hz, 1H, pyridine CH), 7.60 (s, 1H, H-5), 7.75 and 7.95 (t, *J* = 6.4 Hz, each 1H, pyridine CH), 8.05-8.15 (m, 2H, pyridine CH), 8.80 (s, 1H, H-2), 11.60 (bs, 1H, COOH); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 23.52, 42.75, 46.51 (2C), 49.42 (2C), 50.68 (2C), 51.34, 53.00 (2C), 106.67, 107.44 (2C), 108.13, 113.39, 114.49 (2C), 123.15, 132.23, 135.9, 137.9, 142.31, 145.50, 146.31 (2C), 151.06, 152.93, 153.60, 167.33, 176.74. Anal. calcd. for C₃₁H₃₆N₈O₃: C 65.47, H 6.38, N 19.70, found: C 65.40, H 6.70, N 19.55.

2.1.30. 6-Amino-4-Oxo-1-[3-(Pyridin-2-Ylamino)Propyl]-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylic Acid Hydrochloride (7)

The title compound was obtained starting from **35** using Method C (6 h) after purification by treatment with Et₂O in 91% yield: mp 290 °C (c); ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.00-2.10 (m, 2H, CH₂CH₂CH₂), 3.10-3.20 (m, 4H, piperazine CH₂), 3.40-3.50 (m, 2H, NCH₂), 3.90-4.00 (m, 4H, piperazine CH₂), 4.65-4.75 (m, 2H, CH₂NH), 6.75-6.85 and 6.90-7.00 (m, each 1H, pyridine CH), 7.10-7.20 (m, 1H, pyridine CH), 7.35-7.45 (m, 2H, H-8 and pyridine CH), 7.60 (s, 1H, H-5), 7.80-8.00 (m, 3H, pyridine CH), 8.10-8.20 (m, 1H, pyridine CH), 8.75 (s, 1H, H-2), 9.35 (s, 1H, NH); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 28.42, 39.45, 46.81 (2C), 49.62 (2C), 51.63, 106.79, 108.62, 112.39 (2C), 112.98, 113.36 (2C), 123.10, 133.11, 135.91, 137.90, 140.91, 144.44 (2C), 146.00, 146.55, 152.45, 152.96, 167.20, 176.64. Anal. calcd. for C₂₇H₃₀ClN₇O₃:

C 60.50, H 5.64, N 18.29, found: C 60.62, H 5.77, N 18.33.

2.1.31. 6-Amino-4-Oxo-1-(3-(Piperazin-1-Yl)Propyl)-7-(4-(Pyridin2-yl)Piperazin-1-yl)-1,4-Dihydroquinoline-3-Carboxylic Acid (8)

The title compound was obtained starting from **36** using Method C (4 h) after purification by crystallization from DMF in 60% yield: mp > 300 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.00-2.20 (m, 2H, CH₂), 3.00-3.25 and 3.30-3.55 (m, each 8H, piperazine CH₂), 3.90-4.00 and 4.50-4.60 (m, each 2H, CH₂), 6.90 (t, *J* = 6.5 Hz, 1H, pyridine CH), 7.20 (s, 1H, H-8), 7.50 (d, *J* = 9.1 Hz, 1H, pyridine CH), 7.70 (s, 1H, H-5), 7.90-8.00 (m, 2H, pyridine CH), 8.70 (s, 1H, H-2); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 23.48, 42.69, 45.32 (2C), 56.12 (2C), 50.73 (2C), 51.69, 53.25 (2C), 106.18, 107.35, 108.23, 113.45, 114.61, 123.38, 132.91, 136.12, 142.75, 145.56, 146.42, 151.18, 153.06, 167.45, 176.88. Anal. calcd. for C₂₆H₃₃N₇O₃: C 63.52, H 6.77, N 19.95, found: C 63.79, H 7.02, N 20.13.

2.1.32. Ethyl 7-Chloro-4-Oxo-1-[2-(4-(Pyridin2-yl)Piperazin-1-yl)Ethyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylate (40)

The title compound was obtained starting from **37** [32] and [2-(4-(Pyridin2-yl)Piperazin-1-yl)ethyl]amine [33] using Method D (2 h) to give intermediate **38**, followed by Method E in 56% overall yield: mp 123-124° C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.30 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.45-2.55 (m, 4H, piperazine CH₂), 2.75 (t, *J* = 6.7 Hz, 2H, CH₂), 3.40-3.50 (m, 4H, piperazine CH₂), 4.20 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.60 (t, *J* = 6.7 Hz, 2H, CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.80 (d, *J* = 8.7 Hz, 1H, pyridine CH), 7.50-7.60 (m, 1H, pyridine CH), 7.65 (d, *J* = 8.3 Hz, 1H, H-6), 8.10-8.15 (m, 1H, pyridine CH), 8.60 (d, *J* = 8.3 Hz, 1H, H-5), 8.85 (s, 1H, H-2).

2.1.33. Ethyl 7-Chloro-4-Oxo-1-[3-(4-(Pyridin2-yl)Piperazin-1-yl)Propyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylate (41)

The title compound was obtained starting from **37** [32] and [3-(4-(pyridin-2-yl)piperazin-1-yl)propyl]amine [33] using Method D (2 h) to give intermediate **39**, followed by Method E in 59% overall yield: mp 124-125 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.30 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.00 (t, *J* = 6.7 Hz, 2H, NCH₂CH₂), 2.40-2.50 (m, 6H, CH₂ and piperazine CH₂), 3.40-3.50 (m, 4H, piperazine CH₂), 4.25 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.50 (t, *J* = 6.7 Hz, 2H, NCH₂CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.80 (d, *J* = 8.7 Hz, 1H, pyridine CH), 7.50-7.60 (m, 1H, pyridine CH), 7.65 (d, *J* = 8.3 Hz, 1H, H-6), 8.10-8.15 (m, 1H, pyridine CH), 8.55 (d, *J* = 8.3 Hz, 1H, H-5), 8.90 (s, 1H, H-2).

2.1.34. Ethyl 7-[4-(1,3-Benzothiazol-2-Yl)Piperazin-1-Yl]-4-Oxo-1-[2-(4-(Pyridin2-yl)Piperazin-1-yl)Ethyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylate (42)

The title compound was obtained starting from **40** and 1-(1,3-benzothiazol-2-yl)piperazine [34] using Method A (80 °C, 24 h). After cooling the reaction mixture was poured into ice/water and extracted with CH₂Cl₂. The organic layers were

evaporated to dryness affording an oil, which was triturated with Et₂O, giving a solid which was filtered, to afford **42** in 68% yield: mp 124-125 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.25 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.50-2.60 (m, 4H, piperazine CH₂), 2.75 (t, *J* = 6.7 Hz, 2H, CH₂), 3.40-3.50, 3.65-3.75, and 3.80-3.90 (m, each 4H, piperazine CH₂), 4.20 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.75 (t, *J* = 6.7 Hz, 2H, CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.80 (d, *J* = 8.7 Hz, 1H, pyridine CH), 7.00-7.10 (m, 2H, H-6 and benzothiazole CH), 7.25 (t, *J* = 7.4 Hz, 1H, benzothiazole CH), 7.40-7.50 (m, 2H, pyridine CH and benzothiazole CH), 7.75 (d, *J* = 7.9 Hz, 1H, benzothiazole CH), 8.05-8.10 (m, 1H, pyridine CH), 8.20 (d, *J* = 8.8 Hz, H-5), 8.60 (s, 1H, H-2).

2.1.35. Ethyl 7-[4-(1,3-Benzothiazol-2-Yl)Piperazin-1-Yl]-4-Oxo-1-[3-(4-(Pyridin2-yl)Piperazin-1-yl)Propyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylate (43)

The title compound was obtained starting from **41** and 1-(1,3-benzothiazol-2-yl)piperazine [34] using Method A (80 °C, 24 h) in 63% yield: mp 140-141 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.30 (t, *J* = 7.0 Hz, 3H, CH₂CH₃), 2.00 (t, *J* = 6.7 Hz, 2H, NCH₂CH₂), 2.40-2.50 (m, 6H, CH₂ and piperazine CH₂), 3.40-3.50, 3.65-3.75 and 3.85-3.95 (m, each 4H, piperazine CH₂), 4.20 (q, *J* = 7.0 Hz, 2H, CH₂CH₃), 4.40 (t, *J* = 6.7 Hz, 2H, NCH₂CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.80 (d, *J* = 8.7 Hz, 1H, pyridine CH), 7.00-7.10 (m, 2H, H-6 and benzothiazole CH), 7.25 (t, *J* = 7.4 Hz, 1H, benzothiazole CH), 7.40-7.50 (m, 2H, pyridine CH and benzothiazole CH), 7.75 (d, *J* = 7.9 Hz, 1H, benzothiazole CH), 8.05-8.10 (m, 1H, pyridine CH), 8.20 (d, *J* = 8.8 Hz, H-5), 8.60 (s, 1H, H-2).

2.1.36. 7-[4-(1,3-Benzothiazol-2-Yl)Piperazin-1-Yl]-4-Oxo-1-[2-(4-(Pyridin-2-Yl)Piperazin-1-Yl)Ethyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylic Acid (11)

The title compound was obtained starting from **42** using Method C (2 h) after purification by crystallization from DMF in 41% yield: mp 294-295 °C (dec.) °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.50-2.60 (m, 4H, piperazine CH₂), 2.75 (t, *J* = 6.7 Hz, 2H, CH₂), 3.30-3.40, 3.65-3.75, and 3.90-4.00 (m, each 4H, piperazine CH₂), 4.65 (t, *J* = 6.7 Hz, 2H, CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.80 (d, *J* = 8.7 Hz, 1H, pyridine CH), 7.05 (t, *J* = 7.7 Hz, 1H, benzothiazole CH), 7.20-7.30 (m, 2H, H-6 and benzothiazole CH), 7.40-7.50 (m, 2H, pyridine CH and benzothiazole CH), 7.75 (d, *J* = 7.9 Hz, 1H, benzothiazole CH), 8.05-8.10 (m, 1H, pyridine CH), 8.30 (d, *J* = 8.8 Hz, H-5), 8.80 (s, 1H, H-2), 15.50 (s, 1H, COOH); ¹³C NMR (101 MHz, DMSO-*d*₆) δ 44.21 (2C), 45.27, 47.86 (4C), 52.82 (2C), 56.23, 107.61, 107.81, 108.41, 111.50, 113.54, 119.20, 121.75, 121.88, 126.53, 130.90, 136.42, 137.98, 147.99, 149.62, 150.01, 152.80, 159.29, 159.46, 166.63, 168.47, 177.04. Anal. calcd. for C₃₁H₃₂N₈O₃S: C 62.40, H 5.41, N 18.78, found: C 62.25, H 5.70, N 18.65.

2.1.37. 7-[4-(1,3-Benzothiazol-2-Yl)Piperazin-1-Yl]-4-Oxo-1-[3-(4-(Pyridin-2-Yl)Piperazin-1-Yl)Propyl]-1,4-Dihydro-1,8-Naphthyridine-3-Carboxylic Acid (12)

The title compound was obtained starting from **43** using Method C (2 h) after purification by crystallization from DMF

in 16% yield: mp 256-257 °C (dec); ^1H NMR (400 MHz, DMSO- d_6) δ 2.00 (t, J = 6.7 Hz, 2H, CH₂), 2.25-2.35 (m, 6H, CH₂ and piperazine CH₂), 3.25-3.35, 3.65-3.75 and 3.95-4.00 (m, each 4H, piperazine CH₂), 4.50 (t, J = 6.7 Hz, 2H, CH₂), 6.60-6.65 (m, 1H, pyridine CH), 6.75 (d, J = 8.7 Hz, 1H, pyridine CH), 7.05 (t, J = 7.7 Hz, 1H, benzothiazole CH), 7.20-7.30 (m, 2H, H-6 and benzothiazole CH), 7.40-7.50 (m, 2H, pyridine CH and benzothiazole CH), 7.75 (d, J = 7.9 Hz, 1H, benzothiazole CH), 8.05-8.10 (m, 1H, pyridine CH), 8.30 (d, J = 8.8 Hz, H-5), 9.00 (s, 1H, H-2), 15.50 (s, 1H, COOH); ^{13}C NMR (101 MHz, DMSO- d_6) δ 23.62, 42.93, 46.48 (2C), 48.31 (2C), 49.16 (2C), 51.42, 52.87 (2C), 107.53, 108.16, 108.43, 111.62, 113.48, 119.1, 122.15, 122.88, 126.41, 131.10, 136.72, 138.22, 148.34, 150.21, 150.34, 152.98, 159.75, 159.88, 166.89, 169.17, 177.25. Anal. calcd. for C₃₂H₃₄N₈O₃S: C 62.93, H 5.61, N 18.35, found: C 62.98, H 5.60, N 18.35. .

2.2. Biology

2.2.1. In vitro Anti-HIV Assays

The evaluation of the antiviral activity of the target compounds against HIV-1 strain III_B in MT-4 cells was performed using the MTT assay as previously described [35, 36]. Mock-infected cells were used to assess the cytotoxic effects of the test compounds.

2.2.2. Fluorescence Quenching Assay (FQA)

The effect of tested quinolones on the Tat-TAR complex was evaluated using an FQA, a FRET-based competition assay, as previously described [23], using the Tat-derived peptide labelled with the donor fluorescein at its N terminus and the 29 nt wtTAR labeled at its 3'-end with a dabcyI moiety (quencher).

3. RESULTS AND DISCUSSION

3.1. Chemistry

The introduction of the guanidine group at C-6 position in the quinolone derivative **1** was accomplished by reacting **WM5** [20] with *N,N'*-di-(tert-butoxycarbonyl)thiourea in the presence of HgCl₂ followed by Boc deprotection of intermediate **13** under acid conditions Scheme (1).


The synthesis of the C-7 4-carbamimidoylpiperazine derivative **2** started by reacting derivative **14** [28] with methyl imidothiocarbamate sulfate to give intermediate **15**, which,

after reduction of the nitro group accomplished with FeSO₄ and NH₄OH followed by acid hydrolysis of the ethyl ester intermediate, led to the target compound **2** Scheme (2).


The C-7 imidazolinpyridinipiperazine derivative **3** was synthesized, as outlined in Scheme (3), by reacting intermediate **17** [20] with the 1-[4-(4,5-dihydro-1*H*-imidazol-2-yl)pyridin-2-yl] piperazine **18** in DMF, affording intermediate **19**. Nitro derivative **19** was then catalytically reduced to amino derivative **20** and finally hydrolysed under basic conditions to give the target compound **3**. The required building block **18** was in turn prepared through the reaction of nitrile derivative **16** [29] with ethylenediamine in presence of *p*-toluenesulfonic acid.

The synthesis of quinolone derivatives **4** and **6-8** Scheme (4) was accomplished through a cycloaracylation procedure. Thus, acrylate **21** [30] was reacted with (pyridin-2-ylmethyl) amine, (3-bromopropyl) amine, and *N*-(pyridin-2-yl) propane-1,3-diamine in Et₂O/EtOH mixture at room temperature to give intermediates **22-24**, which by treatment with K₂CO₃ in dry DMF at 80 °C afforded synthons **25-27**. These derivatives were then subjected to nucleophilic reaction with 1-(2-pyridinyl) piperazine in dry DMF, to give intermediates **28-31**. In the case of compound **26**, the nucleophilic reaction led to the mixture of the di-substituted **29** and mono-substituted **30**, which were easily isolated by chromatographical purification. Intermediate **30** was then functionalized at the N-1 position by reaction with piperazine in presence of DIPEA in dry DMF affording derivative **32**. Then, the 6-nitro-derivatives **28**, **29**, **31**, and **32** were catalytically reduced to give 6-amino-derivatives **33-36**. Finally, the target compounds **4** and **6-8** were obtained by saponification of the corresponding ethyl esters **33-36**.


By following an analogous synthetic route, 1,8-naphthyridone derivatives **11** and **12** were prepared as presented in Scheme (5). Thus, the reaction of acrylate **37** [32] with [2-(4-(pyridin-2-yl)piperazin-1-yl)ethyl]amine [33] and [3-(4-(pyridin-2-yl)piperazin-1-yl)propyl]amine [33] in Et₂O/ EtOH mixture at room temperature gave intermediates **38** and **39**, which by treatment with K₂CO₃ in dry DMF at 80 °C, afforded synthons **40** and **41**, respectively. The nucleophilic reaction of naphthyridone derivatives **40** and **41** with 1-(1,3-benzothiazol-2-yl) piperazine [34] led to intermediates **42** and **43**. Finally, the target compounds **11** and **12** were obtained by saponification of the corresponding ethyl esters **42** and **43**.




Scheme 1. Reagents and conditions: (i) *N,N'*-di-(tert-butoxycarbonyl)thiourea, HgCl₂, Et₃N, dry DMF, rt; (ii) HCl g, CHCl₃, EtOH, rt.




Scheme 2. Reagents and conditions: (i) methyl imidothiocarbamate sulfate, dry DMSO, 120 °C; (ii) FeSO₄·7H₂O, NH₄OH, H₂O, rt; (iii) 6N HCl, EtOH, reflux.



Scheme 3. Reagents and conditions: (i) ethylenediamine, *p*-toluenesulfonic acid, reflux; (ii) dry DMF, K₂CO₃, 80 °C; (iii) H₂, Raney Ni, DMF rt; (iv) 4% NaOH, reflux.



Scheme 4. Reagents and conditions: (i) (pyridin-2-ylmethyl)amine, (3-bromopropyl)amine, or *N*-(pyridin-2-yl)propane-1,3-diamine, Et₂O/EtOH (3:1)rt; (ii) K₂CO₃, dry DMF, rt, 80 °C; (iii) 1-(2-pyridinyl)piperazine, dry DMF, K₂CO₃, 80 °C; (iv) piperazine, DIPEA, dry DMF, 80 °C; (v) H₂, Raney Ni, DMF; (vi) 4% NaOH, reflux.



Scheme 5. Reagents and conditions: (i) [2-(4-(pyridin-2-yl)piperazin-1-yl)ethyl]amine, [3-(4-(pyridin-2-yl)piperazin-1-yl)propyl]amine, Et₂O/EtOH (3:1), rt; (ii) K₂CO₃, dry DMF, 80°C; (iii) 1-(1,3-benzothiazol-2-yl)piperazine, K₂CO₃, dry DMF, 80°C; (iv) H₂, Raney Ni, DMF rt; (v) 4% NaOH, reflux.

Table 1. Anti-HIV-1 activity and cytotoxicity of quinolones in MT-4 cells and inhibition of Tat-TAR complex formation.

Compd	EC ₅₀ (μM) ^{a,c} HIV-1 (III _B)	CC ₅₀ (μM) ^{b,c}	SI ^d	Ki Tat-TAR (μM) ^e
1	>296	>296	1	N.T. ^f
2	>363	363	1	N.T.
3	>34.9	34.9 ± 1.43	<1	N.T.
4	>19.8	19.8 ± 5.02	<1	N.T.
5^g	5.05 ± 0.41	20.9 ± 3.93	4	1.15 ± 0.55
6	1.21 ± 0.08	5.92 ± 2.19	5	0.91 ± 0.27
7	≥3.42	3.94 ± 0.62	≤1	2.18 ± 0.40
8	70.8 ± 0.71	>254	>4	N.T.
9^g	1.16 ± 0.18	3.97 ± 0.70	3	N.D. ^h
10^g	≥0.18	3.34 ± 1.47	≤18	1.72 ± 0.22
11	3.37 ± 1.62	11.65 ± 1.17	3.5	N.D.
12	≥0.67	10.70 ± 7.76	≤16	N.D.
WM5ⁱ	0.15 ± 0.05	2.21 ± 1.05	15	2.22 ± 0.66

^a EC₅₀: concentration of compound required to achieve 50% protection of MT-4 cells from HIV-1 induced cytopathogenicity, as determined by the MTT method. ^b CC₅₀: concentration of compound that reduces the viability of mock-infected cells by 50%, as determined by the MTT method. ^c All data represent mean values ± standard deviations for three separate experiments. ^d SI: ratio of CC₅₀/EC₅₀. ^e Ki from FQA experiments performed with 10 aa FAM-Tat peptide in Tris 10 mM, pH 7.5, Mg(ClO₄)₂ 1 mM, NaCl 20 mM (TNMg), 0.01% Triton X-100. ^f N.T.: not tested. ^g Reference [25]. ^h N.D.: not determined due to solubility issue. ⁱ Reference [20].

3.2. Biological Evaluation

The newly synthesized compounds were initially evaluated for anti-HIV-1 (III_B) activity in MT-4 cells, determining their cytotoxicity in parallel Table 1. WM5 [20] was evaluated in the same cell lines for comparative purposes.

From the results, it emerged that the introduction of a C-6 guanidine group (compound 1) as well as an amidine moiety and a 4-imidazolinpyridine ring at the C-7 position (compounds 2 and 3) was completely detrimental in the inhibition of HIV replication. However, the low cytotoxicity showed by the compounds could suggest their inability to enter the cells.

The introduction of a 2-pyridinylmethyl moiety at N-1 position (compound 4) did not produce an active derivative (EC₅₀ > 19.8 μM). Conversely, the presence of a (pyridin-2-yl) piperazine spaced by an ethylene or propylene unit permitted to restore the anti-HIV-1 activity in compounds 5 and 6 (EC₅₀ = 5.05 and 1.21 μM, respectively) at no cytotoxic concentrations.

By alternatively deleting the piperazine or pyridine ring in compound 6, no activity at subcytotoxic concentrations was observed for compound 7, while only a weak inhibitory activity was conserved by compound 8. More interesting results were achieved when protonable side chains were inserted at the N-1 position of the 1,8-naphthyridone scaffold. In particular, the

introduction of the (pyridin-2-yl)piperazine spaced by an ethylene or propylene unit gave compounds **11** and **12**, both endowed with good anti-HIV-1 activity ($EC_{50} = 3.37$ and ≥ 0.67 μM) coupled with SI values of 3.5 and ≤ 16 , respectively. The deletion of the piperazine ring, as in compound **9** and **10**, also in this case permitted to maintain anti-HIV-1 ($EC_{50} = 1.16$ and ≥ 0.18 μM , SI = 3 and ≤ 18 , respectively), with compound **10** that emerged as the most potent of the series.

In order to investigate the ability of active anti-HIV-1 compounds (derivative **5-7** and **9-12**) to interfere with the formation of the Tat-TAR complex, they were evaluated in a FQA using a truncated Tat peptide. Unfortunately, solubility issues did not allow the evaluation of compounds **9**, **11** and **12**. Derivatives **5**, **6**, and **10** appeared to be able to inhibit the Tat-TAR complex formation, all presenting K_i values even improved as compared to the values obtained with **WM5** Table 1. In particular, enhancing the distance of the pyridine ring from the quinolone nucleus, the ability of the compounds to inhibit the formation of the complex peptide-TAR increases: K_i **6** < **5** < **7**; moreover, the presence of a piperazine in the side chain (compounds **6** and **5**) has a positive effect on the activity of compounds, probably because it confers more rigidity to the chain and the position of the pyridine ring respect to the quinolone nucleus is fixed. Even if a strict correlation between the K_i values and the anti-HIV activities did not exist, this assay confirmed that the impairment of the Tat-TAR complex formation represents a putative target for the new derivatives.

CONCLUSION

The Tat-mediated transcription, the only phase of the HIV-1 replicative cycle in which viral genome amplification occurs, is a really attractive step that if inhibited by a suitable inhibitor could lead to a functional cure or even HIV infection eradication. For years, the inhibition of the Tat-mediated transcription has been highly pursued through different approaches, but this step of the viral replicative cycle remains untouched by any of the drugs in therapy. However, there are examples of promising Tat-mediated transcription inhibitors worthy to be mentioned, such as triptolide [37] and dCA [38] working on Tat protein.

In the attempt to obtain potent TAR binders, we started from **WM5**, a small molecule endowed with potent anti-HIV and Tat-mediated transcription inhibition activity thanks to the ability to selectively bind the bulge region of TAR; indeed no interaction with TAR-unrelated nucleic acids such as tRNA sequence and single-stranded or double-stranded DNA structures was observed [39].

Starting from **WM5**, various protonable moieties were inserted in different positions of the quinolone scaffold, intended to mime the basic region of Tat involved in the TAR interaction. The modifications introduced in the structure do not result in loss of activity, but rather seem to improve the activity as inhibitors of the Tat-TAR complex formation, leading to the identification of compounds able to recognize TAR better than **WM5**. Important SAR insights were also achieved for the quinolone class of anti-HIV derivatives. In particular, the C-6 and C-7 positions were confirmed as particularly sensitive to structural modifications, while the N-1

emerged as a suitable position to host protonable moieties such as the pyridine-based side chains. This is an important new insight since the most potent quinolones reported until now were all characterized by the presence of small alkyl groups at the N-1 position [22, 40]. In addition, potent anti-HIV compounds were identified, with naphthyridone derivatives **10** and **12** which showed EC_{50} values in the sub-micromolar range.

In conclusion, the addition of protonable chains as tool to improve the TAR binding properties permitted to obtain active compounds, but additional knowledge on the interaction of TAR RNA with viral protein Tat and host Super Elongation Complex (SEC) [41, 42] are still necessary to design compounds able to better recognize the TAR RNA and potently inhibit HIV replication. During the reviewing of this work, two interesting papers were published that could help the design of an effective TAR binder. In particular, Schulze-Gahmen and Hurley solved the crystal structure of the TAR in complex with Tat and the SEC core (CycT1/CDK9/AFF4) [43], while Varani and co-workers reported the NMR structure of TAR RNA in complex with a ultra-potent macrocyclic peptide that mimics the Arginine Rich Motif (ARM) of Tat [44], which however only weakly inhibit the Tat-mediated transcription.

These studies revealed how the direct interaction of Tat-ARM with TAR RNA bulge is required to induce the RNA structural change, but it is the recognition of the loop by the CycT1 that dominantly contributes to the binding energy. Thus, a future TAR binder should be able to target not only the UCU bulge but above all the interaction of CycT1 with the rearranged TAR RNA loop, to achieve an effective Tat-mediated transcription inhibition in cells.

LIST OF ABBREVIATIONS

ARM	=	Arginine Rich Motif
SAR	=	Structure-Activity Relationship
SEC	=	Super Elongation Complex
TAR	=	Transactivation Response Element
FQA	=	Fluorescence Quenching Assay.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable

HUMAN AND ANIMAL RIGHTS

No Animals/Humans were used for studies that are basis of this research.

CONSENT FOR PUBLICATION

Not applicable

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial, or otherwise.

ACKNOWLEDGEMENTS

We thank K. Erven, K. Uyttersprot and C. Heens for

technical assistance with the evaluation of the anti-HIV activity and the assessment of cytotoxicity.

REFERENCES

- [1] Thomas, J.R.; Hergenrother, P.J. Targeting RNA with small molecules. *Chem. Rev.*, **2008**, *108*(4), 1171-1224. [http://dx.doi.org/10.1021/cr0681546] [PMID: 18361529]
- [2] Hermann, T. Small molecules targeting viral RNA. *Wiley Interdiscip. Rev. RNA*, **2016**, *7*(6), 726-743. [http://dx.doi.org/10.1002/wrna.1373] [PMID: 27307213]
- [3] Kolář, M.H.; Tabarrini, O. Halogen Bonding in Nucleic Acid Complexes. *J. Med. Chem.*, **2017**, *60*(21), 8681-8690. [http://dx.doi.org/10.1021/acs.jmedchem.7b00329] [PMID: 28605588]
- [4] Weeks, K.M.; Ampe, C.; Schultz, S.C.; Steitz, T.A.; Crothers, D.M. Fragments of the HIV-1 Tat protein specifically bind TAR RNA. *Science*, **1990**, *249*(4974), 1281-1285. [http://dx.doi.org/10.1126/science.2205002] [PMID: 2205002]
- [5] Aboul-ela, F.; Karn, J.; Varani, G. Structure of HIV-1 TAR RNA in the absence of ligands reveals a novel conformation of the trinucleotide bulge. *Nucleic Acids Res.*, **1996**, *24*(20), 3974-3981. [http://dx.doi.org/10.1093/nar/24.20.3974] [PMID: 8918800]
- [6] Dethoff, E.A.; Petzold, K.; Chugh, J.; Casiano-Negrone, A.; Al-Hashimi, H.M. Visualizing transient low-populated structures of RNA. *Nature*, **2012**, *491*(7426), 724-728. [http://dx.doi.org/10.1038/nature11498] [PMID: 23041928]
- [7] Puglisi, J.D.; Chen, L.; Frankel, A.D.; Williamson, J.R. Role of RNA structure in arginine recognition of TAR RNA. *Proc. Natl. Acad. Sci. USA*, **1993**, *90*(8), 3680-3684. [http://dx.doi.org/10.1073/pnas.90.8.3680] [PMID: 7682716]
- [8] Tao, J.; Chen, L.; Frankel, A.D. Dissection of the proposed base triple in human immunodeficiency virus TAR RNA indicates the importance of the Hoogsteen interaction. *Biochemistry*, **1997**, *36*(12), 3491-3495. [http://dx.doi.org/10.1021/bi962259t] [PMID: 9131998]
- [9] Huthoff, H.; Girard, F.; Wijmenga, S.S.; Berkhout, B. Evidence for a base triple in the free HIV-1 TAR RNA. *RNA*, **2004**, *10*(3), 412-423. [http://dx.doi.org/10.1261/rna.5161304] [PMID: 14970387]
- [10] Mousseau, G.; Valente, S.T. Role of host factors on the regulation of Tat-Mediated HIV-1 transcription. *Curr. Pharm. Des.*, **2017**, *23*(28), 4079-4090. [http://dx.doi.org/10.2174/1381612823666170622104355] [PMID: 28641539]
- [11] Schiralli Lester, G.M.; Henderson, A.J. Mechanisms of HIV transcriptional regulation and their contribution to latency. *Mol. Biol. Int.*, **2012**, *2012*, 614120. [http://dx.doi.org/10.1155/2012/614120] [PMID: 22701796]
- [12] Massari, S.; Sabatini, S.; Tabarrini, O. Blocking HIV-1 replication by targeting the Tat-hijacked transcriptional machinery. *Curr. Pharm. Des.*, **2013**, *19*(10), 1860-1879. [http://dx.doi.org/10.2174/1381612811319100010] [PMID: 23092279]
- [13] Tabarrini, O.; Desantis, J.; Massari, S. Recent advances in the identification of Tat-mediated transactivation inhibitors: progressing toward a functional cure of HIV. *Future Med. Chem.*, **2016**, *8*(4), 421-442. [http://dx.doi.org/10.4155/fmc.16.3] [PMID: 26933891]
- [14] Abulwerdi, F.A.; Le Grice, S.F.J. Recent advances in targeting the HIV-1 Tat/TAR complex. *Curr. Pharm. Des.*, **2017**, *23*(28), 4112-4121. [http://dx.doi.org/10.2174/1381612823666170616081736] [PMID: 28625133]
- [15] Hamy, F.; Felder, E.R.; Heizmann, G.; Lazdins, J.; Aboul-ela, F.; Varani, G.; Karn, J.; Klimkait, T. An inhibitor of the Tat/TAR RNA interaction that effectively suppresses HIV-1 replication. *Proc. Natl. Acad. Sci. USA*, **1997**, *94*(8), 3548-3553. [http://dx.doi.org/10.1073/pnas.94.8.3548] [PMID: 9108013]
- [16] Abulwerdi, F.A.; Shortridge, M.D.; Sztuba-Solinska, J.; Wilson, R.; Le Grice, S.F.; Varani, G.; Schneekloth, J.S., Jr. Development of small molecules with a noncanonical binding mode to HIV-1 trans activation response (TAR) RNA. *J. Med. Chem.*, **2016**, *59*(24), 11148-11160. [http://dx.doi.org/10.1021/acs.jmedchem.6b01450] [PMID: 28002966]
- [17] Davidson, A.; Begley, D.W.; Lau, C.; Varani, G. A small-molecule probe induces a conformation in HIV TAR RNA capable of binding drug-like fragments. *J. Mol. Biol.*, **2011**, *410*(5), 984-996. [http://dx.doi.org/10.1016/j.jmb.2011.03.039] [PMID: 21763501]
- [18] Davidson, A.; Leeper, T.C.; Athanassiou, Z.; Patora-Komisarska, K.; Karn, J.; Robinson, J.A.; Varani, G. Simultaneous recognition of HIV-1 TAR RNA bulge and loop sequences by cyclic peptide mimics of Tat protein. *Proc. Natl. Acad. Sci. USA*, **2009**, *106*(29), 11931-11936. [http://dx.doi.org/10.1073/pnas.0900629106] [PMID: 19584251]
- [19] Sztuba-Solinska, J.; Shenoy, S.R.; Gareiss, P.; Krumpe, L.R.; Le Grice, S.F.; O'Keefe, B.R.; Schneekloth, J.S., Jr. Identification of biologically active, HIV TAR RNA-binding small molecules using small molecule microarrays. *J. Am. Chem. Soc.*, **2014**, *136*(23), 8402-8410. [http://dx.doi.org/10.1021/ja502754f] [PMID: 24820959]
- [20] Cecchetti, V.; Parolin, C.; Moro, S.; Pecere, T.; Filipponi, E.; Calistri, A.; Tabarrini, O.; Gatto, B.; Palumbo, M.; Fravolini, A.; Palù, G. 6-Aminoquinolones as new potential anti-HIV agents. *J. Med. Chem.*, **2000**, *43*(20), 3799-3802. [http://dx.doi.org/10.1021/jm9903390] [PMID: 11020296]
- [21] Tabarrini, O.; Massari, S.; Cecchetti, V. 6-desfluoroquinolones as HIV-1 Tat-mediated transcription inhibitors. *Future Med. Chem.*, **2010**, *2*(7), 1161-1180. [http://dx.doi.org/10.4155/fmc.10.208] [PMID: 21426162]
- [22] Tabarrini, O.; Massari, S.; Daelemans, D.; Meschini, F.; Manfroni, G.; Bottega, L.; Gatto, B.; Palumbo, M.; Pannecouque, C.; Cecchetti, V. Studies of anti-HIV transcription inhibitor quinolones: identification of potent N1-vinyl derivatives. *Chem Med Chem*, **2010**, *5*(11), 1880-1892. [http://dx.doi.org/10.1002/cmdc.201000267] [PMID: 20928882]
- [23] Manfroni, G.; Gatto, B.; Tabarrini, O.; Sabatini, S.; Cecchetti, V.; Giaretta, G.; Parolin, C.; Del Vecchio, C.; Calistri, A.; Palumbo, M.; Fravolini, A. Synthesis and biological evaluation of 2-phenylquinolones targeted at Tat/TAR recognition. *Bioorg. Med. Chem. Lett.*, **2009**, *19*(3), 714-717. [http://dx.doi.org/10.1016/j.bmcl.2008.12.034] [PMID: 19109014]
- [24] Gatto, B.; Tabarrini, O.; Massari, S.; Giaretta, G.; Sabatini, S.; Del Vecchio, C.; Parolin, C.; Fravolini, A.; Palumbo, M.; Cecchetti, V. 2-Phenylquinolones as inhibitors of the HIV-1 Tat-TAR interaction. *ChemMedChem*, **2009**, *4*(6), 935-938. [http://dx.doi.org/10.1002/cmdc.200800437] [PMID: 19283692]
- [25] Donalizio, M.; Massari, S.; Argenziano, M.; Manfroni, G.; Cagno, V.; Cirva, A.; Sabatini, S.; Cecchetti, V.; Lorigian, A.; Cavalli, R.; Lembo, D.; Tabarrini, O. Ethyl 1,8-naphthyridone-3-carboxylates downregulate human papillomavirus-16 E6 and E7 oncogene expression. *J. Med. Chem.*, **2014**, *57*(13), 5649-5663. [http://dx.doi.org/10.1021/jm500340h] [PMID: 24905115]
- [26] Massari, S.; Daelemans, D.; Barreca, M.L.; Knezevich, A.; Sabatini, S.; Cecchetti, V.; Marcello, A.; Pannecouque, C.; Tabarrini, O. A 1,8-naphthyridone derivative targets the HIV-1 Tat-mediated transcription and potentially inhibits the HIV-1 replication. *J. Med. Chem.*, **2010**, *53*(2), 641-648. [http://dx.doi.org/10.1021/jm901211d] [PMID: 19958026]
- [27] Tabarrini, O.; Massari, S.; Sancineto, L.; Daelemans, D.; Sabatini, S.; Manfroni, G.; Cecchetti, V.; Pannecouque, C. Structural investigation of the naphthyridone scaffold: identification of a 1,6-naphthyridone derivative with potent and selective anti-HIV activity. *Chem Med Chem*, **2011**, *6*(7), 1249-1257. [http://dx.doi.org/10.1002/cmdc.201100073] [PMID: 21567967]
- [28] Tabarrini, O.; Massari, S.; Daelemans, D.; Stevens, M.; Manfroni, G.; Sabatini, S.; Balzarini, J.; Cecchetti, V.; Pannecouque, C.; Fravolini, A. Structure-activity relationship study on anti-HIV 6-desfluoroquinolones. *J. Med. Chem.*, **2008**, *51*(17), 5454-5458. [http://dx.doi.org/10.1021/jm701585h] [PMID: 18710207]
- [29] Cecchetti, V.; Clementi, S.; Cruciani, G.; Fravolini, A.; Pagella, P. G.; Savino, A.; Tabarrini, O. 6-Aminoquinolones: A new class of quinolone antibacterials? *J. Med. Chem.*, **1995**, *38*(6), 973-982.
- [30] Anderson, D.; Beutel, B.; Bosse, T. D.; Clark, R.; Cooper, C.; Dandliker, P.; David, C.; Yu-Gui, H.; Todd, M.; Hinman, M.; Calvin, D.; Larson, D. P.; Lynch, L.; Ma, Z.; Motter, C.; Palazzo, F.; Rosenberg, T.; Rehm, T.; Sanders, W.; Tufano, M.; Wagner, R.; Weitzberg, M.; Yong, H.; Zhang, T. Preparation of naphthyridines as antibacterial compounds. *US2003232818* 2003.
- [31] Elliott, R. L.; Hammond, M.; Hank, R. F. 2-Pyridyl-4-heteroaryl-imidazoles for the treatment of obesity. *WO0066578(A1)*, **2000**.
- [32] Ife, R.J.; Catchpole, K.W.; Durant, G.J.; Robin Ganellin, C.; Harvey, C.A.; Meeson, M.L.; Owen, D.A. Parsons, M. E.; Slingsby, B. P.; Theobald, C. J. Non-Basic Histamine H1-Antagonists. I. Synthesis and Biological Evaluation of Some Substituted 2-(2-Pyridylaminoalkylamino) Pyrimidones and Related Compounds. *Eur. J. Med. Chem.*, **1989**, *24*(3), 249-257. [http://dx.doi.org/10.1016/0223-5234(89)90006-8]

- [33] Jung, H.K.; Doddareddy, M.R.; Cha, J.H.; Rhim, H.; Cho, Y.S.; Koh, H.Y.; Jung, B.Y.; Pae, A.N. Synthesis and biological evaluation of novel T-type Ca²⁺ channel blockers. *Bioorg. Med. Chem.*, **2004**, *12*(15), 3965-3970. [http://dx.doi.org/10.1016/j.bmc.2004.06.011] [PMID: 15246072]
- [34] Verderame, M. 1,4-Disubstituted piperazines. 3. Piperazinylbenzothiazoles. *J. Med. Chem.*, **1972**, *15*(6), 693-694. [http://dx.doi.org/10.1021/jm00276a039] [PMID: 4555494]
- [35] Pannecouque, C.; Daelemans, D.; De Clercq, E. Tetrazolium-based colorimetric assay for the detection of HIV replication inhibitors: revisited 20 years later. *Nat. Protoc.*, **2008**, *3*(3), 427-434. [http://dx.doi.org/10.1038/nprot.2007.517] [PMID: 18323814]
- [36] Pauwels, R.; Balzarini, J.; Baba, M.; Snoeck, R.; Schols, D.; Herdewijn, P.; Desmyter, J.; De Clercq, E. Rapid and automated tetrazolium-based colorimetric assay for the detection of anti-HIV compounds. *J. Virol. Methods*, **1988**, *20*(4), 309-321. [http://dx.doi.org/10.1016/0166-0934(88)90134-6] [PMID: 2460479]
- [37] Wan, Z.; Chen, X. Triptolide inhibits human immunodeficiency virus type 1 replication by promoting proteasomal degradation of Tat protein. *Retrovirology*, **2014**, *11*, 88. [http://dx.doi.org/10.1186/s12977-014-0088-6] [PMID: 25323821]
- [38] Mousseau, G.; Clementz, M.A.; Bakeman, W.N.; Nagarsheth, N.; Cameron, M.; Shi, J.; Baran, P.; Fromentin, R.; Chomont, N.; Valente, S.T. An analog of the natural steroidal alkaloid cortistatin A potently suppresses Tat-dependent HIV transcription. *Cell Host Microbe*, **2012**, *12*(1), 97-108. [http://dx.doi.org/10.1016/j.chom.2012.05.016] [PMID: 22817991]
- [39] Richter, S.N.; Gatto, B.; Tabarrini, O.; Fravolini, A.; Palumbo, M. Antiviral 6-amino-quinolones: molecular basis for potency and selectivity. *Bioorg. Med. Chem. Lett.*, **2005**, *15*(19), 4247-4251. [http://dx.doi.org/10.1016/j.bmcl.2005.06.074] [PMID: 16054362]
- [40] Massari, S.; Daelemans, D.; Manfroni, G.; Sabatini, S.; Tabarrini, O.; Pannecouque, C.; Cecchetti, V. Studies on anti-HIV quinolones: new insights on the C-6 position. *Bioorg. Med. Chem.*, **2009**, *17*(2), 667-674. [http://dx.doi.org/10.1016/j.bmc.2008.11.056] [PMID: 19091580]
- [41] Schulze-Gahmen, U.; Echeverria, I.; Stjepanovic, G.; Bai, Y.; Lu, H.; Schneidman-Duhovny, D.; Doudna, J.A.; Zhou, Q.; Sali, A.; Hurley, J.H. Insights into HIV-1 proviral transcription from integrative structure and dynamics of the Tat:AFF4:P-TEFb:TAR complex. *eLife*, **2016**, *5*, e15910. [http://dx.doi.org/10.7554/eLife.15910] [PMID: 27731797]
- [42] Gu, J.; Babayeva, N.D.; Suwa, Y.; Baranovskiy, A.G.; Price, D.H.; Tahirov, T.H. Crystal structure of HIV-1 Tat complexed with human P-TEFb and AFF4. *Cell Cycle*, **2014**, *13*(11), 1788-1797. [http://dx.doi.org/10.4161/cc.28756] [PMID: 24727379]
- [43] Schulze-Gahmen, U.; Hurley, J.H. Structural mechanism for HIV-1 TAR loop recognition by Tat and the super elongation complex. *Proc. Natl. Acad. Sci. USA*, **2018**, *115*(51), 12973-12978. [http://dx.doi.org/10.1073/pnas.1806438115] [PMID: 30514815]
- [44] Shortridge, M.D.; Wille, P.T.; Jones, A.N.; Davidson, A.; Bogdanovic, J.; Arts, E.; Kam, J.; Robinson, J.A.; Varani, G. An ultra-high affinity ligand of HIV-1 TAR reveals the RNA structure recognized by P-TEFb. *Nucleic Acids Res.*, **2018**. [http://dx.doi.org/10.1093/nar/gky1197] [PMID: 30481318]