

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This author's accepted manuscript may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

The full details of the published version of the article are as follows:

TITLE: The utility of diagnostic tests for immune-mediated hemolytic anemia

AUTHORS: Members of the Veterinary and Comparative Clinical Immunology Society Diagnostic Task Force, Amy L. MacNeill, Julien Dandrieux, George Lubas, Davis Seelig, Balázs Szladovits

JOURNAL: Veterinary Clinical Pathology

PUBLISHER: Wiley

PUBLICATION DATE: 9 September 2019

DOI: [10.1111/vcp.12771](https://doi.org/10.1111/vcp.12771)

1 **Title:** The Utility of Diagnostic Tests for Immune-mediated Hemolytic Anemia

2

3 **Short Title:** Diagnosis of IMHA

4

5 **Authors:** Members of the Veterinary and Comparative Clinical Immunology Society Diagnostic Task

6 Force: Amy L. MacNeill,¹ Julien Dandrieux,² George Lubas³ Davis Seelig,⁴ Balázs Szladovits⁵

7

8 ¹Department of Microbiology, Immunology, and Pathology, College of Veterinary Medicine and

9 Biomedical Sciences, Colorado State University, Fort Collins, Colorado, USA

10 ²Faculty of Veterinary and Agricultural Sciences, Translational Research and Animal Clinical Trial Study

11 (TRACTS) Group, The University of Melbourne, Werribee, Victoria, Australia

12 ³Department of Veterinary Sciences, University of Pisa, Pisa, Italy

13 ⁴Department of Veterinary Clinical Sciences, College of Veterinary Medicine, University of Minnesota, St.

14 Paul, Minnesota, USA

15 ⁵Department of Pathobiology and Population Sciences, The Royal Veterinary College, University of

16 London, Hertfordshire, UK

17

18 **Correspondence:** Amy L. MacNeill, 300 W. Drake Rd., Fort Collins, CO, USA 80523.

19 amy.macneill@colostate.edu

20

21 **Abstract**

22 **Background:** A definitive diagnosis of immune-mediated hemolytic anemia (IMHA) can be difficult to
23 make. However, it is critical to differentiate IMHA from other causes of anemia due to the impact on
24 prognosis and outcome for IMHA patients. Recently published American College of Veterinary Internal
25 Medicine recommendations for the diagnosis of IMHA should be followed to concurrently confirm
26 ongoing anemia, verify *in vivo* hemolysis, and detect anti-erythrocyte antibodies. The reliability of
27 immunologic IMHA tests varies depending on which test is used and how it is performed.

28 **Objectives:** Our aims were to determine which tests are currently used in veterinary medicine to
29 diagnose IMHA and to review the utility of assays that have historically been used to diagnose IMHA.

30 **Methods:** A short survey was designed to see which diagnostic tests for IMHA were currently being
31 used by veterinary practices. The survey was distributed *via* list-serves to veterinarians and veterinary
32 technologists. A literature review was performed to report the utility of diagnostic tests for the diagnosis
33 of IMHA.

34 **Results:** Survey respondents indicated variability in test protocols used to diagnose IMHA. Most
35 respondents perform saline agglutination or Coombs' tests to detect anti-erythrocyte antibodies.
36 Additional tests that can be used to support a diagnosis of IMHA are discussed in this review.

37 **Conclusions:** A standardized diagnostic approach should be followed to differentiate IMHA from other
38 causes of anemia. Test methodology can vary from one laboratory to another, and clinicians should be
39 familiar with the procedures used by their laboratory.

40

41 **Key Words:** Agglutination, Anti-erythrocyte antibody, Coombs', Diagnosis, Spherocytosis

42

43 **Introduction**

44 Immune-mediated hemolytic anemia (IMHA) is commonly diagnosed in canine patients with hemolytic
45 anemia. It is a less prevalent, but an equally important cause of hemolytic anemia in cats, horses, and
46 cattle.^{1,2} While there are certain clinicopathologic findings supportive of an IMHA diagnosis (e.g.,
47 peripheral blood spherocytosis, RBC agglutination, demonstration of immunoglobulins attached to RBC
48 membranes), the diagnosis of IMHA is often presumptive. Misdiagnosis of IMHA is problematic because
49 of important differences in treatment decisions and prognostic consequences for patients with IMHA as
50 compared with other causes of anemia.³ Additionally, it is critical to evaluate patients that have IMHA
51 for underlying diseases that drive the immune-mediated response. This article summarizes the results of
52 a short survey on the use of diagnostic tests for IMHA in veterinary medicine and reviews the utility of
53 diagnostic tests for IMHA.

54

55 **Diagnosis of IMHA**

56 In anemic patients, the mechanism causing the anemia (decreased RBC production, hemorrhage, or
57 hemolysis) needs to be determined. A CBC (including quantitative data and a qualitative blood film
58 review) provides fundamental information needed for the assessment of anemia. Anemia can be
59 classified as either regenerative or nonregenerative based on quantification of the reticulocytosis using
60 an absolute reticulocyte concentration or corrected reticulocyte percentage in the sample. If the anemia
61 is regenerative, blood loss or hemolytic anemia should be considered. Blood loss anemia can often be
62 differentiated from hemolytic anemia using clinical examination findings and additional
63 clinicopathologic data. Specifically, acute hemorrhage results in loss of serum proteins and,
64 consequently, decreased serum albumin and globulin concentrations. Although blood loss typically
65 results in a strongly regenerative anemia, chronic hemorrhage can cause a poorly regenerative anemia if
66 blood loss leads to iron depletion, since the resultant iron-restricted erythropoiesis impedes reticulocyte
67 production. It is important to assess patients for occult blood loss as gastrointestinal and urinary blood

68 loss can be easily missed. While the majority of IMHA patients present with a regenerative anemia, this
69 initial classification is not a certainty. IMHA can be nonregenerative or pre-regenerative if there has
70 been insufficient time for a clearly regenerative response to occur (anemia present < 5-7 days)⁴ or if
71 immune-mediated destruction of erythroid precursors is occurring (“precursor-targeted immune-
72 mediated anemia”).⁵

73
74 It is vital to document that true *in vivo* hemolysis is occurring in patients with IMHA. Accordingly, proper
75 collection of blood samples is critical to avoid *ex vivo* hemolysis, which can lead to false decreases in RBC
76 concentration, hematocrit (HCT), and packed cell volume (PCV); false increases in mean cell hemoglobin
77 concentration (MCHC); variable alterations in serum bilirubin concentrations (depending on the
78 methodology used); and can also interfere with several other serum chemistry values.⁶ Diagnostic test
79 results that support *in vivo* hemolysis are provided in **Table 1**.

80
81 To distinguish IMHA from other causes of anemia, an American College of Veterinary Internal Medicine
82 (ACVIM) consensus statement recommends documenting all three of the following diagnostic findings as
83 minimum criteria to diagnose IMHA in dogs:⁷

- 84 1) Decreased packed cell volume (PCV)
- 85 2) At least one of the following abnormalities supportive of hemolysis: erythrocyte ghost cells,
86 hyperbilirubinemia, bilirubinuria, icterus, hemoglobinemia, or hemoglobinuria
- 87 3) A positive saline agglutination test that persists with washing or at least two of the following:
 - 88 a) A positive saline agglutination test without washing
 - 89 b) ≥ 5 spherocytes per 1000 \times microscopic field
 - 90 c) Detection of anti-erythrocyte antibodies by a Coombs’ test or flow cytometry

91 Note that erythrocyte ghost cell numbers can markedly increase *ex vivo*, so examination of a freshly
92 made smear (at the time of blood collection) is recommended to assess this morphologic abnormality.⁶

93

94 A short survey (Appendix A) to determine how tests are being used to diagnose IMHA was sent to an
95 undetermined number of people in Australia, Canada, Europe, and the United States *via* list-serves and
96 email messages. Veterinary practitioners in both general practice and specialty hospitals, veterinary
97 clinical pathologists, and medical technologists and technicians in clinical pathology laboratories were
98 asked to reply. Respondents were not asked to self-identify. Ninety-four people completed at least the
99 first question of the survey.

100

101 The hematology parameters that > 75% of the respondents said were provided with a CBC are listed in
102 Figure 1. Seventy-two people (77%) indicated that, when needed, a saline agglutination test was
103 included with CBC results. An additional 13 respondents (14%) did not include a saline agglutination test
104 as part of a CB, but as an additional test. Seventy-two people (77%) recommended a test to detect anti-
105 erythrocyte antibodies (saline agglutination test, Coombs test, and/or flow cytometry). Additional tests
106 that were recommended by > 15 % of respondents included a biochemistry panel (66%), screening for
107 tick-borne diseases (60%), urinalysis (52%), and diagnostic imaging (34%).

108

109 Several tests have been used to aid in the diagnosis of IMHA, none of which are 100% sensitive and
110 specific. Differences in the protocols used to perform diagnostic tests for IMHA can dramatically affect
111 the performance of the test.^{8,9} Tests for IMHA must account for physical properties associated with
112 antibody-antigen interactions, one of which is temperature-dependent antibody-antigen binding.¹⁰
113 Therefore, the Coombs' test is often carried out at both 37°C and 4°C.¹¹ Additionally, inclusion criteria
114 for IMHA patients vary between references, which can alter the assessment of test performance. Most

115 studies evaluating diagnostic tests for canine IMHA list auto-agglutination or spherocytosis as inclusion
116 criteria. However, Reimer *et al.* (1999) concluded that 2/70 (3%) canine patients with IMHA did not have
117 auto-agglutination or spherocytosis, nor did they have polychromasia.¹²

118

119 Genetics

120 Several dog breeds have a higher incidence of IMHA than the general canine population including;
121 cocker spaniels,¹²⁻¹⁵ English springer spaniels,¹² miniature schnauzers,^{13,15} and old English sheepdogs¹⁶;
122 however, a specific gene association with IMHA has not be found. A few studies have associated the
123 presence of specific major histocompatibility complex alleles in dogs with the occurrence of immune-
124 mediated disease, but the predictive value of these associations remains uncertain.¹⁷⁻¹⁹ The frequency
125 of specific dog erythrocyte antigens found during blood typing was not significantly different in 33 dogs
126 with IMHA as compared to 1,014 dogs without IMHA.²⁰ Therefore, expectedly, genetic testing was not
127 recommended as an additional test for IMHA by survey respondents.

128

129 Non-specific Tests for Antibodies

130 Two proteins, staphylococcal protein A (SpA) and papain, have been used in several papers to document
131 increased immunoglobulin in serum samples. SpA is an immunoglobulin binding protein produced by
132 *Staphylococcus aureus* that has been shown to be more sensitive than indirect antiglobulin tests in dogs
133 with IMHA.²¹ Papain is an enzyme that digests antibodies into three 50 kDa segments comprised of one
134 fragment crystallizable (Fc) and two fragment antigen-binding (Fab) regions. When using a papain test to
135 detect anti-erythrocyte antibodies, false-positive results occur in up to 3/16 (19%) of dogs.²² One study
136 tested 23 papain positive anemic dogs with direct and indirect antiglobulin tests; 8/23 (35%) dogs were
137 positive using a direct test and none were positive using an indirect test.²³ Due to the non-specific

138 nature of SpA and papain reactions, they are no longer recommended as diagnostic tests for IMHA and
139 unsurprisingly were not listed as additional tests used by survey respondents.

140

141 Auto-agglutination

142 Red blood cell agglutination should raise suspicion for IMHA (**Fig. 2**). To determine whether
143 agglutination is likely related to the presence of antibody or complement on the surface of the red blood
144 cell (i.e., auto-agglutination), a saline agglutination test should be performed. It is important to note that
145 artifactual agglutination/rouleaux is not dispersed in all patients using a 1:1 dilution. However, if
146 agglutination persists after washing erythrocytes at least three times in isotonic or phosphate-buffered
147 saline, it is likely that the erythrocyte clumping is true agglutination rather than rouleaux formation. The
148 occurrence of auto-agglutination in dogs with IMHA varies from 42% to 86%.¹²⁻¹⁶ The number of patients
149 with a positive saline agglutination test is decreased when erythrocytes are washed extensively.^{24,25}
150 There is some evidence that the decrease in positive saline agglutination tests with washing is due to a
151 reduction in false-positive results,²⁵ but increased numbers of false-negative results should also be
152 considered.

153

154 Seventy respondents indicated whether they evaluated saline agglutination tests macroscopically or
155 microscopically: 34 looked for microscopic agglutination, 29 looked for macroscopic and microscopic
156 agglutination, and seven looked at the sample macroscopically only. Sixty-three people indicated how
157 they dilute blood samples for a saline agglutination test: 31 performed a 1:4 dilution, 24 performed a 1:1
158 dilution, and eight performed both a 1:1 and a 1:4 dilution of the sample. Of the 28 respondents who
159 indicated whether they wash RBCs for the saline agglutination test, 16 used unwashed RBCs, 11 used
160 washed RBCs, and one used both.

161

162 Erythron

163 Spherocytosis is often used as an inclusion criterion for studies of dogs with IMHA and was included in
164 CBC data by 90/94 (96%) of survey respondents (Fig. 1). However, IMHA also can be non-spherocytic, as
165 the occurrence of spherocytosis in dogs with IMHA ranges from 61% to 95%.^{12,13,15,16,24,26} Spherocytes are
166 generally only recognized in dogs (owing to the pronounced central pallor of their erythrocytes). In
167 contrast, spherocytes are rarely identified with certainty in cats, horses, and cattle since erythrocytes
168 from these animals lack central pallor on blood smears.

169

170 Spherocytosis and some other RBC membrane abnormalities can cause increased erythrocyte fragility.²⁷

171 The osmotic fragility of erythrocytes can be tested by diluting whole blood in progressively decreasing
172 concentrations of sodium chloride (NaCl), incubating the samples for 30 minutes at room-temperature,
173 recording absorbance of the samples at 540 nm, and then creating a data curve assuming that the
174 lowest NaCl concentration causes 100% hemolysis.²⁷ RBC hemolysis occurs as a result of a loss of
175 osmotic regulation and volume control, which is exacerbated in a number of RBC disorders. Increased
176 RBC hemolysis during osmotic fragility testing is commonly reported in spherocytic conditions (e.g.,
177 IMHA), but can be seen in spectrin deficiency,²⁸ hereditary stomatocytocytosis,²⁹ intestinal parasite-
178 associated microcytosis, *Babesia canis* infection,³⁰ or non-hemolytic samples that are lipemic.

179 Erythrocyte fragility testing is not commonly available to practitioners but can provide support for
180 ongoing hemolysis.³¹ One study observed increased erythrocyte fragility in 15/15 (100%) direct
181 antiglobulin test (DAT) positive and 4/12 (33%) DAT negative anemic dogs.²⁵ None (0/91) of the survey
182 respondents indicated that they recommended osmotic fragility testing for patients suspected of having
183 IMHA.

184

185 Most patients with IMHA have a moderate to marked regenerative anemia. A reticulocyte count was
186 listed as a component of a CBC by 77/94 (82%) of the survey respondents (Fig. 1). The occurrence of
187 reticulocytosis in dogs with IMHA ranges from 67% to 82%.^{13,16,26} IMHA patients often have increased
188 numbers of circulating nucleated RBCs and/or Howell-Jolly bodies. Interestingly, reticulocyte
189 hemoglobin (HGB) content was shown to be decreased in 5/14 (36%) dogs with IMHA suggesting that
190 iron-restricted erythropoiesis can be present in some canine IMHA patients.³²

191

192 A diagnosis of erythroid hyperplasia in bone marrow samples is a definitive indication of erythrocyte
193 regeneration. It is noteworthy that Weinkle *et al.* (2005) found 23/45 (51%) dogs with IMHA that
194 underwent bone marrow analysis had erythroid hyperplasia.¹³ Similarly, another study that analyzed
195 bone marrow samples from dogs with IMHA observed erythroid hyperplasia in 6/11 (55%) samples.¹⁵
196 Extramedullary hematopoiesis³³ and secondary myelodysplasia³⁴ also have been reported in dogs with
197 IMHA. Bone marrow evaluation is typically recommended for suspected IMHA patients that have a
198 nonregenerative anemia. Low numbers of survey respondents [6/91 (7%)] indicated that they
199 recommend bone marrow aspiration or biopsy to aid in the diagnosis of IMHA. Two of these
200 respondents specified that this recommendation was warranted in patients with persistent
201 nonregenerative anemia.

202

203 Nonregenerative anemia has been reported in 6/23 (26%) canine IMHA patients in one study²⁴ and 6/20
204 (30%) in another.¹⁵ In a retrospective analysis of dogs with nonregenerative anemia in which a bone
205 marrow sample was clinically indicated, 55/82 (67%) were determined to have IMHA (based on the
206 presence of either Coombs' positivity, auto-agglutination, or > 30% spherocytes).³⁵ In dogs with
207 nonregenerative IMHA, 38/55 (69%) had erythroid hyperplasia, and 17/55 (31%) showed incomplete
208 maturation of the erythroid line.³⁵ In another study of canine patients with a nonregenerative anemia

209 present for more than 5 days, 41/43 (95%) had erythroid hyperplasia in bone marrow samples, 23/43
210 (54%) had a spherocytosis, and 20/35 (57%) were positive by DAT.³⁶ To determine if dogs with a
211 nonregenerative anemia > 5 days duration have precursor-targeted immune-mediated anemia, bone
212 marrow should be evaluated for macrophage phagocytosis of erythroid precursors.⁵ Weiss (2008) also
213 evaluated 57 cats with nonregenerative anemia in which bone marrow analysis was indicated.
214 Approximately half of the cats 28/57 (49%) were determined to have IMHA (based upon measurement
215 of an HCT < 20% and either Coombs' positivity or auto-agglutination).³⁵ In cats with IMHA and
216 nonregenerative anemia, 24/28 (86%) had erythroid hyperplasia and 4/28 (14%) showed maturation
217 arrest of the erythroid line.³⁵ In another study, phagocytosis of erythroid precursors and abnormal
218 presence of stainable iron was documented in the bone marrow of cats with both primary and
219 secondary IMHA.³⁷

220
221 In cases of IMHA with RBC agglutination in the sample, many of the measured or calculated values of the
222 erythron are often erroneous [e.g., RBC concentration, mean cell volume (MCV), HCT, mean cell
223 hemoglobin (MCH), MCHC, red cell distribution width (RDW)] as impedance counters will count
224 erythrocyte clumps as single and large erythrocytes, which leads to a significant reduction in the
225 numbers of RBCs counted and an increase in the mean size of the cells.⁶ In cases of intravascular
226 hemolysis, HGB concentration is not clinically reliable, as it represents a combination of free (plasma)
227 and RBC HGB. If it is available, the determination of the cell hemoglobin concentration mean (CHCM)
228 using an advanced laser cell counter could help assess RBC HGB.

229
230 Leukon
231 Abnormalities in leukocytes are commonly observed in patients with IMHA. Leukocytosis was reported
232 in 43% to 99% of dogs with IMHA (WBC concentrations in these reports ranged from 5,300 cells/ μ L to

233 105,700 cells/ μ L).^{12,15,38,39} In one study, decreased survival time of dogs with IMHA was associated with
234 leukocytosis and lymphopenia,⁴⁰ while lymphocytosis was a positive prognostic factor in cats with
235 IMHA.³⁷ Neutrophil left shifts were noted in up to 16/20 (80%) of dogs with IMHA.¹⁵ One paper
236 observed decreased survival rates in dogs with IMHA with band neutrophil concentrations \geq 3000
237 cells/ μ L.¹³

238
239 IMHA patients with inflammatory leukograms typically have acute patterns such as a neutrophilia with a
240 left shift, lymphopenia, eosinopenia, and monocytosis.¹⁵ In severe inflammatory and erythroid
241 regenerative conditions, a leukoerythroblastic pattern can be observed with a highly acute inflammatory
242 leukogram and a high percentage of nucleated RBCs in different stages of maturation.¹⁵ Rubricytosis
243 causes a false increase in the automated WBC concentration that must be corrected mathematically
244 after the enumeration of nucleated RBCs by blood smear review. This further emphasizes the need for
245 blood smear examination, which can also help with the detection of neutrophil left-shifting and toxicity
246 that can be present in IMHA cases. Sixty-five/94 (69%) survey respondents indicated that blood smear
247 evaluation by a board-certified clinical pathologist was included in CBCs they performed or received (Fig.
248 1).

249

250 Serum Biochemistry and Urinalysis

251 Abnormalities in biochemical parameters have been associated with the clinical outcomes of IMHA
252 patients (**Table 2**), but not all studies report the same findings. An increase in total bilirubin
253 concentrations was observed in 60% to 100% of dogs with IMHA.^{12,15,26} It is important to note that
254 increased conjugated bilirubin can interfere with phosphorus measurements leading to
255 pseudohypophosphatemia in patients with IMHA.⁴¹ Although not linked to decreased survival, Klag et
256 al. (1993) observed hemoglobinemia and/or hemoglobinuria in 4/42 (10%) dogs with IMHA.²⁶ There are

257 also a few case studies of dogs with IMHA with biochemical abnormalities consistent with distal renal
258 tubular acidosis.⁴² In cats with IMHA, hyperglobulinemia is reported to be a positive prognostic factor.³⁷
259 Biochemistry profiles were recommended as additional tests for suspected IMHA patients by 60/91
260 (66%) of survey respondents.

261
262 Additional serum protein parameters have been reported to be altered in dogs with IMHA. For example,
263 cardiac troponin I was > 0.1 ng/mL in 20/27 (74%) dogs with IMHA (authors indicated that < 0.1 ng/mL is
264 expected in healthy dogs, but a true reference interval was not provided).⁴³ C-reactive protein was
265 increased in dogs with IMHA at presentation.⁴⁴⁻⁴⁶ Alpha-1 acid glycoprotein also was increased in dogs
266 with IMHA, while albumin can be decreased at presentation.⁴⁵ Increased serum concentrations of
267 several cytokines have been reported in dogs with IMHA (n = 20) as compared with six healthy dogs.⁴⁶
268 Interleukin-15 (IL-15), IL-18, granulocyte-monocyte colony stimulating factor, and monocyte
269 chemoattractant protein-1 concentrations were increased in animals with IMHA that died ≤ 30 days
270 after hospital admission.⁴⁶ Similarly, IL-2, IL-6, and tumor necrosis factor-α were present at higher
271 concentrations in dogs with primary IMHA (n = 19) when compared with dogs that had other
272 inflammatory diseases (n = 22) or healthy dogs (n = 32).⁴⁷ In question 4, none of the survey respondents
273 indicated that they recommended these protein assays to help diagnose patients with IMHA.

274

275 Thrombon and Coagulation

276 Thrombocytopenia is reported to occur in 29-70% of dogs with IMHA.^{12,14,15,26,38,39} In a study of 151 dogs,
277 a platelet concentration < 150,000 platelets/μL correlated with decreased survival rates.¹³ Also, a
278 decreased mean platelet component concentration was found in dogs with IMHA (n = 95) as compared
279 with healthy dogs (n = 95) or sick canine patients (n = 95)⁴⁸ which could indicate increased platelet
280 activation in IMHA patients.⁴⁹

281
282 Considerations for severe thrombocytopenia include a consumptive process [e.g., disseminated
283 intravascular coagulation (DIC), pulmonary thromboembolism (PTE)] or a concurrent immune-mediated
284 thrombocytopenia (IMT). In humans, concurrent IMHA and IMT have been termed Evan’s Syndrome.
285 This disease process likely occurs in dogs; however, the presence of concurrent anti-erythrocyte and
286 anti-platelet antibodies has rarely been documented in veterinary patients.^{50,51} In 38 dogs with both
287 anemia and thrombocytopenia, 18/38 (47%) of patients were positive by DAT for anti-erythrocyte
288 antibodies.⁵² In a similar study of 21 dogs with concurrent anemia and thrombocytopenia, auto-
289 agglutination that persisted after washing was observed 6/21 (29%) dogs, and two of three dogs tested
290 by DAT were positive.⁵³

291
292 Several studies have assessed coagulation parameters in dogs with IMHA (**Table 3**). Importantly,
293 increased mortality was observed in dogs with IMHA that had thrombocytopenia, prolonged
294 prothrombin time (PT), prolonged activated partial thromboplastin time (APTT), decreased fibrinogen,
295 or DIC.⁴⁰ Reports using thromboelastography determined that 85-100% of dogs with IMHA were
296 hypercoagulable.⁵⁴⁻⁵⁶ Development of DIC is observed in between 10/31 (28%)¹⁴ and 9/20 (45%)¹⁵ dogs
297 with IMHA. One study reported that thromboemboli were found in 20/25 (80%) IMHA dogs at
298 necropsy.¹⁴ The analysis of coagulation parameters is typically recommended in IMHA patients that
299 have clinical signs of coagulopathy. One survey respondent indicated in Question 4 that they
300 recommended measurement of D-dimers in patients suspected of having IMHA.

301 302 Indirect Antiglobulin Tests

303 Indirect antiglobulin tests are not recommended in veterinary species due to low sensitivities and
304 specificities. When these studies were first evaluated for utility in canine patients with IMHA, the

305 sensitivity and specificity of an indirect antiglobulin test were 62.5% and 96.6%, respectively.⁵⁷ The DAT
306 performed by the same laboratory had a 83.3% sensitivity and 98.8% specificity.⁵⁷ None of the survey
307 participants recommended indirect antiglobulin tests in Question 4.

308

309 Direct Antiglobulin Tests (DATs)

310 Various methods for directly detecting RBC surface-bound anti-erythrocyte immunoglobulin (Ig) and
311 opsonizing complement protein (C3) are available. Of the 72 respondents who recommended a test to
312 diagnose IMHA with anti-erythrocyte antibodies, 66 recommended a Coombs' test, three recommended
313 flow cytometry and a Coombs' test, two recommended flow cytometry alone, and one recommended
314 flow cytometry and a Coombs' test at 4°C.

315

316 *Coombs' Tests*

317 Coombs' tests are often performed using a microtiter plate format. Additional methods include gel-
318 based microcolumn, immunochromatographic strip, and capillary DAT assays. Good agreement has been
319 reported between results of the Coombs' test and these methods.²⁵ False-positive DAT results have
320 been reported in anemic dogs. In theory, false-positive results could be due to technical difficulties (e.g.,
321 nonspecific absorption of the antibody, incomplete washing, contamination, assignment of an
322 inappropriate cut off) but patient factors are critical to consider. Dogs that recently received a
323 transfusion can have a positive DAT.^{8,58} Also, dogs with an autoimmune disease that are positive for
324 antinuclear antibodies have been reported to be DAT positive without conclusive evidence of IMHA.⁵⁹
325 Additionally, horses with equine infectious anemia can have a positive Coombs' test.⁶⁰

326

327 The Coombs' test uses species-specific antibodies to detect Ig and/or C3 bound to erythrocytes in a
328 patient blood sample. A positive test results in RBC agglutination. Sixty-five of the people surveyed

329 specified if they recommended a Coombs' test when auto-agglutination was observed, 50 people
330 recommended a Coombs' test if no auto-agglutination was seen (7 specified use of a microtiter plate,
331 and 2 recommended an immunochromatographic strip DAT assay). Fifteen people recommended a
332 Coombs' test with or without auto-agglutination (1 specified the use of a microtiter plate, and another
333 recommended the use of a strip DAT assay). The authors agree with the recent ACVIM consensus
334 statement,⁷ which indicates that a Coombs' test is unnecessary if true auto-agglutination that persists
335 after washing is present.

336

337 When a Coombs' test is warranted, polyvalent and monovalent test antibodies are available. These
338 antibodies are pre-adsorbed onto RBCs from healthy dogs before use in the test. False-negative results
339 can occur with either type of test antibody; therefore, including both polyvalent and monovalent
340 antibodies in a Coombs' test can be beneficial.¹¹ Including both polyvalent and monovalent antibodies
341 increased test performance in a study that reported a sensitivity of 82% and a specificity of 95% when
342 antibodies were combined.⁸

343

344 The antibody binding reaction is temperature-dependent, so it is recommended that testing is
345 performed at both 37°C and 4°C.¹¹ Thirty people specified that they ran a Coombs' test at 37°C, 11
346 respondents performed a Coombs' test at 37°C and 4°C, 1 respondent performed the test at 37°C and
347 room temperature, and (as mentioned above) one person performed flow cytometry plus a Coombs' test
348 at 4°C.

349

350 Also, prozone effects are commonly reported when agglutination is not observed at low serum dilutions
351 (i.e., high antibody concentrations) but is observed at higher serum dilutions (i.e., low antibody
352 concentrations). This is due to the presence of excess immunoglobulins that interfere with agglutination

353 induced by the interactions of the test antibodies with Ig and C3 on the RBCs.⁶¹ This improper test
354 antibody to anti-RBC Ig ratio leads to false-negative results if there are not enough serial serum dilutions
355 tested. A prozone effect was observed in 17/126 (13%) samples tested by Piek *et al.* (2012).⁶²

356
357 The reported performance of Coombs' tests vary, likely due to different samples, protocols, and test
358 reagents that are used at different laboratories. Important positive control samples for the Coombs' test
359 include Ig-coated and complement-coated canine RBCs, but these reagents are not readily available.⁹ It
360 has been reported that the use of whole blood in EDTA or acid citrate dextrose (ACD) yields similar
361 results; however, ACD anticoagulants were preferred in one study because of increased sample
362 hemolysis in EDTA.⁹ False-negative results that reduce the sensitivity, and negative predictive value
363 (NPV) of the Coombs' test can be caused by physical properties of the test antibodies (e.g., low antibody
364 affinity, an inappropriate antibody ratio, steric hindrance), poor technique (e.g., excessive washing,
365 delayed processing, assignment of an inappropriate cut-off), or patient factors (drug-dependent
366 reactions, blood transfusions, steroid administration).⁸

367
368 In dogs with IMHA, one study reported that a low percentage, 17/46 (37%), of patients had positive
369 Coombs' test results,¹² but other studies indicated that 77% of dogs with IMHA were positive.^{13,14} In a
370 small study of 12 dogs, the sensitivity, specificity, positive predictive value (PPV), and NPV of the
371 Coombs' test was 58%, 100%, 100%, and 62%, respectively.⁶³ Similarly, Quigley *et al.* (2001) calculated a
372 PPV of 100% and an NPV of 68%.⁶⁴ In a study of cats with IMHA, 2/89 (2%) healthy cats were reported
373 to have a strongly positive Coombs' test at 37°C.⁶⁵ Another study reported that 0/14 (0%) nonanemic
374 cats and 18/55 (33%) anemic cats were Coombs' positive.⁶⁶ Of the 18 Coombs' positive cats, 15 were
375 diagnosed with primary IMHA, two were feline leukemia virus positive (FeLV), and one had
376 cholangiohepatitis.⁶⁶ An older manuscript indicated a weak positive Coombs' test at 4°C in 9/20 (45%)

377 healthy cats and a positive Coombs' test at 4°C and 37°C in 16/20 (80%) anemic cats (12 of the Coombs'
378 positive anemic cats were FeLV positive).⁶⁷

379

380 It is expected that transfusion reactions can cause positive DAT results.^{58,68,69} Honeckman *et al.*
381 indicated that transfusions given 3 to 21 days before a Coombs' test could be particularly problematic.⁵⁸
382 However, only one study was found that reported DAT test results in seven dogs that had been given a
383 recent transfusion; results of two DAT test kits were reported.⁸ Samples were interpreted as truly
384 positive for five dogs (by at least one Coomb's test kit), falsely positive for one dog, and falsely negative
385 for one dog. This study did not specify the length of time between when the transfusion was given and
386 when the diagnostic testing was done. Interestingly, in humans, blood typing is recommended either
387 prior to transfusion or a minimum of 3 months after transfusion to avoid erroneous results.⁷⁰

388

389 Ninety-one of the 94 survey respondents indicated if they would interpret any diagnostic results with
390 caution following a transfusion. The timeframe of concern for people who interpreted results with more
391 caution varied (Fig. 3). It is evident from this small survey that there is uncertainty if recent transfusion
392 would cause false-positive results in CBC, agglutination, or DAT assays.

393

394 *Flow Cytometry Methods*

395 Flow cytometry can also be used to detect immunoglobulins bound to RBCs. One of the first evaluations
396 of flow cytometry as a diagnostic test for IMHA compared data from 12 dogs and three horses with
397 IMHA to 12 healthy animals from each respective species.⁶³ They reported low specificity of a goat anti-
398 equine IgG but 100% specificity of goat anti-equine IgG F(ab')₂ fragment in their assay.⁶³ In dogs, by
399 pooling anti-canine IgG, IgM, IgA, and C3 antibodies, the sensitivity of the test was 100%, specificity was
400 87.5%, PPV was 92%, and NPV was 100%.⁶³ There was no prozone effect with this assay.⁶³ Quigley et al.

401 (2001) reported a PPV of 100%, and an NPV of 93% when they used flow cytometry to evaluate 13 dogs
402 with IMHA and 13 healthy dogs.⁶⁴ In 2008, Morley *et al.* published an assessment of the utility of flow
403 cytometry to detect anti-erythrocyte antibodies in dogs.⁷¹ They found 26/147 (18%) anemic patients
404 had detectable anti-erythrocyte antibodies. This included 17/22 (77%) IMHA patients, 5/14 (36%) IMT
405 patients, and 3/71 (4%) cancer patients. However, 12/145 (8%) nonanemic dogs also were positive for
406 anti-erythrocyte antibodies, which included 3/5 (60%) patients with infectious disease and 5/81 (6%)
407 cancer patients, and the test had a PPV of 70% and NPV of 95%.

408

409 *Direct Enzyme-Linked Antiglobulin Tests (DELATs)*

410 Immunoglobulins and C3 bound to RBCs are detected by comparing the absorbance of patient samples
411 to healthy control animals in the DELAT. Early evaluation of a DELAT indicated that 1 mg/mL p-
412 nitrophenyl phosphate in carbonate buffer is the preferred substrate for the reaction, however false-
413 positive results were observed in 31/60 (52%) dogs tested.⁷² Another evaluation of DELAT performance
414 yielded comparable results with a Coombs' test; 12/23 (52%) samples were Coombs' positive, while
415 13/23 (57%) were DELAT positive.⁷³ To the authors' knowledge, this test for IMHA is no longer readily
416 available.

417

418 **Guidelines for Performing Diagnostic Tests for IMHA**

419 As mentioned previously, technical difficulties and test protocols can profoundly affect the performance
420 of diagnostic tests for IMHA. Tests performed at 4°C, room temperature, and 37°C can provide different
421 results. This document includes two example protocols for saline agglutination testing (Appendix B) and
422 an example of Coombs' testing (Appendix C). In different diagnostic laboratories, it is expected that
423 these protocols will be performed differently but that proper procedures will yield adequate results.

424

425 **Conclusions**

426 Making a definitive diagnosis of IMHA can be difficult due to the variability in patient presentation and
427 diagnostic test performance. Recommended tests for diagnosing IMHA in anemic patients include: 1) a
428 CBC with verified reticulocyte count, manual PCV with assessment of plasma color, and microscopic
429 examination of a (preferably fresh) blood smear, 2) serum chemistry profile, 3) urinalysis, 4) saline
430 agglutination test (preferably with washing), 5) Coombs' test or flow cytometric analysis if the saline
431 agglutination test is negative, 6) coagulation testing for thrombocytopenic patients and patients with
432 clinical signs of coagulopathy, and 7) tests to determine if any underlying disease is present (e.g., drug or
433 toxin exposure, infection, inflammation, neoplasia, other autoimmune diseases). Our survey proved that
434 clinicians choose different tests to diagnose IMHA and that laboratories perform tests differently.
435 Therefore, it is recommended that veterinarians contact a clinical pathologist or technicians or
436 technologist at the diagnostic laboratory they use to obtain details about the reliability of specific tests
437 being performed to diagnose IMHA.

438

439 **References:**

- 440 1. Day MJ, Stokol T, Seino KK. Chapters 33-35. In: Weiss DJ, Wardrop KJ, eds. *Shalm's Veterinary*
441 *Hematology*. 6th ed. Ames, Iowa: Blackwell Publishing Ltd; 2010.
- 442 2. McCullough S. Immune-mediated hemolytic anemia: Understanding the nemesis. *Vet Clin North*
443 *Am - Small Anim Pract*. 2003;33(6):1295-1315. doi:10.1016/j.cvsm.2003.08.003
- 444 3. Skelly B, Swann J. Canine autoimmune hemolytic anemia: management challenges. *Vet Med Res*
445 *Reports*. 2016;Volume 7:101-112. doi:10.2147/VMRR.S81869
- 446 4. Steele BYBF. The Effects of Blood Loss and Blood Destruction Upon the Erythroid Cells in the
447 Bone Marrow of Rabbits. *Erythroid Cells Bone Marrow Lilly Lab Clin Res Indiana Univ Sch Med*
448 *Indianap*. 1932:881-896.
- 449 5. Lucidi C de A, de Rezende CLE, Jutkowitz LA, Scott MA. Histologic and cytologic bone marrow
450 findings in dogs with suspected precursor-targeted immune-mediated anemia and associated
451 phagocytosis of erythroid precursors. *Vet Clin Pathol*. 2017;46(3):401-415.
452 doi:10.1111/vcp.12502
- 453 6. Stockham S, Scott M. *Fundamentals of Veterinary Clinical Pathology*. 2nd ed. Ames, Iowa:
454 Blackwell Publishing; 2008.
- 455 7. Garden OAO, Kidd L, Mexas AAM, et al. ACVIM consensus statement on the diagnosis of immune-
456 mediated hemolytic anemia in dogs and cats. *J Vet Intern Med*. February 2019.
457 doi:10.1111/jvim.15441
- 458 8. Overmann JA, Sharkey LC, Weiss DJ, Borjesson DL. Performance of 2 microtiter canine Coombs'
459 tests. *Vet Clin Pathol*. 2007;36(2):179-183. doi:10.1111/j.1939-165X.2007.tb00205.x
- 460 9. Jones DRE, Gruffydd-Jones TJ, Stokes CR, Bourne FJ. Investigation into factors influencing
461 performance of the canine antiglobulin test. *Res Vet Sci*. 1990;48:53-58.
- 462 10. Reverberi R, Reverberi L. Factors affecting the antigen-antibody reaction. *Blood Transfus*.

- 463 2007;5(4):227-240. doi:10.2450/2007.0047-07
- 464 11. Warman SM, Murray JK, Ridyard A, Eastwood J, Silva S, Day MJ. Pattern of Coombs' test reactivity
465 has diagnostic significance in dogs with immune-mediated haemolytic anaemia. *J Small Anim*
466 *Pract.* 2008;49(10):525-530. doi:10.1111/j.1748-5827.2008.00641.x
- 467 12. Reimer ME, Troy GC, Warnick LD. Immune-mediated hemolytic anemia: 70 cases (1988-1996). *J*
468 *Am Anim Hosp Assoc.* 1999;35(5):384-391. doi:10.5326/15473317-35-5-384
- 469 13. Weinkle TK, Center S a, Randolph JF, Warner KL, Barr SC, Erb HN. Evaluation of prognostic factors,
470 survival rates, and treatment protocols for immune-mediated hemolytic anemia in dogs: 151
471 cases (1993-2002). *J Am Vet Med Assoc.* 2005;226(11):1869-1880.
472 doi:10.2460/javma.2005.226.1869
- 473 14. Carr AR, Panciera DL, Kidd L. Prognostic factors for mortality and thromboembolism in canine
474 immune-mediated hemolytic anemia: A retrospective study of 72 dogs. *J Vet Intern Med.*
475 2002;16(5):504-509. doi:10.1111/j.1939-1676.2002.tb02378.x
- 476 15. Scott-Moncrieff J, Treadwell N, McCullough S, Brooks M. Hemostatic abnormalities in dogs with
477 primary immune-mediated hemolytic anemia. *J Am Anim Hosp Assoc.* 2001;37(3):220-227.
478 doi:10.5326/15473317-37-3-220
- 479 16. Mills JN, Day MJ, Shaw SE, Penhale WJ. Autoimmune haemolytic anaemia in dogs. *Aust Vet J.*
480 1985;62(4):121-123.
- 481 17. Friedenbergs SG, Buhrman G, Chdid L, et al. Evaluation of a DLA-79 allele associated with multiple
482 immune-mediated diseases in dogs. *Immunogenetics.* 2016;68(3):205-217. doi:10.1007/s00251-
483 015-0894-6
- 484 18. Threlfall AJ, Boag AM, Soutter F, Glanemann B, Syme HM, Catchpole B. Analysis of DLA-DQB1 and
485 polymorphisms in CTLA4 in Cocker spaniels affected with immune-mediated haemolytic anaemia.
486 *Canine Genet Epidemiol.* 2015;2:8. doi:10.1186/s40575-015-0020-y

- 487 19. Kennedy LJ, Barnes A, Ollier WER, Day MJ. Association of a common dog leucocyte antigen class II
488 haplotype with canine primary immune-mediated haemolytic anaemia. *Tissue Antigens*.
489 2006;68(6):502-508. doi:10.1111/j.1399-0039.2006.00715.x
- 490 20. Miller SA, Hohenhaus AE, Hale AS. Case-control study of blood type, breed, sex, and bacteremia
491 in dogs with immune-mediated hemolytic anemia. *J Am Vet Med Assoc*. 2004;224(2):232-235.
- 492 21. Kaplan A V, Quimby FW. A radiolabeled staphylococcal protein a assay for detection of anti-
493 erythrocyte IgG in warm agglutinin autoimmune hemolytic anemia of dogs and man. *Vet*
494 *Immunol Immunopathol*. 1983;4:307-317. doi:10.1016/0926-6410(93)90001-L
- 495 22. Barker RN, Jones DRE. Effects of papain on the agglutination of canine red cells with serum
496 autoantibodies. *Res Vet Sci*. 1993;55(2):156-161. doi:10.1016/0034-5288(93)90075-Q
- 497 23. Jones DR. Use of an enzyme indirect antiglobulin test for the diagnosis of autoimmune
498 haemolytic anaemia in the dog. *Res Vet Sci*. 1986;41(2):187-190.
- 499 24. Engelbrecht R, Kohn B, Leibold W, Giger U. Clinical findings, diagnostics and treatment results in
500 primary and secondary immune-mediated hemolytic anemia in dog. *Kleintierpraxis*.
501 2002;47(5):265-278.
- 502 25. Caviezel LL, Raj K, Giger U. Comparison of 4 direct coombs' test methods with polyclonal
503 antiglobulins in anemic and nonanemic dogs for in-clinic or laboratory use. *J Vet Intern Med*.
504 2014;28(2):583-591. doi:10.1111/jvim.12292
- 505 26. Klag AR, Giger U, Shofer FS. Idiopathic immune-mediated hemolytic anemia in dogs: 42 cases
506 (1986-1990). *J Am Vet Med Assoc*. 1993;202:783-788.
- 507 27. Paes G, Paepe D, Meyer E, et al. The use of the rapid osmotic fragility test as an additional test to
508 diagnose canine immune-mediated haemolytic anaemia. *Acta Vet Scand*. 2013;55(1):74.
509 doi:10.1186/1751-0147-55-74
- 510 28. Slappendel RJ, Van Zwieten R, Van Leeuwen M, Schneijdenberg CTWM. Hereditary spectrin

- 511 deficiency in golden retriever dogs. *J Vet Intern Med.* 2005;19(2):187-192. doi:10.1892/0891-
512 6640(2005)19<187:HSDIGR>2.0.CO;2
- 513 29. Bonfanti U, Comazzi S, Paltrinieri S, Bertazzolo W. Stomatocytosis in 7 related Standard
514 Schnauzers. *Vet Clin Pathol.* 2004;33(4):234-239.
- 515 30. Makinde MO, Bobade PA. Osmotic fragility of erythrocytes in clinically normal dogs and dogs
516 infected with parasites. *Res Vet Sci.* 1994;57(3):343-348.
- 517 31. Paltrinieri S, Comazzi S, Agnes F. Haematological parameters and altered erythrocyte metabolism
518 in anaemic dogs. *J Comp Pathol.* 2000;122(1):25-34. doi:10.1053/jcpa.1999.0339
- 519 32. Schaefer DMW, Stokol T. Retrospective study of reticulocyte indices as indicators of iron-
520 restricted erythropoiesis in dogs with immune-mediated hemolytic anemia. *J Vet Diagnostic*
521 *Investig.* 2016;28(3):304-308. doi:10.1177/1040638715618231
- 522 33. Brenner K, Pohlman L, Muldowney I, Petersen D, Schermerhorn T. Peritoneal EMH in a Dog with
523 Immune-Mediated Hemolytic Anemia. *J Am Anim Hosp Assoc.* 2013;49(5):329-332.
524 doi:10.5326/JAAHA-MS-5869
- 525 34. Weiss DJ, Aird B. Cytologic Evaluation of Primary and Secondary Myelodysplastic Syndromes in
526 the Dog. *Vet Clin Pathol.* 2001;30(2):67-75. doi:10.1111/j.1939-165X.2001.tb00261.x
- 527 35. Weiss DJ. Bone Marrow Pathology in Dogs and Cats with Non-Regenerative Immune-Mediated
528 Haemolytic Anaemia and Pure Red Cell Aplasia. *J Comp Pathol.* 2008;138(1):46-53.
529 doi:10.1016/j.jcpa.2007.10.001
- 530 36. Stokol T, Blue JT, French TW. Nonregenerative Immune-Mediated Anemia in Dogs : 43 Cases (
531 1988 – 1999). *J Am Vet Med Assoc.* 2000;216(9):1429-1436.
- 532 37. Swann JW, Szladovits B, Glanemann B. Demographic Characteristics, Survival and Prognostic
533 Factors for Mortality in Cats with Primary Immune-Mediated Hemolytic Anemia. *J Vet Intern*
534 *Med.* 2016;30(1):147-156. doi:10.1111/jvim.13658

- 535 38. Day MJ. Serial monitoring of clinical, haematological and immunological parameters in canine
536 autoimmune haemolytic anaemia. *J Small Anim Pract.* 1996;37(11):523-534. doi:10.1111/j.1748-
537 5827.1996.tb02313.x
- 538 39. Ridyard AE, Shaw DJ, Milne EM. Evaluation of platelet activation in canine immune-mediated
539 haemolytic anaemia. *J Small Anim Pract.* 2010;51(6):296-304. doi:10.1111/j.1748-
540 5827.2010.00935.x
- 541 40. Griebisch C, Arndt G, Kohn B. Evaluation of different prognostic markers in dogs with primary
542 IMHA. *Berliner und Muenchener tieraerztliche Wochenschrift.* 2010;(123):160-168.
- 543 41. Harkin KR, Braselton WE, Tvedten H. Pseudohypophosphatemia in two dogs with immune-
544 mediated hemolytic anemia. *J Vet Intern Med.* 1998;12(3):178-181. doi:10.1111/j.1939-
545 1676.1998.tb02115.x
- 546 42. Shearer LR, Boudreau AE, Holowaychuck MK. Distal Renal Tubular Acidosis and Immune-
547 Mediated Hemolytic Anemia in 3 Dogs. *J Vet Intern Med.* 2009;23:1284-1288.
548 doi:10.1097/OPX.0000000000001132
- 549 43. Gow DJ, Gow AG, Bell R, et al. Serum cardiac troponin I in dogs with primary immune-mediated
550 haemolytic anaemia. *J Small Anim Pract.* 2011;52(5):259-264. doi:10.1111/j.1748-
551 5827.2011.01061.x
- 552 44. Griebisch C, Arndt G, Raila J, Schweigert FJ, Kohn B. C-reactive protein concentration in dogs with
553 primary immune-mediated hemolytic anemia. *Vet Clin Pathol.* 2009;38(4):421-425.
554 doi:10.1111/j.1939-165X.2009.00146.x
- 555 45. Mitchell KD, Kruth SA, Wood RD, Jefferson B. Serum acute phase protein concentrations in dogs
556 with autoimmune hemolytic anemia. *J Vet Intern Med.* 2009;23(3):585-591. doi:10.1111/j.1939-
557 1676.2009.0282.x
- 558 46. Kjelgaard-Hansen M, Goggs R, Wiinberg B, Chan DL. Use of serum concentrations of interleukin-

- 559 18 and monocyte chemoattractant protein-1 as prognostic indicators in primary immune-
560 mediated hemolytic anemia in dogs. *J Vet.* 2011;25:76-82.
- 561 47. Swann JW, Woods K, Wu Y, Glanemann B, Garden OA. Characterisation of the immunophenotype
562 of dogs with primary immune-mediated haemolytic anaemia. *PLoS One.* 2016;11(12):1-18.
563 doi:10.1371/journal.pone.0168296
- 564 48. Zoia A, Gerou-Ferriani M, Drigo M, Caldin M. Case-control study of plasma mean platelet
565 component concentration and survival analysis for dogs with immune-mediated hemolytic
566 anemia. *J Am Vet Med Assoc.* 2018;252(11). doi:10.2460/javma.252.11.1384
- 567 49. Macey MG, Carty E, Webb L, et al. Use of mean platelet component to measure platelet
568 activation on the ADVIA 120 haematology system. *Cytometry.* 1999;38(5):250-255.
- 569 50. Dircks BH, Schuberth H-J, Mischke R. Underlying diseases and clinicopathologic variables of
570 thrombocytopenic dogs with and without platelet-bound antibodies detected by use of a flow
571 cytometric assay: 83 cases (2004–2006). *J Am Vet Med Assoc.* 2009;235(8):960-966.
572 doi:10.2460/javma.235.8.960
- 573 51. Giannuzzi AP, Simone A De, Ricciardi M, Gernone F. Presumptive Ischemic Brain Infarction in a
574 Dog with Evans ' Syndrome. *Case reports Vet Med.* 2014;2014:1-8. doi:10.1155/2014/456524
- 575 52. Jackson ML, Kruth S a. Immune-mediated Hemolytic Anemia and Thrombocytopenia in the Dog:
576 A retrospective study of 55 cases diagnosed from 1979 through 1983 at the Western College of
577 Veterinary Medicine. *Can Vet J.* 1985;26(8):245-250.
- 578 53. Goggs R, Boag a K, Chan DL. Concurrent IMHA and severe thrombocytopenia in 21 dogs. *Vet Rec.*
579 2008;163:323-327. doi:10.1136/vr.163.11.323
- 580 54. Goggs R, Wiinberg B, Kjølgaard-Hansen M, Chan DL. Serial assessment of the coagulation status
581 of dogs with immune-mediated haemolytic anaemia using thromboelastography. *Vet J.*
582 2012;191(3):347-353. doi:10.1016/j.tvjl.2011.03.015

- 583 55. Fenty RK, deLaforcade AM, Shaw SE, O'Toole TE. Identification of hypercoagulability in dogs with
584 primary immune-mediated hemolytic anemia by means of thromboelastography. *J Am Vet Med*
585 *Assoc.* 2011;238(4):463-467. doi:10.2460/javma.238.4.463
- 586 56. Sinnott VB, Otto CM. Use of thromboelastography in dogs with immune-mediated hemolytic
587 anemia: 39 cases (2000-2008): Retrospective Study. *J Vet Emerg Crit Care.* 2009;19(5):484-488.
588 doi:10.1111/j.1476-4431.2009.00455.x
- 589 57. Quimby FW, Smith C, Brushwein M, Lewis RW. Efficacy of immunoserdiagnostic procedures in
590 the recognition of canine immunologic diseases. *Am J Vet Res.* 1980;41:1622-1666.
- 591 58. Honeckman A, Knapp D, Reagan W. Diagnosis of canine immune-mediated hematologic disease.
592 *Compend.* 1996;18(2):113-127.
- 593 59. Jacobs RM, Murtaugh RJ, Crocker DB. Use of a microtiter Coombs' test for study of age, gender,
594 and breed distributions in immunohemolytic anemia of the dog. *J Am Vet Med Assoc.*
595 1984;185(1):66-69.
- 596 60. McGuire TC, Henson JB, Burger D. Complement (C'3)-coated red blood cells following infection
597 with the virus of equine infectious anemia. *J I.* 1969;103(2):293-299.
- 598 61. Hubinont PO. Prozone Phenomenon in Haemagglutination Reactions. *Bmj.* 2009;2(4733):708-
599 711. doi:10.1136/bmj.2.4733.708
- 600 62. Piek CJ, Teske E, van Leeuwen MW, Day MJ. Good agreement of conventional and gel-based
601 direct agglutination test in immune-mediated haemolytic anaemia. *Acta Vet Scand.*
602 2012;54(1):10. doi:10.1186/1751-0147-54-10
- 603 63. Wilkerson MJ, Davis E, Shuman W, Harkin K, Cox J, Rush B. Isotype-specific antibodies in horses
604 and dogs with immune-mediated hemolytic anemia. *J Vet Intern Med.* 2000;14(2):190-196.
605 doi:10.1111/j.1939-1676.2000.tb02235.x
- 606 64. Quigley KA, Chelack BJ, Haines DM, Jackson ML. Application of a direct flow cytometric

607 erythrocyte immunofluorescence assay in dogs with immune-mediated hemolytic anemia and
608 comparison to the direct antiglobulin test. *J Vet Diagnostic Investig.* 2001;13(4):297-300.
609 doi:10.1177/104063870101300403

610 65. Abrams-Ogg ACG, Lim S, Kocmarek H, et al. Prevalence of antinuclear and anti-erythrocyte
611 antibodies in healthy cats. *Vet Clin Pathol.* 2018;47(1):51-55. doi:10.1111/vcp.12570

612 66. Kohn B, Weingart C, Eckmann V, Ottenjann M, Leibold W. Primary immune-mediated hemolytic
613 anemia in 19 cats: Diagnosis, therapy, and outcome (1998-2004). *J Vet Intern Med.*
614 2006;20(1):159-166. doi:10.1892/0891-6640(2006)20[159:PIHAIC]2.0.CO;2

615 67. Dunn JK, Searcy GP, Hirsch VM. The diagnostic significance of a positive direct antiglobulin test in
616 anemic cats. *Can J Comp Med.* 1984;48(4):349-353.

617 68. Switzer JW, Jain NC. Autoimmune hemolytic anemia in dogs and cats. *Vet Clin North Am Small
618 Anim Pract.* 1981;11(2):405-420.

619 69. Zantek ND, Koepsell SA, Tharp DRJ, Cohn CS. The direct antiglobulin test: A critical step in the
620 evaluation of hemolysis. *Am J Hematol.* 2012;87(7):707-709. doi:10.1002/ajh.23218

621 70. Go RS, Winters JL, Kay NE. How I treat autoimmune hemolytic anemia. *Blood.* 2017;129(22):2971-
622 2979. doi:10.1182/blood-2016-11-693689

623 71. Morley P, Mathes M, Guth A, Dow S. Anti-erythrocyte antibodies and disease associations in
624 anemic and nonanemic dogs. *J Vet Intern Med.* 2008;22(4):886-892. doi:10.1111/j.1939-
625 1676.2008.0112.x

626 72. Jones DR, Gruffydd-Jones TJ, Stokes CR, Bourne FJ. Use of a direct enzyme-linked antiglobulin test
627 for laboratory diagnosis of immune-mediated hemolytic anemia in dogs. *Am J Vet Res.*
628 1992;53(4):457-465.

629 73. Campbell KL, George JW. Application of the Enzyme-linked immunosorbent assay to detect
630 canine erythrocyte antibodies. *Am J Vet Res.* 1984;45(4):747-750.

- 631 74. Holahan ML, Brown AJ, Drobatz KJ. The association of blood lactate concentration with outcome
632 in dogs with idiopathic immune-mediated hemolytic anemia: 173 cases (2003-2006). *J Vet Emerg*
633 *Crit Care*. 2010;20(4):413-420. doi:10.1111/j.1476-4431.2010.00551.x
- 634 75. Weiss DJ, Brazzell JL. Detection of activated platelets in dogs with primary immune-mediated
635 hemolytic anemia. *J Vet Intern Med*. 2006;20(3):682-686. doi:10.1892/0891-
636 6640(2006)20[682:DOAPID]2.0.CO;2
- 637

638 **Table 1.** Clinical pathology findings that support the diagnosis of hemolytic anemia.⁶

| CBC/Blood smear exam | Serum Biochemistry | Urinalysis |
|---|--|-----------------------------|
| Increased polychromasia | Hyperbilirubinemia | Hyperbilirubinuria |
| Reticulocytosis ± rubricytosis ± Howell-Jolly bodies | Hemoglobinemia ⁱ | Hemoglobinuria ⁱ |
| Macrocytosis & anisocytosis | Decreased haptoglobin | Increased urobilinogen |
| Decreased MCHC ^e or increased MCHC ⁱ | concentration ⁱ | |
| Spherocytes ^e ; ghost cells ^l – immune-mediated damage or other cause | Decreased hemopexin concentration ⁱ | |
| Heinz bodies ^e ; eccentrocytes ^l – oxidative damage | | |
| Hemoparasites – direct physical &/or immune-mediated damage | | |
| Schistocytes ^l ; acanthocytes ^l ; keratocytes ^l – direct physical damage | | |

639 e = specifically associated with extravascular hemolysis

640 i = specifically associated with intravascular hemolysis

641 MCHC = mean cell hemoglobin concentration

642

643 **Table 2.** Serum biochemistry results associated with decreased survival in IMHA patients.

| Abnormality | Species | References |
|---------------------------------------|----------------|-------------------|
| Hyperbilirubinemia | Dog | 12-14,40 |
| | Cat | 37 |
| Hyperlactatemia | Dog | 74 |
| Increased alanine aminotransferase | Dog | 40 |
| Increased aspartate aminotransferase | Dog | 40 |
| Increased urea nitrogen | Dog | 40 |
| Increased alkaline phosphatase | Dog | 12 |
| Hypoalbuminemia (< 3.0 mg/dL) | Dog | 13 |
| Hypophosphatemia (< 3.5 mEq/L) | Dog | 13 |
| Increased creatine kinase (> 250 U/L) | Dog | 13 |

644

645 **Table 3.** Evidence of coagulation abnormalities in dogs with IMHA.

| Abnormality | Dogs with IMHA affected | References |
|--|--------------------------------|-------------------|
| Increased prothrombin time | 10-28% | 14,15 |
| Increased activated partial thromboplastic time | 45-47% | 14,15 |
| Increased fibrin degradation products | 57-60% | 14,15 |
| Increased fibrinogen | 17/20 (85%) | 15 |
| Increased D-dimers | 16/20 (80%) | 15 |
| Decreased anti-thrombin III | 10/20 (50%) | 15 |
| Increased Russell viper venom time | 7/20 (35%) | 15 |
| Increased von Willebrand factor associated antigen | 9/20 (45%) | 15 |
| Increased Kaolin clotting time | 3/20 (15%) | 15 |
| Increased P-selectin* | 15/20 (75%) | 75 |
| Hypercoagulability using thromboelastography | 85-100% | 54-56 |
| Disseminated intravascular coagulation | 28-45% | 14,15 |
| Thromboemboli found at necropsy | 20/25 (80%) | 14 |

646 * A second paper saw no increase in P-selectin.¹³

647 **Figure Legends:**

648 **Figure 1.** Proportions (above 75%) of hematologic parameters reported or received on the CBC results of
649 patients suspected of having IMHA by survey respondents (n=94).

650

651 **Figure 2.** Microscopic evidence of agglutination in a wet-mount saline agglutination test. Peripheral
652 blood in EDTA (0.15%) was diluted 1:4 in isotonic saline (0.9% NaCl). A drop of the mixture was placed
653 on a glass slide, and a coverslip was placed over the drop. Grape-like aggregates of erythrocytes can be
654 observed (unstained, 200× magnification).

655

656 **Figure 3.** Survey responses indicating the post-transfusion time-frame in days (d) during which
657 respondents would cautiously interpret hematologic test results that support a diagnosis of IMHA
658 (n=94).

Survey of Tests Used to Diagnose Immune Mediated Hemolytic Anemia (IMHA) in Veterinary Medicine
Designed by the Diagnostic Task Force of the Veterinary and Comparative Clinical Immunology Society
Data from this survey will be presented at the ACVP/ASVCP Meeting in November 2018. Thank you for taking the time to answer these questions about the diagnosis of IMHA.

Question Title

1. Which parameters do you report or receive as part of a complete blood count when an animal is suspected of having IMHA? Please check all that apply.

- Packed cell volume (PCV)
- Plasma Total Protein (TP)
- Fibrinogen using heat precipitation
- Fibrinogen using a turbidimetric assay
- Hemolysis
- Icterus
- Lipemia
- Red blood cell (RBC) count
- Hematocrit
- Hemoglobin concentration
- Mean corpuscular volume (MCV)
- Mean corpuscular hemoglobin (MCH)
- Mean corpuscular hemoglobin content (MCHC)
- Red cell distribution width (RDW)
- Spherocytes
- Ghost cells
- Acanthocytes
- Shistocytes
- Keratocytes
- Hemoparasites
- Relative numbers of nucleated RBCs (nRBCs/100 WBCs)
- Absolute nRBC count
- Manual reticulocyte count using New Methyene Blue
- Manual reticulocyte count using Brilliant Cresyl Blue
- Automated hematology instrument reticulocyte count
- Average mean reticulocyte volume
- Average reticulocyte hemoglobin concentration
- Average hemoglobin content of reticulocytes
- White blood cell count

- Automated hematology instrument differential count
- Manual 200-cell differential count
- Manual 100-cell differential count
- Blood smear review by a board-certified clinical pathologist
- Saline agglutination test at a 1:1 dilution
- Saline agglutination test at a 1:4 dilution
- Saline agglutination test without extensive washing
- Saline agglutination test with extensive washing
- Saline agglutination test by macroscopic evaluation
- Saline agglutination test by microscopic evaluation

Question Title

2. If not already included in the CBC, what additional testing do you recommend for IMHA? Please check all that apply.

- Saline agglutination test at a 1:1 dilution
- Saline agglutination test at a 1:4 dilution
- Saline agglutination test without extensive washing
- Saline agglutination test with extensive washing
- Saline agglutination test by macroscopic evaluation
- Saline agglutination test by microscopic evaluation
- Direct antigen testing using Coombs' test only if there is no auto-agglutination
- Direct antigen testing using Coombs' test with or without auto-agglutination
- Direct antigen testing using Coombs' test at 37 degrees C
- Direct antigen testing using Coombs' test at 4 degrees C
- Direct antigen testing using a Coombs' test microtiter assay
- Direct antigen testing using a Coombs' test gel microcolumn assay
- Direct antigen testing using a Coombs' test immunochromatographic strip assay
- Direct antigen testing using a Coombs' test capillary assay
- Flow cytometry to detect anti-RBC antibodies
- Biochemistry panel
- Urinalysis
- Bone marrow aspiration needle biopsy
- Bone marrow core biopsy
- Infectious disease screening for tick-borne diseases
- Infectious disease screening for viral diseases
- Infectious disease screening for a large panel of different organisms

- Anti-nuclear antigen testing
- Osmotic fragility testing
- Diagnostic imaging

Question Title

3. If the animal was transfused recently, do you interpret test results more cautiously? Please check all that apply.

- Yes, if the transfusion occurred within the last 24 hours
- Yes, if the transfusion occurred 1-6 days ago
- Yes, if the transfusion occurred 7-14 days ago
- No

Question Title

4. Please indicate if you use a test method or report out a parameter that you feel supports a diagnosis of IMHA, but was not listed in Question 1 or Question 2.

Appendix B

Saline agglutination:

The following is a very basic protocol for testing for agglutination.¹ This protocol is likely to give some false positive results.² Reagents needed include isotonic saline, a dropper, a clean glass microscope slide, a wooden applicator stick, and a microscope. Results using fresh EDTA blood from the patient should be compared to fresh EDTA blood from a normal animal of the same species.

- Place one drop (50 μ L) of saline onto a glass slide
- Place an applicator stick in a tube of well-mixed whole blood and stir
- Tap the blood-filled applicator stick into the drop of saline and stir
 - This should produce a 1:4 to 1:10 dilution of blood³
 - If needed, a more accurate dilution of blood can be tested
 - Add 5-10 μ L of whole blood to the drop of saline and stir
- Evaluate the slide for agglutination macroscopically and microscopically at 100 \times magnification
 - Agglutination appears as grape-like clusters of RBCs (Fig. 2) and must be distinguished from rouleaux where erythrocytes are arranged in linear stacks.
 - If agglutination is present, the result is positive and supports a diagnosis of IMHA

A more stringent protocol for testing for agglutination is listed below.¹ This protocol can give some false negative results. Reagents needed include isotonic saline, a clean glass microscope slide, micropipettes, a centrifuge, and a microscope. Results using fresh EDTA blood from the patient should be compared to fresh EDTA blood from a normal animal of the same species.

- Wash RBCs
 - Dilute 100 μ L of whole blood in 900 μ L of isotonic saline to wash the RBCs
 - Centrifuge the sample for 5 minutes at 1500 \times g to pellet the RBCs

- Remove the supernatant
- Re-suspend the RBC pellet in 900 μ L of isotonic saline
- Wash the sample in saline (as outlined above) at least 2 additional times
- Place 50 μ L of the re-suspended sample onto a glass slide
- Evaluate the slide for agglutination macroscopically and microscopically at 100 \times magnification
 - Agglutination appears as grape-like clusters of RBCs (Fig. 2) and must be distinguished from rouleaux where erythrocytes are arranged in linear stacks.
 - If agglutination is present, the result is positive and supports a diagnosis of IMHA

1. Harvey J. Hematology Procedures. In: Harvey J, ed. *Veterinary Hematology: A Diagnostic Guide and Color Atlas*. St. Louis: Elsevier Saunders; 2012:13-14.
2. Caviezel LL, Raj K, Giger U. Comparison of 4 direct coombs' test methods with polyclonal antiglobulins in anemic and nonanemic dogs for in-clinic or laboratory use. *J Vet Intern Med*. 2014;28(2):583-591. doi:10.1111/jvim.12292
3. Stockham S, Scott M. *Fundamentals of Veterinary Clinical Pathology*. 2nd ed. Ames, Iowa: Blackwell Publishing; 2008:136.

Appendix C

Coombs' test:

Test kits for Coombs' tests are available for dogs, cats, and horses.¹⁻³ Instructions provided in the kit should be followed. Reagents typically include rabbit antiserum that has been adsorbed with normal species-specific RBCs so that the reagents are unlikely to react with erythrocytes from healthy animals. Blood collected in EDTA, ACD, or heparin can be used. Ideally, results using fresh whole blood from the patient should be compared to fresh whole blood from a normal animal of the same species. However, samples can be stored at 4°C for up to 7 days before testing.⁴

- Wash RBCs
 - Centrifuge blood for 5 minutes at 1500 × g
 - Remove 100 µL of packed RBCs
 - Add 4.9 mL phosphate buffered saline (PBS) and mix
 - Centrifuge the sample for 5 minutes at 1500 × g
 - Remove the supernatant
 - Re-suspend the RBC pellet in 4.9 mL PBS
- Wash the sample in PBS (as outlined above) at least 2 additional times
- Dilute the Coombs' test reagent
 - Label 2 sets of 10 test tubes: 1:2, 1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256, auto, neg
 - Note that the use of microwell titer plates in place of test tubes can dramatically reduce the amount of serum needed.
 - Higher dilutions are recommended based upon the findings of Piek *et al.*⁵
 - Add 100 µL PBS to each test tube
 - Add 100 µL species-specific Coombs' reagent to the test tube labeled 1:2
 - Transfer 100 µL solution from tube 1:2 to tube 1:4

- Transfer 100 μ L solution from tube 1:4 to tube 1:8
- Transfer 100 μ L solution from tube 1:8 to tube 1:16
- Transfer 100 μ L solution from tube 1:16 to tube 1:32
- Transfer 100 μ L solution from tube 1:32 to tube 1:64
- Transfer 100 μ L solution from tube 1:64 to tube 1:128
- Transfer 100 μ L solution from tube 1:128 to tube 1:256
- Perform the Coombs' test
 - Add 100 μ L of washed RBCs from the patient to test tubes 1:2, 1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256, and auto
 - Add 100 μ L of washed RBCs from the normal animal to test tube neg
 - Incubate one set of 10 test tubes for 30 minutes at 37°C
 - Incubate the other set of 10 test tubes for 30 minutes at 4°C
 - Centrifuge for 1 minute at 1500 \times g
 - Hold each tube up to the light at a 45° angle and mix by tapping
 - If RBCs stay in a pellet or break off in clumps, the sample is positive for macro-agglutination
- Evaluate the slides for agglutination macroscopically and microscopically at 100 \times magnification
 - Place 50 μ L each sample onto glass slides
 - Agglutination appears as grape-like clusters of RBCs and must be distinguished from rouleaux where erythrocytes are arranged in linear stacks (Fig. 2).
 - If agglutination is present, the result is positive and supports a diagnosis of IMHA

Accessed May 10, 2019.

2. ALVEDIA. Alice Veterinary Diagnostics. <http://www.alvedia.com/quicktest-dat-canine/>. Accessed May 15, 2019.
3. VMRD. Veterinary Medical Research and Development. <https://www.vmr.com/reagents/detail/canine-coombs-reagent-5-ml/>. Accessed May 15, 2019.
4. Caviezel LL, Raj K, Giger U. Comparison of 4 direct coombs' test methods with polyclonal antiglobulins in anemic and nonanemic dogs for in-clinic or laboratory use. *J Vet Intern Med*. 2014;28(2):583-591. doi:10.1111/jvim.12292
5. Piek CJ, Teske E, van Leeuwen MW, Day MJ. Good agreement of conventional and gel-based direct agglutination test in immune-mediated haemolytic anaemia. *Acta Vet Scand*. 2012;54(1):10. doi:10.1186/1751-0147-54-10