



Bilinear local controllability to the trajectories of the Fokker-Planck equation with a localized control

Michel Duprez, Pierre Lissy

► To cite this version:

Michel Duprez, Pierre Lissy. Bilinear local controllability to the trajectories of the Fokker-Planck equation with a localized control. 2019. hal-02280501

HAL Id: hal-02280501

<https://hal.archives-ouvertes.fr/hal-02280501>

Submitted on 6 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bilinear local controllability to the trajectories of the Fokker-Planck equation with a localized control

MICHEL DUPREZ* PIERRE LISSY †

September 6, 2019

Abstract

This work is devoted to the control of the Fokker-Planck equation, posed on a bounded domain of \mathbb{R}^d ($d \geq 1$). More precisely, the control is the drift force, localized on a small open subset. We prove that this system is locally controllable to regular nonzero trajectories. Moreover, under some conditions on the reference control, we explain how to reduce the number of controls around the reference control. The results are obtained thanks to a linearization method based on a standard inverse mapping procedure and the fictitious control method. The main novelties of the present article are twofold. Firstly, we propose an alternative strategy to the standard fictitious control method: the algebraic solvability is performed and used directly on the adjoint problem. Secondly, we prove a new Carleman inequality for the heat equation with one order space-varying coefficients: the right-hand side is the gradient of the solution localized on a subset (rather than the solution itself), and the left-hand side can contain arbitrary high derivatives of the solution.

Keywords: Controllability, Parabolic equations, Carleman estimates, Fictitious control method, Algebraic solvability.

2010 MSC: 93B05, 93B07, 93B25, 93C10, 35K40.

Contents

1	Introduction and main results	2
1.1	Introduction	2
1.2	Mains results	3
1.2.1	Controls with d components	4
1.2.2	Controllability acting through a control operator	4
2	Null controllability of the linearized system	6
2.1	Carleman estimates	7
2.2	Algebraic resolubility	16
2.3	Regular control	19
3	Controllability to the trajectories	22

*CEREMADE, Université Paris-Dauphine & CNRS UMR 7534, Université PSL, 75016 Paris, France. E-mail: duprez@ceremade.dauphine.fr.

†CEREMADE, Université Paris-Dauphine & CNRS UMR 7534, Université PSL, 75016 Paris, France. E-mail: lissy@ceremade.dauphine.fr.

1 Introduction and main results

1.1 Introduction

Let $T > 0$ and let Ω be a bounded domain in \mathbb{R}^d ($d \in \mathbb{N}^*$), regular enough (for example of class \mathcal{C}^∞). Denote by $Q_T := (0, T) \times \Omega$ and $\Sigma_T := (0, T) \times \partial\Omega$. We consider the following system

$$\begin{cases} \partial_t y &= \Delta y + \operatorname{div}(uy) & \text{in } Q_T, \\ y &= 0 & \text{on } \Sigma_T, \\ y(0, \cdot) &= y^0 & \text{in } \Omega, \end{cases} \quad (1.1)$$

where $y^0 \in L^2(\Omega)$ is the initial data and $u = (u_1, \dots, u_d) \in L^\infty((0, T) \times \Omega)^d$ is the control.

It is well-known (see for instance [23, Theorem and Proposition 3.1]) that for every initial data $y^0 \in L^2(\Omega)$ and every control $u \in L^\infty((0, T) \times \Omega)^d$, there exists a unique solution y to System (1.1) in the space $W(0, T)$, where

$$W(0, T) := L^2((0, T), H_0^1(\Omega)) \cap H^1((0, T), H^{-1}(\Omega)) \hookrightarrow \mathcal{C}^0([0, T]; L^2(\Omega)).$$

Equation (1.1), introduced in [30], is called the Fokker-Planck equation. In the case where the Fokker-Planck equation is posed on the whole space \mathbb{R}^d , it is strongly related to the stochastic differential equation (SDE)

$$\begin{cases} dX_t &= \sum_{i=1}^d u_i(X_t) dt + dW_t & \text{in } (0, T) \times \mathbb{R}^d, \\ X(0, \cdot) &= X^0 & \text{in } \mathbb{R}^d, \end{cases} \quad (1.2)$$

where W_t is the standard multi-dimensional Brownian motion starting from 0. System (1.2) describes the movement of a particule of negligible mass, with constant and isotropic diffusion, under the action of a force field $u = (u_1, \dots, u_d)$.

Under some regularity conditions on the drift term U , it is well-known that, by the Itô Lemma, the probability density function p associated to (1.2) verifies

$$\begin{cases} \partial_t p &= \frac{1}{2} \Delta p + \operatorname{div}(up) & \text{in } (0, T) \times \mathbb{R}^d, \\ p(0, \cdot) &= p^0 & \text{in } \mathbb{R}^d, \end{cases} \quad (1.3)$$

where p^0 is some initial probability density function (see *e.g.* [41, Section 5.3]). By definition of a probability measure, we have $p^0 \geq 0$ a.e. and $\int_{\mathbb{R}^d} p^0 = 1$. It is then very easy to prove that these properties are preserved during time: any solution p of System (1.3) verifies also $p(t, \cdot) \geq 0$ a.e. and $\int_{\mathbb{R}^d} p(t, \cdot) = 1$, for any $t \in [0, T]$ and hence remains a probability measure. We refer to [42] for more explanations on the Fokker-Planck equation, notably in the case of nonlinear drift terms or non-constant and anisotropic diffusion.

However, in the case where we impose Dirichlet boundary conditions as in (1.1), the derivation of the Fokker-Planck equation from a SDE is more difficult: the Brownian motion has to be replaced by an “absorbed” or “killed” Brownian motion, see *e.g.* [11, pp. 31-60]. Moreover, the total mass of the initial condition is not conserved anymore, meaning that the probability of remaining inside Ω decreases in time, and the solution to (1.1) is not a probability density function anymore. We refer to [23, Section 2] for a discussion on the relevance of Dirichlet boundary conditions in this context. Neumann boundary conditions (that would restore the conservation of mass) seem to be beyond the scope of the present article.

While the controllability properties of the scalar linear heat equation in the case of internal control and Dirichlet boundary condition are now well-understood (see notably [32] and [24]), bilinear controllability seems to have been less explored. Equation (1.1) has been studied in [8], in the whole space and with controls localized everywhere in space and time. Concerning bilinear control in the case where the bilinear term $\operatorname{div}(uy)$ is replaced by uy with $u \in L^\infty((0, T) \times \Omega)$, we refer to [9, 10, 27, 28, 26, 29, 34, 40, 44, 45].

Let us mention that bilinear optimal control of parabolic equations has previously been studied. A first result was proved in [1], where a close forth-order in time model is investigated, with controls depending only on time. This result has been extended to second-order parabolic equations firstly in [4] in the one-dimensional case, then in [5] in the multi-dimensional case, still for time-varying controls. For equation (1.1) (in a slightly more general form), the case of space and time-varying controls is treated in [23]. Notably, for a drift term that is affine in the control, the authors prove the existence of optimal controls for general cost functionals, and derive first-order necessary optimality conditions using an adjoint state. The controllability of the continuity equation, *i.e.* System (1.1) without diffusion, has been investigated in [19, 20].

The structure of the article is as follows. In Section 1.2, we give the mains results of the article (Theorem 1.1, resp. Theorem 1.2, which gives a result of local controllability to the trajectories with d components, resp. a reduced number of controls around the reference control) and some remarks. Section 2 is devoted to studying a linearized version of (1.1). In Section 2.1, we prove a new Carleman estimate (Proposition 2.1) for solutions of the linear backward heat equation with one-order terms. The main novelty is that the local observation term is the gradient of the solution of the adjoint problem (2.4). This has already been proved in [17] for constant coefficients. Moreover, we are able to put as many derivatives as we want in the left-hand side of our Carleman estimate, which will be needed for the rest of the proof. In Section 2.2, we explain how to remove some components of the gradient in the Carleman inequality. This is performed by using what we call an argument of “algebraic solvability” (as introduced in [12] in the context of the stabilization of ODEs and in [16] for the study of coupled systems of PDEs), based on ideas developed by Gromov in [25, Section 2.3.8]. This procedure has already been used successfully in [2, 17, 18, 15, 33, 43]. The main novelty compared to the existing literature is that the algebraic solvability is performed directly on the dual problem. Moreover, we are able to get rid of the high order derivatives of the right in order to obtain the final Carleman estimate (2.34). In Section 2.3, we use some arguments coming from optimal control theory in order to derive from our observability inequality the existence of regular enough controls, with a special form, in appropriate weighted spaces. In Section 3, we go back to the nonlinear problem by using a standard strategy coming from [37] together with some adapted inverse mapping Theorem. To finish, in Section 4, we give an example of trajectory for which the local controllability does not hold with a reduced number of controls.

1.2 Mains results

Let (\bar{y}, \bar{u}) be a trajectory of (1.1), *i.e.* verifying

$$\begin{cases} \partial_t \bar{y} &= \Delta \bar{y} + \operatorname{div}(\bar{u} \bar{y}) & \text{in } Q_T, \\ \bar{y} &= 0 & \text{on } \Sigma_T, \\ \bar{y}(0, \cdot) &= \bar{y}^0 \in L^2(\Omega) \setminus \{0\} & \text{in } \Omega. \end{cases} \quad (1.4)$$

1.2.1 Controls with d components

We first state a result of local controllability to the trajectories to System (1.4) with a control containing d components:

THEOREM 1.1. *Let ω be any nonempty open subset of Ω . Assume that the trajectory (\bar{y}, \bar{u}) with $\bar{u} = (\bar{u}_1, \dots, \bar{u}_d)$ of System (1.4) is regular enough (for example of class C^∞ on $(0, T) \times \Omega$), and that there exists some open subset $\omega_{\bar{u}}$, strongly included in Ω , such that the support of \bar{u} is included in $[0, T] \times \omega_{\bar{u}}$.*

Then, System (1.1) is locally controllable with localized controls, in the following sense: for every $\varepsilon > 0$ and every $T > 0$, there exists $\eta > 0$ such that for any $y^0 \in L^2(\Omega)$ verifying

$$\|y^0 - \bar{y}(0)\|_{L^2(\Omega)} \leq \eta, \quad (1.5)$$

there exists a trajectory (y, u) to System (1.1) such that

$$\begin{cases} y(T) & = \bar{y}(T), \\ u & = \bar{u} + v \text{ for some } v \in L^\infty((0, T) \times \Omega)^d, \\ \text{Supp}(v) & \subset (0, T) \times \omega, \\ \|v\|_{L^\infty((0, T) \times \Omega)^d} & \leq \varepsilon, \\ \|y - \bar{y}\|_{W(0, T)} & \leq \varepsilon. \end{cases}$$

Remark 1. • The regularity assumptions on (\bar{y}, \bar{u}) can be improved, notably it is enough that the reference trajectory is C^r for some $r \in \mathbb{N}^*$ large enough, on an open subset of $(0, T) \times \Omega$.

- If $y^0 = 0$, the only solution to (1.1) is $y \equiv 0$, whatever u is, so that the only reachable state at time T is 0. As a consequence, $\eta > 0$ has notably to be chosen small enough such that $y^0 \neq 0$.
- From the results given in [7], as soon as $y^0 \geq 0$, then any trajectory to System (1.1) remains non-negative (see also [23]). This fact differs from the usual linear heat equation with internal control (see [38]).

1.2.2 Controllability acting through a control operator

In this section, we give a result of local controllability to the trajectories to System (1.4) with a control acting through a control operator $B \in \mathcal{M}_{d, m}(\mathbb{R})$ with $m \in \mathbb{N}^*$ such that $m \leq d$.

We first introduce some notations. Let $q \in \mathbb{N}$ and consider the following set

$$\mathcal{E}(m, q) = \{(\alpha_1, \dots, \alpha_m) \in \mathbb{N}^m \mid 0 < \alpha_1 + \dots + \alpha_m \leq q\},$$

with the convention that $\mathcal{E}(m, q) = \emptyset$ if $q = 0$. Note that by an elementary computation,

$$\#\mathcal{E}(m, q) = \frac{(q+m)!}{m!q!} - 1 =: N(m, q).$$

For $v \in \mathbb{R}^m$, we write $Bv = (B_1v, \dots, B_dv) \in \mathbb{R}^d$. For $j \in \{1, \dots, m\}$, we write

$$(B_m^* \cdot \nabla) : \psi \in C^\infty(\mathbb{R}^d) \mapsto B_m^*(\nabla \psi) \in C^\infty(\mathbb{R}^d).$$

For $(\alpha_1, \dots, \alpha_m) \in \mathcal{E}(m, q)$, we introduce the following operator:

$$(B^* \cdot \nabla)^{\alpha_1, \dots, \alpha_m} : \psi \in C^\infty(\mathbb{R}^d) \mapsto \underbrace{(B_1^* \cdot \nabla) \dots (B_1^* \cdot \nabla)}_{\alpha_1 \text{ times}} \dots \underbrace{(B_m^* \cdot \nabla) \dots (B_m^* \cdot \nabla)}_{\alpha_m \text{ times}} \psi \in C^\infty(\mathbb{R}^d).$$

We introduce the following matrix:

$$M_q(\bar{u}) = \begin{pmatrix} & B_1^* & & \\ & \vdots & & \\ & B_m^* & & \\ (B^* \cdot \nabla)^{1,0,\dots,0} \bar{u}_1 & \dots & (B^* \cdot \nabla)^{1,0,\dots,0} \bar{u}_d & \\ (B^* \cdot \nabla)^{0,1,\dots,0} \bar{u}_1 & \dots & (B^* \cdot \nabla)^{0,1,\dots,0} \bar{u}_d & \\ \vdots & \ddots & \vdots & \\ (B^* \cdot \nabla)^{0,\dots,0,q} \bar{u}_1 & \dots & (B^* \cdot \nabla)^{0,\dots,0,q} \bar{u}_d & \end{pmatrix} \in \mathcal{M}_{N(m,q)+m,d}(\mathbb{R}). \quad (1.6)$$

We have the following controllability result.

THEOREM 1.2. *Let $m \in \mathbb{N}^*$ (with possibly $m < d$). Under the hypothesis of Theorem 1.1, assume that there exists $q \in \mathbb{N}$ and some $(t_0, x_0) \in (0, T) \times \omega$ such that*

$$\text{Rank}(M_q(\bar{u})(t_0, x_0)) = d. \quad (1.7)$$

Then, System (1.1) is locally controllable with localized controls, in the following sense: for every $\varepsilon > 0$ and every $T > 0$, there exists $\eta > 0$ such that for any $y^0 \in L^2(\Omega)$ verifying

$$\|y^0 - \bar{y}(0)\|_{L^2(\Omega)} \leq \eta,$$

there exists a trajectory (y, u) to System (1.1) such that

$$\begin{cases} y(T) & = \bar{y}(T), \\ u & = \bar{u} + Bv \text{ for some } v \in L^\infty((0, T) \times \Omega)^m, \\ \text{Supp}(v) & \subset (0, T) \times \omega, \\ \|v\|_{L^\infty((0, T) \times \Omega)^m} & \leq \varepsilon, \\ \|y - \bar{y}\|_{W(0, T)} & \leq \varepsilon. \end{cases}$$

Remark 2. • Remark that if $B = I_d$ (i.e. we control every component of the gradient of u), condition (1.7) is automatically verified for $q = 0$, whatever \bar{u} is. Hence Theorem 1.2 contains the result given in Theorem 1.1. Thus we will only give a proof of Theorem 1.2.

- Condition (1.7) notably implies that q has to be chosen large enough such that $N(m, q) \geq d - m$.
- Assumption (1.7) is generic, in the following sense: if $C^\infty((0, T) \times \omega)^2$ is endowed with the C^q topology, the sets of the functions $(\bar{y}, \bar{u}) \in C^\infty((0, T) \times \omega)^2$ verifying (1.7) is an dense open set.
- In Section 4, we give an example of trajectory which does not satisfy condition (1.7) and for which the local controllability to the trajectories does not hold. It highlights that Condition (1.7) is not artificial. Even if the authors think that Condition 1.7 is optimal, find a necessary and sufficient condition remains on open problem.

Exemple 1.1. We give an explicit example, in order to explain better condition (1.7). Let us assume

that we want to control only the $m(< n)$ first components of the gradient, *i.e.*

$$B = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 \\ 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \end{pmatrix} \in \mathcal{M}_{n,m}(\mathbb{R}).$$

Then for any $q \in \mathbb{N}$ such that $N(m, q) \geq d - m$, we have

$$M_q(\bar{u}) = \begin{pmatrix} 1 & 0 & \dots & \dots & \dots & \dots & 0 \\ 0 & 1 & 0 & & & & 0 \\ \vdots & \ddots & \ddots & \ddots & & & \vdots \\ 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ \partial_{x_1} \bar{u}_1 & \partial_{x_1} \bar{u}_2 & \dots & \dots & \partial_{x_1} \bar{u}_{m+1} & \dots & \partial_{x_1} \bar{u}_d \\ \vdots & & & & \vdots & & \vdots \\ \partial_{x_m} \bar{u}_1 & \dots & \dots & \dots & \partial_{x_m} \bar{u}_{m+1} & \dots & \partial_{x_m} \bar{u}_d \\ \partial_{x_1}^2 \bar{u}_1 & \dots & \dots & \dots & \partial_{x_1}^2 \bar{u}_{m+1} & \dots & \partial_{x_1}^2 \bar{u}_d \\ \vdots & & & & \vdots & & \vdots \\ \partial_{x_m}^q \bar{u}_1 & \dots & \dots & \dots & \partial_{x_m}^q \bar{u}_{m+1} & \dots & \partial_{x_m}^q \bar{u}_{m+1} \end{pmatrix} \in \mathcal{M}_{N(m,q)+m,d}(\mathbb{R}).$$

We observe that $M_q(\bar{u})$ is of maximal rank d if and only if the following matrix:

$$\widetilde{M}_q(\bar{u}) = \begin{pmatrix} \partial_{x_1} \bar{u}_{m+1} & \dots & \partial_{x_1} \bar{u}_d \\ \vdots & & \vdots \\ \partial_{x_m} \bar{u}_{m+1} & \dots & \partial_{x_m} \bar{u}_d \\ \partial_{x_1}^2 \bar{u}_{m+1} & \dots & \partial_{x_1}^2 \bar{u}_d \\ \vdots & & \vdots \\ \partial_{x_m}^q \bar{u}_{m+1} & \dots & \partial_{x_m}^q \bar{u}_{m+1} \end{pmatrix} \in \mathcal{M}_{N(m,q),d-m}(\mathbb{R}),$$

is of maximal rank $d - m$.

2 Null controllability of the linearized system

In what follows, we always assume that the trajectory (\bar{y}, \bar{u}) of (1.4) verifies the hypothesis of Theorem 1.1. Consider the following linear parabolic system

$$\begin{cases} \partial_t y & = \Delta y + \operatorname{div}(\bar{u}y) + \operatorname{div}(\theta u) & \text{in } Q_T, \\ y & = 0 & \text{on } \Sigma_T, \\ y(0, \cdot) & = y^0 & \text{in } \Omega, \end{cases} \quad (2.1)$$

where $y^0 \in L^2(\Omega)$ and $\theta \in C^\infty(\bar{\Omega})$ is such that

$$\begin{cases} \text{Supp}(\theta) & \subseteq \omega, \\ \theta & \equiv 1 & \text{in } \omega_0, \\ 0 & \leq \theta \leq 1 & \text{in } \Omega, \end{cases} \quad (2.2)$$

for some non-empty open subset ω_0 which is strongly included in ω . The goal of this section is to prove the null controllability of System (2.1), with less controls than equations and regular enough controls in a special form.

Remark 3. Note that the null controllability of (2.1) is equivalent to the null controllability of the “real” linearized version of (1.1) around (\bar{y}, \bar{u}) given by

$$\begin{cases} \partial_t y & = \Delta y + \text{div}(\bar{u}y) + \text{div}(\bar{y}\tilde{u}) & \text{in } Q_T, \\ y & = 0 & \text{on } \Sigma_T, \\ y(0, \cdot) & = y^0 & \text{in } \Omega. \end{cases} \quad (2.3)$$

Indeed, by unique continuation of the solution \bar{y} of (1.4), as soon as $y^0 \neq 0$, since \bar{y} cannot vanish on a subset of $(0, T) \times \Omega$ of positive measure (see [6]) and \bar{y} is in $C^\infty((0, T) \times \Omega)$, there exists some subset $(T_1, T_2) \times \tilde{\omega}_0$ of $(0, T) \times \omega_0$ such that $|\bar{y}| \geq C > 0$ on $(T_1, T_2) \times \omega_0$, that we can assume to be exactly $(0, T) \times \omega_0$ without loss of generality. Hence, for any $i \in \{1, \dots, d\}$, one can solve (in \tilde{u}_i) the equation $\theta u_i = \bar{y}\tilde{u}_i$ by posing

$$\tilde{u}_i = \frac{\theta u_i}{\bar{y}}.$$

Remark that \tilde{u}_i enjoys the same regularity properties as u_i .

2.1 Carleman estimates

Let us consider the following adjoint system associated to System (2.1)

$$\begin{cases} -\partial_t \psi & = \Delta \psi + \bar{u} \cdot \nabla \psi & \text{in } Q_T, \\ \psi & = 0 & \text{on } \Sigma_T, \\ \psi(T, \cdot) & = \psi^0 & \text{in } \Omega. \end{cases} \quad (2.4)$$

First of all, we will introduce some notations. We denote by $|\cdot|$ the euclidean norm on \mathbb{R}^M , whatever $M \in \mathbb{N}^*$ is. For $s, \lambda > 0$ and $p \geq 1$, let us define the two following functions:

$$\alpha(t, x) := \frac{\exp((2p+2)\lambda\|\eta^0\|_\infty) - \exp[\lambda(2p\|\eta^0\|_\infty + \eta^0(x))]}{t^p(T-t)^p} \quad (2.5)$$

and

$$\xi(t, x) := \frac{\exp[\lambda(2p\|\eta^0\|_\infty + \eta^0(x))]}{t^p(T-t)^p}. \quad (2.6)$$

Here, $\eta^0 \in C^\infty(\bar{\Omega})$ is a function satisfying

$$|\nabla \eta^0| \geq \kappa \text{ in } \Omega \setminus \omega_1, \quad \eta^0 > 0 \text{ in } \Omega \quad \text{and} \quad \eta^0 = 0 \text{ on } \partial\Omega,$$

with $\kappa > 0$ and ω_1 some open subset verifying $\omega_1 \subset\subset \omega_0$. The proof of the existence of such a function η^0 can be found in [24, Lemma 1.1, Chap. 1] (see also [13, Lemma 2.68, Chap. 2]). We will use the two notations

$$\alpha^*(t) := \max_{x \in \Omega} \alpha(t, x) \quad \text{and} \quad \xi^*(t) := \min_{x \in \Omega} \xi(t, x), \quad (2.7)$$

for all $t \in (0, T)$. Note that these maximum and minimum are reached at the boundary $\partial\Omega$. For $s, \lambda > 0$, let us define

$$I(s, \lambda; u) := s^3 \lambda^4 \iint_{Q_T} e^{-2s\alpha} \xi^3 u^2 dxdt + s \lambda^2 \iint_{Q_T} e^{-2s\alpha} \xi |\nabla u|^2 dxdt. \quad (2.8)$$

Let us now give some useful auxiliary results that we will need in our proofs. The first one is a Carleman estimate which holds for solutions of the heat equation with non-homogeneous Neumann boundary conditions:

LEMMA 2.1. *There exists a constant $C > 0$ such that for any $u^0 \in L^2(\Omega)$, $f_1 \in L^2(Q_T)$ and $f_2 \in L^2(\Sigma_T)$, the solution to the system*

$$\begin{cases} -\partial_t u - \Delta u & = f_1 & \text{in } Q_T, \\ \frac{\partial u}{\partial n} & = f_2 & \text{on } \Sigma_T, \\ u(T, \cdot) & = u^0 & \text{in } \Omega \end{cases}$$

satisfies

$$I(s, \lambda; u) \leq C \left(s^3 \lambda^4 \iint_{(0, T) \times \omega_1} e^{-2s\alpha} \xi^3 u^2 dxdt + s \lambda \iint_{\Sigma_T} e^{-2s\alpha} \xi^* f_2^2 d\sigma dt + \iint_{Q_T} e^{-2s\alpha} f_1^2 dxdt \right),$$

for all $\lambda \geq C$ and $s \geq C(T^p + T^{2p})$.

Lemma 2.1 is proved in [22, Theorem 1] in the case $p = 1$. However, following the steps of the proof given in [22], one can prove exactly the same inequality for any $p \in \mathbb{N}^*$.

From Lemma 2.1, one can deduce the following result:

LEMMA 2.2. *Let $f \in L^2(\Sigma_T)$, $G = (g_1, \dots, g_d) \in L^\infty(Q_T)^d$ and $h \in L^2(Q_T)$. Then there exists a constant $C > 0$ such that for every $\varphi^T \in L^2(\Omega)$, the solution φ to the system*

$$\begin{cases} -\partial_t \varphi & = \Delta \varphi + G \cdot \nabla \varphi + h & \text{in } Q_T, \\ \frac{\partial \varphi}{\partial n} & = f & \text{on } \Sigma_T, \\ \varphi(T, \cdot) & = \varphi^T & \text{in } \Omega \end{cases}$$

satisfies

$$I(s, \lambda; \varphi) \leq C \left(s^3 \lambda^4 \iint_{(0, T) \times \omega_1} e^{-2s\alpha} \xi^3 \varphi^2 dxdt + s \lambda \iint_{\Sigma_T} e^{-2s\alpha} \xi^* f^2 d\sigma dt + \iint_{Q_T} e^{-2s\alpha} h^2 dxdt \right),$$

for every $\lambda \geq C$ and $s \geq s_0 = C(T^p + T^{2p})$.

The proof of Lemma 2.2 is standard and is left to the reader (one just has to apply Lemma 2.1 and absorb the remaining lower-order terms thanks to the left-hand side).

We will also need the following estimates.

LEMMA 2.3. *Let $r \in \mathbb{R}$. Then there exists $C := C(r, \omega_1, \Omega) > 0$ such that, for every $T > 0$ and every $u \in L^2((0, T), H^1(\Omega))$,*

$$s^{r+2} \lambda^{r+2} \iint_{Q_T} e^{-2s\alpha} \xi^{r+2} u^2 dxdt \leq C \left(s^r \lambda^r \iint_{Q_T} e^{-2s\alpha} \xi^r |\nabla u|^2 dxdt + s^{r+2} \lambda^{r+2} \iint_{(0, T) \times \omega_1} e^{-2s\alpha} \xi^{r+2} u^2 dxdt \right),$$

for every $\lambda \geq C$ and $s \geq C(T^{2p})$.

The proof of this lemma can be found for example in [14, Lemma 3] in the case $p = 9$. However, following the steps of the proof given in [14], one can prove exactly the same inequality for any $p \in \mathbb{N}^*$.

In order to deal with more regular solutions, one needs the following lemma.

LEMMA 2.4. *Let $z_0 \in H_0^1(\Omega)$, $G \in C^\infty(Q_T)^d$ and $f \in L^2(Q_T)^m$. Let us denote by $\mathcal{R} := -\Delta - G \cdot \nabla$ and consider the solution z to the system*

$$\begin{cases} \partial_t z &= \Delta z + G \cdot \nabla z + f & \text{in } Q_T, \\ z &= 0 & \text{on } \Sigma_T, \\ z(0, \cdot) &= z_0 & \text{in } \Omega. \end{cases}$$

Let $n \in \mathbb{N}$. Let us assume that $z_0 \in H^{2n+1}(\Omega)$, $f \in L^2((0, T), H^{2n}(\Omega)) \cap H^n((0, T), L^2(\Omega))$ and satisfy the following compatibility conditions:

$$\begin{cases} g_0 := z_0 \in H_0^1(\Omega), \\ g_1 := f(0, \cdot) - \mathcal{R}g_0 \in H_0^1(\Omega), \\ \vdots \\ g_d := \partial_t^{n-1} f(0, \cdot) - \mathcal{R}g_{d-1} \in H_0^1(\Omega). \end{cases} \quad (2.9)$$

Then $z \in L^2((0, T), H^{2n+2}(\Omega)) \cap H^{n+1}((0, T), L^2(\Omega))$ and we have the estimate

$$\|z\|_{L^2((0, T), H^{2n+2}(\Omega)) \cap H^{n+1}((0, T), L^2(\Omega))} \leq C(\|f\|_{L^2((0, T), H^{2n}(\Omega)) \cap H^n((0, T), L^2(\Omega))} + \|z_0\|_{H^{2n+1}(\Omega)}).$$

It is a classical result that can be easily deduced for example from [21, Th. 6, p. 365].

We are now able to prove the following crucial inequality:

PROPOSITION 2.1. *Let $N \in \mathbb{N}$ with $N \geq 3$. Then, there exists $C > 0$ such that for every $\psi^0 \in L^2(\Omega)$, the corresponding solution ψ to System (2.4) satisfies*

$$\begin{aligned} & \lambda^2 \iint_{Q_T} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi) |\nabla^{N+1} \psi|^2 dx dt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi)^{2N+1} |\nabla \psi|^2 dx dt \\ & \quad + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^* - 2\mu s \alpha^*} (s\xi^*)^{2N+1} |\psi|^2 dx dt \\ & \leq C \lambda^{2N+2} \iint_{(0, T) \times \omega_0} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi)^{2N+1} |\nabla \psi|^2 dx dt \end{aligned} \quad (2.10)$$

for every $\lambda \geq C$ and $s \geq s_0 = C(T^p + T^{2p})$.

Such a Carleman inequality seems new to the authors in the context of non-constant coefficients (it was proved in [17] in the case of constant coefficients). The main improvement comes from the fact that the observation is a gradient of the solution ψ on ω_0 (and not the solution itself). We are also able to introduce as many derivatives of ψ as we want in the left-hand side, as soon as \bar{u}_i is regular enough.

Remark 4. • Note that the proof proposed here relies on the fact that the lower-order terms in equation (2.4) are of order 1, and would fail in the presence of lower-order terms of order 0. Indeed, in the first step of our proof (inequality (2.13)), some term that cannot be absorbed will appear.

- Note that inequality (2.10) automatically implies that any solution ψ of (2.4) lives in high order weighted Sobolev spaces. This is not a surprise since we know that away from the final time $t = T$, any solution of (2.4) is regular.

Proof of Proposition 2.1.

The proof is inspired by [14] and is quite similar to [17]. Let $\mu > 0$. In all what follows, $C > 0$ is a constant that does not depend on s or λ (but that might depend on the other parameters, notably p, N, η, T, μ) and that might change from inequality to inequality. We assume without loss of generality that N is odd (the case N even can be treated similarly).

Let ψ the solution to System (2.4). We introduce the following auxiliary functions:

$$\rho_0^* := e^{-\mu s \alpha^*}, \quad \psi_1 := \rho_0^* \psi. \quad (2.11)$$

Then ψ_1 is solution of

$$\begin{cases} -\partial_t \psi_1 &= \Delta \psi_1 + \bar{u} \cdot \nabla \psi_1 - \partial_t \rho_0^* \psi & \text{in } Q_T, \\ \psi_1 &= 0 & \text{on } \Sigma_T, \\ \psi_1(T, \cdot) &= 0 & \text{in } \Omega. \end{cases} \quad (2.12)$$

We remark that $\phi := \nabla^N \psi_1$ (the operator ∇ applied N times, or in other words, all the derivatives of order N of ψ_1 , ordered for example lexicographically) satisfies the system

$$\begin{cases} -\partial_t \phi = \Delta \phi + \sum_{i=1}^N G_i \cdot \nabla^i \psi_1 + \bar{u} \cdot \nabla \phi - \partial_t \rho_0^* \nabla^N \psi & \text{in } Q_T, \\ \frac{\partial \phi}{\partial n} = \frac{\partial \phi}{\partial n} & \text{on } \Sigma_T, \\ \phi(T, \cdot) = 0 & \text{in } \Omega, \end{cases}$$

where, for any $i \in \{1, \dots, N\}$, G_i is an essentially bounded tensor of appropriated size, whose coefficients are depending only on \bar{u}_i and its derivatives in space up to the order i . Applying Lemma 2.2 to the different components of ϕ , we obtain the following estimate

$$\begin{aligned} I(s, \lambda; \phi) \leq C & \left(s^3 \lambda^4 \iint_{(0, T) \times \omega_1} e^{-2s\alpha} \xi^3 |\phi|^2 dx dt + s \lambda \iint_{\Sigma_T} e^{-2s\alpha^*} \xi^* \left| \frac{\partial \phi}{\partial n} \right|^2 d\sigma dt \right. \\ & \left. + \iint_{Q_T} e^{-2s\alpha} \sum_{i=1}^N |\nabla^i \psi_1|^2 dx dt + \iint_{Q_T} e^{-2s\alpha} |\partial_t \rho_0^* \nabla^N \psi|^2 dx dt \right). \end{aligned} \quad (2.13)$$

The rest of the proof is divided into four steps:

- In a first step, we will estimate the boundary term appearing in the right-hand side of (2.13) by some global interior term involving ψ_1 , which will be absorb later on (in the last step). We will also absorb the last term of the right-hand side under some condition on p .
- In a second step, we will estimate the last term in the right-hand side of (2.13) by some local terms involving $\nabla \psi_1$ and its derivatives on ω_1 , and get rid of the third term of the right-hand side.
- In a third step, we will estimate the high-order local terms created at the previous step by some local terms involving only $\nabla \psi_1$ on ω_0 .
- In a last step, we will use some Poincaré-like inequality in order to recover the variable ψ in the left-hand side and bound the global interior term of the right-hand side involving ψ_1 by an interior term involving $\nabla \psi$. We will conclude by coming back to the original variable ψ , in order to establish (2.10).

Step 1: Let $\tilde{\theta} \in C^2(\bar{\Omega})$ a function satisfying

$$\frac{\partial \tilde{\theta}}{\partial n} = \tilde{\theta} = 1 \text{ on } \partial\Omega.$$

An integration by parts of the boundary term leads to

$$\begin{aligned} s\lambda \int_0^T e^{-2s\alpha^*} \xi^* \int_{\partial\Omega} \left| \frac{\partial \phi}{\partial n} \right|^2 d\sigma dt &= s\lambda \int_0^T e^{-2s\alpha^*} \xi^* \int_{\partial\Omega} \frac{\partial \phi}{\partial n} \nabla \phi \cdot \nabla \tilde{\theta} d\sigma dt \\ &= s\lambda \int_0^T e^{-2s\alpha^*} \xi^* \int_{\Omega} \Delta \phi \nabla \phi \cdot \nabla \tilde{\theta} dx dt + s\lambda \int_0^T e^{-2s\alpha^*} \xi^* \int_{\Omega} \nabla(\nabla \tilde{\theta} \cdot \nabla \phi) \cdot \nabla \phi dx dt. \end{aligned}$$

Hence

$$s\lambda \int_0^T e^{-2s\alpha^*} \xi^* \int_{\partial\Omega} \left| \frac{\partial \phi}{\partial n} \right|^2 d\sigma dt \leq C\lambda \int_0^T e^{-2s\alpha^*} s\xi^* \|\psi_1\|_{H^{N+2}(\Omega)} \|\psi_1\|_{H^{N+1}(\Omega)} dt.$$

Using the interpolation inequality

$$\|\psi_1\|_{H^{N+2}(\Omega)} \leq C \|\psi_1\|_{H^{N+1}(\Omega)}^{1/2} \|\psi_1\|_{H^{N+3}(\Omega)}^{1/2}$$

and Young's inequality $ab \leq \frac{a^q}{q} + \frac{b^{q'}}{q'}$ ($\frac{1}{q} + \frac{1}{q'} = 1$) for $a, b \geq 0$ and $q = 4$, we deduce that for any $c \in \mathbb{R}$, we have

$$\begin{aligned} \lambda \int_0^T e^{-2s\alpha^*} s\xi^* \int_{\partial\Omega} \left| \frac{\partial \phi}{\partial n} \right|^2 d\sigma dt &\leq C\lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^c \|\psi_1\|_{H^{N+3}(\Omega)}^{1/2} (s\xi^*)^{(1-c)} \|\psi_1\|_{H^{N+1}(\Omega)}^{3/2} dt \\ &\leq C\lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^{4c} \|\psi_1\|_{H^{N+3}(\Omega)}^2 dt \\ &\quad + C\lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}} \|\psi_1\|_{H^{N+1}(\Omega)}^2 dt. \end{aligned} \tag{2.14}$$

Consider the function $\psi_2 := \rho_1^* \psi_1$, where

$$\rho_1^* := (s\xi^*)^{\frac{2(1-c)}{3}} e^{-s\alpha^*}. \tag{2.15}$$

The function ψ_2 is solution to the system

$$\begin{cases} -\partial_t \psi_2 &= \Delta \psi_2 + \bar{u} \cdot \nabla \psi_2 - \partial_t(\rho_1^*) \psi_1 - \rho_1^* \partial_t(\rho_0^*) \psi & \text{in } Q_T, \\ \psi_2 &= 0 & \text{on } \Sigma_T, \\ \psi_2(T, \cdot) &= 0 & \text{in } \Omega. \end{cases}$$

Using Lemma 2.4 for ψ_2 (remark that the compatibility conditions (2.9) are verified, since $\psi_2(T, \cdot) = 0$ and \bar{u} has spatial support strongly included in Ω), we deduce that

$$\|\psi_2\|_{L^2((0,T), H^{2n+2}(\Omega)) \cap H^{n+1}((0,T), L^2(\Omega))} \leq C \|\partial_t(\rho_1^*) \psi_1 + \rho_1^* \partial_t(\rho_0^*) \psi\|_{L^2((0,T), H^{2n}(\Omega)) \cap H^n((0,T), L^2(\Omega))}, \tag{2.16}$$

for $n = 1, 2, \dots, (N+1)/2$. The definitions of ξ^* and α^* given in (2.7), the definition of ρ_0^* given in

(2.11), and the definition of ρ_1^* given in (2.15) lead to

$$\left\{ \begin{array}{l} |\partial_t \rho_0^*| \leq C(s\xi^*)^{1+\frac{1}{p}} e^{-\mu s\alpha^*}, \\ \vdots \\ |\partial_t^{\frac{N+3}{2}} \rho_0^*| \leq C(s\xi^*)^{\frac{N+3}{2}+\frac{N+3}{2p}} e^{-\mu s\alpha^*}, \\ |\partial_t \rho_1^*| \leq C(s\xi^*)^{\frac{2(1-c)}{3}+1+\frac{1}{p}} e^{-s\alpha^*}, \\ \vdots \\ |\partial_t^{\frac{N+3}{2}} \rho_1^*| \leq C(s\xi^*)^{\frac{2(1-c)}{3}+\frac{N+3}{2}+\frac{N+3}{2p}} e^{-s\alpha^*}. \end{array} \right. \quad (2.17)$$

Remark that for any $k \leq l$, we have

$$|\partial_t^k \rho_0^*| \leq C |\partial_t^l \rho_0^*|. \quad (2.18)$$

Combining (2.16) for $n = (N-1)/2$, (2.17), (2.18) and the equations satisfied by ψ and ψ_1 , we obtain

$$\begin{aligned} & \lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}} \|\psi_1\|_{H^{N+1}(\Omega)}^2 dt \leq C\lambda \left(\int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi_1\|_{L^2(\Omega)}^2 dt \right. \\ & \quad \left. + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi\|_{L^2(\Omega)}^2 dt \right. \\ & \quad \left. + \int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+2+\frac{2}{p}} \|\psi_1\|_{H^{N-1}(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+2+\frac{2}{p}} \|\psi\|_{H^{N-1}(\Omega)}^2 dt \right). \end{aligned} \quad (2.19)$$

In the right-hand side of (2.19), we would like to estimate the terms

$$\int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+2+\frac{2}{p}} \|\psi_1\|_{H^{N-1}(\Omega)}^2 dt \quad \text{and} \quad \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+2+\frac{2}{p}} \|\psi\|_{H^{N-1}(\Omega)}^2 dt.$$

This can be done using exactly the same processus by introducing some appropriate auxiliary weight that multiplies ψ or ψ_1 as in (2.15), using Lemma 2.4 successively for $n = (N-1)/2, \dots, 0$, (2.17) and (2.18). At the end, by gathering all the inequalities, we obtain

$$\begin{aligned} & \lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}} \|\psi_1\|_{H^{N+1}(\Omega)}^2 dt \\ & \leq C\lambda \left(\int_0^T (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi_1\|_{L^2(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi\|_{L^2(\Omega)}^2 dt \right). \end{aligned} \quad (2.20)$$

Applying the same technique also leads to

$$\begin{aligned} & \lambda \int_0^T e^{-2s\alpha^*} (s\xi^*)^{4c} \|\psi_1\|_{H^{N+3}(\Omega)}^2 dt \\ & \leq C\lambda \left(\int_0^T (s\xi^*)^{4c+N+3+\frac{N+3}{p}} \|\psi_1\|_{L^2(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{4c+N+3+\frac{N+3}{p}} \|\psi\|_{L^2(\Omega)}^2 dt \right). \end{aligned} \quad (2.21)$$

From (2.14), (2.20) and (2.21), we deduce that

$$\begin{aligned} & \lambda \int_0^T e^{-2s\alpha^*} s\xi^* \int_{\partial\Omega} \left| \frac{\partial\phi}{\partial n} \right|^2 d\sigma dt \\ \leq & C\lambda \left(\int_0^T (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi_1\|_{L^2(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{4(1-c)}{3}+N+1+\frac{N+1}{p}} \|\psi\|_{L^2(\Omega)}^2 dt \right. \\ & \left. + \int_0^T (s\xi^*)^{4c+N+3+\frac{N+3}{p}} \|\psi_1\|_{L^2(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{4c+N+3+\frac{N+3}{p}} \|\psi\|_{L^2(\Omega)}^2 dt \right). \end{aligned} \quad (2.22)$$

Since we would like the powers in the right-hand side to be equal, it is natural to impose that

$$4c + N + 3 + \frac{N + 3}{p} = \frac{4(1 - c)}{3} + N + 1 + \frac{N + 1}{p},$$

i. e.

$$c = \frac{-3 - p}{8p}. \quad (2.23)$$

Thus, using (2.22) and (2.23), we deduce that

$$\begin{aligned} & \lambda \int_0^T e^{-2s\alpha^*} s\xi^* \int_{\partial\Omega} \left| \frac{\partial\phi}{\partial n} \right|^2 d\sigma dt \\ \leq & C\lambda \left(\int_0^T e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} \|\psi_1\|_{L^2(\Omega)}^2 dt + \int_0^T e^{-2(1+\mu)s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} \|\psi\|_{L^2(\Omega)}^2 dt \right). \end{aligned} \quad (2.24)$$

From (2.13), (2.24), the first line of (2.17) and the definition of ψ_1 given in (2.11), we already deduce that

$$\begin{aligned} I(s, \lambda; \phi) \leq & C \left(s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dx dt + \lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dx dt \right. \\ & \left. + \iint_{Q_T} e^{-2s\alpha} \sum_{i=1}^N |\nabla^i \psi_1|^2 dx dt + \iint_{Q_T} e^{-2s\alpha} (s\xi^*)^{2+\frac{2}{p}} |\nabla^N \psi_1|^2 dx dt \right). \end{aligned}$$

By definition the definition of ξ^* given in (2.7), it is clear that $\xi^* \leq \xi$. Hence, taking p large enough such that $2 + \frac{2}{p} \leq 3$ (*i. e.* $p \geq 2$), s, λ large enough and using the definition of $I(s, \lambda; \phi)$ given in (2.8), we deduce that we can absorb the last term of the right-hand-side, so that we obtain

$$\begin{aligned} I(s, \lambda; \phi) \leq & C \left(s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dx dt + \lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dx dt \right. \\ & \left. + \iint_{Q_T} e^{-2s\alpha} \sum_{i=1}^N |\nabla^i \psi_1|^2 dx dt \right). \end{aligned} \quad (2.25)$$

Step 2: We apply Lemma 2.3 successively with

$$(u, r) = (\nabla^{N-1} \psi_1, 3), \dots, (u, r) = (\nabla \psi_1, 2N - 1).$$

We obtain a sequence of inequalities of the form

$$\begin{aligned}
& s^5 \lambda^6 \iint_{Q_T} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt \leq C \left(s^3 \lambda^4 \iint_{Q_T} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt \right. \\
& \left. + s^5 \lambda^6 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt \right), \\
& \dots \\
& s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \leq C \left(s^{2N-1} \lambda^{2N} \iint_{Q_T} e^{-2s\alpha} \xi^{2N-1} |\nabla^2 \psi_1|^2 dxdt \right. \\
& \left. + s^{2N+1} \lambda^{2N+2} \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right).
\end{aligned}$$

We deduce by starting from the last inequality and using in cascade the other ones that

$$\begin{aligned}
& s^5 \lambda^6 \iint_{Q_T} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \\
& \leq C \left(s^3 \lambda^4 \iint_{Q_T} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt + s^5 \lambda^6 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt \right. \\
& \left. + \dots + s^{2N+1} \lambda^{2N+2} \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right). \tag{2.26}
\end{aligned}$$

Combining (2.25), (2.26) and using the definition of $I(s, \lambda, \phi)$ given in (2.8), we deduce that we can absorb the first term on the right-hand side of (2.26) and obtain

$$\begin{aligned}
& s \lambda^2 \iint_{Q_T} e^{-2s\alpha} \xi |\nabla^{N+1} \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \\
& \leq C \left(\lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dxdt + \iint_{Q_T} e^{-2s\alpha} \sum_{i=1}^{N-1} |\nabla^i \psi_1|^2 dxdt \right. \\
& \left. + s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt + s^5 \lambda^6 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^5 |\nabla^N \psi_1|^2 dxdt + \dots \right. \\
& \left. + s^{2N+1} \lambda^{2N+2} \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right).
\end{aligned}$$

Absorbing the second term of the right-hand side, we deduce that for s, λ large enough, we have

$$\begin{aligned}
& s \lambda^2 \iint_{Q_T} e^{-2s\alpha} \xi |\nabla^{N+1} \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \\
& \leq C \left(\lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dxdt \right. \\
& \left. + s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N} \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right). \tag{2.27}
\end{aligned}$$

Step 3: Now, we consider some open subset ω_2 such that $\omega_1 \subset\subset \omega_2 \subset\subset \omega_0$. We consider some function $\tilde{\theta} \in C^\infty(\Omega, \mathbb{R})$ such that:

- $\text{Supp}(\tilde{\theta}) \subset \omega_2$,
- $\tilde{\theta} = 1$ on ω_1 ,
- $\tilde{\theta} \in [0, 1]$.

Some integrations by parts give

$$\begin{aligned} s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt &\leq s^3 \lambda^4 \iint_{(0,T) \times \omega_2} \theta e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt \\ &\leq C s^3 \lambda^4 \iint_{(0,T) \times \omega_2} (|\nabla(\theta e^{-2s\alpha} \xi^3)| \cdot |\nabla^N \psi_1| \cdot |\nabla^{N-1} \psi_1| + |\theta e^{-2s\alpha} \xi^3| \cdot |\nabla^{N+1} \psi_1| \cdot |\nabla^{N-1} \psi_1|) dxdt. \end{aligned}$$

From the definition of ξ and α given in (2.5) and (2.6), we deduce that

$$|\nabla(\theta e^{-2s\alpha} \xi^3)| \leq C s \lambda e^{-2s\alpha} \xi^4. \quad (2.28)$$

Combining this estimate with Young's inequality, we obtain that for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that for any s and λ large enough, we have

$$\begin{aligned} s^3 \lambda^4 \iint_{(0,T) \times \omega_1} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt &\leq C \left(\varepsilon s^3 \lambda^4 \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^3 |\nabla^N \psi_1|^2 dxdt \right. \\ &\left. + \varepsilon s \lambda^2 \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi |\nabla^{N+1} \psi_1|^2 dxdt + C_\varepsilon s^5 \lambda^6 \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt \right). \end{aligned} \quad (2.29)$$

Combining (2.27) and (2.29), we can absorb the local terms in $|\nabla^{N+1} \psi_1|^2$ and $|\nabla^N \psi_1|^2$ to deduce

$$\begin{aligned} s \lambda^2 \iint_{Q_T} e^{-2s\alpha} \xi |\nabla^{N+1} \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \\ \leq C \left(\lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dxdt + s^5 \lambda^6 \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt \right. \\ \left. + \dots + s^{2N+1} \lambda^{2N+2} \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right). \end{aligned}$$

We can perform exactly the same procedure on the terms

$$s^5 \lambda^6 \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^5 |\nabla^{N-1} \psi_1|^2 dxdt, \dots, s^{2N-1} \lambda^{2N-2} \iint_{(0,T) \times \omega_2} e^{-2s\alpha} \xi^{2N-1} |\nabla^2 \psi_1|^2 dxdt$$

in order to obtain the following estimate:

$$\begin{aligned} s \lambda^2 \iint_{Q_T} e^{-2s\alpha} \xi |\nabla^{N+1} \psi_1|^2 dxdt + \dots + s^{2N+1} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \\ \leq C \left(\lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(r+1)+5r+3}{2r}} |\psi_1|^2 dxdt \right. \\ \left. + s^{2N+1} \lambda^{2N+2} \iint_{(0,T) \times \omega_0} e^{-2s\alpha} \xi^{2N+1} |\nabla \psi_1|^2 dxdt \right). \end{aligned} \quad (2.30)$$

Step 4:

Since the weight $(s\xi^*)^{2N-1}$ does not depend on the space variable, and using the definition of α^* and ξ^* given in (2.7), the following Poincaré's inequality holds:

$$\begin{aligned} \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{2N+1} |\psi_1|^2 dxdt &\leq C \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{2N+1} |\nabla \psi_1|^2 dxdt \\ &\leq C \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} (s\xi)^{2N+1} |\nabla \psi_1|^2 dxdt. \end{aligned} \quad (2.31)$$

Combining (2.30) and (2.31), we deduce that for s large enough

$$\begin{aligned}
& \lambda^2 \iint_{Q_T} e^{-2s\alpha} s\xi |\nabla^{N+1} \psi_1|^2 dxdt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} (s\xi)^{2N+1} |\nabla \psi_1|^2 dxdt \\
& \quad + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{2N+1} |\psi_1|^2 dxdt \\
& \leq C \left(\lambda \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} |\psi_1|^2 dxdt + \lambda^{2N+2} \iint_{(0,T) \times \omega_0} e^{-2s\alpha} (s\xi)^{2N+1} |\nabla \psi_1|^2 dxdt \right). \tag{2.32}
\end{aligned}$$

We now fix $p \geq 2$ large enough such that

$$\frac{2N(p+1) + 5p + 3}{2p} < 2N + 1,$$

which is clearly possible since $\frac{2N(p+1)+5p+3}{2p} \rightarrow N + \frac{5}{2}$ as $p \rightarrow \infty$ and $N \geq 3$ (so that $N + 5/2 < 2N + 1$).

Using that $e^{-2s\alpha^*} (s\xi^*)^{\frac{2N(p+1)+5p+3}{2p}} \leq C e^{-2s\alpha} (s\xi)^{2N+1}$, we deduce by absorbing the first term of the right-hand side of (2.32) that

$$\begin{aligned}
& \lambda^2 \iint_{Q_T} e^{-2s\alpha} (s\xi) |\nabla^{N+1} \psi_1|^2 dxdt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha} (s\xi)^{2N+1} |\nabla \psi_1|^2 dxdt \\
& + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^*} (s\xi^*)^{2N+1} |\psi_1|^2 dxdt \leq C \lambda^{2N+2} \iint_{(0,T) \times \omega_0} e^{-2s\alpha} (s\xi)^{2N+1} |\nabla \psi_1|^2 dxdt.
\end{aligned}$$

Going back to ψ thanks to (2.12), we deduce (2.10). ■

2.2 Algebraic resolubility

In this section, we will derive a new Carleman inequality, adapted to the control problem with less controls we want to prove. We assume here that $q \in \mathbb{N}^*$ (if $q = 0$, necessarily, by condition (1.7), we have $m = d$ and we can take $\mathcal{M} = (B^*)^{-1}$ and $\mathcal{M}_2 = 0$ in the following Lemma).

LEMMA 2.5. *Let $m \in \mathbb{N}^*$ such that $m \leq d - 1$. Assume that the \bar{u} is regular enough (for example of class C^∞).*

Consider two partial differential operators $\mathcal{L}_1 : C^\infty(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}^d)^m$ and $\mathcal{L}_2 : C^\infty(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}^d)$ defined for every $\varphi \in C^\infty(\mathbb{R}^d)$ by

$$\mathcal{L}_1 \varphi := B^*(\nabla \varphi) \text{ and } \mathcal{L}_2 \varphi := \partial_t \varphi + \Delta \varphi + (\bar{u} \cdot \nabla) \varphi.$$

Assume that (1.7) holds.

There exists an open subset $(t_1, t_2) \times \tilde{\omega}$ of $(0, T) \times \omega$ and there exist two partial differential operators $\mathcal{M}_1 : C^\infty(\mathbb{R}^d)^m \rightarrow C^\infty(\mathbb{R}^d)^d$ (of order 1 in time and $q + 1$ in space) and $\mathcal{M}_2 : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R}^d)^d$ (of order 0 in time and q in space) such that

$$\mathcal{M}_1 \circ \mathcal{L}_1 + \mathcal{M}_2 \circ \mathcal{L}_2 = \nabla \text{ in } C^\infty((t_1, t_2) \times \tilde{\omega}). \tag{2.33}$$

Proof of Lemma 2.5:

Since (1.7) is verified, there exists an open subset $(t_1, t_2) \times \tilde{\omega}$ of $(0, T) \times \omega$ and a constant $C > 0$ such that $|\det(M)| > C$ on $(t_1, t_2) \times \tilde{\omega}$. Let $j \in \{1, \dots, m\}$. We call \mathcal{L}_1^j the j -th line of \mathcal{L}_1 . We remark that

$$\begin{aligned}
(B_j^* \cdot \nabla) \mathcal{L}_2 \varphi - (\partial_t + \Delta) \mathcal{L}_1^j \varphi - (\bar{u} \cdot \nabla) \mathcal{L}_1^j \varphi &= (B_j^* \cdot \nabla)(\bar{u} \cdot \nabla) \varphi - (\bar{u} \cdot \nabla)(B_j^* \cdot \nabla) \varphi \\
&= (\bar{u} \cdot \nabla)(B_j^* \cdot \nabla) \varphi + \sum_{k=1}^d ((B_j^* \cdot \nabla) \bar{u}_k) \partial_k \varphi \\
&\quad - (\bar{u} \cdot \nabla)(B_j^* \cdot \nabla) \varphi \\
&= \sum_{k=1}^d ((B_j^* \cdot \nabla) \bar{u}_k) \partial_k \varphi \\
&=: \mathcal{L}_3.
\end{aligned}$$

Now, for some $l \in \{1, \dots, m\}$, the same computations easily give

$$(B_l^* \cdot \nabla) \mathcal{L}_3 \varphi - \sum_{k=1}^d ((B_j^* \cdot \nabla) \bar{u}_k) \partial_k \mathcal{L}_1^l \varphi = \sum_{k=1}^d ((B_l^* \cdot \nabla)(B_j^* \cdot \nabla) \bar{u}_k) \partial_k \varphi =: \mathcal{L}_4 \varphi.$$

Continuing this procedure, we can easily create two partial differential operators $\widetilde{\mathcal{M}}_1$ (of order 1 in time and $q+1$ in space) and $\widetilde{\mathcal{M}}_2$ (of order 0 in time and q in space) such that

$$\widetilde{\mathcal{M}}_1(\mathcal{L}_1(\varphi)) + \widetilde{\mathcal{M}}_2(\mathcal{L}_2(\varphi)) = M_q(\nabla \varphi),$$

where M_q is defined in (1.6). Under condition (1.7), M_q is of maximal rank on $(t_1, t_2) \times \tilde{\omega}$, so that it admits a left inverse at any point of on $(t_1, t_2) \times \tilde{\omega}$. We call $M_q(\bar{u})^{-1}$ any of its left inverses. Then, it is clear that $\mathcal{M}_1 := M_q^{-1} \widetilde{\mathcal{M}}_1$ and $\mathcal{M}_2 := M_q^{-1} \widetilde{\mathcal{M}}_2$ verify (2.33) and have C^∞ coefficients on $(t_1, t_2) \times \tilde{\omega}$. \blacksquare

We now have all the tools to deduce our final Carleman inequality:

PROPOSITION 2.2. *Assume that Condition (1.7) and the hypotheses of Proposition 2.1 hold. Then, for all $\eta \in (0, 1)$, there exists $p \geq 2$, $C > 0$ and $K > 0$ such that for every $\psi^0 \in L^2(\Omega)$, the corresponding solution ψ to System (2.4) satisfies*

$$\begin{aligned}
\int_{\Omega} \psi(0)^2 dx + \iint_{Q_T} e^{\frac{-2K}{\eta(T-t)^p}} \{ \psi^2 + |\partial_t \psi|^2 + \dots + |\partial_{t \dots t}^{\lfloor \frac{N+1}{2} \rfloor} \psi|^2 + |\nabla \psi|^2 + \dots + |\nabla^{N+1} \psi|^2 \} dx dt \\
\leq C e^{K/T^p} \iint_{(0,T) \times \omega_0} e^{\frac{-2K}{(T-t)^p}} |B^*(\nabla \psi)|^2 dx dt.
\end{aligned} \tag{2.34}$$

Proof of Proposition 2.2. We assume that $q \in \mathbb{N}^*$. Let ω_1 some open subset strongly included in ω_0 . Combining Proposition 2.1, Lemma 2.5 (that is still true by replacing ω_0 by ω_1), and the fact that any solution ψ of (2.4) verifies by definition $\mathcal{L}_2 \psi = 0$, we deduce that, for any $\psi^0 \in L^2(\Omega)$, the corresponding solution ψ to System (2.4) satisfies

$$\begin{aligned}
\lambda^2 \iint_{Q_T} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi) |\nabla^{N+1} \psi|^2 dx dt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi)^{2N+1} |\nabla \psi|^2 dx dt \\
+ \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^* - 2\mu s \alpha^*} (s\xi^*)^{2N+1} |\psi|^2 dx dt \\
\leq C \lambda^{2N+2} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha - 2\mu s \alpha^*} (s\xi)^{2N+1} |\mathcal{M}_1 B^*(\nabla \psi)|^2 dx dt,
\end{aligned}$$

where \mathcal{M}_1 is a linear partial differential operator of order 1 in time and $q + 1$ in space, and $\tilde{\theta} \in C^\infty(\Omega, \mathbb{R})$ such that:

- $\tilde{\theta} = 1$ on ω_1 ,
- $\text{Supp}(\tilde{\theta}) \subset \omega_0$,
- $\tilde{\theta} \in [0, 1]$.

We first remark that

$$\begin{aligned} & \lambda^{2N+2} \iint_{Q_T} \theta e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} |\mathcal{M}_1 B^*(\nabla\psi)|^2 dxdt \\ & \leq C\lambda^{2N+2} \iint_{Q_T} \theta e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} \left(\sum_{i=0}^{q+1} (|\nabla^i B^* \nabla\psi|^2 + |\partial_t \nabla^i B^* \nabla\psi|^2) \right) dxdt. \end{aligned}$$

Using that ψ verifies (2.4), we can deduce that

$$\begin{aligned} & \lambda^{2N+2} \iint_{Q_T} \theta e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} |\mathcal{M}_1 B^*(\nabla\psi)|^2 dxdt \\ & \leq C\lambda^{2N+2} \iint_{Q_T} \theta e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} \left(\sum_{i=0}^{q+3} |\nabla^i B^* \nabla\psi|^2 \right) dxdt. \end{aligned}$$

Some integrations by parts give

$$\begin{aligned} & \lambda^{2N+2} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} |\nabla B^*(\nabla\psi)|^2 dxdt \\ & \leq C\lambda^{2N+2} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} |B^*(\nabla\psi)| |\nabla^3\psi| dxdt \\ & + C\lambda^{2N+2} \iint_{Q_T} |\nabla(\tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1})| |B^*(\nabla\psi)| |\nabla^2\psi| dxdt. \end{aligned}$$

Let $\varepsilon > 0$. Young's inequality gives

$$\begin{aligned} & \lambda^{2N+2} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1} |B^*(\nabla\psi)| |\nabla^3\psi| dxdt \\ & \leq C_\varepsilon \lambda^{2N+6} \iint_{(0,T) \times \omega_0} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+5} |B^*(\nabla\psi)|^2 dxdt \\ & + \varepsilon \lambda^{2N-2} \iint_{(0,T) \times \omega_0} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N-3} |\nabla^3\psi|^2 dxdt \end{aligned}$$

and also, by (2.28),

$$\begin{aligned} & \lambda^{2N+2} \iint_{Q_T} |\nabla(\tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+1})| |B^*(\nabla\psi)| |\nabla^2\psi| dxdt \\ & \leq C\lambda^{2N+3} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+2} |B^*(\nabla\psi)| |\nabla^2\psi| dxdt \\ & \leq C_\varepsilon \lambda^{2N+6} \iint_{(0,T) \times \omega_0} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N+5} |B^*(\nabla\psi)| dxdt \\ & + \varepsilon \lambda^{2N} \iint_{(0,T) \times \omega_0} e^{-2s\alpha-2\mu s\alpha^*} (s\xi)^{2N-1} |\nabla^3\psi| dxdt. \end{aligned}$$

Thus, by taking ε small enough, we deduce that

$$\begin{aligned} & \lambda^2 \iint_{Q_T} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi) |\nabla^{N+1}\psi|^2 dxdt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi)^{2N+1} |\nabla\psi|^2 dxdt \\ & \quad + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^* - 2\mu s\alpha^*} (s\xi^*)^{2N+1} |\psi|^2 dxdt \\ & \leq C\lambda^{2N+6} \iint_{(0,T)\times\omega_0} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi)^{2N+5} |B^*(\nabla\psi)|^2 dxdt \\ & \quad + C\lambda^{2N+2} \iint_{Q_T} \tilde{\theta} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi)^{2N+1} \left(\sum_{i=2}^{q+3} |\nabla^i B^* \nabla\psi|^2 \right) dxdt. \end{aligned}$$

By iterating this process for $i = 2, \dots, q+3$, we can get rid of the sum in the right-hand side and obtain

$$\begin{aligned} & \lambda^2 \iint_{Q_T} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi) |\nabla^{N+1}\psi|^2 dxdt + \dots + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi)^{2N+1} |\nabla\psi|^2 dxdt \\ & \quad + \lambda^{2N+2} \iint_{Q_T} e^{-2s\alpha^* - 2\mu s\alpha^*} (s\xi^*)^{2N+1} |\psi|^2 dxdt \\ & \leq C \left(\lambda^{2N+2+4(q+2)} \iint_{(0,T)\times\omega_0} e^{-2s\alpha - 2\mu s\alpha^*} (s\xi)^{2N+1+4(q+2)} |B^*(\nabla\psi)|^2 dxdt \right). \end{aligned}$$

Inequality (2.34) is easily deduced by replacing the space-dependent weights by their infimum in space in the left-hand-side and their supremum in the right-hand side, fixing s and λ large enough, then choosing μ large enough (depending on $\|\eta_0\|_{\infty(\bar{\Omega})}$) with respect to the parameter $\eta \in (0, 1)$, applying usual energy estimates and remarking that the fact that ψ verifies (2.4) enables us to add all the derivatives in time on the left-hand side. ■

2.3 Regular control

Our goal in this section is to construct regular enough controls. Remind that θ is defined in (2.2).

PROPOSITION 2.3. *Let $r \in \mathbb{N}$. Assume that Condition (1.7) holds.*

Under the hypotheses of Proposition 2.1, System

$$\begin{cases} \partial_t y = \Delta y + \operatorname{div}(\bar{u}y) + \operatorname{div}(\theta Bv) & \text{in } Q_T, \\ y = 0 & \text{on } \Sigma_T, \\ y(0, \cdot) = y^0 & \text{in } \Omega, \end{cases} \quad (2.35)$$

is null controllable at time T , i.e. for every $y^0 \in L^2(\Omega)$, there exists a control $v \in L^2(Q_T)^m$ such that the solution z to System (2.35) satisfies $z(T) \equiv 0$ in Ω . Moreover, we can choose $u \in L^2((0, T), H^{2r+2}(\Omega))^m \cap H^{r+1}((0, T), L^2(\Omega))^m$ with

$$\|v\|_{L^2((0,T), H^{2r+2}(\Omega))^m \cap H^{r+1}((0,T), L^2(\Omega))^m} \leq C e^{K/T^p} \|y^0\|_{L^2(\Omega)},$$

where K is the constant in (2.34).

Proof of Proposition 2.3. Let $k \in \mathbb{N}^*$ and let us consider the following optimal control problem

$$\begin{cases} \text{minimize } J_k(v) := \frac{1}{2} \|\tilde{\rho}^{-1/2} v\|_{L^2(Q_T)^m}^2 + \frac{k}{2} \int_{\Omega} |z(T)|^2 dx, \\ v \in \mathcal{U} := \{w \in L^2(Q_T)^m : \tilde{\rho}^{-1/2} w \in L^2(Q_T)^m\}, \end{cases} \quad (2.36)$$

where $\tilde{\rho} := e^{\frac{-2K}{(T-t)^p}}$ (for the $K > 0$ given by Proposition 2.2 with N an even number to be chosen later and some fixed $\eta \in (1/2, 1)$) and z is the solution in $W(0, T)$ to

$$\begin{cases} \partial_t z = \mathcal{A}z + \mathcal{B}v & \text{in } Q_T, \\ y = 0 & \text{on } \Sigma_T, \\ y(0, \cdot) = y^0 & \text{in } \Omega, \end{cases}$$

where

$$\begin{cases} \mathcal{A} := \Delta + \operatorname{div}(\bar{u} \cdot), \\ \mathcal{B} := \operatorname{div}(B\theta \cdot). \end{cases} \quad (2.37)$$

Here, \mathcal{U} is endowed with its natural weighted L^2 -norm.

The functional $J_k : \mathcal{U} \rightarrow \mathbb{R}^+$ is differentiable, coercive and strictly convex on the space \mathcal{U} . Therefore, following [35, [p. 116], there exists a unique solution to the optimal control problem (2.36) and the optimal control v_k is characterized thanks to the solution z_k of the primal system by

$$\begin{cases} \partial_t z_k = \mathcal{A}z_k + \mathcal{B}v_k & \text{in } Q_T, \\ z_k = 0 & \text{on } \Sigma_T, \\ z_k(0, \cdot) = y^0 & \text{in } \Omega, \end{cases} \quad (2.38)$$

the solution φ_k to the dual system

$$\begin{cases} -\partial_t \varphi_k = \mathcal{A}^* \varphi_k & \text{in } Q_T, \\ \varphi_k = 0 & \text{on } \Sigma_T, \\ \varphi_k(T, \cdot) = kz_k(T, \cdot) & \text{in } \Omega \end{cases} \quad (2.39)$$

and the relation

$$\begin{cases} v_k = -\tilde{\rho} \mathcal{B}^* \varphi_k & \text{in } Q_T, \\ v_k \in \mathcal{U}. \end{cases} \quad (2.40)$$

The characterization (2.38), (2.39) and (2.40) of the minimizer v_k of J_k in \mathcal{U} leads to the following computations

$$\begin{aligned} J_k(v_k) &= -\frac{1}{2} \langle \mathcal{B}^* \varphi_k, v_k \rangle_{L^2(Q_T)^m} + \frac{1}{2} \langle z_k(T), \varphi_k(T) \rangle_{L^2(\Omega)} \\ &= -\frac{1}{2} \int_0^T \langle \varphi_k, \mathcal{B}v_k \rangle_{L^2(\Omega)} dt + \frac{1}{2} \int_0^T \{ \langle z_k, \partial_t \varphi_k \rangle_{L^2(\Omega)} + \langle \partial_t z_k, \varphi_k \rangle_{L^2(\Omega)} \} dt \\ &\quad + \frac{1}{2} \langle y^0, \varphi_k(0, \cdot) \rangle_{L^2(\Omega)} \\ &= \frac{1}{2} \langle y^0, \varphi_k(0, \cdot) \rangle_{L^2(\Omega)}. \end{aligned} \quad (2.41)$$

Moreover, using (2.34) with $N = 2s$ and the expression of $\tilde{\rho}$, we infer

$$\|\varphi_k(0, \cdot)\|_{L^2(\Omega)} \leq C e^{K/T^p} \|\tilde{\rho}^{-1/2} v_k\|_{L^2(Q_T)^m}. \quad (2.42)$$

Now, using the definition of J_k , the expression (2.41), the inequality (2.42) and the Cauchy-Schwartz inequality, we infer

$$\|\varphi_k(0, \cdot)\|_{L^2(\Omega)}^2 \leq C e^{2K/T^p} J_k(v_k) \leq C e^{2K/T^p} \|\varphi_k(0, \cdot)\|_{L^2(\Omega)} \|y^0\|_{L^2(\Omega)},$$

from which we deduce

$$\|\varphi_k(0, \cdot)\|_{L^2(\Omega)} \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}. \quad (2.43)$$

Then, using (2.41) and (2.43), we deduce

$$J_k(v_k) \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2. \quad (2.44)$$

Furthermore, we have (see [35, p. 116])

$$\begin{aligned} \|z_k\|_{W(0,T)} &\leq C (\|\mathcal{B}v_k\|_{L^2((0,T),H^{-1}(\Omega))} + \|y^0\|_{L^2(\Omega)}), \\ &\leq C (\|\tilde{\rho}^{-1/2}v_k\|_{L^2(Q_T)^m} + \|y^0\|_{L^2(\Omega)}), \\ &\leq C(1 + C e^{K/T^p}) \|y^0\|_{L^2(\Omega)}, \end{aligned} \quad (2.45)$$

where C does not depend on y^0 and k . Then, using inequalities (2.44) and (2.45), we deduce that there exist subsequences, which are still denoted v_k, z_k , such that the following weak convergences hold:

$$\begin{cases} v_k \rightharpoonup v & \text{in } \mathcal{U}, \\ z_k \rightharpoonup z & \text{in } W(0, T), \\ z_k(T) \rightarrow 0 & \text{in } L^2(\Omega). \end{cases}$$

Passing to the limit in k , z is solution to System (2.37). Moreover, using the expression of J_k given in (2.36) and inequality (2.44), we deduce by letting k going to ∞ that $z(T) \equiv 0$ in Ω . Thus the solution z to System (2.37) with control $v \in \mathcal{U}$ satisfies $z(T) \equiv 0$ in Ω and using (2.44), we obtain the inequality

$$\|v\|_{\mathcal{U}}^2 \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2.$$

Since $\rho^{-1} \geq 1$, using the definition of the norm on \mathcal{U} , we also deduce that

$$\|v\|_{L^2(Q_T)^m}^2 \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2.$$

Now, let us explain why the controls are more regular. First of all, using the fact that φ_k verifies (2.39), we deduce that

$$\|\mathcal{B}^* \varphi_k\|_{L^2(Q_T)}^2 \leq C \|\partial_t \varphi_k\|_{L^2(Q_T)}^2.$$

Hence, for each $i \in \{1, \dots, \frac{N}{2} - 1\}$ and $k \in \mathbb{N}$, using inequalities similar to (2.17) and (2.18), we deduce that for any $\varepsilon > 0$, there exists $C > 0$ such that

$$\begin{aligned} \|\partial_t^i v_k\|_{L^2(Q_T)^m}^2 &= \iint_{Q_T} \partial_t^i (|\tilde{\rho} \mathcal{B}^* \varphi_k|)^2 \\ &\leq C \iint_{Q_T} \tilde{\rho}^{2-2\varepsilon} |\partial_t^{i+1} \varphi_k|^2 \\ &\leq C \iint_{Q_T} \tilde{\rho}^{2-2\varepsilon - \frac{1}{\eta}} \tilde{\rho}^{\frac{1}{\eta}} |\partial_t^{i+1} \varphi_k|^2. \end{aligned} \quad (2.46)$$

Now, we fix $\varepsilon > 0$ small enough (with respect to η) such that $2 - 2\varepsilon - \frac{1}{\eta} \geq 0$. With this choice of ε , we infer that $\tilde{\rho}^{2-2\varepsilon - \frac{1}{\eta}} \leq 1$. Hence, using (2.46) together with (2.34) and (2.44), we deduce that for any $i \in \{0, \dots, \frac{N}{2} - 1\}$, $\|\partial_t^i v_k\| \in L^2(Q_T)$ and

$$\begin{aligned} \|\partial_t^i v_k\|_{L^2(Q_T)^m}^2 &\leq C \iint_{Q_T} \tilde{\varepsilon}^{\frac{-2K}{\eta(T-i)^p}} |\partial_t^{i+1} \varphi_k|^2 \\ &\leq C \iint_{Q_T} e^{\frac{-2K}{(T-i)^p}} |\theta \mathcal{B}^*(\varphi_k)|^2 \\ &\leq C \|v_k\|_{\mathcal{U}}^2 \\ &\leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2. \end{aligned}$$

Thus, extracting one more time a subsequence if necessary and letting k go to $+\infty$, we deduce that for each $i \in \{1, \dots, \frac{N}{2} - 1\}$,

$$\|\partial_i^i v\|_{L^2(Q_T)^m} \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2.$$

We similarly deduce that, for each $i \in \{1, \dots, N - 2\}$,

$$\|\nabla^i v\|_{L^2(Q_T)^{m \times i \times d}} \leq C e^{2K/T^p} \|y^0\|_{L^2(\Omega)}^2.$$

The proof is completed by setting $r = \frac{N}{2} + 1$. ■

3 Controllability to the trajectories

Let $r \in \mathbb{N}$. We use the strategy developed in [37], modifying it slightly to fit our case. Usual interpolation estimates (see [36, Section 13.2, p. 96]) show that

$$L^2((0, T), H^{2r+2}(\Omega)) \cap H^{r+1}((0, T), L^2(\Omega)) \hookrightarrow L^2((0, T), H^{2r+2}(\Omega)) \cap H^1((0, T), H^{2r}(\Omega)),$$

from which we deduce

$$L^2((0, T), H^{2r+2}(\Omega)) \cap H^{r+1}((0, T), L^2(\Omega)) \hookrightarrow L^\infty((0, T), H^{2r}(\Omega)).$$

Now, there exists $R > 0$ large enough such that by Sobolev embeddings, we have

$$L^2((0, T), H^{2R+2}(\Omega)) \cap H^{R+1}((0, T), L^2(\Omega)) \hookrightarrow L^\infty((0, T), W^{1,\infty}(\Omega)).$$

Hence, from Proposition 2.3 and Remark 3, for any $y^0 \in L^2(\Omega)$, there exists a control $v \in L^\infty((0, T), W^{1,\infty}(\Omega))^m$ such that the solution y to System (2.3) satisfies $y(T) \equiv 0$ in Ω and

$$\|v\|_{L^\infty((0, T), W^{1,\infty}(\Omega))^m} \leq C e^{K/T^p} \|y^0\|_{L^2(\Omega)},$$

where $K > 0$ is the constant given by Proposition 2.2 with $N = 2R$ and $p \geq 2$ is given in Proposition 2.1.

Letting the system evolve freely a little bit if needed, we may assume without loss of generality that $y^0 - \bar{y}^0 \in H_0^1(\Omega)$. Indeed, by the regularizing effect, it is very easy to deduce that for any solution (\bar{y}, \bar{u}) to (1.4), there exists some $C(T) > 0$ such that for any solution $(y, 0)$ to (1.1) on $[0, \frac{T}{2}]$, we have $y(\frac{T}{2}) - \bar{y}(\frac{T}{2}) \in H_0^1(\Omega)$ and

$$\|y\left(\frac{T}{2}\right) - \bar{y}\left(\frac{T}{2}\right)\|_{H^1(\Omega)} \leq C(T) \|y^0 - \bar{y}^0\|_{L^2(\Omega)}.$$

Hence, if $\|y^0 - \bar{y}^0\|_{L^2(\Omega)}$ is small, so is $\|y(\frac{T}{2}) - \bar{y}(\frac{T}{2})\|_{H^1(\Omega)}$, so that the condition (1.5) is sufficient for our argument to be valid.

Following [37, p. 24], we introduce the cost of controllability given by

$$\gamma(t) = C e^{K/t^p}, \quad t \in (0, T),$$

and the following weight functions

$$\rho_F(t) = e^{-\frac{\alpha}{(T-t)^{p+1}}}, \quad t \in [0, T]$$

and

$$\rho_0(t) = e^{\frac{\kappa}{(q-1)(T-t)^p} - \frac{\alpha}{q^{2p+2}(T-t)^{p+1}}}, \quad t \in \left[T \left(1 - \frac{1}{q^2} \right), T \right],$$

extended on $[0, T \left(1 - \frac{1}{q^2} \right)]$ by

$$\rho_0(t) = \rho_0 \left(T \left(1 - \frac{1}{q^2} \right) \right), \quad t \in [0, T \left(1 - \frac{1}{q^2} \right)],$$

for some parameters $q > 1$ and $\alpha > 0$ to be chosen later on.

We remark that ρ_F and ρ_0 are non-increasing, verify $\rho_F(T) = \rho_0(T) = 0$ and are related by the relation

$$\rho_0(t) = \rho_F(q^2(T-t) + T)\gamma((q-1)(T-t)), \quad t \in \left[T \left(1 - \frac{1}{q^2} \right), T \right].$$

We introduce for some $\beta > 0$ the weight function

$$\rho(t) = e^{-\frac{\beta}{(T-t)^{p+1}}}.$$

We remark that

$$\rho_F \leq C\rho, \quad \rho_0 \leq C\rho, \quad |\rho'| \rho_0 \leq C\rho^2,$$

as soon as $\beta > 0$ is chosen small enough, precisely

$$\beta < \frac{\alpha}{q^{2p+2}}. \quad (3.1)$$

We introduce the following spaces:

$$\mathcal{F} = \{f \in L^2((0, T) \times \Omega) \text{ such that } \frac{f}{\rho_F} \in L^2((0, T) \times \Omega)\},$$

$$\mathcal{U} = \{u \in L^\infty((0, T), W^{1,\infty}(\Omega))^m \text{ such that } \frac{u}{\rho_0} \in L^\infty((0, T), W^{1,\infty}(\Omega))^m\},$$

$$\begin{aligned} \mathcal{Z} &= \{z \in C^0([0, T], H_0^1(\Omega)) \cap L^2((0, T), H^2(\Omega) \cap H_0^1(\Omega)) \cap H^1((0, T), L^2(\Omega)) \\ &\quad \text{such that } \frac{z}{\rho} \in C^0([0, T], H_0^1(\Omega)) \cap L^2((0, T), H^2(\Omega) \cap H_0^1(\Omega)) \cap H^1((0, T), L^2(\Omega))\}, \end{aligned}$$

endowed with the weighted Sobolev norms naturally induced by the definition of these spaces.

Following [37, Proofs of Propositions 2.5, 2.8] in the spirit of [31, Section 7.2 and Appendix 5], it is easy to obtain the following result.

PROPOSITION 3.1. *For any $z^0 \in H_0^1(\Omega)$ and any $f \in \mathcal{F}$, there exists $v \in \mathcal{U}$ such that the solution z of*

$$\begin{cases} \partial_t z &= \Delta z + \operatorname{div}(\bar{u}z) + \operatorname{div}(\theta \bar{y} B v) + f & \text{in } Q_T, \\ z &= 0 & \text{on } \Sigma_T, \\ z(0, \cdot) &= z^0 & \text{in } \Omega, \end{cases}$$

verifies $z \in \mathcal{Z}$ (and hence $z(T) = 0$).

To conclude, we use the following inverse mapping theorem:

THEOREM 3.1 (see [3]). *Let \mathcal{X} and \mathcal{Y} be Banach spaces and let $M : \mathcal{X} \mapsto \mathcal{Y}$ be a C^1 mapping. Let us assume that the derivative $M'(0) : \mathcal{X} \mapsto \mathcal{Y}$ is onto and let us set $y_0 = M(x_0)$ with $x_0 \in \mathcal{X}$ and $y_0 \in \mathcal{Y}$. Then there exist $\eta > 0$, a mapping $W : B_\eta(y_0) \subset \mathcal{Y} \mapsto \mathcal{X}$ and a constant $K > 0$ satisfying:*

$$\begin{cases} W(z) \in \mathcal{X} \text{ and } M(W(z)) = z \quad \forall z \in B_\eta(y_0), \\ \|W(z) - x_0\|_{\mathcal{X}} \leq K \|z - y_0\|_{\mathcal{Y}} \quad \forall z \in B_\eta(y_0). \end{cases}$$

Proof of Theorem 1.2. We are looking for a solution in the form

$$y(x, t) = \bar{y}(x, t) + w(x, t), \quad u(x, t) = \bar{u}(x, t) + \theta(x)Br(x, t),$$

where (y, u) and (\bar{y}, \bar{u}) are solution to the Systems (1.1) and (1.4), respectively. Then (w, r) is solution to

$$\begin{cases} N(w, r) := \partial_t w - \Delta w - \operatorname{div}(\bar{u}w + \theta Br\bar{y} + \theta Brw) = 0 & \text{in } Q_T, \\ w = 0 & \text{on } \Sigma_T, \\ w(0, \cdot) = y^0 - \bar{y}^0 & \text{in } \Omega. \end{cases}$$

We introduce the following spaces:

$$\mathcal{X} := \{(w, r) \in \mathcal{Z} \times \mathcal{U} \text{ such that } \partial_t w - \Delta w - \operatorname{div}(\bar{u}w + \theta Br\bar{y}) \in \mathcal{F}\},$$

endowed with the norm

$$\|(w, r)\|_{\mathcal{X}} = \|w\|_{\mathcal{Z}} + \|r\|_{\mathcal{U}} + \|\partial_t w - \Delta w - \operatorname{div}(\bar{u}w + \theta Br\bar{y})\|_{\mathcal{F}},$$

and the space

$$\mathcal{Y} = \mathcal{F} \times H_0^1(\Omega),$$

endowed with the norm

$$\|(f, z^0)\|_{\mathcal{Y}} := \|f\|_{\mathcal{F}} + \|z^0\|_{H^1(\Omega)}.$$

Introduce the mapping M given by

$$\begin{aligned} M : \quad \mathcal{X} &\rightarrow \mathcal{Y} \\ (w, r) &\mapsto (N(w, r), w(0, \cdot)). \end{aligned}$$

Let us determine what are the conditions on q, α, β ensuring that M is well-defined. It is clear that

$$\|w(0, \cdot)\|_{H^1(\Omega)} \leq \|w\|_{C^0([0, T], H_0^1(\Omega))} \leq C \left\| \frac{w}{\rho} \right\|_{C^0([0, T], H_0^1(\Omega))} \leq \|(w, r)\|_{\mathcal{X}}.$$

Now, we remark that by definition of the space \mathcal{X} , we have

$$\|\partial_t w - \Delta w - \operatorname{div}(\bar{u}w + \theta Br\bar{y})\|_{\mathcal{F}} \leq \|(w, r)\|_{\mathcal{X}}.$$

Hence, the only difficulty is to treat the bilinear part $\operatorname{div}(\theta w Br)$. We remark that

$$\left\| \frac{\operatorname{div}(\theta w Br)}{\rho_F} \right\|_{L^2((0, T) \times \Omega)} \leq C \left\| \frac{r}{\rho_F^{\frac{1}{2}}} \right\|_{L^\infty((0, T), W^{1, \infty}(\Omega))} \left\| \frac{w}{\rho_F^{\frac{1}{2}}} \right\|_{L^2((0, T), H^1(\Omega))}.$$

We can impose that $\rho^2 \leq C\rho_F$ and $\rho_0^2 \leq C\rho_F$ as soon as

$$\alpha < 2\beta \quad \text{and} \quad q^{2p+2} < 2. \tag{3.2}$$

Remark that these conditions are compatible with condition (3.1).

Hence, under conditions (3.1) and (3.2), we deduce that

$$\left\| \frac{\operatorname{div}(wBq)}{\rho^F} \right\|_{L^2((0,T) \times \Omega)} \leq C \left\| \frac{r}{\rho_0} \right\|_{L^\infty((0,T), W^{1,\infty}(\Omega))} \left\| \frac{w}{\rho} \right\|_{L^2((0,T), H^1(\Omega))} \leq C \|(w, r)\|_{\mathcal{X}}^2.$$

We conclude that under these conditions, M is indeed well-defined and continuous. Moreover, we remark that $M(0,0) = (0,0)$ and M is of class \mathcal{C}^1 as a sum of a linear continuous function and a quadratic continuous function. Furthermore, Proposition 3.1 exactly means that $M'(0,0)$ is onto (see Remark 3). Theorem 3.1 leads to the conclusion. ■

4 Example of a non-controllable trajectory with a reduced number of controls

In this section, we give an example of trajectory which does not satisfy condition (1.7) and for which the local controllability to the trajectories does not hold.

Consider $\bar{u} \in L^\infty(Q_T)^m$ which will be determined later on. Assume that for each $\bar{y}^0 \in L^2(\Omega) \setminus \{0\}$ the following system is locally controllable to the trajectories with a control operator B

$$\begin{cases} \partial_t \bar{y} &= \Delta \bar{y} + \operatorname{div}(\bar{u}\bar{y}) & \text{in } Q_T, \\ \bar{y} &= 0 & \text{on } \Sigma_T, \\ \bar{y}(0, \cdot) &= \bar{y}^0 & \text{in } \Omega. \end{cases}$$

Then for each $\varepsilon \in (0,1)$ small enough, there exists $u \in L^\infty(Q_T)^m$ such that

$$\begin{cases} \partial_t y &= \Delta y + \operatorname{div}(uy) & \text{in } Q_T, \\ y &= 0 & \text{on } \Sigma_T, \\ y(0, \cdot) &= (1 - \varepsilon)\bar{y}^0 & \text{in } \Omega, \\ y(T, \cdot) &= \bar{y}(T) & \text{in } \Omega, \end{cases}$$

where $u = \bar{u} + Bv$ with $\operatorname{Supp}(v) \subset (0,T) \times \omega$. We remark that $(z, w) := (y - \bar{y}, yv)$ is solution to

$$\begin{cases} \partial_t z &= \Delta z + \operatorname{div}(\bar{u}z) + \operatorname{div}(Bw) & \text{in } Q_T, \\ z &= 0 & \text{on } \Sigma_T, \\ z(0, \cdot) &= \varepsilon\bar{y}^0 & \text{in } \Omega, \\ z(T, \cdot) &= 0 & \text{in } \Omega. \end{cases} \quad (4.1)$$

We deduce that System (4.1) is null controllable at time $T > 0$, then approximately controllable at time $T > 0$. It is well known that the approximate controllability of System (4.1) on $(0,T)$ implies the following property, called the Fattorini-Hautus test (see *e.g.* [39]) : for every $s \in \mathbb{C}$ and every $\varphi \in D(\Delta)$,

$$\left. \begin{array}{l} -\Delta\varphi - \bar{u} \cdot \nabla\varphi = s\varphi \quad \text{in } \Omega \\ B^*\nabla\varphi = 0 \quad \text{in } \omega \end{array} \right\} \Rightarrow \varphi = 0. \quad (4.2)$$

We now give an explicit \bar{u} in contradiction with (4.2).

Let $d = 2$, $\Omega = (0, \pi)^2$, $\omega = (5\pi/24, 7\pi/24)^2$ and $B = (1, 0)$.

Consider $\varphi(x_1, x_2) = \varphi_1(x_1)\varphi_2(x_2)$, where $\varphi_1 \in C^\infty([0, \pi], \mathbb{R})$ and $\varphi_2 \in C^\infty([0, \pi], \mathbb{R})$ any two functions verifying

$$\varphi_1(x_1) = \begin{cases} \sin(x_1) & \text{in } [0, \pi] \setminus (\pi/6, 2\pi/6), \\ 1 & \text{in } \omega, \\ \varphi_1 \neq 0 & \text{on } (0, \pi) \end{cases}$$

and

$$\varphi_2(x_2) = \begin{cases} \sin(x_2) & \text{in } [0, \pi] \setminus (\pi/6, 2\pi/6), \\ \sin(2x_2) & \text{in } \omega, \\ \varphi_2'(x_2) \neq 0 & \text{in } [\pi/6, 5\pi/24] \cup [7\pi/24, 2\pi/6]. \end{cases}$$

Remark that it is possible to impose the last condition, since $\varphi_2(\pi/6) \neq \varphi_2(5\pi/24)$ and $\varphi_2(7\pi/24) \neq \varphi_2(2\pi/6)$. Now, we introduce

$$\bar{u} := \begin{cases} (0, 0) & \text{in } (0, \pi/6) \cup (5\pi/24, 7\pi/24) \cup (2\pi/6, \pi), \\ (0, -\frac{2\varphi + \Delta\varphi}{\partial_{x_2}\varphi}) & \text{otherwise.} \end{cases}$$

Remark that \bar{u} is well-defined. Indeed, by construction, $\partial_{x_2}\varphi \neq 0$ in $[\pi/6, 5\pi/24] \cup [7\pi/24, 2\pi/6]$ and $\varphi_1 \neq 0$ on $[0, \pi]$. Moreover, by construction, $2\varphi + \Delta\varphi = 0$ on $(0, \pi/6) \cup (5\pi/24, 7\pi/24) \cup (2\pi/6, \pi)$, so that the extension by 0 of the function $\frac{2\varphi + \Delta\varphi}{\partial_{x_2}\varphi}$ defined on $(0, \pi/6) \cup (5\pi/24, 7\pi/24) \cup (2\pi/6, \pi)$ is of class C^∞ on Ω . We deduce that \bar{u} is of class C^∞ on Ω . To conclude, we remark that by construction,

$$\begin{cases} -\Delta\varphi - \bar{u} \cdot \nabla\varphi = 2\varphi & \text{in } \Omega, \\ \partial_{x_1}\varphi = 0 & \text{in } \omega, \\ \varphi \neq 0. \end{cases}$$

Thus, we obtain a contradiction with (4.2).

Acknowledgements

Pierre Lissy is partially supported by the project IFSMACS (ANR-15-CE40-0010) funded by the french Agence Nationale de la Recherche, 2015-2019.

References

- [1] A. Addou and A. Benbrik. Existence and uniqueness of optimal control for a distributed-parameter bilinear system. *J. Dynam. Control Systems*, 8(2):141–152, 2002.
<https://link.springer.com/article/10.1023/A:1015372725255>.
- [2] F. Alabau-Boussouira, J.-M. Coron, and G. Olive. Internal controllability of first order quasi-linear hyperbolic systems with a reduced number of controls. *SIAM J. Control Optim.*, 55(1):300–323, 2017.
<https://epubs.siam.org/doi/pdf/10.1137/15M1015765>.
- [3] V. M. Alekseev, V. M. Tikhomirov, and S. V. Fomin. *Optimal control*. Contemporary Soviet Mathematics. Consultants Bureau, New York, 1987. Translated from the Russian by V. M. Volosov.
- [4] M. Annunziato and A. Borzi. Optimal control of probability density functions of stochastic processes. *Math. Model. Anal.*, 15(4):393–407, 2010.
<https://www.tandfonline.com/doi/abs/10.3846/1392-6292.2010.15.393-407>.
- [5] M. Annunziato and A. Borzi. A Fokker-Planck control framework for multidimensional stochastic processes. *J. Comput. Appl. Math.*, 237(1):487–507, 2013.
<https://www.sciencedirect.com/science/article/pii/S0377042712002543>.
- [6] J. Apraiz, L. Escauriaza, G. Wang, and C. Zhang. Observability inequalities and measurable sets. *J. Eur. Math. Soc. (JEMS)*, 16(11):2433–2475, 2014.
https://www.ems-ph.org/journals/show_abstract.php?issn=1435-9855&vol=16&iss=11&rank=4.

- [7] D. G. Aronson. Non-negative solutions of linear parabolic equations. *Ann. Scuola Norm. Sup. Pisa (3)*, 22:607–694, 1968.
http://www.numdam.org/article/ASNSP_1968_3_22_4_607_0.pdf.
- [8] A. Blaqui ere. Controllability of a Fokker-Planck equation, the Schr odinger system, and a related stochastic optimal control (revised version). *Dynam. Control*, 2(3):235–253, 1992.
<https://link.springer.com/article/10.1007/BF02169515>.
- [9] P. Cannarsa, G. Floridia, and A. Y. Khapalov. Multiplicative controllability for semilinear reaction-diffusion equations with finitely many changes of sign. *J. Math. Pures Appl. (9)*, 108(4):425–458, 2017.
<https://www.sciencedirect.com/science/article/pii/S0021782417300892>.
- [10] P. Cannarsa and A. Khapalov. Multiplicative controllability for reaction-diffusion equations with target states admitting finitely many changes of sign. *Discrete Contin. Dyn. Syst. Ser. B*, 14(4):1293–1311, 2010.
<https://aimsciences.org/article/doi/10.3934/dcdsb.2010.14.1293>.
- [11] K. L. Chung and Z. X. Zhao. *From Brownian motion to Schr odinger’s equation*, volume 312 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1995.
- [12] J.-M. Coron. Global asymptotic stabilization for controllable systems without drift. *Math. Control Signals Systems*, 5(3):295–312, 1992.
<https://link.springer.com/article/10.1007/BF01211563>.
- [13] J.-M. Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [14] J.-M. Coron and S. Guerrero. Null controllability of the N -dimensional Stokes system with $N - 1$ scalar controls. *J. Differential Equations*, 246(7):2908–2921, 2009.
<https://www.sciencedirect.com/science/article/pii/S0022039608004348>.
- [15] J.-M. Coron and J.-P. Guilleron. Control of three heat equations coupled with two cubic nonlinearities. *SIAM J. Control Optim.*, 55(2):989–1019, 2017.
<https://epubs.siam.org/doi/pdf/10.1137/15M1041201>.
- [16] J.-M. Coron and P. Lissy. Local null controllability of the three-dimensional Navier-Stokes system with a distributed control having two vanishing components. *Invent. Math.*, 198(3):833–880, 2014.
<https://link.springer.com/article/10.1007/s00222-014-0512-5>.
- [17] M. Duprez and P. Lissy. Indirect controllability of some linear parabolic systems of m equations with $m - 1$ controls involving coupling terms of zero or first order. *J. Math. Pures Appl. (9)*, 106(5):905–934, 2016.
<https://www.sciencedirect.com/science/article/pii/S002178241630023X>.
- [18] M. Duprez and P. Lissy. Positive and negative results on the internal controllability of parabolic equations coupled by zero- and first-order terms. *J. Evol. Equ.*, 18(2):659–680, 2018.
<https://link.springer.com/article/10.1007/s00028-017-0415-1>.
- [19] M. Duprez, M. Morancey, and F. Rossi. Approximate and exact controllability of the continuity equation with a localized vector field. *SIAM J. Control Optim.*, 57(2):1284–1311, 2019.
<https://epubs.siam.org/doi/pdf/10.1137/17M1152917>.
- [20] M. Duprez, M. Morancey, and F. Rossi. Minimal time problem for crowd models with a localized vector field. Submitted,
<https://hal.archives-ouvertes.fr/hal-01676679>, 2019.
- [21] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [22] E. Fern andez-Cara, M. Gonz alez-Burgos, S. Guerrero, and J.-P. Puel. Null controllability of the heat equation with boundary Fourier conditions: the linear case. *ESAIM Control Optim. Calc. Var.*, 12(3):442–465 (electronic), 2006.
<https://www.esaim-cocv.org/articles/cocv/abs/2006/03/cocv0515/cocv0515.html>.
- [23] A. Fleig and R. Guglielmi. Optimal control of the Fokker-Planck equation with space-dependent controls. *J. Optim. Theory Appl.*, 174(2):408–427, 2017.
<https://link.springer.com/article/10.1007%2Fs10957-017-1120-5>.
- [24] A. V. Fursikov and O. Y. Imanuvilov. *Controllability of evolution equations*. Number 34. Seoul National University, 1996.

- [25] M. Gromov. *Partial differential relations*, volume 9 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1986.
- [26] A. Y. Khapalov. Global non-negative controllability of the semilinear parabolic equation governed by bilinear control. *ESAIM Control Optim. Calc. Var.*, 7:269–283, 2002.
<https://www.esaim-cocv.org/articles/cocv/abs/2002/01/cocvVo17-11/cocvVo17-11.html>.
- [27] A. Y. Khapalov. On bilinear controllability of the parabolic equation with the reaction-diffusion term satisfying Newton’s law. *Comput. Appl. Math.*, 21(1):275–297, 2002. Special issue in memory of Jacques-Louis Lions.
- [28] A. Y. Khapalov. Controllability of the semilinear parabolic equation governed by a multiplicative control in the reaction term: a qualitative approach. *SIAM J. Control Optim.*, 41(6):1886–1900, 2003.
<https://epubs.siam.org/doi/abs/10.1137/S0363012901394607>.
- [29] A. Y. Khapalov. *Controllability of partial differential equations governed by multiplicative controls*, volume 1995 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2010.
- [30] A. Kolmogoroff. Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung. *Math. Ann.*, 104(1):415–458, 1931.
<https://link.springer.com/article/10.1007/BF01457949>.
- [31] K. Le Balc’h. Local controllability of reaction-diffusion systems around nonnegative stationary states. *Submitted*, 2018.
<https://arxiv.org/abs/1809.05303>.
- [32] G. Lebeau and L. Robbiano. Contrôle exact de l’équation de la chaleur. *Comm. Partial Differential Equations*, 20(1-2):335–356, 1995.
<https://www.tandfonline.com/doi/abs/10.1080/03605309508821097?journalCode=lpde20>.
- [33] T. Liard and P. Lissy. A Kalman rank condition for the indirect controllability of coupled systems of linear operator groups. *Math. Control Signals Systems*, 29(2):Art. 9, 35, 2017.
<https://link.springer.com/article/10.1007/s00498-017-0193-x>.
- [34] P. Lin, Z. Zhou, and H. Gao. Exact controllability of the parabolic system with bilinear control. *Appl. Math. Lett.*, 19(6):568–575, 2006.
<https://www.sciencedirect.com/science/article/pii/S0893965905002697>.
- [35] J.-L. Lions. *Contrôle optimal de systèmes gouvernés par des équations aux dérivées partielles*. Avant propos de P. Lelong. Dunod, Paris, 1968.
- [36] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 2*. Travaux et Recherches Mathématiques, No. 18. Dunod, Paris, 1968.
- [37] Y. Liu, T. Takahashi, and M. Tucsnak. Single input controllability of a simplified fluid-structure interaction model. *ESAIM Control Optim. Calc. Var.*, 19(1):20–42, 2013.
http://www.numdam.org/item/COCV_2013__19_1_20_0/.
- [38] J. Lohéac, E. Trélat, and E. Zuazua. Minimal controllability time for the heat equation under unilateral state or control constraints. *Math. Models Methods Appl. Sci.*, 27(9):1587–1644, 2017.
<https://www.worldscientific.com/doi/abs/10.1142/S0218202517500270>.
- [39] G. Olive. Boundary approximate controllability of some linear parabolic systems. *Evol. Equ. Control Theory*, 3(1):167–189, 2014.
<https://aimsciences.org/article/doi/10.3934/eect.2014.3.167>.
- [40] M. Ouzahra, A. Tsouli, and A. Boutoulout. Exact controllability of the heat equation with bilinear control. *Math. Methods Appl. Sci.*, 38(18):5074–5084, 2015.
<https://onlinelibrary.wiley.com/doi/full/10.1002/mma.3428>.
- [41] S. Primak, V. Kontorovich, and V. Lyandres. *Stochastic methods and their applications to communications*. John Wiley & Sons, Inc., Hoboken, NJ, 2004. Stochastic differential equations approach.
- [42] H. Risken. *The Fokker-Planck equation*, volume 18 of *Springer Series in Synergetics*. Springer-Verlag, Berlin, second edition, 1989. Methods of solution and applications.
- [43] D. Steeves, B. Gharesifard, and A.-R. Mansouri. Controllability of coupled parabolic systems with multiple underactuators, parts 1 and 2. *Submitted*, 2017.
<https://arxiv.org/abs/1710.01397>.
- [44] A. Tsouli and A. Boutoulout. Controllability of the parabolic system via bilinear control. *J. Dyn. Control Syst.*, 22(1):35–44, 2016.
<https://link.springer.com/article/10.1007/s10883-014-9247-2>.

- [45] Z. Zhou, P. Lin, and H. Gao. Some results on exact controllability of parabolic systems. *Taiwanese J. Math.*, 12(3):635–648, 2008.
https://www.jstor.org/stable/43833940?seq=1#metadata_info_tab_contents.