

Pure

Scotland's Rural College

Non CO2 greenhouse gas sources from managed and natural soils - fluxes and mitigation

Skiba, U; Rees, RM; Siong, J; Sentian, J

Published in:

Journal of Oil Palm and the Environment

DOI:

<http://dx.doi.org/10.5366/jope.2012.11>

First published: 01/01/2012

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Skiba, U., Rees, RM., Siong, J., & Sentian, J. (2012). Non CO2 greenhouse gas sources from managed and natural soils - fluxes and mitigation. *Journal of Oil Palm and the Environment*, 3, 107 - 113.
<https://doi.org/10.5366/jope.2012.11>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Non CO₂ greenhouse gas sources from managed and natural soils - fluxes and mitigation

Ute Skiba*¹, Robert Rees², Jambery Siong³, Justin Sentian³

1. Introduction

This paper outlines the current knowledge of the processes controlling nitrous oxide (N₂O) and methane (CH₄) fluxes, methods of measurement, mitigation options and models designed to simulate N₂O and CH₄ fluxes. Natural and managed soils are globally important sources and sinks of the main non-CO₂ greenhouse gases (GHG) N₂O and CH₄. Compared to CO₂ their global warming potential over a 100 year period is 298 and 25 times, respectively, larger than that of CO₂.

Keywords: nitrous oxide, methane, land use change, fertilization, microbial processes
JOPE 2012,3:107-113

Soils are the single largest source of N₂O, accounting for over 60% of the global total annual N₂O budget. Tropical forest soils are the largest natural soil source of N₂O (2.11 Tg N₂O y⁻¹) because the wet, warm environment provides optimal conditions for N₂O production. Nitrogen-fertilised agricultural soils, especially in warm wet climates are the largest anthropogenic source of N₂O (4.4 Tg N₂O y⁻¹) (Figure 1).

Global CH₄ production is dominated by emissions from wetlands, ruminants and gas and coal mining. Natural wetlands account for 26% of the global production and rice paddy fields for a further 5%. Soils are also a small, but significant sink for CH₄ (30 Tg y⁻¹). Natural forest soils are believed to be the largest sinks, but their sink strength can change from sink to

source, depending on the soil moisture conditions and land management (Figure 1).

2. Microbial processes leading to N₂O and CH₄ production and consumption

In soils and waters the microbial processes, 'nitrification', 'denitrification', 'methanogenesis' and 'CH₄ oxidation', are responsible for the soil atmosphere exchange of N₂O and CH₄ everywhere on our planet. In order to model and mitigate N₂O and CH₄ emissions, one needs to understand how these processes operate and how they are influenced by climate, ecosystem type and human intervention (Skiba and Smith 2000).

Email: ums@ceh.ac.uk

* Corresponding Author

Address: ¹ Centre for Ecology and Hydrology, Penicuik, UK

² SRUC, Edinburgh, UK

³ Universiti Malaysia Sabah, Malaysia

Published: 28 December 2012

Received: 29 August 2012

Accepted: 24 November 2012

© 2012 Ute Skiba

This is an Open Access article which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure 1: Global N₂O and CH₄ sources (IPCC, 2007; Bousquet et al., 2007).

2.1 Nitrification and denitrification

Nitrification is the oxidation of ammonium to nitrate and requires oxygen and a source of ammonium. This process is a common source of N₂O in the top few mm of the soil and dominates N₂O production when soils are relatively dry. Large rates of N₂O production are associated with denitrification, the reduction of nitrate to N₂O or N₂. This process requires nitrate and simple organic carbon

compounds (Bremner, 1997; Davidson, 1991). Complete denitrification to N₂ only occurs in the absence of oxygen, whereas in the presence of small amounts of oxygen, N₂O is the dominant denitrification product. Optimal conditions for denitrification are realised in waterlogged soils or in soils with high oxygen demand. The latter conditions can be achieved by decaying organic matter, root exudates or additions of manure (Wrage et al., 2001).

Nitrification or denitrification can only proceed in the presence of an adequate supply of mineral nitrogen. Most common sources of mineral nitrogen are (i) nitrogen fertilisers, (ii) microbial mineralisation of added organic nitrogen, for example excrete from grazing animals, leaf litter, (iii) mineralisation of stored soil carbon compounds and (iv) atmospheric nitrogen deposition.

The third source of nitrogen is an interesting one. Soils are great sinks for carbon, stored as organic matter. However, when soil is disturbed or drained the organic matter and a lot of the nitrogen locked up in organic soil compounds are released by the process 'mineralisation'. Land use change from forest to arable land is a typical example of soil disturbance and its impact on N₂O production and emission is the least understood.

2.2 Methanogenesis and methane oxidation

Under strictly anaerobic conditions CH₄ is formed during the consumption of simple carbon compounds by very specific groups of microbes. Methanogens are responsible for the large CH₄ emissions observed from rice paddies, peat wetlands, termite mounds and ruminants. In wetlands the key controllers of CH₄ emissions rates are water table height, temperature and plant communities (Laanbroek et al., 2010). A large part of the CH₄ produced in the anaerobic soil layers is oxidised by methanotrophs, which live at the interface of aerobic and anaerobic environments. They use CH₄ as carbon and energy source and require aerobic conditions. Another group of methanotrophs is present in dry soils, where CH₄ production and emission are negligible. This group of methanotrophs used the CH₄ from the atmosphere. They are very sensitive to disturbance, consequently CH₄ oxidation rates are reduced by land use change, drainage, ploughing or nitrogen additions (MacDonald et al., 1997; Zhang et al., 2008, Laanbroek et al., 2010). Undisturbed forests, such as primary tropical forests, with good well-aerated soil structures are the largest sinks for CH₄.

Our understanding of the variables that control N₂O and CH₄ production, emission and uptake is predominately based on measurements of N₂O emissions from nitrogen fertilised soils in temperate, managed ecosystems and of CH₄ emissions from rice paddies and high latitude wetlands. These numerous studies have shown very large spatial and temporal variability, especially for N₂O and CH₄ emissions, hence detailed information of physical and chemical characteristics of the

soil, the climate and soil management practices is required to upscale to annual and regional fluxes and identify mitigation options.

3. Methods to measure soil N₂O and CH₄ fluxes

Nitrous oxide and CH₄ fluxes can be measured at scales ranging from a few grams of soil to several km. Our global understanding of N₂O and CH₄ fluxes and their control by physical, chemical and microbial processes has largely arisen from static flux chamber measurements (Clayton et al., 1994). Recent development of high frequency instruments, that detect very small concentration changes, has improved our knowledge of N₂O and CH₄ biosphere atmosphere exchange at the field/landscape scale and at a high temporal resolution (i.e. Fowler et al., 2011). Advancements in satellite technology have made it possible to measure CH₄ and N₂O concentrations from space (i.e. Schneising et al., 2011).

4. Modelling N₂O and CH₄ fluxes

Signatories of the Kyoto Protocol are obliged to submit annual accounts of their anthropogenic greenhouse gas emissions, which include N₂O and CH₄, to the UNFCCC. The reporting follows internationally agreed protocols (IPCC, 2006), mostly using simple equations (emission factors), represented by a Tier 1 methodology. The Tier 1 methodology is very simplistic and is designed to calculate emissions based on easily available data. For example N₂O emissions from fertilised grasslands and arable land are assumed to be 1% of the mineral fertiliser N input and CH₄ emissions from rice without organic amendments 20 g m⁻² (IPCC, 2006).

The Tier 1 emission factor approach assumes a linear relationship between N input and N₂O emission. The emission factor for N additions from (i) mineral fertilisers, (ii) organic amendments and crop residues or (iii) N mineralised from mineral soil as a result of loss of soil carbon, is 1% of N applied, with an uncertainty range of 0.3 to 3%. The reason for the large uncertainty range is that not only N input, but also soil conditions, agronomy and climate are important drivers of N₂O emission. Soil and fertiliser type in particular have been shown to influence N₂O fluxes; but the magnitude of this effect is very varied and cannot be quantified by simple regression equations.

Some countries are currently developing emission factors specific to their growing conditions, management or crop type (Tier 2 methodology). The most complex, but most targeted approach uses process based dynamic models (Tier 3 methodology). Such models use mathematical equations to define the rates of the individual source processes, allowing the model to take a detailed account of site conditions. Climate is a particularly important driver and the models often run on daily or sub daily time steps. In particular for N₂O the denitrification and decomposition model 'DNDC' (Li et al., 1992) has been developed to simulate N₂O emissions. This model also appears to work well for CH₄ emissions from rice paddies (Babu et al., 2009).

5. Effects of land use change from forests on N₂O and CH₄ fluxes

Globally deforestation is continuing at a rate of 12.9 million ha⁻¹ y⁻¹, primarily as a result of converting forests to agricultural land in South America, Africa and Asia (FAO, 2006). Land use change from forestry invariably requires logging, sometimes tree stump removal, drainage, and ploughing. In the short term these destructive activities will increase microbial decomposition and mineralisation and lead to increased emissions of CO₂, N₂O and CH₄ (Yashiro et al., 2008). It has been estimated that humid tropical primary forests grown on mineral soil lose about 30% of their carbon stock when converted to oil palm plantations or agricultural land (Murty et al., 2002). On peat such conversion results in much larger carbon losses, due to its high organic matter content. Land use change from forestry invariably increases N₂O emissions and reduces CH₄ oxidations rates; and in some cases turns CH₄ oxidation to emissions. The rate of change and length of increased flux rates depends on soil type, drainage, rainfall and land management. However relatively few studies have been undertaken, and there are high levels of uncertainty associated with estimates of greenhouse gas fluxes during land use change.

Interests in growing biofuel as a 'low carbon energy' source have increased conversion of forests and permanent grasslands to growing annual and perennial bioenergy crops across the globe. Calculating the carbon footprint for biofuel production must include soil emissions of N₂O and CH₄ (Crutzen et al. 2008), as these emissions can offset or reduce carbon

savings from replacing fossil fuel with renewable bioenergy based fuel (Drewer et al. 2011). Due to lack of sufficient N₂O and CH₄ flux data, carbon footprint analyses of land use change to bioenergy crops are highly uncertain.

6. A case study: Land use change from primary forest to oil palm

The Centre for Ecology and Hydrology and the Universiti Malaysia Sabah (UMS), Malaysia participated in the project 'Oxidant and Particle Photochemical Processes above a South-East Asian tropical rain forest' (OP3) funded by the UK Natural Environment Research Council, UK and measured soil N₂O and CH₄ fluxes from primary, secondary forests at Danum Valley and from an oil palm plantation near Lahad Datu, Sabah, Malaysia. The conventional static chamber method was used to measure fluxes (Figure 2). We observed typically large spatial and temporal variability for both gases. Land use change did not significantly influence CH₄ fluxes (Figure 3). Small rates of CH₄ uptake and release were observed at all sites, but they were smaller than those reported for a Chinese tropical primary forest and rubber plantation (Werner et al., 2006). Nitrous oxide fluxes increased with disturbance, and were enormous (6566 µg N₂O-N m⁻² y⁻¹) near the oil palm tree trunks, where fertiliser was applied into 4 holes around the stem (Figure 3). However, taking into account the much lower emissions from the unfertilised areas of the plantation, suggests an annual emission rate for the entire plantation of 4.4 ± 3.5 kg N₂O-N ha⁻¹y⁻¹.



Figure 2: A static chamber measuring N₂O and CH₄ fluxes on an oil palm plantation near Lahad Datu, Borneo. The chamber is sealed for 60 minutes. Air samples were withdrawn at intervals and sent to CEH, Edinburgh for analysis by gas chromatography.




Figure 3: Average N₂O and CH₄ fluxes measured from Forests (Danum Valley) and an oil palm plantation (Lahad Datu) Sabah, Borneo between April and July 2008. Oil palm (f) is the 2 m area where fertiliser was spread; the flux was 6566 µg N₂O-N m⁻² y⁻¹. Oil palm (nf) is the area on the plantation that was not fertilised.

This emission rate is of the same order of magnitude as emissions reported for an oil palm plantation grown on peat in Sarawak, Malaysia (1.2 kg N₂O-N ha⁻¹y⁻¹, Melling et al., 2007). These two examples both show enhanced N₂O emissions when converting forests to oil palm plantations. Their annual flux calculations are based on many assumptions and more research is required to improve the carbon footprint of land use change from forest to oil palm.

7. Mitigating N₂O and CH₄ emissions

Reducing N₂O emissions from oil palm plantations is critically important in reducing the carbon footprint of the end product, but because mitigation strategies often require increased nutrient use efficiency, there are both environmental and economic benefits (Rees et al. 2012). Nitrous oxide emissions from managed land can be reduced by

- i. Optimising application rates of N, P, K and micro-nutrients for crop type, soil nutrient status, soil type, climate and management. Such management should take full account of any nutrients that are contained in manures or other organic substrates applied to the land. Over- and under-fertilisation, or imbalance in N, P, K micro-nutrients can increase soil N₂O emissions when expressed on a unit product basis (Mosier et al., 1998).

- ii. Avoid N fertilisation in high rainfall periods, as this can increase N₂O emissions (Skiba et al., 2012).
- iii. Use slow release fertilisers or apply nitrification inhibitors, both reduce N₂O emissions (de Klein and Ledgard, 2005).
- iv. Maintain good soil conditions, this includes good soil structure, avoiding compaction and the installation of drainage, where soils retain excessive water (Ball et al., 1999).
- v. Use alternatives to synthetic N fertilisers, such as legumes which can be undersown with oil palm, and provide an input of biologically fixed N (Mosier et al., 1998)
- vi. Optimise crop management and aim for maximum yield. Thereby less land needs to be cultivated and converted from forests or natural grasslands, avoiding greenhouse gas fluxes as a result of land use change.
- vii. Convert grasslands, rather than forest to cropland or oil palm. Land use change from grassland to cropland has a much lower carbon footprint (Germer & Sauerborn, 2007).

Methane emission and oxidation rates tend to be small from managed soils. Land use change often reduces the soil CH₄ sink strength, which however can be improved by

maintaining good soil structure, avoid carbon losses and compaction and minimise tillage.

References

1. Bousquet,P., Ciais,P., Miller,J.B., Dlugokencky,E.J., Hauglustaine,D.A. et al. 2006. Contribution of anthropogenic and natural sources to atmospheric methane variability. *Nature*, 443, 439–443.
2. Bremner,J.M. 1997. Sources of nitrous oxide in soils. *Nutrient Cycling in Agroecosystems*, 49, 7-16.
3. Clayton,H., Arah,J.R.M. and Smith,K.A. 1994. Measurements of nitrous oxide emissions from fertiliser grassland using closed chambers. *Journal of Geophysical Research* 99, 16599-16607.
4. Crutzen,P.J., Mosier,A.R., Smith,K.A., Winiwarter,W. 2008. N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels. *Atmospheric Chemistry and Physics*, 8, 389-395.
5. Davidson,E.A., 1991. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In: Rogers JE and Whitman WB (eds.) *Microbial Production and Consumption of Greenhouse Gases: Methane, Nitrogen Oxides and Halomethanes*, pp. 219–234.
6. de Klein,C.A.M. and Ledgard,S.F. 2005. Nitrous oxide emissions from New Zealand agriculture - key sources and mitigation strategies. *Nutrient Cycling in Agroecosystems* 72, 77-85.
7. Drewer,J., Finch,J., Lloyd,C.R., Baggs,E.M., Skiba,U., 2012. How do soil emissions of N₂O, CH₄ and CO₂ from perennial bioenergy crops differ from arable annual crops? *Global Change Biology, Bioenergy*. 10.1111/j.1757-1707.2011.01136.
8. FAO, 2006: *Global Forest Resources Assessment, 2005. Progress towards sustainable forest management*. FAO Forestry Paper 147, 320 pp.
9. Fowler,D., Nemitz,E., Misztal,P., diMarco,C., Skiba,U.,Nicholas Hewitt, C., 2011. Effects of land use on surface-atmosphere exchanges of trace gases and energy in Borneo: Comparing fluxes over oil palm plantations and a rainforest. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366, 3196-3209.
10. Germer,J. and Sauerborn,J., 2007. Estimation of the impact of oil palm plantation establishment on greenhouse gas balance. *Environmental Development and Sustainability* DOI 10.1007/s10668-006-9080-1.
11. IPCC 2006. *IPCC Guidelines for National Greenhouse Gas Inventories*; Prepared by the National Greenhouse Gas Inventories Programme. Japan: IPCC; 2006.
12. IPCC 2007. *Climate change 2007: the physical science basis*.Contribution of the Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK: Cambridge University Press.
13. Laanbroek, H.J., 2010. Methane emission from natural wetlands: interplay between emergent macrophytes and soil microbial processes. A mini-review. *Annals of Botany* 105, 141-153.
14. Jagadeesh Babu,, Y., Li,C., Froelking,S., Nayak,D.R., Adhya,T.K., 2012. Field Validation of DNDC Model for Methane and Nitrous Oxide Emissions from Rice-based Production Systems of India. *Nutrient Cycling in Agroecosystems* 74,157-174.
15. Li,C.S., Froelking,S. and Froelkin,T.A. (1992) A model of nitrous-oxide evolution from soil driven by rainfall events.1. model structure and sensitivity. *Journal of Geophysical Research-Atmospheres* 97, 9759-9776.MacDonald,J.A., Eggleton,P., Bignell,D.E., Forzi,F. and Fowler,D., 1998. Methane emission by termites and oxidation by soils, across a forest disturbance gradient in the Mbalmayo Forest reserve, Cameroon, *Global Change Biology* 4, 409 - 418.
16. Melling,L., Hatano,R., Goh,K.J. 2007. Nitrous oxide emissions from three ecosystems in tropical peatland of Sarawak, Malaysia. *Soil Science and Plant Nutrition*, 53, 792-805.
17. Mosier, A.R., Duxbury, J.M., Freney, J.R., Heinemeyer,O. and Minami,K., 1998. Assessing and mitigating N₂O emissions from agricultural soils. *Climate Change* 40, 7-38.
18. Murty,D.,Kirschbaum,M.U.F., McMurtrie,R.E., & McGilvray,A., 2002. Does conversion of forest to agricultural land change soil carbon and nitrogen? A review of the literature. *Global Change Biology*, 8, 105-123.
19. Rees,R.M., Baddeley,J.A., Bhogal,A., Ball,B.C., Chadwick,D.R., MacLeod,M., Lilly,A., Pappa,V.A., Thorman,R., Watson,C.A. and Williams,J.R., 2012. Nitrous oxide mitigation in UK agriculture. *Soil Science and Plant Nutrition*, In Press.

20. Schneising, O., Buchwitz, M., Reuter, M., Heymann, J., Bovensmann, H., Burrows, J.P. 2011. Long-term analysis of carbon dioxide and methane column-averaged mole fractions retrieved from SCIAMACHY. *Atmospheric Chemistry & Physics*, 11, 2863–2880.
21. Siong, J. 2012. Greenhouse gas emission in SEA: soil methane (CH₄) and nitrous oxide (N₂O) fluxes in selected forest stand and oil palm plantation in Sabah. Masters Thesis, University of Sabah, Malaysia.
22. Skiba, U. and Smith, K.A., 2000. The control of nitrous oxide emissions from agricultural and natural soils. *Chemosphere* 2: 379–386.
23. Skiba, U., Jones, S., Dragosits, U., Drewer, J., Fowler, D., Rees, R., Pappa, V., Cardenas, L., Chadwick, D., Yamulki, S. 2012. UK emissions of the greenhouse gas nitrous oxide. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367:1175-1185.
24. Werner, C., [Zheng, X.](#), [Tang, J.](#), [Xie, B.](#), [Liu, C.](#), [Kiese, R.](#), [Butterbach-Bahl, K.](#) 2006. N₂O, CH₄ and CO₂ emissions from seasonal tropical rainforests and a rubber plantation in Southwest China. *Plant and Soil* 289, 335-353.
25. Wrage, N., Velthof, G.L., van Beusichem, M.L. and Oenema, O. 2001. Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biology and Biochemistry* 33: 1723–1732.
26. Yashiro, Y., Kadir, W.R., Okuda, T., Koisumi, H., 2008. The effects of logging on soil greenhouse gas (CO₂, CH₄, N₂O) flux in a tropical rain forest, Peninsular Malaysia. *Agricultural and Forest Meteorology*, 148, 799-806.
27. Zhang, W., Mo, J., Zhou, G., Gundersen, P., Fang, Y., Lu, X., Zhang, T., and Dong, S. 2008. Methane uptake responses to nitrogen deposition in three tropical forests in southern China. *Journal of Geophysical Research*, 113, D11116, doi:10.1029/2007JD009195.