# Pure

## Scotland's Rural College

## Management and personality in Labrador Retriever dogs

Lofgren, SE; Wiener, P; Blott, SC; Sanchez-Molano, E; Woolliams, JA; Clements, DN; Haskell, MJ

Published in: **Applied Animal Behaviour Science** 

DOI: 10.1016/j.applanim.2014.04.006

Print publication: 01/01/2014

**Document Version** Peer reviewed version

Link to publication

*Citation for pulished version (APA):* Lofgren, SE., Wiener, P., Blott, SC., Sanchez-Molano, E., Woolliams, JA., Clements, DN., & Haskell, MJ. (2014). Management and personality in Labrador Retriever dogs. *Applied Animal Behaviour Science*, *156*, 44 - 53. https://doi.org/10.1016/j.applanim.2014.04.006

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
   You may freely distribute the URL identifying the publication in the public portal ?

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

| 1  | Management and Personality in Labrador Retriever dogs  |
|----|--|
| 2  |  |
| 3  | Sarah E. Lofgren <sup>1</sup> , Pamela Wiener <sup>2</sup> , Sarah C. Blott <sup>3</sup> , Enrique Sanchez-Molano <sup>2</sup> , John A. |
| 4  | Woolliams <sup>2</sup> , Dylan N. Clements <sup>1,2</sup> , Marie J. Haskell <sup>4</sup>  |
| 5  |  |
| 6  | 1. Royal (Dick) School of Veterinary Science, University of Edinburgh, Easter Bush,  |
| 7  | Scotland, UK   |
| 8  | 2. Roslin Institute, Royal (Dick) School of Veterinary Science, University of Edinburgh,   |
| 9  | Easter Bush, Scotland, UK  |
| 10 | 3. Animal Health Trust, Newmarket, UK  |
| 11 | 4. Scotland's Rural College, West Mains Road, Edinburgh, Scotland, UK  |

12

13

#### 14 Abstract

15 Canine personality is of keen interest to dog owners and researchers alike. The 16 regular human contact with them makes dogs an ideal species to use in the investigation of 17 animal personality. This study specifically focused on Labrador Retrievers, consistently one 18 of the most popular breeds both in the UK and around the world. Using surveys completed 19 by dog owners, data was gathered on the behaviour of the dogs, in addition to the physical 20 characteristics and management characteristics of the dogs (n=1978). Twelve personality 21 traits were identified and investigated for associations with the demographic data. It was 22 found that the working status of the dog was more commonly associated with differences in 23 personality than other analyzed factors. Gundogs had higher scores for 'fetching tendency' 24 and 'trainability' than Showdogs or Pets (P<0.05). Chocolate dogs were more 'agitated 25 when ignored' and showed more 'excitability' than black dogs, and lower 'trainability' and 26 'noise fear' than both yellow and black dogs (all P<0.05). Dogs exercised for longer 27 periods showed less aggression, less fear of humans and objects and lower separation 28 anxiety than dogs that were not as active. The effects observed in this study may be due to 29 the experience and training of the dogs, the work-related genetic strain of Labrador 30 Retriever or most likely, a combination of both influences.

31

32 Keywords: Labrador Retriever, Management, Personality, Demographics, C-BARQ,
33 canine

#### 35 **1. Introduction**

It is commonly observed that individual animals show consistency in the way they respond to situations, and that the intensity of the response varies between individuals. In farm animals this phenomenon is termed temperament (Burrow and Corbet, 2000, Hoppe *et al.*, 2010). However, in dogs it is often called personality (Svartberg *et al.* 2005, Ley *et al.*, 2008), and this is the convention that we will follow for this paper.

41 An animal's personality arises from the influences of both genetics and its 42 environment, including previous experience. Prenatal experience has been shown to have 43 long term effects on personality and other traits. Zebra finch eggs injected with testosterone 44 produced birds that habituated quicker to novel food (Tobler and Sandell, 2007). Sows born 45 to mothers that experienced social stress during pregnancy show more restlessness and 46 aggression toward their own piglets (Jarvis, et al., 2006). There are also many postnatal 47 influences that determine an animal's personality. Critical periods in early life are known to 48 affect the long-term behaviour of the dog (Scott and Marston, 1950). The time at which a 49 puppy is introduced to humans is critical, with earlier introduction resulting in more 50 positive reactions towards humans in adulthood (Freedman *el al.*, 1961). Svartberg *et al.* 51 (2005) also found that dogs' reactions to some tests changed following later repetition, such 52 as tests intended to provoke aggression using unusual stimuli. Although the individual dogs' 53 reactions changed, the relative ranking of the dogs remained the same. Since personality is 54 unique to each individual animal, it can be influenced by other factors and experiences in 55 the animals' life history. Kutsumi et al., (2013) found that puppy training classes improved 56 long term obedience as well as response to strangers. McMillan et al., (2013) found that 57 puppies obtained from pet stores scored less favorably on a personality assessment than

58 puppies from non-commercial breeders, such that pet store dogs showed higher aggression 59 and separation-related problems than dogs purchased from breeders. Later retesting 60 produced similar results, showing that early experience has a long-term effect on the 61 personality of the dogs.

62 The genetic influence on animal personality has often been studied in terms of breed 63 differences. Differences in temperament were found between breeds of cattle which were 64 raised in identical environments (Hoppe *et al.*, 2010). These differences are presumably due 65 to genetic differences, since other variation had been removed. Dog breeds are well known 66 to show differences in personality (Hart and Hart, 1995). Dachshunds and Chihuahuas have 67 shown higher aggression toward humans, while Akitas and Pit Bull Terriers show higher 68 dog-directed aggression (Duffy et al., 2008). Personality traits, including aggression, have 69 also been shown to be heritable in dogs in a number of studies (Liinamo *et al.*, 2007; 70 Mackenzie, et al., 1986; Goddard and Beilharz, 1983; Saetre et al., 2006), which could 71 have implications for breeding programmes. This is especially true for working dogs, since 72 an appropriate personality is important to fulfilling their duties. Additionally, Svartberg et 73 al., (2006) found that recent selection pressures have affected personality, with personality 74 being more highly correlated with the current role of the dogs than with the breed's original 75 purpose. For instance, breeds that are currently popular as house pets show higher 76 playfulness regardless of the breeds' original purpose. The same experience is likely to 77 affect genetically different individuals in different ways (Stamps and Groothius, 2010). 78 As personality traits have been shown to be influenced by both genetic and non-79 genetic ('environmental') factors, it is of interest to determine the relative importance of 80 these different factors. In this case, 'environment' is defined as the management and

81 housing conditions experienced by domestic dogs. 'Physical' traits, such as age, sex and 82 bodyweight, are also likely to influence personality. Therefore the aim of this study was to 83 determine how personality traits are affected by physical and management factors in dogs. 84 In order to account for the complexity of the study a large sample size was needed. In order 85 to accomplish this, surveys were sent to several thousand dog owners. The Canine 86 Behaviour and Research Questionnaire (C-BARQ), developed at the University of 87 Pennsylvania, was used for this study (http://vetapps.vet.upenn.edu/cbarq/). Originally 88 developed as a method for evaluating and predicting the success of guide dogs (Serpell and 89 Hsu, 2001), this survey can be filled out by any dog owner. It covers many behavioural 90 responses which are categorized into different aspects of animal personality. The survey 91 responses are recorded on a 1 to 5 scale of the intensity of behavioural response to various 92 situations. This is very similar to the approach of Svartberg and Forkman (2002), except the 93 ratings are made by owners instead of a separate observer, and the behaviours recorded are 94 elicited by normal interactions instead of induced by the test setup. The C-BARQ has been 95 translated and used successfully in Japan (Nagasawa, et al., 2011), Taiwan (Hsu and Sun, 96 2010), and the Netherlands (van den Berg *et al.*, 2006), further demonstrating its generality. 97 It has been used in the past to identify problematic behaviours being exhibited by individual 98 dogs (Hsu and Serpell, 2003). It has also been used to study variation in specific traits 99 among dogs. Using the C-BARQ, Duffy et al., (2008) found that levels of aggression 100 towards people versus aggression towards dogs varies within and between breeds. 101 For this study the issue of between-breed variation was eliminated by only studying 102 a single breed, Labrador Retrievers registered with the UK Kennel Club. The overall aim of

103 the study was to test for associations between the animal's physical characteristics,

104 lifestyle, potential genetic differences, and personality.

105

#### 106 **2. Materials and Methods**

#### 107 2.1. Surveys

A survey was created to gather demographic and management data on the dogs participating in a larger study investigating the factors associated with canine hip dysplasia. It included 38 questions on physical traits such as weight, coat colour and health, as well as management data related to activities, housing, management and feeding (further details given below).

113 The C-BARQ questionnaire consists of 102 questions pertaining to dog behaviour,

114 divided into seven sections. The sections pertain to Training and obedience (8 questions),

115 Aggression (25), Fear and anxiety (19), Separation-related behaviour (8), Excitability (6),

116 Attachment and attention seeking (6), and Miscellaneous (Barking, chasing, unusual

117 behaviours, etc.) (28).

The demographic survey was sent by the UK Kennel Club to the owners of 12,408 registered Labrador Retrievers which had known hip scores. Of these, 3071 surveys were completed and returned. The Canine Behavioural Assessment and Research Questionnaire (C-BARQ) surveys were distributed to the 2974 of those who had completed the first survey and also agreed to take part in the personality assessment. C-BARQ surveys were received for 2020 dogs.

124

125 **2.2. Personality trait analysis** 

| 126 | C-BARQ responses were recorded as letters A-E, with A representing a low or                      |
|-----|--|
| 127 | infrequent display of the behaviour in question, and E representing a high or frequent           |
| 128 | response. The C-BARQ data was transformed to numerical values, with A=1, B=2, C=3,               |
| 129 | D=4, E=5, and non-responses (N/A or Unanswered). Histograms were plotted for each                |
| 130 | question which were used to examine response variation for each question. Values for             |
| 131 | questions 6-8 (regarding disobedience) were reversed so 'desirable' was represented by a         |
| 132 | high score and 'undesirable' by a low score to be consistent with the other questions in that    |
| 133 | section. A description of the seven categories of behaviour are shown in Table 1.                |
| 134 | In order to define distinct personality traits, we investigated whether some of the              |
| 135 | questions referred to the same or closely related behavioural characteristics. A Principle       |
| 136 | Components Analysis (PCA) for correlation was run in Minitab 16 (Minitab Inc.) to                |
| 137 | determine whether the answers to individual questions related to each other. Questions with      |
| 138 | 100 or more missing values were removed from analysis. This included questions 23, 32-           |
| 139 | 36, 42, 50, 66, 71, 79, 87, and 103. These questions largely concerned multi-dog                 |
| 140 | households or other situations that many owners and dogs had never been exposed to.              |
| 141 | Surveys missing any responses from the remaining questions were removed due to the               |
| 142 | constraints of the PCA. This resulted in a final sample size of 1077 surveys covering 89         |
| 143 | questions. The first three components of the PCA accounted for 13%, 6.6%, and 3.9% of            |
| 144 | the variation, respectively. On the basis of the clustering of the question level traits (Figure |
| 145 | 1), they were combined into 'personality traits' by taking the mean of the responses to          |
| 146 | questions within each PCA-defined group of traits, with the added constraint that the            |
| 147 | questions were from the same category in the C-BARQ questionnaire (Hsu & Serpell,                |
| 148 | 2003; http://vetapps.vet.upenn.edu/cbarq/). For example, the trait of Owner aggression was 7     |

149 calculated by averaging the values of all the questions that pertained to aggressive

150 behaviour directed at the owner of the dog (Questions 10,14,15,18,20,26,31).

- 151 Urinating/defecating was removed due to very low variation (all dogs had low scores). The
- 152 correlations between the 18 remaining personality traits (Supplementary Table 1) were then
- 153 calculated with the aim of further combining highly correlated traits (>0.4), again with the
- 154 constraint that the questions were from the same category in the questionnaire
- 155 (Supplementary Table 2). All questions included in the highly correlated traits were
- averaged to create the new trait in the same process as described above. If a survey had
- 157 missing values for more than half the questions used to make up a personality trait, that
- 158 individual did not receive a value for that trait.
- 159 This analysis resulted in a final group of 12 personality traits: Agitated when
- 160 Ignored, Attention Seeking, Barking Tendency, Excitability, Fetching, Human and Object

161 Fear, Noise Fear, Non-Owner Aggression, Owner Aggression, Separation Anxiety,

162 Trainability, and Unusual Behaviour (Supplementary Table 3).

163

#### 164 2.3. Management/Physical Traits

Following a quality control procedure to remove questionnaires missing key data, complete C-BARQ surveys and management and physical characteristics data were available for 1978 dogs. Age at the time of survey completion was calculated from the date of birth and the survey received date. Age was then rounded to the nearest 0.5 years. Dogs were aged between 2 and 9.5 years. Measurements made in Imperial units were converted to SI units. Body mass index (BMI) was calculated as Girth divided by Length squared. Dogs were separated into categories based on their Working Status. These categories were 172 Pets, Gundogs, and Showdogs. Dogs that were reported as 'Other' were either reassigned 173 based on information provided in the comments or removed, resulting in a final sample size 174 of 1,978. Dogs that were reported as being both Pets and Gundogs were classified as 175 Gundogs, while dogs classified as Showdogs and Pets were grouped with Showdogs. Dogs 176 were classified as living Indoor, Outdoor, or Indoor/Outdoor based on where they were 177 reported to spend most of their time throughout the year. For instance, if a dog spent most 178 of its time in a run, outdoor kennel, or yard, it was classified as Outdoor. If it spent most of 179 its time in a house or garage, it was classified as Indoor. Dogs that were classified as Indoor 180 for one half of the year and Outdoor for the other comprise the Indoor/Outdoor category. 181 Gender Status was used to combine the Gender and Neuter responses. This resulted in four 182 possible categories: Entire Males (EM) for uncastrated dogs, Entire Females (EF) for 183 unneutered bitches, Neutered Males (NM) for castrated dogs, and Neutered Females (NF) 184 for neutered bitches. Coat Colour was limited to the three main colours of Black, 185 Chocolate, and Yellow. The small number of dogs that reported a coat colour of Fox Red 186 (5), Liver (17), or Black and Tan (2) were grouped with the three categories Yellow, 187 Chocolate, and Black, respectively. Health Status was determined by the presence of a 188 disease or veterinary condition, with dogs either identifying as healthy (0) or having a 189 significant health problem (1). No single health problem occurred with high enough 190 frequency to be examined independently. Exercise was categorized into 1 (up to one hour 191 per day), 2(1-2 hours), 3(2-4 hours), or 4 (more than 4 hours). 192 Sire ID was used to identify full or half siblings and was used to account for any 193 variation due to family relationships. Of the study dogs, 693 had sires which had no other 194 progeny in our study. The remaining 1285 dogs had sires that had between 2 and 37 9

progeny in our sample (mean family size=1.91, median=1). In summary, eight factors were
extracted from the demographic survey for use in subsequent analysis: Age, BMI, Coat
Colour, Gender Status, Health Status, Indoor/Outdoor Housing, Exercise, and Working
Status.

199

#### 200 **2.4. Statistical model-building**

201 Models were analyzed in Genstat 15 (VSN International, 2000-2013) using the 202 General Linear Mixed Models option with Sire Identity as the random term. The binomial 203 variable of Health Status was analyzed using a Binomial model with a binomial total of 2 204 and a Logit link function. Variables that had a normal distribution were analyzed using a 205 Normal model with an Identity link function. Variables where the distribution of responses 206 was right-skewed were analyzed using a Poisson model with a Logarithm link function. 207 The single variable where the distribution was left-skewed (Fetching) was a single-question 208 personality trait and was therefore analyzed using a Binomial model with a binomial total

209 of 5 and a Logit link function.

210

For constructing the models we followed a set of rules designed to determine the explanatory variables that influenced each of the twelve response variables. The eight explanatory variables were all included in the model together. The variables with the highest p-values were then removed singly until all variables in the model had p-values  $\leq$ 0.200

216

## 217 **2.4.1. Interactions**

218 All two-way interactions between the demographic factors were checked by 219 including only the two independent variables and the interaction between them in the 220 model. Only interactions which returned a p-value of 0.05 or lower during this analysis 221 were included in the next step. These interactions were added simultaneously to the 222 previously established model of demographic factors. Those with the highest p-values were 223 removed until all interactions in the model had a p-value  $\leq 0.05$ , leading to the final model. 224 Effect size was determined by taking the largest difference between means for a 225 single factor in the final model, e.g. the difference between Chocolate and Black dogs for 226 Agitated when Ignored. Average effect size is the mean of all significant effects within a 227 factor.

228

229 3. Results

230 A different full model was used for each response variable. Variables included in 231 the models and their significance are shown in Table 2; interactions present in the model 232 are shown in Table 3, and effect sizes are shown in Table 4.

233

#### 234 **3.1. Working Status**

235 The factor significantly associated with the most response variables was the 236 Working Status of the dogs (Figure 2). Working Status featured in all models except 237 Separation Anxiety, and was significantly associated with all response variables except 238 Separation Anxiety and Owner Aggression. There was an average effect size of 0.33 over 239 all significant associations, the highest of all factors. Pets and Gundogs were more Agitated 240 when Ignored than Showdogs. Gundogs exhibited more Attention Seeking than Showdogs

and Pets. Pets showed greater Barking Tendency and Excitability than Gundogs. Gundogs
showed higher Fetching Tendency than Showdogs or Pets, and this factor had the largest
effect size (1.32). Pets and Gundogs exhibited more Human and Object Fear than
Showdogs. Pets showed greater Noise Fear than Gundogs or Showdogs. Pets showed more
Non-Owner Aggression than Showdogs. Gundogs exhibited greater Trainability than Pets,
and both were greater than Showdogs. Finally, Showdogs and Pets were more likely to
exhibit Unusual Behaviours than Gundogs.

## **3.2. Coat Colour**

Coat Colour was also shown to be associated with several response variables, and
had an average effect size of 0.19. Chocolate dogs were more Agitated when Ignored than
Black dogs. Chocolate dogs showed more Excitability than Black dogs. Black dogs showed
a higher Fetching Tendency than Chocolate dogs. Black and Yellow dogs showed higher
Noise Fear than Chocolate. Yellow dogs showed more Separation Anxiety than Black dogs.
Chocolate dogs exhibited lower Trainability and a higher incidence of Unusual Behaviour
than Black or Yellow dogs.

257

## **3.3. Exercise**

259 The amount dogs were exercised was significantly associated with several

260 personality traits and had the third highest average effect size (0.17). Dogs exercised <1

hour/day were more likely to become Agitated when Ignored than dogs exercised 1-4

262 hours/day. Dogs exercised <1 hour/day had a greater Barking Tendency and greater Human

and Object Fear than those exercised 4+ hours/day. Dogs exercised <1 hour/day were more

264 Excitable than others. Dogs exercised 1-2 hours/day were more likely to show Non-Owner 265 Aggression than dogs exercised 2+ hours/day. Dogs exercised 1-2 hours/day were more 266 likely to show Owner Aggression than dogs exercised 2-4 hours/day. Dogs exercised <1 or 267 2-4 hours/day showed more Separation Anxiety than those exercised 4+ hours/day. Dogs 268 exercised 1+ hour/day had higher Trainability than dogs exercised <1 hour/day. Dogs 269 exercised <1 hour/day were more likely to exhibit Unusual Behaviour than others, and dogs 270 exercised <2 hours/day were more likely to exhibit these behaviours than dogs exercised 4+ 271 hours/day.

272

#### **3.4. Housing**

Housing had an average effect size of 0.13. Dogs kept Indoor/Outdoor were more
likely than Outdoor dogs to become Agitated when Ignored, although Indoor dogs were not
significantly different from either group. Outdoor dogs showed less Excitability and
Human and Object Fear than others, and were less likely to show Noise Fear than Indoor
dogs. Dogs kept Indoor/Outdoor were more likely to show Non-Owner Aggression than
other dogs.

280

#### 281 **3.5. Gender Status**

Gender Status also had an average effect size of 0.13, and played a significant role in nine traits (Attention Seeking, Excitability, Human and Object Fear, Noise Fear, Non-Owner Aggression, Owner Aggression, Separation Anxiety, Trainability, and Unusual Behaviour) (Figure 3). Entire dogs showed more Attention Seeking and Excitability, and lower Human and Object Fear and Noise Fear, than Neutered Females. All Females showed 13 287 higher Non-Owner Aggression than Entire Males, and Entire Males showed higher Owner

288 Aggression than females. Entire Males showed higher Separation Anxiety than all other

289 categories. Entire Females had higher Trainaibility than Neutered Females. All Females

290 showed a higher incidence of Unusual Behaviour than Entire Males.

291

#### 292 3.6. Health, Age, BMI

293 Health Status had an average effect size of 0.069, the lowest of the categorical 294 variables. Healthy dogs were more likely to exhibit Attention Seeking. Younger dogs were 295 more likely to show Human and Object Fear. Dogs with a lower BMI were more likely to 296 show Non-Owner Aggression.

297

#### 298 4. Discussion

299 Given that the survey data by owners is subjective in nature, there is potential for 300 inaccuracy. The large sample size of this study may counter potential imprecision in 301 judgment. Furthermore, multiple questions targeting similar personality traits were 302 grouped, further reducing the role of individual inaccuracies. The survey data does not 303 allow us to directly investigate the causal relationship between variables. Therefore, these 304 results must be discussed in terms of associations and causal relationships can only be 305 hypothesized in most cases.

306

#### 307 **4.1.** Personality trait groupings

308 The traits were grouped according to the PCA results and correlations. The most 309 distinct trait group was that for Trainability which appeared separately from the others in 14

310 the PCA (Figure 1). The questions from the Aggression section were consolidated into two 311 traits: Non-owner and Owner Aggression. Stranger, Dog, and Animal Aggression were 312 highly correlated with each other, but not with Owner Aggression. This indicates that there 313 are some fundamental differences between the expression of aggression towards human 314 owners and other people and animals in the Labrador Retriever. Previous studies with the 315 C-BARQ in a range of breeds have separated aggression into three categories, towards 316 owners, strangers, and dogs (Hsu and Serpell, 2003; Nagasawa et al., 2011; Serpell and 317 Hsu, 2005). In a study comparing aggression in a large number of breeds, Labrador 318 Retrievers were shown to exhibit below-average levels of aggression towards owners, dogs 319 and strangers (Duffy et al., 2008).

320

#### 321 **4.2. Effects of genetics and lifestyle**

Genetic and experiential differences are known causes of personality variation in dogs (Podberscek and Serpell, 1996). The variation in genetics and lifestyle was primarily examined through Working Status, Coat Colour, and Exercise. Other sources of variation were accounted for in the models, including Indoor/Outdoor Housing, Health Status and BMI, but are not discussed in detail because of their limited impact in the statistical analysis.

#### 328 4.2.1. Working Status

Working Status was significantly associated with 10 out of 12 personality traits and had the largest average effect sizes, making it the most influential factor. There are a number of possible explanations for these effects: genetic differences between the Working Status categories, difference in management and/or training between the categories or a

combination of the two influences. Additionally, differences in the effects of Working
Status may be exaggerated by the movement of dogs from one category to another if their
behaviour is inappropriate (i.e. a dog that does not perform well as a show dog becomes a
pet). We will refer to this phenomenon as 'category shift'.

337 In terms of the genetic influence, our results may reflect a known division in this 338 breed. Labrador Retriever breeders and dog researchers recognize two types of Labrador 339 Retrievers, which are referred to as "conformation"-bred and "field" Labrador Retrievers in 340 the U.S. (http://en.wikipedia.org/wiki/Labrador\_Retriever, accessed 22/11/13; Duffy et al., 341 2008) or 'show' and 'working' strains in the UK (Craig, 2011). The former are generally 342 seen in dog shows while the latter are the type generally used as gundogs in the UK. 343 Genetic differences between the two strains may be the result of breeding animals for good 344 'performance' in either Showdog or Gundog roles, where performance in either category is 345 likely to be related to the dog's behaviour and personality. Gundogs are working dogs and 346 are expected to be responsive and obedient throughout long periods where activity 347 (fetching/retrieval) is interspersed with periods of inactivity (waiting for the next shoot to 348 take place), unlike Pets and Showdogs that are not relied upon to complete specific tasks. 349 The increased Attention Seeking, Fetching, and Trainability may relate to being attentive to 350 the commands of the handler, performing the retrieval task reliably and being easily trained 351 for all required tasks. Similarly the decreased Barking Tendency and Noise Fear of 352 Gundogs may be attributed to their requirement to be quiet whilst working and between 353 shoots, and unafraid of gunshot. Gundogs were also less likely to show Unusual 354 Behaviours, which are often labeled as 'stereotypies'. It has been shown that stereotypies are 355 negatively associated with stimulation and engagement (Sergiel et al., 2012). The lowered 16

tendency to exhibit Unusual Behaviour in Gundogs may be due to the increased

357 environmental complexity that is associated with being a working dog, although category

358 shift could also explain the observed pattern.

Showdogs have to tolerate distracting environments with many people and animals in close proximity and occasional physical handling by unfamiliar people, which may explain their lower scores for Agitated when Ignored, Human and Object Fear, Noise Fear and Non-owner aggression than Pets. The breeding of successful showdogs may have promoted these personality traits.

364 However, previous training and experience may also explain some of these 365 differences in personality traits. The behavioural phenotype recorded in the questionnaire 366 may be influenced by the training or management regime of show and working dogs. 367 Whilst the differences in Trainability could be due to deliberate breeding strategies in 368 Gundogs, a Gundog will also undergo intensive training for its role, often by highly 369 experienced trainers. Therefore, the behavioural phenotype that was recorded in the 370 questionnaire may be influenced by this training, as has been shown in other studies 371 (Svartberg, 2002; Kutsumi et al., 2013). Similarly, Showdogs may have been become 372 desensitized to the multiple distractions of the dog show environment, and therefore react 373 less to these stimuli (Kubinyi et al., 2009), and thus score lower in Excitability. Those that 374 did not adapt well to either activity may have been removed due to category shift. 375 It is likely that both genetic and training/experience influence the personality traits 376 documented by the questionnaire in this population of dogs. An experimental approach

377 would be required to disentangle these factors, in which behavioural outcomes are recorded

from Show and Gundog strains that are managed and trained in pet, show and gundogenvironments.

380 **4.2.2. Coat Colour** 

381 Chocolate Labrador Retrievers were different from Black and Yellow dogs for 382 several traits. Chocolate dogs had lower Noise Fear and Trainability, and exhibited more 383 Unusual Behaviour than yellow or black dogs. Additionally, Chocolate dogs were more 384 Agitated when Ignored, more Excitable, and had lower Fetching than black dogs. There are 385 two possible explanations for the differences. Firstly, the genes responsible for chocolate 386 coat colour could be genetically linked to the genes responsible for these personality traits, 387 which would make these characteristics likely to co-occur. The inheritance pattern of coat 388 colour has been studied in Labrador Retrievers and other breeds, and mutations in the 389 tyrosine related protein 1 gene (TYRP1) have been shown to be responsible for brown coat 390 colour in dogs (Templeton et al., 1977; Schmutz et al., 2002), however, genetic associations 391 between this gene and behavioural traits have yet to be investigated. Another explanation is 392 that in the attempts of dog breeders to produce a high frequency of Chocolate dogs, the 393 gene pool of dogs carrying the alleles for a Chocolate coat may have become somewhat 394 separated from that of the other Labrador retrievers. By chance, traits other than coat colour 395 may have higher frequencies in this gene pool, which could explain the differences in 396 personality. Although the main difference was between Chocolate dogs and other dogs, 397 there were also some differences between Black and Yellow Labrador Retrievers (i.e. 398 Separation Anxiety). 399 4.2.3. Exercise

The level of exercise and stimulation a dog gets impacts its health and mental wellbeing (Sergiel *et al.*, 2012). The amount of time the dogs were exercised was associated with 8 of the 12 personality traits, and it had the second highest effect sizes. Associations with personality traits could be due to the level of exercise directly influencing the expression of certain behaviours, or it could be that dogs showing unwanted traits are not taken out by owners as much as other dogs, for fear of an inability to control them leading to embarrassment or harm, which we will call 'Behavioural Deterrence'.

407 Dogs exercised less had higher Excitability, lower Trainability, and exhibited more 408 Unusual Behaviour. The association between high levels of exercise and Trainability is 409 likely in part due to an increased exposure to training during activity. Dogs exercised more 410 showed less Unusual Behaviour, supporting the idea that higher exercise levels are good for 411 the mental health of the dogs. The increased stimulation from human interaction and time 412 outside in novel environments may help to reduce the incidence of stereotypies (Menor-413 Campos et al., 2011). This however is inconsistent with the results of Clark et al. (1997) 414 who found exercise had little effect on behaviour. The only behavioural difference they 415 found was an increase in barking among dogs exercised with a conspecific. However, the 416 periods of exercise used in their study were much shorter (20 minutes, 3 times a week) than 417 those reported here, suggesting that the quantity of exercise is important.

Dogs exercised more showed lower Non-Owner Aggression. This could be due to increased exposure to unknown stimuli during prolonged activity. Frequent contact would help familiarize the dog to strange people, animals and environments, and reduce the likelihood of an aggressive response to novelty. This is supported by dogs exercised less also showing higher Human and Object Fear, since fear and aggression have been shown to

be correlated in other studies (Duffy *et al.*, 2008). Behavioural Deterrence may also account
for these patterns, such that dogs showing non-owner aggression are not walked as often in
order to avoid awkward situations for the owner.

Dogs exercised less showed higher Barking Tendency and Owner Aggression. This may again be due to Behavioural Deterrence, or they may be barking to attract the attention of owners or as an outlet for boredom. This is supported by the finding that dogs exercised less also showed more Attention Seeking and less Separation Anxiety. Boredom may lead to frustration, manifested as aggression towards the people within the household.

431

## 432 **4.3. Gender and Age**

Gender status was significant in relation to 9 out of the 12 traits. Gender status is a combination of the sex and neuter status of the dogs, and both of these may have had an effect on their personality. Gender status has previously been shown to be associated with personality traits in dogs (Wilsson and Sundgren, 1997; Svartberg, 2002), including aggressiveness and boldness. Personality differences between the groups are likely to be due to hormonal differences.

Entire Males and Entire Females differed significantly from Neutered Females for
four traits. Neutered Females showed less Attention Seeking and Excitability, and showed
more Human and Object Fear and Noise Fear than the Entire dogs. Neutered Males were in
the middle and not significantly different from either group for all four of these traits.
Excitability may be higher in Entire than Neutered Females due to differences in hormone
levels.

445 Females showed more Non-Owner Aggression than Entire Males, and Entire Males 446 showed more Owner Aggression and less Unusual Behaviour than all Females, and higher 447 Separation Anxiety than all groups. Higher aggression among Males has been reported 448 previously (Hart and Hart, 1985, Wilsson and Sundgren, 1997). Castration of male dogs has 449 been shown to reduce aggressive dominance to some extent, but not territorial aggression 450 (Hart and Eckstein, 1997). The latter category may contain components of the Non-owner 451 Aggression category from the present study. However, higher Non-Owner Aggression from 452 Females was not been reported in either of these studies. This may represent a form of 453 territorial aggression, but levels of all aggression from this breed were low in this study. 454 The age of the dog is one of the easier factors to understand. The age of the animal 455 relates to its past experience, and therefore has an influence on its personality (Stamps and 456 Groothius, 2012). Older dogs showed significantly less Human and Object Fear. This is 457 possibly because experience has led them to discriminate between actual threats and 458 innocuous things. This is supported by older dogs showing less Separation Anxiety than 459 younger dogs, although this effect was not statistically significant. Older dogs have learned 460 their owners' routines, and are less concerned about prolonged absences.

461

#### 462 **5. Conclusions**

463 This large-scale study of behavioural characteristics in Labrador Retrievers revealed 464 a number of associations between physical, lifestyle and management characteristics of the 465 dogs and personality traits. The explanatory factor with the largest overall effect was the 466 working status of the dog, where pets showed dispositions that are generally considered 467 less desirable than those of gundogs and showdogs. The mechanism by which working 21 468 status could affect behaviour is not yet known, but it is likely to involve both genetic and

469 environmental factors. Further research is required to disentangle these factors. There were

- 470 also significant associations between personality traits and other factors considered,
- 471 including coat colour, levels of exercise, age, sex, neuter status and housing.
- 472

#### 473 Acknowledgments:

- 474 We are very grateful to Professor James Serpell, of the University of Pennsylvania, for
- 475 allowing us to use C-BARQ in our study and for supplying additional information about the
- 476 questionnaire for Table 1. We would like to thank the UK Kennel Club for initially
- 477 contacting the dog owners. We are very grateful to all the owners of Labrador Retrievers
- 478 who completed and returned the surveys and made this study possible. Funding was
- 479 provided by the UK Biotechnology and Biological Sciences Research Council (grant
- 480 BB/H019073/1 and core funding to the Roslin Institute).
- 481

#### 482 **References:**

- 483 Burrow, H.M., Corbet, N.J., 2000. Genetic and environmental factors affecting
- 484 temperament of zebu and zebu-derived beef cattle grazed at pasture in the tropics.
- 485 Australian Journal of Agricultural Research 51, 155-162.
- 486 Clark, J.D., Rager, D.R., Crowell-Davis, S., Evans, D.L., 1997. Housing and exercise of
- 487 dogs: effects on behavior, immune function, and cortisol concentration. Laboratory Animal
- 488 Science 47, 500-510.
- 489 Craig, D., 2011. The Labrador Retriever. Pet Book Publishing Company. PO Box Lydney
- 490 Gloucestershire GL15 6YD, United Kingdom.

- 491 Duffy, D.L., Hsu, Y.Y., Serpell, J.A., 2008. Breed differences in canine aggression.
- 492 Applied Animal Behaviour Science 114, 441-460.
- 493 Freedman, D.G., Elliot, O., King, J.A., 1961. Critical period in social development of dogs.
- 494 Science 133, 1016-1017.
- 495 Goddard, M.E., Beilharz, R.G., 1983. Genetics of traits which determine the suitability of
- 496 dogs as guide-dogs for the blind. Applied Animal Ethology 9, 299-315.
- 497 Hart, B.L., Hart, L.A., 1985. Selecting pet dogs on the basis of cluster-analysis of breed
- 498 behavior profiles and gender. Journal of the American Veterinary Medical Association 186,
- 499 1181-1185.
- 500 Hart, B.L. and Eckstein, R.A., 1997. The role of gonadal hormones in the occurrence of
- 501 objectionable behaviours in dogs and cats. Applied Animal Behaviour Science, 52, 331-
- 502 344.
- 503 Hoppe, S., Brandt, H.R., Konig, S., Erhardt, G., Gauly, M., 2010. Temperament traits of
- 504 beef calves measured under field conditions and their relationships to performance. Journal
- 505 of Animal Science 88, 1982-1989.
- 506 Hsu, Y.Y., Serpell, J.A., 2003. Development and validation of a questionnaire for
- 507 measuring behavior and temperament traits in pet dogs. Journal of the American Veterinary
- 508 Medical Association 223, 1293-1300.
- 509 Hsu, Y.Y., Sun, L.C., 2010. Factors associated with aggressive responses in pet dogs.
- 510 Applied Animal Behaviour Science 123, 108-123.
- 511 Jarvis, S., Moinard, C., Robson, S.K., Baxter, E., Ormandy, E., Douglas, A.J., Seckl, J.R.,
- 512 Russell, J.A., Lawrence, A.B., 2006. Programming the offspring of the pig by prenatal
- 513 social stress: Neuroendocrine activity and behaviour. Hormones and Behavior 49, 68-80.

- 514 Kubinyi, E., Turcsan, B., Miklosi, A., 2009. Dog and owner demographic characteristics
- and dog personality trait associations. Behavioural Processes 81, 392-401.
- 516 Kutsumi, A., Nagasawa, M., Ohta, M., Ohtani, N., 2013. Importance of puppy training for
- 517 future behavior of the dog. The Journal of veterinary medical science / the Japanese Society
- 518 of Veterinary Science 75, 141-149.
- 519 Ley, J., Bennett, P., Coleman, G., 2008. Personality dimensions that emerge in companion
- 520 canines. Applied Animal Behaviour Science 110, 305-317.
- 521 Liinamo, A.-E., van den Berg, L., Leegwater, P.A.J., Schilder, M.B.H., van Arendonk,
- 522 J.A.M., van Oost, B.A., 2007. Genetic variation in aggression-related traits in Golden
- 523 Retriever dogs. Applied Animal Behaviour Science 104, 95-106.
- 524 Mackenzie, S.A., Oltenacu, E.A.B., Houpt, K.A., 1986. Canine behavioral genetics-A
- 525 review. Applied Animal Behaviour Science 15, 365-393.
- 526 McMillan, F.D., Serpell, J.A., Duffy, D.L., Masaoud, E., Dohoo, I.R., 2013. Differences in
- 527 behavioral characteristics between dogs obtained as puppies from pet stores and those
- 528 obtained from noncommercial breeders. Journal of the American Veterinary Medical
- 529 Association 242, 1359-1363.
- 530 Menor-Campos, D.J., Molleda-Carbonell, J.M., Lopez-Rodriguez, R., 2011. Effects of
- 531 exercise and human contact on animal welfare in a dog shelter. Veterinary Record 169,
- 532 388.
- 533 Nagasawa, M., Tsujimura, A., Tateishi, K., Mogi, K., Ohta, M., Serpell, J.A., Kikusui, T.,
- 534 2011. Assessment of the Factorial Structures of the C-BARQ in Japan. Journal of
- 535 Veterinary Medical Science 73, 869-875.

- 536 Podberscek, A.L., Serpell, J.A., 1996. The English Cocker Spaniel: Preliminary findings on
- 537 aggressive behaviour. Applied Animal Behaviour Science 47, 75-89.
- 538 Schmutz, S.M., Berryere, T.G., Goldfinch, A.D., 2002. TYRP1 and MC1R genotypes and
- their effects on coat color in dogs. Mammalian Genome 13, 380-387.
- 540 Saetre, P., Strandberg, E., Sundgren, P.E., Pettersson, U., Jazin, E., Bergstrom, T.F., 2006.
- 541 The genetic contribution to canine personality. Genes Brain and Behavior 5, 240-248.
- 542 Scott, J.P., Marston, M.V., 1950. Critical periods affecting the development of normal and
- 543 mal-adjustive social behavior of puppies. Journal of Genetic Psychology 77, 25-60.
- 544 Sergiel, A., Maslak, R., Kusznierz, J., Pasko, L., 2012. Stereotypies: Development and
- 545 effects. Medycyna Weterynaryjna 68, 402-405.
- 546 Serpell, J.A., Hsu, Y.Y., 2001. Development and validation of a novel method for
- 547 evaluating behavior and temperament in guide dogs. Applied Animal Behaviour Science
- 548 72, 347-364.
- 549 Serpell, J.A., Hsu, Y.Y., 2005. Effects of breed, sex, and neuter status on trainability in
- 550 dogs. Anthrozoos 18, 196-207.
- 551 Stamps, J., Groothuis, T.G.G., 2010. The development of animal personality: relevance,
- 552 concepts and perspectives. Biological Reviews 85, 301-325.
- 553 Svartberg, K., 2002. Shyness-boldness predicts performance in working dogs. Applied
- 554 Animal Behaviour Science 79, 157-174.
- 555 Svartberg, K., Forkman, B., 2002. Personality traits in the domestic dog (Canis familiaris).
- 556 Applied Animal Behaviour Science 79, 133-155.
- 557 Svartberg, K., Tapper, I., Temrin, H., Radesater, T., Thorman, S., 2005. Consistency of
- 558 personality traits in dogs. Animal Behaviour 69, 283-291.

- 559 Svartberg, K., 2006. Breed-typical behaviour in dogs Historical remnants or recent
- 560 constructs? Applied Animal Behaviour Science 96, 293-313.
- 561 Templeton, J.W., Stewart, A.P., Fletcher, W.S., 1977. Coat color genetics in Labrador
- 562 Retriever. Journal of Heredity 68, 134-136.
- 563 Tobler, M., Sandell, M.I., 2007. Yolk testosterone modulates persistence of neophobic
- responses in adult zebra finches, Taeniopygia guttata. Hormones and Behavior 52, 640-645.
- van den Berg, L., Schilder, M.B.H., Knol, B.W., 2003. Behavior genetics of canine
- 566 aggression: Behavioral phenotyping of golden retrievers by means of an aggression test.
- 567 Behavior Genetics 33, 469-483.
- 568 Wilsson, E. and Sundgren P-E., 1997. The use of a behaviour test for the selection of dogs
- 569 for service and breeding, I: Method of testing and evaluating test results in the adult dog,
- 570 demands on different kinds of service dogs, sex and breed differences. Applied Animal
- 571 Behaviour Science 53: 279-295.
- 572
- 573
- 574
- 575
- 576
- 577
- 578
- 579
- 580
- 581

| 583 | Figure legends   |
|-----|--|
| 584 | Figure 1: Scatterplot of PC1 vs PC2 for C-BARQ responses. The different symbols refer to |
| 585 | the sections within C-BARQ (1: Training and obedience; 2: Aggression; 3: Fear and        |
| 586 | anxiety; 4: Separation-related behavior; 5: Excitability; 6: Attachment and attention-   |
| 587 | seeking; 7: Miscellaneous; Hsu & Serpell, 2003; http://vetapps.vet.upenn.edu/cbarq/).    |
| 588 |  |
| 589 | Figure 2: Adjusted means for the three Working Status categories (Gundog, Pet, Showdog)  |
| 590 | for the 12 personality traits described in the text (Agitated when Ignored, Attention    |
| 591 | Seeking, Barking Tendency, Excitability, Fetching, Human and Object Fear, Noise Fear,    |
| 592 | Non-Owner Aggression, Owner Aggression, Separation Anxiety, Trainability, and Unusual    |
| 593 | Behaviour).  |
| 594 |  |
| 595 | Figure 3: Adjusted means for the four gender/neuter status categories (EF=entire female; |
| 596 | EM=entire male; NF=neutered female; NM=neutered male) for the 12 personality traits      |
| 597 | described in the text (Agitated when Ignored, Attention Seeking, Barking Tendency,       |
| 598 | Excitability, Fetching, Human and Object Fear, Noise Fear, Non-Owner Aggression,         |
| 599 | Owner Aggression, Separation Anxiety, Trainability, and Unusual Behaviour).              |