

Pure

Scotland's Rural College

Sensitivity of the integrated Welfare Quality® scores to changing values of individual dairy cattle welfare measures

de Graaf, S; Bart, A; Buijs, S; Andreasen, S; de Boyer des Roches, A; Haskell, MJ; Kirchner, M; Mounier, L; Radeski, M; Winckler, C; Bijttebier, J; Lauwers, L; Verbeke, W; Tuytens, F

Published in:
Animal Welfare

DOI:
[10.7120/09627286.27.2.157](https://doi.org/10.7120/09627286.27.2.157)

First published: 01/05/2018

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

de Graaf, S., Bart, A., Buijs, S., Andreasen, S., de Boyer des Roches, A., Haskell, MJ., ... Tuytens, F. (2018). Sensitivity of the integrated Welfare Quality® scores to changing values of individual dairy cattle welfare measures. *Animal Welfare*, 27(2), 157 - 166. <https://doi.org/10.7120/09627286.27.2.157>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Animal Welfare

Sensitivity of the integrated Welfare Quality® scores to changing values of individual dairy cattle welfare measures

Journal:	<i>Animal Welfare</i>
Manuscript ID	F2001.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	09-Nov-2017
Complete List of Authors:	de Graaf, Sophie; ILVO, Farm animal welfare and behaviour Ampe, Bart; Institute for Agricultural and Fisheries Research (ILVO), Animal Sciences Buijs, Stephanie Andreasen, Sine; SEGES P/S, Danish Pig Research Center de Boyer des Roches, Alice; Université de Lyon, VetAgro Sup, UMR 1213 Herbivores; INRA, UMR1213 Herbivores van Eerdenburg, Frank; Faculty of Veterinary Medicine, Farm Animal Health Haskell, Marie; SRUC, Animal and Veterinary Sciences Kirchner, Marlene; University of Copenhagen, Large Animals Mounier, Luc; INRA, UMR1213 Herbivores, ; Université de Lyon, VetAgro Sup, UMR1213 Herbivores, Radeski, Miroslav; Ss. Cyril and Methodius University Winckler, Christoph; University of Natural Resources and Life Sciences, Department of Sustainable Agricultural Systems Bijttebier, Jo; Institute for Agricultural and Fisheries Research (ILVO), Social Sciences Unit Lauwers, Ludwig; Institute for Agricultural and Fisheries Research (ILVO) , Social Sciences Unit Verbeke, Wim; Universiteit Gent, Faculty of Bioscience Engineering, Department of Applied Biosciences Tuytens, Frank; ILVO, Animal Sciences Unit
Keywords:	animal welfare, animal-based welfare indicators, dairy cattle, integrated welfare index, sensitivity analysis, Welfare Quality

SCHOLARONE™
Manuscripts

1 Dairy cattle welfare assessment using Welfare Quality[®]
2 **Sensitivity of the integrated Welfare Quality[®] scores to changing values of individual**
3 **dairy cattle welfare measures**

4
5 S. de Graaf^{1,10}, B. Ampe¹, S. Buijs¹, S.N. Andreasen², A. De Boyer Des Roches^{3,7}, F.J.C.M.
6 van Eerdenburg⁴, M.J. Haskell⁵, M.K. Kirchner⁶, L. Mounier³, M. Radeski⁸, C. Winckler⁹, J.
7 Bijttebier¹, L. Lauwers¹, W. Verbeke¹⁰, F.A.M. Tuytens^{1*}

8
9 ¹*Institute for Agricultural and Fisheries Research (ILVO), Burgemeester van Gansberghelaan*
10 *92, 9820 Merelbeke, Belgium*

11 ²*Department of Large Animal Sciences, Faculty of Health and Medical Sciences, University*
12 *of Copenhagen, Groennegaardsvej 8, DK-1870 Frederiksberg, Denmark*

13 ³*Université de Lyon, VetAgro Sup, UMR1213 Herbivores, 69280 Marcy-L'Étoile, France*

14 ⁴*Department of Herd Animal Health, Utrecht University, 3508 TD Utrecht, The Netherlands*

15 ⁵*SRUC, West Mains Road, Edinburgh EH9 3JG, Scotland, United Kingdom*

16 ⁶*University of Copenhagen, Dept. of Veterinary and Animal Sciences, Section of Animal*
17 *Welfare and Disease Control, Grønnegårdsvej 8, 1870 Frederiksberg Copenhagen, Denmark*

18 ⁷*Institut National de la Recherche Agronomique, UMR1213 Herbivores, Equipe*

19 *Comportement Animal, Robustesse et Approche Intégrée du Bien-Etre, 63122 Saint Genes*

20 *Champanelle, France*

21 ⁸*Animal Welfare Center, Faculty of Veterinary Medicine, Ss. Cyril and Methodius University*
22 *in Skopje, Lazar Pop-Trajkov 5-7, 1000 Skopje, Republic of Macedonia*

23 ⁹*Division of Livestock Sciences, Department of Sustainable Agricultural Systems, University*
24 *of Natural Resources and Life Sciences, Gregor-Mendel Straße 33, 1180 Vienna, Austria*

1
2
3 25 ¹⁰Faculty of Bioscience Engineering, Ghent University, Coupure links 653, 9000 Ghent,
4
5 26
6
7
8
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27 * Contact for correspondence and requests for reprints: frank.tuytens@ilvo.vlaanderen.be
28

28 Abstract

29 The Welfare Quality[®] (WQ) protocol for on-farm dairy cattle welfare assessment describes 33
30 measures and a step-wise method to integrate the outcomes into 12 criteria scores, grouped
31 into four principle scores and into an overall welfare categorization with four possible levels.
32 The relative contribution of various welfare measures to the integrated scores has been
33 contested. Using a European dataset (491 herds), we investigated 1) variation in sensitivity of
34 integrated outcomes to extremely low and high values of measures, criteria and principles by
35 replacing each actual value with minimum and maximum observed and theoretically possible
36 values and 2) the reasons for this variation in sensitivity. As intended by the WQ consortium,
37 the sensitivity of integrated scores depends on 1) the observed value of the specific
38 measures/criteria, 2) whether the change was positive/negative, and 3) the relative weight
39 attributed to the measures. Additionally, two unintended factors of considerable influence
40 appear to be side-effects of the complexity of the integration method. Namely 1) the number
41 of measures integrated into criteria and principle scores, and 2) the aggregation method of the
42 measures. Therefore, resource-based measures related to drinkers, of which validity to assess
43 absence of prolonged thirst was criticized, have a much larger influence on integrated scores
44 than health-related measures like 'mortality rate' and 'lameness score'. Hence, the integration
45 method of the WQ protocol for dairy cattle should be revised to ensure that the relative
46 contribution of the various welfare measures to the integrated scores more accurately reflect
47 their relevance for dairy cattle welfare.

48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 49 **Keywords:** animal welfare, animal-based welfare indicators, dairy cattle, integrated welfare
4
5 50 index , sensitivity analysis, Welfare Quality®
6
7
8

9 51 **Introduction**

10 52 Accurate welfare assessment is vital for improving animal welfare. In dairy cattle, measures
11
12 53 have been developed and validated for a wide variety of both negative and positive aspects of
13
14 54 welfare. However, only a few protocols exist that aggregate the scores of multiple welfare
15
16 55 measures into one score or index reflecting the overall welfare status of a given herd. Such an
17
18 56 overall welfare status score might be used for example in the communication with consumers
19
20 57 (food labelling), as an incentive for on-farm welfare improvements and as regulative target
21
22 58 (Blokhuis et al 2010). Examples of schemes that calculate an overall welfare status of dairy
23
24 59 cattle are a protocol by Whay et al (2003) based on the “Five Freedoms” (Farm Animal
25
26 60 Welfare Council 1992) which generates a ranking of herds’ welfare status. The Animal Needs
27
28 61 Index (ANI) produces an overall welfare score based on integrating mostly resource-based
29
30 62 measures (measures of environmental aspects that affect welfare) (Bartussek et al 2000).
31
32 63 Finally, the Welfare Quality® (WQ) protocol categorizes overall welfare status of a herd as
33
34 64 ‘excellent’, ‘enhanced’, ‘acceptable’ or ‘not classified’ based on a step-wise integration
35
36 65 procedure (Welfare Quality® 2009). The current study focuses on the WQ protocol, as this is
37
38 66 the only protocol that predominantly uses animal-based measures to calculate an integrated
39
40 67 welfare index. Such measures are generally preferred over resource-based measures as the
41
42 68 latter tend to reflect risk factors for welfare impairments instead of directly measuring welfare
43
44 69 (Blokhuis et al 2003, 2010).
45
46
47
48
49
50

51 70

52
53
54 71 In the EU project Welfare Quality® (WQ), protocols for the welfare assessment of the main
55
56 72 types of farm animals (cattle, pigs and chickens) were proposed. The dairy cattle protocol
57
58 73 describes 33 welfare measures performed on-farm by means of behavioural observations,
59
60

1
2
3 74 qualitative behaviour assessment, an avoidance distance test, a management questionnaire, a
4
5 75 resource checklist and clinical scoring (Table 1). Subsequently, three steps are used to
6
7 76 integrate separate measures into one overall welfare category. Measures are first integrated
8
9 77 into criteria scores on a scale of 0 – 100 which are in turn collated into four welfare principles
10
11 78 ('good feeding', 'good housing', 'good health' and 'appropriate behaviour'). These principle
12
13 79 scores are then used to determine herds' overall welfare category (Welfare Quality® 2009).
14
15
16 80 Integration methods are intended to limit compensation of poor scores with better scores on
17
18 81 other welfare aspects (Veissier et al 2011). Expert opinion of social and animal scientists and
19
20 82 stakeholders was used to determine weights for the integration method (Botreau et al 2007).
21
22 83 Additionally, the protocols were designed with the intention of modifying and updating
23
24 84 assessment methods according to advances in animal welfare science
25
26
27 85 (www.welfarequalitynetwork.net/network/45848/7/0/40).
28
29
30 86

31
32 87 Discussion has arisen recently about WQ's **measures and** integration methods. Some of the
33
34 88 measures have been criticised for their poor or undocumented reliability, validity or feasibility
35
36 89 (Knierim and Winckler, 2009; de Vries et al., 2013; de Jong et al., 2015; Tuytens et al., 2015;
37
38 90 de Graaf et al., in press). In addition, studies have indicated that a few, resource-based
39
40 91 measures have a disproportionately large influence on the overall welfare category (Heath et
41
42 92 al 2014; de Vries et al 2013). Both critical findings may harm the credibility and validity of
43
44 93 the WQ protocol in assessing herd welfare. To further examine the functioning of the WQ
45
46 94 protocol for dairy cattle, the aim of the current study was to examine 1) if there is variation in
47
48 95 sensitivity of integrated outcomes (criteria and principle scores and overall welfare category)
49
50 96 to extremely low and high values of measures, criteria and principles and 2) the reasons for
51
52 97 this variation in sensitivity. More specifically, we aimed to critically evaluate whether
53
54 98 differences in sensitivity appear to be deliberate and justifiable rather than unintentional side-
55
56
57
58
59
60

1
2
3 99 effects of the complex integration method. To this end, we performed a sensitivity analysis by
4
5 100 replacing individual observed values for a given herd with both the theoretically possible and
6
7 101 the actually observed worst and best values. The latter values were based on a large database
8
9 102 of WQ data that reflect a wide range of herd types in Europe and thereby ensuring a
10
11 103 substantial but realistic spread in observed values.
12
13
14

15 16 104 **Materials and methods**

17 18 105 *WQ protocol*

19
20
21 106 Only a brief description of the integration method of the WQ protocol for on-farm dairy cattle
22
23 107 welfare assessment is given here. The full protocol can be found at
24
25 108 <http://www.welfarequalitynetwork.net/>.

26 27 28 29 30 110 *Step 1: from measures to criteria scores*

31
32 111 Aggregation starts by combining 33 measures into 11 rather than 12 criteria (Table 1),
33
34 112 because no data is collected on-farm for the criterion 'thermal comfort'. Because the
35
36 113 recording scales of measures differ, various aggregation methods are used. For categorical
37
38 114 measures, decision trees are used resulting in a score between 0 – 100 where 100 indicates the
39
40 115 best possible score. Other measures are converted to ordinal scores where required (e.g.
41
42 116 scores within 'comfort around resting' are converted into three categories: normal, moderate
43
44 117 problem or serious problem using thresholds in seconds for time needed to lie down and
45
46 118 percentages of cows for the other measures) and then combined into index values using
47
48 119 weighted sums. Spline functions are used to re-weight these sums based on their severity
49
50 120 according to expert opinion. Finally, when multiple spline functions were used, Choquet
51
52 121 integrals are used to combine these functions into criteria scores on a scale of 0 – 100
53
54
55 122 (Botreau et al 2007). These algorithmic operators calculate the criteria scores in such a way
56
57
58
59
60

1
2
3 123 that a poor score cannot be fully compensated for by a better score in another measure
4
5 124 (Botreau et al 2007). Consequently, poor scores will have a bigger influence on the integrated
6
7 125 scores than good scores. Using Choquet integrals, the weight given to each element (measures
8
9 126 or criteria) depends on its value relative to the other elements, where the poorest score always
10
11 127 gets the highest weight (Botreau et al., 2008; Welfare Quality 2009).
12
13

14 128

15
16 129 *Step 2: from criterion scores to principle scores*

17
18 130 To integrate criterion scores into principle scores, Choquet integrals are used (Welfare Quality
19
20 131 2009). The resulting principle scores range from 0 (worst) to 100 (best). Because no data is
21
22 132 collected on-farm for the criterion 'thermal comfort', this criterion score is replaced with the
23
24 133 best score among 'comfort around resting' and 'ease of movement'.
25
26
27

28 134

29
30 135 *Step 3: from principle scores to overall welfare category*

31
32 136 The third and final integration step is from principle scores to overall welfare category. Dairy
33
34 137 welfare in a herd is considered 'excellent' when it scores >50 for each principle and >75 on
35
36 138 two of them. When a herd scores >15 on each principle and >50 on at least two of them, it is
37
38 139 classified as 'enhanced'. 'Acceptable' herds score >5 for all principles and >15 for at least
39
40 140 three principles. Herds that do not reach the thresholds for the category 'acceptable' are
41
42 141 considered 'not classified' (Botreau et al 2009).
43
44

45
46
47 142 <Table 1>
48

49 143 ***Data collection and collation***

50
51
52 144 To reflect the current range present in Europe across various herding systems, pre-existing
53
54 145 research datasets of assessments using the WQ protocol for on-farm dairy cattle welfare were
55
56 146 collated from seven European research institutes and included data from 10 countries. The
57
58
59
60

1
2
3 147 collected samples were selected by the research institutes to be representative for 1) small
4
5 148 scale dairy herds in Macedonia (n = 12); 2) non-organic and non-tie stall dairy herds in The
6
7 149 Netherlands (n = 60) and France (n = 128); 3) random herds with individual Somatic Cell
8
9 150 Count data available (SCC, to be able to calculate WQ scores) in Belgium (n = 140), Scotland
10
11 151 (n = 16) and Denmark (n = 42); 4) typical herds for the regional low-input herding systems in
12
13 152 Romania, Northern Ireland and Spain (n = 30); and 5) loose housed dairy herds with at least
14
15 153 20 cows in Austria (n = 65). The total number of herds in the collated database was 491. To
16
17 154 ensure a homogenous integration method for all data, integrated WQ scores were calculated
18
19 155 from raw data using a custom-made integration procedure programmed in R 3.2.2 (R
20
21 156 Foundation for Statistical Computing, Vienna, Austria). The R integration programme is
22
23 157 available on request. The results were checked for coherence with the INRA WAFA webtool
24
25 158 (<http://www1.clermont.inra.fr/wq/>), in which WQ measure values can be entered (for dairy
26
27 159 cows, fattening pigs, growing pigs and broilers), and WQ criteria, principle and classification
28
29 160 scores can be calculated.
30
31
32
33

34 161 *Sensitivity analysis*

35
36
37 162 In order to investigate the extent to which values for separate measures affected the criteria
38
39 163 and principle scores and the overall welfare category, each herd-level observation for each
40
41 164 measure and each herd was replaced one by one with both the theoretically possible and the
42
43 165 observed (of the entire dataset of 491 herds) worst and best values. This was repeated for
44
45 166 individual criteria and principle scores to assess the impact of criteria and principle scores on
46
47 167 the overall welfare category. For these calculations, farms that were already in the highest or
48
49 168 lowest overall welfare category were excluded. This decision was made because these
50
51 169 excluded farms were not able to shift categories, therefore retaining them would give a
52
53 170 distorted picture of the results. Subsequently, the median increase and decrease in criteria and
54
55 171 principle scores and the percentage of herds that shifted to a lower or higher overall welfare
56
57
58
59
60

1
2
3 172 category were quantified for each replacement by the theoretically and observed worst and
4
5 173 best values.
6
7 174
8
9 175 For most measures, values that were altered were scored as either percentage of cows (e.g. %
10
11 176 of severely lame cows) or ‘yes’ and ‘no’ (e.g. for cleanliness of drinkers). However, for some
12
13 177 measures (avoidance distance at the feed rack (ADF), lameness and integument alterations)
14
15 178 the aggregated measure indexes rather than individual percentages were replaced with worst
16
17 179 and best scores. Because these measures together add up to 100% of animals, changing
18
19 180 percentages within these could create an impossible situation (i.e. percentages would add up
20
21 181 to over 100%). In addition, the theoretical best score for the measures ‘length of drinking
22
23 182 trough’ and ‘number of drinking bowls’ depends on the average number of cows on the herd.
24
25 183 Therefore, we replaced these with scores that would meet the requirements for all herds in the
26
27 184 dataset (10,000 cm for drinking trough length and 100 for number of drinking bowls) as best
28
29 185 scores. For the measures of dehorning and tail docking, we replaced the actual methods used
30
31 186 at each herd with the methods which would generate the best (i.e. no dehorning, no tail
32
33 187 docking respectively) and the worst score (i.e. dehorning using surgery with no anaesthetics
34
35 188 or analgesics, tail docking using a rubber band without anaesthetics and analgesics,
36
37 189 respectively).

190 **Results**

191 None of the 491 herds were originally (i.e. before replacement with worst/best scores) in the
192 ‘excellent’ category, 174 (35%) were in the ‘enhanced’ category, 308 (63%) in the
193 ‘acceptable’ category and nine (2%) in the ‘not classified’ category. For eight of the nine ‘not
194 classified’ herds, classification was due to a ‘good feeding’ principle score below 5 (the
195 threshold for the not-classified category). The median, minimum, and maximum scores are
196 given at the measure (Table 2) and principle and criterion level (Table 4). For several

1
2
3 197 measures, the observed range spanned the entire theoretical range (i.e. 0 – 100 for
4
5 198 percentages, 0 – 24 for hours and 0 – 365 for days). However, for several other measures (18
6
7 199 out of 33), criteria (6 out of 12) and principles (3 out of 4), the observed data range was
8
9 200 narrower than was theoretically possible (Tables 2 and 3). Only 5% of herds were not
10
11 201 dehorned or disbudded, 18% were disbudded using caustic paste, 76% using thermocautery,
12
13 202 and 1% was dehorned using surgery. Analgesics and/or anaesthetics were used during these
14
15 203 procedures in 24% and 60% of the herds, respectively. Only 5 (ca. 1%) herds were tail-
16
17 204 docked (3 by rubber ring and 2 by surgery). Analgesics were never used during tail docking
18
19 205 whilst anaesthetics were used in two herds.
20
21
22
23
24

207 *Sensitivity analysis using observed values: measurement level*

208 *Sensitivity of the overall welfare category*

209 When separate measure values were increased to the observed maximum value (i.e. to the
210 level of the herd that scored best for that specific measure) fewer herds shifted between
211 overall categories than when separate scores were decreased to the observed minimum value
212 (Table 2). Regarding the overall welfare categories between which the shifts occurred, for
213 most measures, the highest percentage of shifts occurred between the ‘enhanced’ and
214 ‘acceptable’ category (percentage of shifts ranging from 0 – 99%). However, for increases in
215 some measures (‘% of lean cows’, ‘number of water bowls’, ‘cleanliness of drinker’ and
216 ‘loose versus tied housing’) highest % of shifts to a higher category were between ‘not
217 classified’ and ‘acceptable’ (percentage of shifts ranging from 22 - 100%).
218

219 Replacements of measure values only rarely led to negative shifts of more than one category
220 and never to positive shifts of more than one category (Table 2). The effects of replacing a
221 measure often differed greatly, even between measures that belong to the same principle.

1
2
3 222 'Good health' was the only principle for which changing the values of any of its underlying
4
5 223 measures did not result in a substantial (>10%) effects on herd classification. All measures
6
7 224 that were the only measure of a certain criterion caused a relatively high percentage of herds
8
9 225 to shift category: '% of lean cows', 'loose or tied housing' and the 'QBA index' when
10
11 226 replaced with the worst possible score, with the exception of the 'ADF index'. Although
12
13 227 seemingly combined with many other measures, most measures of the criterion 'absence of
14
15 228 prolonged thirst' had a relatively large influence as well. Most upgrades to a higher overall
16
17 229 welfare category were achieved by increasing (to the observed maximum levels) 'number of
18
19 230 water bowls', 'trough length', and to a lesser extent '% of cows colliding'. Within the two
20
21 231 criteria that contained most measures, either sensitivity was very low for all measures
22
23 232 ('absence of disease') or sensitivity was greater for those measures that were attributed the
24
25 233 highest weight (i.e. within 'comfort around resting', the measures for resting behaviour are
26
27 234 given a higher weight than cleanliness).

28
29
30
31
32
33 235 <Table 2>

34
35
36 236 *Sensitivity of the principles and criteria scores*

37
38
39 237 The sensitivity analysis of the effect of changes in separate measure values on the principles
40
41 238 scores and on the criteria scores (Table 3) showed the same pattern as the sensitivity analysis
42
43 239 of the overall welfare category. The decrease caused by changing a measure to the lowest
44
45 240 observed value was usually greater than the increase caused by changing the same measure to
46
47 241 its highest observed value. Exceptions to this trend often concerned measures of which the
48
49 242 observed values were very poor. Furthermore, measures that caused the greatest difference
50
51 243 tended to belong to criteria that contain few other measures. Exceptions to this trend once
52
53 244 again concerned most measures within 'absence of prolonged thirst' and the measure '% of
54
55 245 cows colliding with housing'. There was a difference in the sensitivity of the principles and
56
57
58
59
60

1
2
3 246 the criteria in that measure values have a more direct influence on criteria scores, and
4
5 247 therefore had a greater influence on criteria scores than on principle scores.
6
7
8

9 248 <Table 3>
10
11 249

12
13 250 ***Sensitivity analysis using observed values: criteria and principle level***

14
15 251 Of all principles, alteration of ‘good feeding’ led to the highest number of negative shifts as
16
17 252 well as positive shifts (Table 4). Moreover, replacing the ‘good feeding’ score to the lowest
18
19 253 observed score in the database caused all ‘enhanced’ herds to be re-categorised as ‘non-
20
21 254 classified’. Alterations to the other principle scores never caused a change of more than one
22
23 255 overall welfare category. Alteration of the ‘good housing’ principle caused the fewest positive
24
25 256 shifts of all principles, as most farms already scored relatively high for this principle (median
26
27 257 score of 54).
28
29
30

31 258

32
33 259 Of all criteria, replacement with the lowest observed score was most effective in generating
34
35 260 negative shifts for ‘absence of prolonged hunger’ followed by ‘absence of prolonged thirst’.
36
37 261 Replacement with the highest observed score was most effective in generating a positive shift
38
39 262 for ‘absence of prolonged thirst’. Both criteria within the principle ‘good housing’ (‘comfort
40
41 263 around resting’ and ‘ease of movement’) caused 27% of herds to be downgraded when
42
43 264 replaced by the observed minimum. Effects of replacing criteria scores within the ‘good
44
45 265 health’ and ‘appropriate behaviour’ principles varied considerably between criteria.
46
47
48
49

50
51 266 <Table 4>
52

53 267 ***Differences between replacement with observed and theoretically possible scores***

54
55 268 For several measures, criteria and principles, the observed range did not span the entire
56
57 269 theoretical range. For three measures (‘lameness index’, ‘head butts/cow/15 min’ and ‘ADF
58
59
60

1
2
3 270 index'), four criteria ('absence of injuries', 'absence of diseases', and 'absence of pain
4
5 271 induced by management procedures') and three principles ('good housing', 'good health' and
6
7 272 'appropriate behaviour'), replacement with the theoretically possible scores instead of the
8
9 273 observed scores resulted in a higher % of herds shifting between overall welfare categories
10
11 274 (Table 5). For four measures ('% lean cows', 'lameness index', 'number of coughs/cow/15
12
13 275 min.', '% cows with hampered respiration' and 'ADF index), this resulted in a higher median
14
15 276 increase or decrease of the principle and criteria scores than when worst or best observed
16
17 277 scores were used (Table 6).

18
19
20
21
22 278 <Table 5>

23
24
25
26 279 <Table 6>

27 280 **Discussion**

28
29
30
31
32
33 281 This study investigated the sensitivity of the integrated scores of the WQ protocol for on-farm
34
35 282 dairy cattle welfare assessment to extreme changes in individual measure, criterion and
36
37 283 principle scores. The impact of one by one replacement of observed herd-level measure,
38
39 284 criteria and principle scores by extremely low or high values had variable effects on the more
40
41 285 highly integrated scores and on the overall welfare category. Investigation into what type of
42
43 286 replacements have a large versus negligible impact suggests that a considerable part of this
44
45 287 variation appears to be an unwanted side-effect of the complex step-wise integration method
46
47 288 rather than being intentional or justifiable.

48
49
50 289

51 290 *Sensitivity analysis using observed values: measurement level*

52
53
54
55 291 Generally, the impact of a replacement with an extremely low score was bigger than
56
57 292 replacement with an extremely high score. This reflects the intention of the WQ integration
58
59
60

1
2
3 293 method to limit compensation of poor scores with better scores on other welfare aspects
4
5 294 (Veissier et al 2011). The effect of replacing observed measure scores with extreme values on
6
7 295 more highly integrated scores (criteria and principles) and on the overall welfare category was
8
9 296 very variable and seemed to depend on various aspects. Replacements of the measures ‘% of
10
11 297 lean cows’, ‘loose/tied housing’, the ‘QBA index’, ‘drinker trough length’ and ‘cleanliness of
12
13 298 drinkers’, had a bigger impact on overall classification compared to other measures
14
15
16 299 (particularly when substituted by observed worst scores). The common feature shared by the
17
18 300 first three measures is that they are the only measure of the criterion they belong to (‘absence
19
20 301 of prolonged hunger’, ‘ease of movement’ and ‘positive emotional state’, respectively). One
21
22 302 other criterion is also documented by a single measure, namely ‘expression of other normal
23
24 303 behaviour’ measured with the ADF-test. This measure had less impact compared with the
25
26 304 aforementioned three measures, presumably because the ADF-index was already poor for
27
28 305 most farms to begin with (so the change by replacing the actual score with the worst possible
29
30 306 score was often very small) .
31
32
33
34
35

36 308 The relatively large impact of drinker space and cleanliness of drinkers is in accordance with
37
38 309 previous findings for both the dairy cattle protocol (de Vries et al 2013; Heath et al 2014) and
39
40 310 the WQ broiler chicken protocol (Buijs et al 2016). This seems to be caused by a combination
41
42 311 of factors. First, these measures both belong to the criterion of ‘absence of prolonged thirst’
43
44 312 which contains few measures that matter for calculating the criterion scores (in the decision
45
46 313 tree only number/length of drinkers and cleanliness are taken into account). The other
47
48 314 measures are either prerequisites for the required number/length of drinkers and therefore less
49
50 315 directly influence criterion scores (‘water flow’), or are related to the number of drinkers (‘at
51
52 316 least 2 drinkers/cow’). Second, the principle ‘good feeding’ contains only one other criterion
53
54
55 317 apart from ‘absence of prolonged thirst’, whereas most other principles are composed of more
56
57
58
59
60

1
2
3 318 criteria. It could be argued that the large impact of these measures is not necessarily
4
5 319 problematic if they are valid indicators of an important welfare problem. However, as
6
7 320 resource-based measures, drinker space and cleanliness would appear to be potential risk
8
9 321 factors rather than direct measures of thirst (Sprenger et al 2009; Vanderhasselt et al 2014).
10
11 322 Moreover, to our knowledge, the validity of these measures of thirst has not yet been tested.
12
13 323 Therefore, the finding that these measures have a relatively large influence on integrated
14
15 324 scores can be considered problematic. Animal-based indicators of thirst have been developed,
16
17 325 such as blood sodium concentrations, plasma osmolality (Reece, 2009; Vanderhasselt et al.,
18
19 326 2013) and voluntary water consumption (in broiler chickens; Sprenger et al., 2009;
20
21 327 Vanderhasselt et al., 2014). Whereas blood parameters are too invasive to perform in on-farm
22
23 328 welfare monitoring, it could be promising to develop voluntary water consumption tests
24
25 329 further. Identifying the most reliable, valid and feasible measure of prolonged thirst in dairy
26
27 330 cattle should be a priority in future animal welfare assessment research.
28
29
30
31

32 331
33
34 332 Replacements of measures within the principle 'good health' with the best or worst scores had
35
36 333 little influence on principle and criterion scores and on overall classification, in accordance
37
38 334 with previous results (de Vries et al 2013; Heath et al 2014; Nielsen et al 2015). This is
39
40 335 remarkable because it includes measures which indicate important welfare problems in dairy
41
42 336 cattle according to many experts, such as mortality, mastitis and lameness (Nielsen et al 2014;
43
44 337 Lievaart and Noordhuizen, 2011). In addition, Tuytens et al (2010) reported that both
45
46 338 consumers and farmers rank health aspects as the most important for farm animal welfare.

47
48 339 The very limited effect of extreme changes in measures within the criterion 'absence of
49
50 340 diseases' on integrated WQ scores seems to be caused, at least partially, by the aggregation
51
52 341 method of this criterion. In this aggregation, prevalence of symptoms of diseases is compared
53
54 342 to warning and alarm thresholds (e.g. warning threshold for nasal discharge is 5% of cows and
55
56
57
58
59
60

1
2
3 343 alarm threshold 10% of cows). Subsequently, a weighted sum is calculated of warnings and
4
5 344 alarms, with a weight of 1 for warnings and 3 for alarms, which is computed into the criterion
6
7 345 score using a spline function. Because of this method, increasing prevalences that were
8
9 346 already above the alarm threshold (or decreasing those that were already below the threshold)
10
11 347 will not affect classification at all. Also, when the prevalence of one disease symptom
12
13 348 changes, it has only a limited effect on the criterion scores because it is aggregated with many
14
15 349 other disease symptoms.
16

17
18
19 350

20
21 351 Similarly to measures within ‘absence of diseases’, measures within ‘absence of injuries’ also
22
23 352 had a small impact on the integrated scores. However, a different method is used to integrate
24
25 353 the measures within ‘absence of injuries’ to one score. Partial scores for lameness and
26
27 354 integument alterations are first calculated using weighted sums and i-spline curves, and are
28
29 355 then combined using a Choquet integral. The lameness index had most influence, but still
30
31 356 caused only 10% of herds to be downgraded when replaced with the theoretically worst
32
33 357 possible score (i.e. 100% severely lame cows). This surprisingly low impact seems to be due
34
35 358 to the large number of criteria within the principle ‘good health’, and to the observation that
36
37 359 herds often score relatively low for these criteria. Therefore, changing another score within
38
39 360 this principle to a low score is likely to have a smaller effect than when it is done for a score
40
41 361 in another principle with fewer criteria such as ‘good feeding’. Due to the limited impact of
42
43 362 good health measures on overall welfare categorisation, in theory a situation could occur
44
45 363 where farms categorised as ‘acceptable’ or better have 100% severely lame animals, while
46
47 364 this may obviously be considered a major welfare problem.
48
49
50

51
52 365

53
54 366 Regarding positive shifts, the percentage of cows colliding with housing had a relatively large
55
56 367 positive impact when replaced with best observed score. This is likely because a large
57
58
59
60

1
2
3 368 proportion of farms (55%) were classified as having a serious problem for this measure to
4
5 369 begin with, so for many farms a vast improvement was possible (compared to 37% for % of
6
7 370 cows laying out and 28% which were above the threshold value of 6.3 seconds for mean time
8
9 371 needed to lie down).

10
11 372

12
13
14 373 *Sensitivity analysis using observed values: criteria and principle level*

15
16 374 There are two, three, or four criteria per principle. This difference in the number of criteria is
17
18 375 reflected in the results of the sensitivity analysis: replacement with the worst criteria scores
19
20 376 within the principle ('good feeding') containing only two criteria ('absence of prolonged
21
22 377 hunger' and 'absence of prolonged thirst') generated most shifts towards a different welfare
23
24 378 category. The principle 'good housing' also consists of only two criteria for which measures
25
26 379 have been developed (for its third criterion 'thermal comfort' no measure is available). The
27
28 380 impact of both criteria are smaller compared to the two criteria of 'good feeding'. However,
29
30 381 even though for 'thermal comfort' no data are collected, the missing criterion score is
31
32 382 replaced with the best score among 'comfort around resting' and 'ease of movement'. This
33
34 383 dilutes the effect of a very low score on either of these two criteria. Although some validated
35
36 384 measures for thermal comfort exist for dairy cattle (e.g. respiration rate, Schutz et al., 2010),
37
38 385 inclusion of such measures may complicate timing of farm visits, as the **outcomes of these**
39
40 386 measures are highly **influenced by** ambient temperature **and humidity**. **Therefore, climatic**
41
42 387 **conditions should be similar during farm visits to capture farm-level differences in thermal**
43
44 388 **comfort rather than differences based on ambient weather conditions**. Further research on how
45
46 389 to deal with these complexities in the WQ protocol is necessary, or removal of 'thermal
47
48 390 comfort' as a criterion for dairy cattle welfare should be considered.

49
50
51
52
53
54 391

1
2
3 392 In line with the criteria, of all principles, alteration of ‘good feeding’ led to the most negative
4
5 393 and positive shifts when replaced with observed worst and best scores. For negative shifts this
6
7 394 was because ‘good feeding’ was the only principle for which scores <5 were observed, which
8
9 395 automatically categorizes a herd as ‘not classified’. For positive shifts, this was because this
10
11 396 principle caused more ‘not classified’ and ‘acceptable’ categorizations than any other
12
13 397 principle (as 131 farms originally had a score between 5 and 15 for this principle, as opposed
14
15 398 to 9 for housing, 3 for health and 23 for behaviour). Therefore, more positive shifts could
16
17 399 occur when ‘good feeding’ was altered than when the other principles were replaced with
18
19 400 observed maximum scores.
20
21
22
23

401

402 *Differences between replacement with observed and theoretically possible scores*

403 As the sample size in the current study was large and contained a wide variety of herds (given
404 the different sampling aims), we can draw some conclusions about the observed scores in
405 relation to theoretical possible scores. For most measures, observed scores spanned the entire
406 theoretical range. This means that for the dairy cattle protocol, most limits set by WQ seem
407 realistically attainable. For some measures however, observed scores were less extreme than
408 the theoretically possible scores. In most cases, this did not affect criterion scores as these
409 were within the criterion ‘absence of diseases’, where warning and alarm thresholds are used
410 to integrate scores. For lameness index and ADF index however, fewer shifts of the overall
411 welfare category were observed when replaced with the observed scores. This was also
412 reflected in the corresponding criteria and principle scores, of which the worst possible score
413 never occurred. This is one of the reasons that the principles ‘good health’ and ‘appropriate
414 behaviour’ never caused herds to be categorized as ‘not classified’ when replaced by the
415 observed minimum score.

416 **Conclusion**

417 The results of the current study provide insight into the functioning of the integration methods
418 for the dairy cattle WQ protocol. Findings indicate that the sensitivity of integrated scores to
419 replacement of individual scores by extreme scores is dependent on a number of factors which
420 were intended by the WQ protocol: 1) the observed value of the specific measure (or
421 criterion), relative to the values of the other measure in the same criterion (or principle); 2)
422 whether the values were replaced by an extremely low or an extremely high value (more
423 impact of the former); 3) the relative weight WQ attributes to the measures. However, two
424 other factors that were not intended and appear to be unwanted side-effects of the complexity
425 of the step-wise integration method also had considerable influence. These factors were: 1)
426 the number of measures that are integrated into criteria and principle scores; and 2) the
427 aggregation method of the measures (e.g. decision trees or weighted sums). The effect of both
428 integration method and grouping is problematic, as it should be the severity of the welfare
429 problem that affects the overall category. As a result, sensitivity is highest for changes in
430 measures of the ‘good feeding’ principle, of which a large proportion of the measures are
431 criticized for their validity (i.e. measures of ‘absence of prolonged thirst’). On the contrary,
432 measures within the principle ‘good health’ have the lowest impact while some of these
433 measures are considered to most severely affect dairy cattle welfare. For instance, a farm in
434 the ‘acceptable’ category or higher could theoretically have 100% severely lame animals. The
435 unwanted side-effects of the current WQ integration methods shown in this study warrant
436 research to develop and evaluate alternative integration methods.

437

438 *Animal welfare implications*

439 This study indicates that the WQ integration method does not adequately balance the relative
440 importance of all welfare measures that are included in order to adhere to the multi-

1
2
3 441 dimensional nature of animal welfare. Therefore, using the current integrated WQ scores
4
5 442 could lead to a focus on a limited set of (often resource-based) measures which is hard to
6
7 443 justify. As this harms the credibility of the assessment protocol, we recommend a revision of
8
9 444 the integration method, so that the relative contribution of the various welfare measures to the
10
11 445 integrated scores more correctly reflects their relevance for dairy cattle welfare.
12
13
14
15

16 446 **References**

- 17
18 447 **Bartussek H, Leeb C and Held S** 2000 Animal Needs Index for Cattle. ANI 35, L/2000.
19
20 448 <http://www.bartussek.at/veroeffentlichungen/511134991b0db8204/index.html>
21
22
23 449 **Blokhuis HJ, Jones RB, Geers R, Miele M and Veissier I** 2003 Measuring and monitoring animal
24
25 450 welfare: Transparency in the food product quality chain. *Animal Welfare* 12: 445-455
26
27
28 451 **Blokhuis HJ, Veissier I, Miele M and Jones B** 2010 The Welfare Quality® project and beyond:
29
30 452 Safeguarding herd animal well-being. *Acta Agriculturae Scand* 60: 129-140
31
32
33 453 **Buijs S, Ampe B and Tuytens FAM** 2016 Sensitivity of the Welfare Quality® Broiler chicken
34
35 454 protocol to differences between intensively reared indoor flocks: which factors explain overall
36
37 455 classification? *Animal* 15: 1-10
38
39
40 456 **Botreau R, Veissier I, Butterworth A, Bracke MBM and Keeling LJ** 2007 Definition of criteria for
41
42 457 overall assessment of animal welfare. *Animal Welfare* 16: 225-228
43
44
45 458 **Botreau R, Capdeville J, Perny P, and Veissier I** 2008 Multicriteria evaluation of animal welfare at
46
47 459 farm level: An application of MCDA methodologies. *Foundation of Computing and Decision*
48
49 460 *Sciences* 33: 1-18
50
51
52 461 **de Graaf S, Ampe B and Tuytens FAM** in press Assessing dairy cow welfare at the beginning and
53
54 462 end of the indoor period using the Welfare Quality® protocol. *Animal Welfare*.
55
56
57
58
59
60

- 1
2
3 463 **de Vries M, Bokkers EAM, van Schaik G, Botreau RI, Engel B, Dijkstra T and de Boer IJM**
4
5 464 2013 Evaluating results of the Welfare Quality multi-criteria evaluation model for
6
7 465 classification of dairy cattle welfare at the herd level. *Journal of dairy science* 96: 6264-6273
8
9
10 466 **Herd Animal Welfare Council** 1992 FAWC updates the five freedoms. *Veterinary Record*
11
12 467 17: 357
13
14 468 **Heath CAE, Browne, WJ, Mullan S, Main DCJ** 2014 Navigating the iceberg: reducing the number
15
16 469 of parameters within the Welfare Quality® assessment protocol for dairy cows. *Animal* 8:
17
18 470 1978-1986
19
20
21 471 **Lievaart JJ and Noordhuizen JPTM** 2011 Ranking experts' preferences regarding measures and
22
23 472 methods of assessment of welfare in dairy herds using Adaptive Conjoint Analysis. *Journal of*
24
25 473 *dairy science* 94: 3420-3427.
26
27
28 474 **Nielsen BH, Angelucci A, Scalvenzi A, Forkman B, Fusi F, Tuytens F, Houe H, Blokhuis H,**
29
30 475 **Sørensen JT, Rothmann J, Matthews L, Mounier L, Bertocchi L, Richard M, Donati M,**
31
32 476 **Nielsen PP, Salini R, de Graaf S, Hild S, Messori S, Nielsen SS, Lorenzi V, Boivin X and**
33
34 477 **Thomsen PT** 2014 Use of animal based measures for the assessment of dairy cow welfare-
35
36 478 ANIBAM. *EFSA External scientific report*.
37
38
39 479 **Reece WO 2009**. Functional anatomy and physiology of domestic animals. John Wiley & Sons. Iowa,
40
41 480 USA.
42
43
44 481 **Schütz KE, Rogers AR, Poulouin YA, Cox NR, and Tucker CB** 2010 The amount of shade
45
46 482 **influences the behavior and physiology of dairy cattle. *Journal of dairy science* 93: 125–133.**
47
48
49
50 483 **Sprenger M, Vangestel C and Tuytens FAM** 2009 Measuring thirst in broiler chickens. *Animal*
51
52 484 *Welfare* 18: 553-560
53
54
55 485 **Tuytens FAM, Vanhonacker F, Van Poucke E and Verbeke W** 2010 Quantitative verification of
56
57 486 the correspondence between the Welfare Quality® operational definition of herd animal
58
59
60

- 1
2
3 487 welfare and the opinion of Flemish herders, citizens and vegetarians. *Livestock Science* 131:
4
5 488 108-114
6
7
8 489 **Vanderhasselt RF, Buijs S, Sprenger M, Goethals K, Willemsen H, Duchateau L and Tuytens**
9
10 490 **FAM** 2013 Dehydration indicators for broiler chickens at slaughter. *Poultry Science* 92: 612-
11
12 491 619.
13
14
15 492 **Vanderhasselt RF, Goethals K, Buijs S, Federici JF, Sans ECO, Molento CFM, Duchateau L**
16
17 493 **and Tuytens, FAM** 2014 Performance of an animal-based test of thirst in commercial broiler
18
19 494 chicken herds. *Poultry science* 93:1327-1336.
20
21
22 495 **Veissier I, Jensen KK, Botrea R and Sandøe P** 2011 Highlighting ethical decisions underlying the
23
24 496 scoring of animal welfare in the Welfare Quality® scheme. *Animal Welfare* 20:89-101
25
26
27 497 **Welfare Quality® Consortium** 2009 Welfare Quality® Assessment Protocol for Cattle. Lelystad,
28
29 498 The Netherlands
30
31 499 **Whay HR, Main DCJ, Webster AJF and Green LE** 2003 Assessment of the welfare of dairy cattle
32
33 500 using animal based measurements: direct observations and investigation of herd records. *The*
34
35 501 *Veterinary Record* 153: 197-202
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

502 **Appendix**

503 Percentages of herds¹ (n = 491) that were downgraded and upgraded by 1 or 2 overall welfare
 504 categories when individual values at measure level within the criterion ‘absence of diseases’ were
 505 replaced with theoretical worst and best values per measure

Measures	Observed worst score	Observed best score
	% downgraded 1 category	% upgraded 1 category
Number of coughs/cow/minute	2	0
% cows with nasal discharge	2	0
% cows with ocular discharge	2	0
% cows with hampered respiration	1	0
% cows with diarrhoea	2	0
% cows with vulvar discharge	2	0
% cows with SCC >400.000	2	1
% cows mortality	2	1
% calvings with dystocia	1	0
% downer cows	1	1

506 ¹Percentages were based on the herds that were actually able to shift one or two categories. For
 507 downgrades of 1 category n = 482, for downgrades of 2 categories n = 174. For upgrades of 1
 508 category n = 491.

509

Table 1: All principles, the corresponding criteria and indicators used in the Welfare Quality® assessment protocol for dairy cattle welfare

Principles	Criteria	Measures	Aggregation method measures
Good feeding	Absence of prolonged hunger	Body Condition Score (% very lean animals)	Spline curve fitting
	Absence of prolonged thirst	Availability & cleanliness water	Decision tree
Good housing	Comfort around resting	Lying down duration; collisions during lying down ; on edge/outside of lying area; cleanliness	Converted to ordinal scores, combined in weighted sums and spline curve fitting
	Thermal comfort Ease of movement	No measure for dairy cattle Free stalls or presence of tethering and exercise	Decision tree
Good health	Absence of injuries	Lameness; integument alterations	Combined in weighted sums, spline curve fitting and Choquet integration
	Absence of disease	Respiration/digestive diseases; mastitis; mortality; dystocia, downer cows	Converted to ordinal scores, combined in weighted sums and spline curve fitting
	Absence of pain induced by management procedures	Mutilations (dehorning; tail docking; use of anaesthetics/analgesics)	Decision tree
Appropriate behaviour	Expression of social behaviour	Incidence agonistic interactions	Combined in weighted sums and spline curve fitting
	Expression of other behaviours	Access to pasture	Spline curve fitting
	Good human-animal relationship	Avoidance distance at feeding place	Combined in weighted sums and spline curve fitting
	Positive emotional state	Qualitative Behavioural Assessment	Combined in weighted sums and spline curve fitting

Table 2: Percentages of herds¹ (n = 491) that were downgraded or upgraded 1 or 2 overall welfare categories when individual values at measure level (continuous and binary) were replaced with observed worst and best values per measure

Principles	Criteria, Continuous measures	Observed median, min - max	Observed worst score		Observed best score
			% downgraded 1 category	% downgraded 2 categories	
			% downgraded 1 category	% downgraded 2 categories	% upgraded 1 category
Good feeding	Absence of prolonged hunger				
	% of lean cows ²	4, 0 – 88	53	0	5
Good housing	Comfort around resting				
	Mean time needed to lie down (s)	6, 3 – 20	10	0	6
	% of cows colliding with housing	33, 0 – 100	5	0	12
	% of cows lying outside of lying area	0, 0 – 73	11	0	8
	% of cows with dirty flanks	64, 0 – 100	0	0	7
	% of cows with dirty lower legs	80, 0 – 100	2	0	7
	% cows with a dirty udder	37, 0 – 100	2	0	7
Good health	Absence of injuries				
	Lameness index	88, 37 – 100	6	0	5
	Integument alterations index	53, 0 – 100	2	0	4
	Absence of diseases				
	Range of all disease-measures ²	-	1-2	0	0-1
Appropriate behaviour	Expression of social behaviour				
	Head butts/cow/15 min.	0.5, 0 – 7	13	0	1
	Displacements/cow/15 min.	0.4, 0 – 5	16	0	4
	Expression of other normal behaviour				
	Number of hours on pasture)	7.5, 0 - 24	9	0	1
	Number of days on pasture	175, 0 - 365	9	0	1
	Human-animal interaction				
	ADF index	67, 23 – 100	13	0	6
Positive emotional state					
	QBA index	0.3, -11 – 5	24	1	7
	Criteria, Binary measures	% farms with best score			
Good feeding	Absence of prolonged thirst				
	Water flow	82	22	3	3
	Trough length	18	26	1	19
	Number of water bowls		11	1	20
	Drinker cleanliness	76	23	0	8
	At least 2 drinkers/cow	84	9	0	1
Good	Ease of movement				

housing	Loose or tied housing	93	38	2	3
Good health	Absence of pain induced by management procedures				
	Dehorning method	5	9	0	3
	Tail docking method	95	8	0	0

¹Percentages were based on the herds that were actually able to shift one or two categories. For downgrades of 1 category n = 482, for downgrades of 2 categories n = 174. For upgrades of 1 category n = 491.

²As absence of disease contains a very high number of measures with a very small range of shifts, we present only the range here. All separate measures can be found in the Appendix.

For Review Only

Table 3: Median (min – max) decrease and increase in principle and criteria scores when measure scores were replaced with worst and best observed measure scores

Principles / Criteria	Measures	Changes in principles scores		Changes in criteria scores	
		Median decrease in worst scenario	Median increase in best scenario	Median decrease in worst scenario	Median increase in best scenario
Good feeding					
Absence of prolonged hunger	% lean cows	24 (0 – 71)	5 (0 – 69)	67 (0 - 98)	30 (0 - 98)
Absence of prolonged thirst	Water flow	11 (0 – 85)	0 (0 – 85)	29 (0 - 97)	0 (0 - 0)
	Trough length	25 (0 – 85)	0 (0 – 85)	29 (0 - 97)	0 (0 - 97)
	Number of water bowls	0 (0 – 85)	10 (0 – 85)	0 (0 - 97)	12 (0 - 97)
	Drinker cleanliness	12 (0 – 60)	0 (0 – 60)	40 (0 - 68)	0 (0 - 68)
	At least 2 drinkers per animal	0 (0 – 35)	0 (0 – 35)	20 (0 – 97)	0 (0 - 40)
Good housing					
Comfort around resting	Mean time to lie down	6 (0 – 20)	5 (0 – 20)	10 (0 – 32)	8 (0 - 31)
	% cows colliding with housing	0 (0 – 19)	11 (0 – 17)	0 (0 - 32)	18 (0 - 27)
	% cows lying outside of lying area	10 (0 – 20)	0 (0 – 29)	16 (0 – 32)	0 (0 – 46)
	% cows with dirty flanks	0 (0 – 5)	4 (0 – 14)	0 (0 – 12)	6 (0 – 22)
	% cows with dirty lower legs	0 (0 – 9)	4 (0 – 12)	0 (0 – 15)	6 (0 – 18)
	% cows with a dirty udder	0 (0 - 9)	4 (0 – 8)	0 (0 – 15)	6 (0 – 18)
Ease of movement	Loose or tied housing	24 (0 – 37)	0 (0 – 40)	66 (0 – 66)	0 (0 – 85)
Good health					
Absence of injuries	Lameness index	13 (0 – 37)	5 (0 – 35)	27 (3 – 69)	33 (0 – 57)
	Integument alteration index	4 (0 – 24)	5 (0 – 26)	10 (0 – 44)	26 (0 – 42)
Absence of disease	Number of coughs/cow/minute	0 (0 – 0)	0 (0 – 0)	0 (0 – 0)	0 (0 – 0)
	% cows with nasal discharge	1 (0 – 12)	0 (0 – 10)	8 (0 – 35)	0 (0 – 21)
	% cows with ocular discharge	1 (0 – 12)	0 (0 – 8)	8 (0 – 35)	0 (0 – 35)
	% cows with hampered respiration	1 (0 – 5)	0 (0 – 1)	4 (0 – 14)	0 (0 – 14)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	% cows with diarrhoea	2 (0 – 12)	0 (0 – 10)	9 (0 – 35)	0 (0 – 35)
	% cows with vulvar discharge	3 (0 – 12)	0 (0 – 7)	10 (0 – 35)	0 (0 – 24)
	% cows with SCC >400.000	2 (0 – 12)	1 (0 – 12)	8 (0 – 35)	4 (0 – 35)
	% cows mortality	2 (0 – 11)	0 (0 – 12)	8 (0 – 35)	0 (0 – 35)
	% calvings with dystocia	1 (0 – 12)	0 (0 – 13)	7 (0 – 35)	4 (0 – 35)
	% downer cows	2 (0 – 12)	1 (0 – 13)	0 (0 – 35)	3 (0 – 35)
Absence of pain induced by management procedures	Dehorning method (none, surgery)	15 (0 – 35)	6 (0 – 40)	50 (0 – 89)	48 (0 – 98)
	Tail docking method (none, ring)	14 (0 – 34)	0 (0 – 6)	6 (0 – 89)	0 (0 – 0)
Appropriate behaviour					
Expression of social behaviour	Head butts/cow/15 min.	13 (0 – 37)	1 (0 – 16)	69 (0 – 100)	8 (0 – 49)
	Displacements/cow/15 min.	16 (0 – 44)	2 (0 – 30)	69 (0 – 100)	19 (0 – 93)
Expression of other behaviour	Number of hours on pasture	15 (0 – 38)	0 (0 – 34)	64 (1 – 100)	0 (0 – 85)
	Number of days on pasture	15 (0 – 38)	1 (0 – 24)	64 (1 – 100)	15 (0 – 86)
Good human-animal relationship	ADF index	10 (0 – 37)	9 (0 – 37)	31 (0 – 87)	56 (0 – 87)
Positive emotional state	QBA index	20 (0 – 50)	7 (0 – 44)	52 (0 – 93)	40 (0 – 93)

Table 4: Percentages of herds¹ (n = 491) that shifted into a different overall welfare category when individual scores were replaced with observed worst and best criteria or principle scores (observed median, min. and max. score given in column b)

Principles, Criteria	Original observed median, min - max	Observed worst score		Observed best score	
		% farms downgraded 1 category	% farms downgraded 2 categories	% farms upgraded 1 category	% farms upgraded 2 categories
Good feeding	40, 4 – 100	64	100	36	1
Absence of prolonged hunger	70, 3 – 100	59	0	6	0
Absence of prolonged thirst	60, 3 – 100	35	3	30	1
Good housing	54, 6 – 86	37	0	13	0
Comfort around resting	27, 0 – 80	27	0	13	0
Ease of movement	100, 15 – 100	27	0	0	0
Good health	34, 8 – 86	37	0	23	0
Absence of injuries	35, 4 – 100	21	0	8	0
Absence of diseases	40, 12 – 100	4	0	7	0
Absence of pain induced by management procedures	52, 2 – 100	9	0	3	0
Appropriate behaviour	35, 6 – 86	37	0	25	0
Expression of social behaviour	69, 0 – 100	16	0	5	0
Expression of other normal behaviour	64, 0 -100	9	0	8	0
Good human-animal relationship	44, 13 – 100	14	0	8	0
Positive emotional state	53, 0 – 93	24	1	7	0

¹Percentages were based on the herds that were actually able to shift one or two categories. For downgrades of 1 category n = 482, for downgrades of 2 categories n = 174. For upgrades 1 category n = 491, for upgrades of 2 categories n = 317.

Table 5: Percentages of herds¹ (n = 491) that shifted into a different overall welfare category when scores at the **measure, criterion, and principle level**² were replaced with theoretically possible¹ worst and best scores

	Worst score		Best score
	% downgraded 1 category	% downgraded 2 categories	% upgraded 1 category
<i>Measures</i> ¹			
Lameness index ³	10	0	5
Head butts/cow/15 min. ³	16	0	1
ADF index ³	20	0	6
<i>Criteria</i> ¹			
Absence of injuries ⁴	29	1	8
Absence of diseases ⁴	36	1	7
Absence of pain induced by management procedures ⁴	12	0	3
Good human-animal relationship ⁴	23	0	8
<i>Principles</i> ¹			
Good housing ⁴	64	100	13
Good health ⁴	64	100	23
Appropriate behaviour ⁴	64	100	25

¹Percentages were based on the herds that were actually able to shift one or two categories. For downgrades of 1 category n = 482, for downgrades of 2 categories n = 174. For upgrades of 1 category n = 491.

²Scores shown are of those measures, criteria and principles where replacement with theoretical score generated different results than when replaced with observed score.

³ theoretical possible worst score was 100, **theoretical best score was 0**

⁴ theoretical possible worst score was 0, **theoretical best score was 100**

Table 6: Median (min – max) decrease and increase in principle and criterion scores when measures were replaced with worst and best theoretically possible values

Principles, criteria	Measures	Change in principle scores		Change in criteria scores	
		Median decrease in worst scenario	Median increase in best scenario	Median decrease in worst scenario	Median increase in best scenario
Good feeding ¹					
Absence of prolonged hunger	% lean cows ²	25 (2 – 73)	5 (0 – 69)	69 (2 – 100)	30 (0 – 98)
Good health ¹					
Absence of injuries	Lameness index ³	15 (2 – 39)	5 (0 – 35)	27 (3 – 69)	33 (0 – 57)
Absence of disease	Number of coughs/cow/15 min. ²	4 (0 – 12)	0 (0 – 0)	10 (5 – 35)	0 (0 – 0)
	% cows with hampered respiration ²	4 (1 – 12)	0 (0 – 1)	10 (6 – 35)	0 (0 – 14)
Appropriate behaviour ¹					
Good human-animal relationship	ADF index ²	46 (11 – 82)	9 (0 – 37)	44 (13 – 100)	55 (0 – 87)

¹ Scores shown are of those where replacement with theoretical score generated different results than when replaced with observed score

² theoretical possible worst score was 100, theoretical best score was 0

³ theoretical possible worst score was 0, theoretical best score was 100