

Scotland's Rural College

Modelling farmer decision-making to anticipate tradeoffs between provisioning ecosystem services and biodiversity

Guillem, EE; Murray-Rust, D; Robinson, DT; Barnes, AP; Rounsevell, MDA

Published in:
Agricultural Systems

DOI:
[10.1016/j.agry.2015.03.006](https://doi.org/10.1016/j.agry.2015.03.006)

Print publication: 01/01/2015

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (APA):

Guillem, EE., Murray-Rust, D., Robinson, DT., Barnes, AP., & Rounsevell, MDA. (2015). Modelling farmer decision-making to anticipate tradeoffs between provisioning ecosystem services and biodiversity. *Agricultural Systems*, 137, 12 - 23. <https://doi.org/10.1016/j.agry.2015.03.006>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Modelling farmer decision-making to anticipate tradeoffs between provisioning**
2 **ecosystem services and biodiversity**

3
4
5
6 E.E. Guillem^{a,b,*}, D. Murray-Rust^b, D.T. Robinson^c, A. Barnes^a, and M.D.A.
7 Rounsevell^b

8
9
10 ^aLand Economy and Environment Research Group, Scottish Rural College (SRUC),
11 King Buildings, West Mains Road, Edinburgh, EH9 3JG, UK; ^bGeography and the
12 Lived Environment, School of Geosciences, University of Edinburgh, Drummond
13 Street, Edinburgh, EH8 9XP, UK; ^c Department of Geography and Environmental
14 Management, University of Waterloo, 200 University Avenue West, Waterloo,
15 Ontario, N2L 3G1

16
17
18 * Corresponding author. Email: eleonore.guillem@gmail.com

19 Postal address: LPED – UMR 151, Université Aix-Marseille, Case 10, 13331
20 Marseille Cedex 3, France

21
22
23 **Abstract**

24
25 In this paper, an agent-based model of heterogeneous farmer decision-making was
26 coupled with an individual-based model of skylark breeding populations, and applied
27 to a small intensive arable catchment in Scotland. The impacts of farmer decisions on
28 a tradeoff between food and bioenergy production, and skylark numbers, were
29 simulated under the assumptions of three socio-economic scenarios until the year
30 2050. Bioenergy and food production had a significant negative effect on adult and
31 fledgling skylarks. In a business-as-usual context, the production of food and
32 bioenergy increases smoothly, and the number of skylarks is more stable over time
33 than in other scenarios. Food production was higher in an economic liberalisation
34 scenario, due to intensive management and higher yield performance. This explained
35 the low average number of skylarks found at the landscape level in this scenario. The
36 number of skylarks was highest in a sustainability-oriented scenario, but a sharp

37 decrease was observed from 2035 onwards due to the large area planted with
38 bioenergy crops. The different values for economic, environmental and social
39 attributes of farmer decisions played an important role in the land use mosaic, the
40 implementation of ecologically-related actions and on the provision of ecosystem
41 services and biodiversity. Overall, results suggest that a re-assessment of policy
42 targets and design is necessary to maximise environmental management efficiency at
43 the catchment level by taking into account the heterogeneity in farmer objectives and
44 the tradeoffs in ecosystem services provision. The novel approach of coupling an
45 ABM with an IBM is encouraged in further land use related studies.

46

47

48

49 **Keywords:** agent-based model; bioenergy crops; farmer behaviour; food production;
50 land use change; skylark

51

52

53

54

55

56 **1. Introduction**

57 Land use and cover change (LUCC) is a major concern for the sustainability
58 of farming areas, biodiversity levels and the provision of ecosystem services
59 responsible for human welfare. Agricultural landscapes are largely shaped by human
60 actions driven by socio-political and environmental stimuli (Antle et al., 2001;
61 Lambin et al., 2001), and host a number of species that underpin the provision of
62 ecosystem services. These species are under constant threat following changes in
63 farming practices and management styles.

64 Land-related policies have been modified to prevent environmental
65 degradation, but the reforms have created unexpected issues undetected in common
66 *ex-ante* analysis, i.e. land abandonment and intensification of arable land use after the
67 Fischler Reforms in 2005 (Acs et al., 2010; Holland et al., 2011; Doxa et al., 2012). In
68 the near future, the Common Agricultural Policy (CAP) will tend towards
69 liberalisation, which will create increasing reliance on fluctuating commodity prices
70 and a possible switch from food to non-food production (Tranter et al., 2007), and
71 lead to uncertain impacts on the long-term economic and ecological sustainability of
72 farming areas (European Commission, 2010). The anticipation of consequences due
73 to changing conditions (i.e. market, policy, climate) can be improved through the

74 understanding of how actors within the system make decisions and when changes will
75 occur.

76 Indeed, the heterogeneity of land-use activities and management observed at
77 the landscape level has relevance in *ex-ante* analysis, but cannot be explained by
78 common methodologies (i.e. linear programming). In the Agent-Based Modelling
79 (ABM) approach, this landscape heterogeneity is seen from a bottom-up point of view
80 where each actor (i.e. each farmer) is considered to react autonomously and
81 cognitively to external pressures (e.g. Janssen et al., 2000; Berger, 2001; Murray-Rust
82 et al., 2011). In the same way, ecological, individual-based models (IBM) can
83 simulate species population from the behaviour and life cycles of the individuals
84 under different LUCC scenarios (e.g. De Angelis et al., 1998; Topping et al., 2003;
85 McLane et al., 2011).

86 Too often, the impacts of policy on farmer decisions and LUCC (explored via
87 ABMs), and the effect of LUCC on biodiversity and ecosystem services (explored via
88 IBMs) are studied separately. In general, the current ABMs and IBMs lack
89 transparency in some of the component sub-models that drive simulation outcomes.
90 This can be improved by integration, or coupling, of an ABM of LUCC with an IBM,
91 which offers greater potential to understand processes and feedbacks between human
92 and natural systems (Luus et al., 2011) and to study the indirect effect of policy on
93 ecosystem services through farmer decision making (Milner-Gulland, 2012;
94 Sutherland and Freckleton, 2012). Only a few studies have presented results from
95 such a combination (Jepsen et al., 2005; Bithell and Brasington, 2009; Verburg and
96 Overmars, 2009), but the decision maker agents were not heterogeneous, which limits
97 the relevance of such models since not all land managers react similarly to policies
98 (Beilin et al., 2012). Indeed, the nonlinear interactions between farmer decisions and
99 the ecosystem, often acting at different spatio-temporal scales, cannot be considered
100 independently since they involve feedbacks. In particular, these feedbacks occur in
101 respect of a wide variety of ecosystem services and on species by providing or
102 removing habitats (Antle et al., 2001; Liu et al., 2007). For instance, farmland
103 specialist bird species (e.g. skylark, lapwing, yellowhammer), which require specific
104 farmland habitat to nest and to feed, have decreased faster than other types of birds
105 and drastically since the 1970s due to the intensification of agricultural land use
106 (Siriwardena et al., 1998; Donald et al., 2002). Simultaneously, intensive agriculture
107 allows a larger production of food, which is an important ecosystem service.
108 Therefore tradeoffs between several services and with biodiversity levels must be
109 considered.

110 This article reports on the integration of an agent-based model of farmer
111 decision-making with an individual-based model of skylarks applied to a spatial

112 (Geographic Information System (GIS)) database representing a Scottish intensive
113 arable catchment. The model represents relationships between external pressures
114 (market, climate, and policy), heterogeneous farmer decisions about farming
115 practices, and the effects of these on provisioning services (food production,
116 renewable energy), and an indicator of biodiversity (skylark local population). A set
117 of simulation experiments was carried out based on three socio-economic scenarios to
118 test the adaptation and responses of agents to changing contexts and the effects of this
119 on provisioning services and biodiversity.

120

121

122 **2. Materials and Methods**

123 **2.1 Study site**

124 The study area comprises 132 km² of a mostly arable catchment in the Tayside
125 region, East Scotland (Figure 1). 115 active farmers manage the land with a mix of
126 land use activities, essentially cereals and root crops (65%), and grasslands (35%)
127 (Scottish Government, 2007). The study area is one of the few places in Scotland
128 where intensive cropping occurs due to a relatively flat and fertile soil. Intensive
129 cropping takes place on 9% of Scottish agricultural land and generates 34% of
130 agricultural outputs (Scotland's Environment, 2014). Farmers in the catchment share
131 similar biophysical conditions, agricultural activities and market prospects, while
132 avoiding the problem arising from variations observed at larger scales.

133 This site has been intensively studied as it represents an example of a
134 catchment with a number of typical indicators for Scottish farming and shows
135 fragility in terms of water and air quality (Vinten et al., 2009). Since 2003, the
136 catchment has been designated as a Nitrate Vulnerable Zone (NVZ)¹, which puts
137 constraints on how farmers manage their land (Scottish Executive, 2003).

138 The catchment also includes a Site of Special Scientific Interest (SSSI) under
139 the Nature Conservation (Scotland) Act 2004 (Rescobie and Balgavies Lochs), active
140 fisheries, and the Balgavies Scottish Wildlife Trust reserve. In addition, the catchment
141 forms part of the Scottish Environment Protection Agency's Monitored Priority
142 Catchment Project, which aims to establish monitored baselines against which the
143 effectiveness of diffuse pollution mitigation measures can be assessed (Vinten et al.,

¹ The Environment Agency has designated conservation zones, the NVZs, to reduce the risk of nitrate polluted waters (EU Nitrate Directive 91/676/EEC and the EU Water Framework Directive 2000/60/EC). Restrictions include reduction of the amount of fertiliser used and limited fertiliser and animal waste application periods.

144 2009). Thus, the catchment and the broader region are of particular interest to policy
145 makers.

146

147 <FIGURE 1>

148

149 **2.2 Model Development**

150 The integrated ABM/IBM comprises four components (see Figure 2):

151 1) An agent-based model of farmer decision-making for land uses, named “Aporia”²
152 (Robinson et al., 2011; Fontaine et al., 2013; Murray-rust et al., 2014; Guillem et al.,
153 in review);

154 2) An individual-based model of breeding skylarks;

155 3) A vegetation model within which the Aporia model and the skylark IBM are
156 coupled;

157 4) A sub-model that quantifies the provisioning ecosystem services (food and biofuel
158 energy).

159

160 <FIGURE 2>

161

162 **2.2.1 Agent-based model of farmer decision-making for land uses**

163 The model represents heterogeneity in decision-making in terms of farm
164 strategies, i.e. land use regimes per farm. A farmer agent chooses a regime, i.e. crop
165 rotation, for each of the parcels that compose its farm, the management style
166 associated with it (intensive or extensive) and whether an agri-environmental measure
167 or the conversion to bioenergy crops is applied. It is assumed that these choices are
168 based on attitudes and preference structures for the sustainability principles, i.e.
169 economic viability, environmental quality, and social feedback (Murray-Rust et al.,
170 2014; Guillem et al., in review).

171 A sample of farmers within the Lunan catchment was selected for a phone
172 interview and the results used to obtain three attitudinal clusters of respondents:
173 Profit-oriented (38%), Multifunctionalist (25%), and Traditionalist (36%) (Guillem et
174 al., 2012). The proportion of each farm type was randomly allocated and associated
175 with farm parcels within the catchment.

176 In Aporia a set of alternative regimes are evaluated and ranked in order for the
177 farmer agents to select the one that maximises their utility (Murray-Rust et al., 2014).
178 This method computes an economic (difference in gross margins), environmental

²The model framework, and the software and its guidance are available freely at
<http://www.wiki.ed.ac.uk/display/Aporia>

179 (land use cover, nitrogen use and diversity) and social (access to green space and
180 tradition) attributes' score for each regime (Murray-Rust et al., 2014; Guillem et al.,
181 in review). Simultaneously, each farmer type responds to a specific aggregative
182 nonlinear utility function in which the preferences values for these regime attributes
183 was elicited from a choice-based conjoint survey (*ibid*).

184 To anticipate tradeoffs between provisioning ecosystem services and a
185 biodiversity indicator, the change in land use in the Lunan catchment was explored in
186 different socio-economic contexts using three hypothetical scenarios from the
187 Assessing LARge-scale Risks for biodiversity with tested Methods (ALARM) project
188 (Bohunovsky et al., 2011; Settele et al., 2012; Spangenberg et al., 2012): BAMBU
189 (Business-As-Might-Be-Usual) represents the current economic and policy situation
190 with a progressive shift of funds from the CAP pillar 1 (production) to pillar 2
191 (environmental enhancement); GRAS (GRowth Applied Strategy) is characterised by
192 economic liberalism, free trade and international competitiveness - Neither direct
193 payments nor rural development funds are proposed; SEDG (Sustainable European
194 Development Goal) portrays environmental and social development where farmers
195 are encouraged through financial incentives to grow bioenergy crops, to use more
196 extensive management and to apply agri-environmental measures. The scenarios'
197 narratives were adapted to the case study and changing factors were attributed to
198 define market prices, subsidy levels and yield performance over time (initial values
199 taken from SAC (2000 to 2008), and assumptions and forecasted values from
200 Abildtrup et al., 2006)³.

201

202 **2.2.2 Skylark individual-based model**

203 The IBM was designed to estimate the number of skylarks within the Lunan
204 catchment emerging from individual breeding behaviour. Skylark nest suitability and
205 number of brood per year depend mainly on vegetation structure (Chamberlain et al.,
206 1999), which is influenced by crop type.

207 *Behavioural rules* (Figure 3): When entering the breeding period (from April
208 to July), each modelled skylark male “scanned” a territory search space within the
209 virtual GIS-based landscape, and selected a bird territory (i.e. a circular space which
210 is suitable for a nest and a foraging area) until a maximum carrying capacity of the
211 landscape was reached. The territories were suitable for nesting when vegetation
212 height was comprised between 10 to 120 centimetres (Table 1). The maximum
213 capacity was determined by multiplying the area of crops in the search space by their

³ A list of policy instruments and market prices used to define the scenarios is given in the supplementary materials.

214 specific territory density (*Ibid*). Territory densities were upgraded by 20% when a
215 crop was extensively managed or associated with grass margins to represent less
216 dense structure and higher availability of feeding resources for chicks (Henderson et
217 al., 2009). If the number of territories occupied did not exceed the maximum capacity,
218 the male set its nest in a suitable place and attracted a female. Once a male had
219 selected a site, the site remained occupied until the male or its partner dies. In the
220 same manner, if the vegetation structure changed and was no longer suitable, the pair
221 sought another site or became “floaters”, i.e. non-reproductive flock of birds.

222 When a pair established a nest, mating occurred followed by egg laying. The
223 behavioural rules applied to the young stages, i.e. egg, nestling and fledgling, were
224 limited to “Start” and “Die”. In winter, the birds floated randomly in the catchment
225 until a new breeding season started.

226

227 <FIGURE 3>

228 <TABLE 1>

229

230 *Variables* (Table 2): Individual skylarks were characterised by a set of
231 dynamic variables related to their life-cycle stages and recorded daily throughout the
232 simulations: eggs, nestlings, fledglings, adults. Mortality rates are given for each life-
233 cycle stages from empirically-determined means with environmental fluctuations
234 simulated using a daily modifier of 0.1% (adapted from Topping et al., 2005). The
235 number of individual floaters was not initially set but emerged from simulations when
236 some adults were unable to find a nest or a partner (due to the depletion of suitable
237 territory or to the death of a mate).

238

239 <TABLE 2>

240

241 **2.2.3 Vegetation model and coupling of ABM/IBM**

242 A vegetation model (*DefaultVegetationModel*) was used to provide, for each
243 farm parcel, a daily update of vegetation height and a yearly harvestable biomass
244 based on crop types (for yield calculation, see Murray-Rust et al., 2014)⁴. For
245 vegetation height, the *DefaultVegetationModel* uses different equations depending on
246 land use. For crops, a daily growth curve was used based on empirical information
247 collected in the Lunan catchment (own unpublished data; Figure 4). The growth was

⁴Only the harvestable biomass increased across time due to technological improvements in each of the socio-economic scenarios. The height of vegetation is assumed to remain the same.

248 initiated at time of “sowing” and fell to 0 at time of “cutting”. The annual timing of
249 these actions was crop-specific and the same each year. If a parcel was abandoned, a
250 natural succession of shrub vegetation took place, for which the height of vegetation
251 H (in centimetres) at time t (in day) was modelled using the Chapman-Richards
252 equation (Equation 1).

253

$$254 \quad H(t) = A + \frac{(K - A)}{(1 + Q \cdot e^{-B(t-M)})^{1/\nu}} \quad (1)$$

255
256 , with A and K respectively the lower and upper asymptote ($A=0$, $K=150\text{cm}$), B is the
257 growth rate ($B=0.02 \text{ cm.day}^{-1}$), ν is the nearest line between lower and upper
258 asymptote ($\nu=0.5$), Q depends on the value at $H(0)$ and M is the time of maximum
259 growth when $Q=\nu$.

260

261 <FIGURE 4>

262

263 The vegetation model was the connecting interface by which the ABM of
264 farmer decision-making is coupled with the skylark IBM. Indeed, the spatial
265 resolution of both models was the parcel level, delimited by boundaries and attached
266 to a given farmer identity. The environmental factors involved in the skylark IBM
267 (i.e. vegetation heights and territory density) are therefore directly driven by farmers’
268 choices of land use managements and regimes. However, the ABM and IBM are only
269 loosely coupled since the time-step of a changing state was asynchronous (Antle et
270 al., 2001; Bithell and Brasington, 2009): farmer attributes and decisions, and crop
271 yields, were updated annually while skylark behaviour, life-cycle characteristics and
272 vegetation heights were simulated daily.

273

274 **2.2.4 Food and bioenergy production**

275 The harvesting of food for human consumption (i.e. vegetables, potatoes,
276 cereals, beef⁵) and bioenergy crops (i.e. willow and miscanthus) was converted at
277 each annual time-step into energy produced from the whole catchment. This was done
278 by multiplying the amount of commodity harvested (in tonnes) by the energy value
279 for human consumption and renewable energy using FAO and USDA conversion

⁵We assumed that grassland biomass is used to rear beef cattle, and thus the biomass of grass was converted into tons of beef (see supplementary material for details).

280 coefficients (Table 3). The simulation outputs gave a cumulative sum of energy
281 produced in the catchment.

282

283 <TABLE 3>

284

285 **2.3 Initialisation and analysis of simulation results**

286 The model was initialised with the historical spatial arrangement of land use
287 from 2000 to 2008 using Spatially Integrated Administration and Control System
288 (SIACS) data and run for a period of 50 years. The initial population of skylarks was
289 estimated from the carrying capacity of the 2007 historical landscape.

290 Because the model included a stochastic component (i.e. mortality rates of
291 individual skylarks), multiple simulations were performed; 10 simulations for each
292 scenario, applied to four cases of farmer agent populations: ALL, a proportion of
293 farmer types corresponding to the results of the social survey; Multifunctional, a
294 population exclusively composed of multifunctionalist farmers; Profit, a population of
295 profit-oriented farmers; and Traditional, a population of traditionalist farmers.

296 For the ALL simulations, a time series (2008 to 2050) of the proportion of
297 land use types found in the Lunan catchment is given for each scenario. In addition, a
298 time series of the cumulative sum of energy produced, averaged over the 10 multiple
299 simulation runs, and of the average number of adult and fledgling skylarks, were
300 compared across each scenario.

301 The geometric means over 10 simulations from the year 2008 onwards of
302 adult skylarks was used to compare skylark populations in a landscape managed
303 exclusively by a single farmer agent type. Kruskal-Wallis tests were carried out on the
304 null hypothesis that skylark numbers were statistically similar across farmer types.

305 Finally, model outcomes were analysed to test the relationships between the
306 production of food as well as bioenergy (in constant energy units, megajoules (MJ))
307 against the adult and fledgling population of skylarks, using a linear mixed model to
308 account for temporal autocorrelation, i.e. 30 points, related to the 10 simulations for
309 three scenarios, were clustered per year, giving 42 groups (i.e. the 42 groups were the
310 42 years of simulations) for 1260 observations. The model was computed in R using
311 the “nlme” package (Pinheiro et al., 2009). The linear mixed model had the following
312 form (Laird and Ware, 1982):

313

$$314 \mathbf{A}_{i,j} = \beta_1 \cdot \mathbf{x}_{1,i,j} + \dots + \beta_n \cdot \mathbf{x}_{n,i,j} + \mathbf{t}_{i,1} \cdot \mathbf{z}_{1,i,j} + \dots + \mathbf{t}_{i,p} \cdot \mathbf{z}_{p,i,j} + \boldsymbol{\varepsilon}_{i,j} \quad (2)$$

315

$$316 \mathbf{t}_{i,k} \sim N\left(0, \boldsymbol{\psi}_k^2\right), \text{Cov}\left(\mathbf{t}_k, \mathbf{t}_{k'}\right) = \boldsymbol{\psi}_{k,k'} \quad (3)$$

317

$$318 \quad \varepsilon_{i,j} \sim N(0, \theta^2 \cdot \lambda_{i,j}), \text{Cov}(\varepsilon_{i,j}, \varepsilon_{i,j'}) = \theta^2 \cdot \lambda_{i,j,j'} \quad (4)$$

319

320

321 where $A_{i,j}$ is the resulting number of skylarks for observation j ($j = 30$) of cluster i (i

322 $= 42$), $\beta_1 \dots \beta_n$ are the fixed effect coefficients constant across clusters,

323 $\alpha_{1,i,j} \dots \alpha_{n,i,j}$ are the fixed effect regression coefficients, $t_{i,1} \dots t_{i,p}$ are the random

324 effect of time coefficients of cluster i , $z_{1,i,j} \dots z_{p,i,j}$ are the random effects regression

325 coefficients, $\varepsilon_{i,j}$ is the error term, $\psi_{k,k'}$ are the covariances among the random

326 effects and are constant across clusters, $\theta^2 \cdot \lambda_{i,j,j'}$ are the covariances between errors

327 in cluster i .

328

329

330 3. Results

331 3.1 Temporal effects of socio-economic scenarios on farmers' decision, 332 provisioning services and skylark number

333 In BAMBU, the proportion of crop types changes noticeably at each decade
334 (Figure 5), with an increase in root crops due to higher yielding performance, loss of
335 set-aside and grassland⁶. The level of cereals is higher than in the other scenarios and
336 the area planted with miscanthus remains low. The population of adult skylarks
337 increases until a plateau is reached between 2020 and 2040, followed by a small
338 decrease afterwards (Figure 6a). In this scenario the energy produced from
339 miscanthus is the lowest, and does not exceed 10 terajoules (TJ), while energy from
340 food is intermediate compared with other scenarios (Figure 6c and d).

341 In GRAS, the area grown under cereals is cut by 35% by 2050 compared to
342 2030's levels, which is replaced with root and bioenergy crops (Figure 5). Yield
343 improvement and the resulting response from low input and output prices in GRAS
344 allow more land to be converted to bioenergy crops without diminishing food
345 production. Indeed, the production of food energy is the highest compared to the other
346 scenarios, while the adult skylark population is the lowest (until around 2040).

347 In SEDG, the land cultivated for bioenergy crops rise from 2040 (Figure 5),
348 leading to the highest production of bioenergy across the scenarios, which accounts
349 for more than 50 TJ in 2050, and the lowest production of food (Figure 6c and d). The

⁶GIS-based maps showing the simulated distribution of land-uses in the study area in two time slices, 2025 and 2050, under the assumptions of three scenarios GRAS, BAMBU, SEDG, are provided as a supplementary material.

350 number of adult skylarks reaches a maximum level in SEDG around 2030 while the
351 most abrupt decrease is observed afterwards (Figure 6a). The decrease in adult and
352 fledgling skylarks is initiated before the amount of bioenergy produced goes beyond
353 10 TJ and is very abrupt, as opposed to the GRAS scenario where the decrease starts
354 later and is smoother (Figure 6a and b).

355 Figure 6b shows that the number of skylark fledglings produced diminishes in
356 all scenarios over the whole period. A small increase is observed from 2020 in GRAS
357 and SEDG when direct payments start to be reduced (drastically in GRAS and more
358 progressively in SEDG). The only difference found in 2020 between GRAS, SEDG
359 and BAMBU, is a greater diversity of crop types in GRAS and SEDG, i.e. presence of
360 leguminous crops and miscanthus (Figure 5).

361

362 <FIGURE 5>

363

364 <FIGURE 6>

365

366 **3.2 Effects of farmer behaviour on skylarks' number**

367 The mean density of skylark territories over the period 2008-2050 was 0.13
368 per hectare and there were no significant differences between scenarios. However,
369 Kruskal-Wallis tests were performed to test the distribution of adult skylarks across
370 different landscapes virtually managed by each farmer type separately. The average
371 number of skylarks over the period 2008-2050 was significantly different across the
372 three types of landscapes (BAMBU: $p=0.007$, GRAS: $p=0.000$, SEDG: $p=0.002$)
373 (Figure 7).

374 In a landscape managed exclusively by traditionalist farmers, the number of
375 adult skylarks remains the same in the three scenarios, while there are some variations
376 in the case of profit-oriented and multifunctionalist farmers. For profit-oriented
377 farmers, the average number of skylarks is the highest in BAMBU, but the lowest in
378 GRAS. For multifunctionalist farmers, the abundance is similar to the traditionalists
379 in BAMBU and GRAS, but decreases in SEDG.

380 Multifunctionalist farmers generally apply environmentally-friendly practices,
381 i.e. grass margins and spring cereals, but they also adopt newer land use such as
382 bioenergy crops (Guillem et al., in review). This could explain the low abundance
383 found in the SEDG scenario after 2030, in which subsidies allow bioenergy crops to
384 be viable. The profit-oriented farmers grow cereals in BAMBU, but they manage their
385 land more intensively and the crop mosaic is less diverse in GRAS. This type of
386 farmer was the most proficient in adapting to rapidly changing market conditions to
387 maximise profit. Traditionalist farmers maintained intensive regimes in all scenarios,

388 but they use longer and more diverse crop rotations (Guillem et al., in review). In
389 addition this type of farmer was the least likely to apply bioenergy crops. The average
390 number of skylarks in a landscape managed by all types of farmers was very similar
391 to those for the profit-oriented types for both BAMBU and GRAS.

392
393
394

<FIGURE 7>

395 **3.3 Tradeoffs between food production, bioenergy and skylark number**

396 The linear mixed model shows that both bioenergy and food production
397 have a negative fixed effect on the number of skylarks and fledglings when
398 considering potential variation due to time (random effect) (Figure 8). The fixed
399 effect of the explanatory variables, food and bioenergy production, is the average
400 effect over all years of the simulation. The fixed effect of biofuel production against
401 adult and fledgling numbers is significant (respectively, t (Df=1246) = -3.785,
402 $p < 0.001$ and t (Df=1246) = -6.783, $p < 0.001$), with a negative effect occurring when
403 the production exceeds approximately 10 terajoules. Similarly, the linear relationship
404 between food production and adult and fledgling skylark numbers is also significant (t
405 (Df=1246) = -4.053, $p < 0.001$ and t (Df=1246) = -3.868, $p < 0.001$), though the fitted
406 regression line is less abrupt than for bioenergy.

407
408
409
410

<FIGURE 8>

411 **4. Discussion**

412 **4.1 Impacts of socio-economic contexts on farmer behaviour and skylark number**

413 In all scenarios, an increase in skylark numbers is observed at least until 2030.
414 This is explained by the choices most farmers make to increase the cultivation of
415 cereals compared with the area planted in the baseline year 2008. Cereal crops have
416 been defined as “the single most important habitat for skylarks in the UK in terms of
417 the overall number of breeding pairs they support” (Donald and Vickery, 2000). In
418 BAMBU, land uses are not changing as much as in GRAS and SEDG, and therefore
419 the population of adult skylarks is relatively stable. Without subsidies, as is the case
420 in GRAS, land uses change according to commodity price fluctuations, and the land is
421 managed intensively. This has a negative effect on skylark numbers since, on average,
422 these numbers are the lowest compared with the other scenarios. Economic
423 liberalisation therefore brings uncertainty for the viability of farmland bird
424 populations since impacts are dependent on market forces rather than on policy

425 intervention. In SEDG, extensive regimes and grass margins, which are beneficial to
426 skylarks, are encouraged by substantial environmental payments and one would
427 expect an increase in the population of skylarks. However, while the number of
428 skylarks is the highest until 2035 compared with the other scenarios, a sharp decrease
429 was observed afterwards that can be explained by the large expansion of bioenergy
430 cropping occurring in this scenario. Other simulation studies based on Lucc
431 scenarios have shown the negative impact of bioenergy crops on wildlife at different
432 spatial levels (Eggers et al., 2009; Gevers et al., 2011). In the latter study, an
433 individual based model of skylark was used and the effect of land use scenarios was
434 analysed. Gevers et al. (2011) found that skylark numbers were affected by the loss of
435 crop heterogeneity when more than 13% of the land was replaced with maize, but it
436 was also largely explained by the loss of set-aside replaced with these crops. In this
437 study, static land use scenarios were used that did not simulate explicitly any possible
438 lag effect that might occur in real world situations (Liu et al., 2007). We found that
439 the negative effect of bioenergy production on skylark abundance occurred at
440 different times in SEDG and GRAS. Two conclusions can be drawn from this
441 observation. First, since the same area grown with miscanthus produces less energy in
442 SEDG than in GRAS, due to the difference in yield performance, the amount of
443 bioenergy becomes a poor indicator for assessing the impact on skylarks under a
444 given renewable energy target as opposed to an area. Second, the low production of
445 food energy in SEDG could also increase risks for the skylark population, despite the
446 negative relationship described in Section 3.3. This indicates that a possible minimum
447 threshold of food production as well as a maximum proportion of land converted to
448 bioenergy crops are required to sustain skylark populations.

449 The overall decrease in fledgling numbers could be an effect of the population
450 equilibrium state; e.g. when the number of adults increases, less fledglings are
451 produced. However, from 2040 onwards both the number of adults and fledglings
452 decreases. Likewise, it has been found that as the territory density of the overall
453 landscape increases, with a large area being planted with cereals, the size of territory
454 shrinks resulting in lower reproductive success (Both and Visser, 2003). This trend
455 implies the presence of an ecological trap, which often leads to population extinction
456 (Battin, 2004), possibly explaining why the number of skylarks decreases after 2040
457 in all scenarios. However, in this model, the environment has closed boundaries,
458 which does not allow the population to diffuse to surrounding landscapes. This leads
459 to individual skylarks using the landscape to its maximum carrying capacity,
460 establishing nests in sub-optimal conditions (e.g. use of habitat with minimum and
461 maximum vegetation height). Secondly, food availability to skylark was not explicitly
462 modelled and this could have resulted in an overestimation of the number of skylarks,

463 especially in the economic liberalisation scenario, where intensive management
464 reduce significantly the presence of invertebrates for young skylarks (Topping et al.,
465 2005).

466

467 **4.2 Importance of farmer heterogeneous decision-making on ecosystem services** 468 **and biodiversity delivery**

469 The crop mosaic, intensity pressures and provision of ecosystem services in a
470 landscape arise from the decisions of individual farmers. The proportion of farmer
471 behavioural types in the Lunan catchment had an effect on the provision of food and
472 bioenergy, and on skylark abundance. There was however a dominant effect of the
473 way profit-oriented farmers manage their farms in both BAMBU and GRAS,
474 neutralising the positive environmental outcomes expected from other farmer types.
475 The profit-oriented farmers are the most represented in the population of farmers
476 (38%) and they favour the economic viability of the business over the enhancement of
477 habitats for farmland birds (Guillem et al., 2012; Guillem and Barnes, 2013). In
478 SEDG, the aggregate effect of heterogeneous farmer decision-making leads to higher
479 skylark abundance than would be expected in simulations with exclusive farm types.
480 This is possibly a result of the combination of high uptake of agri-environmental
481 measures and extensive regimes up to 2025, and of a variety of farming objectives,
482 which have a cumulative beneficial effect on skylarks; as opposed to BAMBU and
483 GRAS where production and intensification dominate. In Guillem et al. (in review),
484 the consequences of the SEDG scenario on LUCC and management styles were
485 greatly influenced by farmers' environmental and social values. Therefore, farmer
486 (positive) values for the environment, when they are encouraged appropriately, are
487 important to ensure skylark abundance and probably other ecologically-related
488 aspects of the landscape.

489 Nevertheless, a positive attitude towards birds and socio-environmental
490 objectives do not always benefit skylarks. For instance, bioenergy crops, which
491 scored the highest for the environmental attribute in the model (i.e. do not require
492 large amounts of nitrogen and provide a winter cover against soil erosion (see
493 Guillem et al., in review)), were applied by the multifunctionalist farmers to a large
494 area because they wish to maximise environmental benefits over the farm, but had a
495 deleterious effect on skylarks. This highlights the importance of appropriate
496 information on the ecological risks associated with bioenergy cropping, which are
497 advertised as environmentally-friendly.

498

499 **4.3 Negative effect of food and bioenergy production on skylarks**

500 The study revealed a negative effect of bioenergy and food production on
501 adult and fledgling skylarks. In mid-May, during the middle of the breeding period,
502 the height of miscanthus is no longer suitable and the birds have to seek other
503 territories (see Figure 4). It is possible that, at this period, most of the adjacent fields
504 are already occupied leading these birds to become non-reproductive floaters. This
505 was verified by the more severe decrease in fledgling numbers when the production of
506 bioenergy increases, meaning that the breeding period is shortened and less breeding
507 attempts will occur. However, previous field studies related to bird and bioenergy
508 crops showed that miscanthus supports a higher density of breeding skylarks than
509 other arable crops, but at an early stage of crop establishment when the vegetation
510 does not exceed a maximum threshold (Semere and Slater, 2007; Bellamy et al.,
511 2009; Sage et al., 2010). The high skylark density found in the literature was
512 explained by a significant proportion of bare ground and the presence of weeds on
513 which adults feed. Hence, if bioenergy cropping becomes increasingly viable, there is
514 a risk that improved technology aiming at maximising yields will lead to the loss of
515 these benefits. Since high density of skylarks only occurs at the beginning of the
516 breeding season in miscanthus, it is also evident that a certain degree of crop diversity
517 should be maintained for the birds to continue breeding in adjacent fields (Chaney et
518 al., 1997).

519 The provision of food is also shown to have a negative impact on skylarks. In
520 contrast to bioenergy, this relationship is not a function of the area planted with food
521 crops. A large area planted with food crops is in fact advantageous for skylarks, but
522 the intensity at which these crops are managed has more impacts. Donald et al. (2002)
523 found a negative relationship between yield improvement and population trends of
524 farmland bird. This is difficult to measure in ecology-based studies since food crops
525 are very diverse and offer a variety of habitats. Nevertheless, it is particularly relevant
526 to test the effect of policy targets, in particular food security, by quantifying both the
527 level of food and energy required at the European and regional levels, and the
528 variations this induces in the abundance of birds. With further intensification and an
529 increase in yield performance due to technology and climate change, the risk
530 increases for the viability of skylark populations.

531

532 **4.4 Reflection on the approach**

533 The coupled ABM/IBM allowed the study of provisioning ecosystem services
534 and of skylark numbers at landscape level that emerge from farmers' individual
535 valuation of sustainability. This means that qualitative and quantitative case-specific

536 information on various “agents” or “individuals” that act at different spatio-temporal
537 and organisational scales can be linked within a single dynamic process. Hence, the
538 ability of an ABM to simulate LUCC is extended to new functionalities such as the
539 simulation of changes in ecosystem services and biodiversity levels. This is of great
540 importance to, on one hand, quantify dynamically the human decisions’ outcomes
541 (provision of ecosystem services and biodiversity), and thus anticipate the impacts of
542 changing and uncertain circumstances. On the other hand, tradeoffs between different
543 ecosystem services and biodiversity levels can be assessed, which will allow efficient
544 policy making (An et al., 2014).

545 This approach simulates empirically the so-called “Coupled Human-Nature
546 Systems” (CHANS) with its complexity (An et al., 2014), i.e. heterogeneity (of
547 farmer behaviour), emergence (from individual farmers and skylarks), non-linearity
548 (e.g. utility function) and feedbacks (e.g. farmers’ adaptation and learning from the
549 impacts of their practices on biodiversity, see Figure 2). In the coupled model
550 presented here, the feedback processes are not yet implemented but are of high
551 interest to policy makers, especially for the development of instruments such as
552 payment-by-results agri-environmental supports, and adaptive co-management
553 (Goldman et al., 2007; Schwarz et al., 2008; Polasky et al., 2011). Indeed, in this
554 version of the model, farmers chose regimes as a function of their economic,
555 environmental and social values that are computed using a simple scoring system (see
556 Guillem et al., in review). However, the scores are static over time and do not
557 consider bi-directional feedbacks (see Figure 2) that could emerge from the skylark
558 IBM and impel farmer agents to re-consider their choices. For example, the uptake
559 and outcomes of per-clutch payments (Verhulst et al., 2007) or sward height measures
560 (SNH, 2005) could be explored, but would necessitate the estimation of the utility of
561 an attribute of decisions specific to bird impacts.

562 The model presented here has some limitations in terms of predictability and
563 concept. If the model were fully predictive, the ABM/IBM coupling could be used to
564 answer specific questions about assigning proportions and combinations of land uses
565 to enhance ecosystem services and biodiversity. The issue with coupling the ABM of
566 farmer decision-making with the IBM of skylarks was the spatial scale. Farmers are
567 indeed easily contained within a virtual catchment as they interact essentially within
568 their household and farm parcels. For skylarks this is unlikely, i.e. there is a spatial
569 diffusion to areas outside the case study, and assumptions must be made at this point.

570 Some other aspects in the ABM must be improved (see Murray-Rust et al.,
571 2014). The difference in crop height should be related with the improvement of
572 technology stipulated in the socio-economic scenarios. In the same manner, a gross to

573 net factor has to be applied for the calculation of gross margins. Indeed, we can
574 expect a difference in tax level across time and scenarios.

575 The aggregate, or emergent, effect of heterogeneous farmer decisions was
576 assessed on a small number of ecosystem services, essentially the provisioning (food
577 and bioenergy) and on a unique indicator of biodiversity (skylarks). We have
578 demonstrated the negative relationship between bioenergy and food production on
579 skylark number, but one can ask what would be the impacts on other ecosystem
580 services or biodiversity indicators. For instance, while cereal cropping maximises the
581 production of food and the availability of nesting habitat for skylark, it does not
582 induce a high level of carbon storage (compared with grassland) or the accessibility to
583 recreational assets (see additional examples in Bennett et al., 2009; Power, 2010;
584 Setälä et al., 2014). The Aporia framework implements additional ecosystem services
585 assessors such as landscape aesthetics, carbon storage and nitrogen cycle (Murray-
586 Rust et al., 2014), but these have not been applied to the Lunan catchment yet.

587 In parallel and adversely to the requirements for increased level of complexity
588 enumerated above, generalisation could also be addressed in future development. The
589 tradeoffs between ecosystem services are global issues (e.g. the necessity to provide
590 food to developing countries and escalating population while maintaining a
591 sustainable environmental level) and policies are usually designed at large scale
592 (regional, national, continental). This alternative approach to the model development
593 will however imply the loss of details in data and require modification in model
594 concept.

595

596 **5. Conclusion**

597 Through the coupling of an ABM of farmer decision-making with an IBM of
598 skylarks, we have shown that the viability of the local population of skylarks and the
599 provision of food and bioenergy are intrinsically related to the landscape level
600 arrangement of crop types and management styles. Simultaneously, it is individual
601 farmers with differing values for the sustainability principles that decide on crop types
602 and management styles. Economic liberalisation is not a good option for sustaining
603 farmland birds since it encourages most farmers to produce intensively in accordance
604 with market signals and to abandon agri-environmental measures. Farmers who have
605 environmental objectives play an important role in the preservation of farmland birds,
606 but this requires substantial reward, especially if other policy goals have to be met
607 (food security and bioenergy target). For that reason, single ecosystem services should
608 not be assessed and targeted in isolation, and careful information should be passed to
609 farmers on the possible tradeoffs that exist between services and biodiversity
610 indicators. The formulation of policies should strategically take account of tradeoffs

611 between ecosystem service and biodiversity indicators, as proposed by Haughton et
612 al. (2009) and by the European Environmental Agency (EEA, 2007), but in a dynamic
613 manner, and should, we argue, also include farmer heterogeneity in decision-making.
614 This could be achieved through collaborative plans at the scale of several farm units.
615 Each decision maker within this spatial scale would have different functions
616 depending on their interests, skills and other objectives. An alternative implies the
617 collaboration of farmers with similar goals to achieve targets that are realizable at
618 larger scales than the farm and in a complementary manner (Pelosi et al., 2010).
619 The novel approach presented here has proven effective in the advancement of
620 simulation models of land use dynamics and policy-making. Improvements of this
621 method as well as applications to other case studies are worthwhile for further
622 research.

623

624

625

626 **Acknowledgements**

627 This research was funded by the Rural and Environment Research and Analysis
628 Directorate (RERAD) of the Scottish Government. The authors would also like to
629 thank the RSPB Scotland, in particular Professor Jeremy Wilson for his help on
630 skylark ecology.

631

632 **References**

633

634 Acs, S., Hanley, N., Dallimer, M., Gaston, K.J., Robertson, P., Wilson, P., and
635 Armsworth, P.R., 2010. The effect of decoupling on marginal agricultural
636 systems: Implications for farm incomes, land use and upland ecology. *Land*
637 *Use Policy* 27, 550-563.

638 An, L., Zvoleff, A., Liu, J., and Axinn, W., 2014. Agent-Based Modeling in Coupled
639 Human and Natural Systems (CHANS): Lessons from a Comparative
640 Analysis. *Annals of the Association of American Geographers* 104(4), 723-
641 745.

642 Antle, J.M., Capalbo, S.M., Elliott, E.T., Hunt, H.W., Mooney, S., and Paustian, K.H.,
643 2001. Research needs for understanding and predicting the behavior of
644 managed ecosystems: Lessons from the study of agroecosystems. *Ecosystems*
645 4, 723-735.

646 Battin, J., 2004. When good animals love bad habitats: Ecological traps and the
647 conservation of animal populations. *Conservation Biology* 18, 1482-1491.

648 Beilin, R., Sysak, T., and Hill, S., 2012. Farmers and perverse outcomes: The quest
649 for food and energy security, emissions reduction and climate adaptation.
650 *Global Environmental Change* 22(2), 463-471.

651 Bellamy, P.E., Croxton, P.J., Heard, M.S., Hinsley, S.A., Hulmes, L., Hulmes, S.,
652 Nuttall, P., Pywell, R.F., and Rothery, P., 2009. The impact of growing

653 miscanthus for biomass on farmland bird populations. *Biomass & Bioenergy*
654 33, 191-199.

655 Bennett, E.M., Peterson, G.D., and Gordon, L.J., 2009. Understanding relationships
656 among multiple ecosystem services. *Ecology Letters* 12, 1394-1404.

657 Berger, T., 2001. Agent-based spatial models applied to agriculture: a simulation tool
658 for technology diffusion, resource use changes and policy analysis.
659 *Agricultural Economics* 25, 245-260.

660 Bithell, M. and Brasington, J., 2009. Coupling agent-based models of subsistence
661 farming with individual-based forest models and dynamic models of water
662 distribution. *Environmental Modelling & Software* 24, 173-190.

663 Bohunovsky, L., Jager, J., and Omann, I., 2011. Participatory scenario development
664 for integrated sustainability assessment. *Regional Environmental Change* 11,
665 271-284.

666 Bondeau, A., Smith, P.C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten,
667 D., Lotze-Campen, H., Muller, C., Reichstein, M., and Smith, B., 2007.
668 Modelling the role of agriculture for the 20th century global terrestrial carbon
669 balance. *Global Change Biology* 13, 679-706.

670 Both, C. and Visser, M.E., 2003. Density dependence, territoriality, and divisibility of
671 resources: From optimality models to population processes. *American*
672 *Naturalist* 161, 326-336.

673 Browne, S., Vickery, J., and Chamberlain, D., 2000. Densities and population
674 estimates of breeding Skylarks *Alauda arvensis* in Britain in 1997. *Bird Study*
675 47, 52-65.

676 Chamberlain, D.E., and Crick, H.Q.P., 1999. Population declines and reproductive
677 performance of skylarks *Alauda arvensis* in different regions and habitats of
678 Great Britain. *Ibis* 141, 38-51.

679 Chamberlain, D.E., Wilson, A.M., Browne, S.J., and Vickery, J.A., 1999. Effects of
680 habitat type and management on the abundance of skylarks in the breeding
681 season. *Journal of Applied Ecology* 36, 856-870.

682 Chaney, K., Evans, S.A., and Wilcox, A., 1997. Effect of cropping practice on skylark
683 distribution and abundance. Brighton crop protection conference: weeds.
684 Proceedings of an international conference, Brighton, UK, 17-20 November
685 1997, 1173-1178.

686 DeAngelis, D.L., Gross, L.J., Huston, M.A., Wolff, W.F., Fleming, D.M., Comiskey,
687 E.J., and Sylvester, S.M., 1998. Landscape modeling for everglades ecosystem
688 restoration. *Ecosystems* 1, 64-75.

689 Delius, J.D., 1965. A population study of skylarks *Alauda arvensis*. *Ibis* 107, 466-
690 491.

691 Donald, P.F., and Vickery, J.A., 2000. The importance of cereal fields for breeding
692 and wintering skylarks *Alauda arvensis* in the UK. In Aebischer, N.J., Grice,
693 P.V., Evans, A.D. & Vickery, J.A. (eds) *Ecology and Conservation of*
694 *Farmland Birds*, British Ornithologists' Union, 140-150.

695 Donald, P.F., Evans, A.D., Muirhead, L.B., Buckingham, D.L., Kirby, W.B., Schmitt,
696 S.I.A. 2002. Survival rates, causes of failure and productivity of Skylark
697 *Alaudaarvensis* nests on lowland farmland. *Ibis* 144, 652-664.

698 Dougall, T.W., 1997. Biometrics and sex ratios of skylarks *Alauda arvensis* in winter
699 in south-east Scotland. *Ringing & Migration* 18, 37-49.

700 Doxa, A., Paracchini, M.L., Pointereau, P., Devictor, V., and Jiguet, F., 2012.
701 Preventing biotic homogenization of farmland bird communities: The role of

702 High Nature Value farmland. *Agriculture, Ecosystems and Environment* 148,
703 83-88.

704 EEA. 2007. Estimating the environmentally compatible bioenergy potential from
705 agriculture. EEA Technical Report Number 12/2007, Copenhagen, Denmark.

706 Eggers, J., Troltsch, K., Falcucci, A., Maiorano, L., Verburg, P.H., Framstad, E.,
707 Louette, G., Maes, D., Nagy, S., Ozinga, W.A., and Delbaere, B., 2009. Is
708 biofuel policy harming biodiversity in Europe? *Global Change Biology*
709 *Bioenergy* 1, 18-34.

710 European Commission. 2010. The CAP towards 2020: Meeting the food, natural
711 resources and territorial challenges of the future. European Commission
712 Communication, [http://ec.europa.eu/agriculture/cap-post](http://ec.europa.eu/agriculture/cap-post-2013/communication/index_en.htm)
713 [2013/communication/index_en.htm](http://ec.europa.eu/agriculture/cap-post-2013/communication/index_en.htm)

714 Field, R.H., Kirby, W.B., and Bradbury, R.B., 2007. Conservation tillage encourages
715 early breeding by Skylarks *Alauda arvensis*: Capsule Increased crop residue
716 on the soil surface, left as a result of conservation tillage, encouraged earlier
717 nesting in winter wheat fields. *Bird Study* 54(1), 137-141.

718 Fontaine, C.M., Dendoncker, N., De Vreese, R., Mortelmans, D., Jacquemin, I.,
719 François, L., Marek, A., Devillet, G., Van Herzele, A., Mortelmans' D., and François,
720 L., 2013. Towards participatory integrated valuation and modelling of ecosystem
721 services under land-use change. *Journal of Land Use Science* 9(3), 278-303.

722 Gevers, J., Hoye, T.T., Topping, C.J., Glemnitz, M., and Schroder, B.,
723 2011. Biodiversity and the mitigation of climate change through bioenergy:
724 impacts of increased maize cultivation on farmland wildlife. *Global Change*
725 *Biology Bioenergy* 3, 472-482.

726 Goldman, R.L., Thompson, B.H., and Daily, G.C., 2007. Institutional incentives for
727 managing the landscape: Inducing cooperation for the production of
728 ecosystem services. *Ecological Economics* 64, 333-343.

729 Guillem, E.E., Murray-Rust, D., Robinson, D.T., Barnes, A., and Rounsevell, M.D.A.
730 2011. The effects of farmers perceptions and objectives on land use change
731 and ecosystem services: The case of a Scottish arable catchment. *Proceedings*
732 *of the 18th annual UK conference of the International Association for*
733 *Landscape Ecology (IALE-UK), Wolverhampton University, UK, 6-8*
734 *September 2011.*

735 Guillem, E.E., Barnes, A.P., Rounsevell, M.D.A., and Renwick, A., 2012. Refining
736 perception-based farmer typologies with the analysis of past census data.
737 *Journal of Environmental Management* 110, 226-235.

738 Guillem, E.E., and Barnes, A.P., 2013. Farmer perceptions of bird conservation and
739 intended farming management at a catchment level. *Land Use Policy* 31, 565-
740 575.

741 Guillem, E.E., Murray-Rust, D., Robinson, D.T., Barnes, A., Acosta-Michlik, L.,
742 Karali, E., and Rounsevell, M.D.A. Heterogeneous farm strategies and
743 sustainable trade-offs: an empirically-grounded agent-based model.
744 *Agriculture, Ecosystems & Environment*, in review.

745 Haughton, A.J., Bond, A.J., Lovett, A.A., Dockerty, T., Sunnenberg, G., Clark, S.J.,
746 Bohan, D.A., Sage, R.B., Mallott, M.D., Mallott, V.E., Cunningham, M.D.,
747 Riche, A.B., Shield, I.F., Finch, J.W., Turner, M.M., and Karp, A., 2009. A
748 novel, integrated approach to assessing social, economic and environmental
749 implications of changing rural land-use: a case study of perennial biomass
750 crops. *Journal of Applied Ecology* 46, 315-322.

- 751 Henderson, I.G., Ravenscroft, N., Smith, G., and Holloway, S., 2009. Effects of crop
752 diversification and low pesticide inputs on bird populations on arable land.
753 *Agriculture Ecosystems & Environment* 129, 149-156.
- 754 Holland, J. P., Morgan-Davies, C., Waterhouse, T., Thomson, S., Midgley, A., and
755 Barnes, A., 2011. An Analysis of the Impact on the Natural Heritage of the
756 Decline in Hill Farming in Scotland. Scottish Natural Heritage Commissioned
757 Report No. 454.
- 758 Janssen, M.A., Walker, B.H., Langridge, J., and Abel, N., 2000. An adaptive agent
759 model for analysing co-evolution of management and policies in a complex
760 rangeland system. *Ecological Modelling* 131, 249-268.
- 761 Jepsen, J.U., Topping, C.J., Odderskaer, P., and Andersen, P.N., 2005. Evaluating
762 consequences of land-use strategies on wildlife populations using multiple-
763 species predictive scenarios. *Agriculture, Ecosystems & Environment* 105,
764 581-594.
- 765 Laird, N.M. and Ware, J.H., 1982. Random-Effects Models for Longitudinal
766 Data. *Biometrics* 38, 963-974.
- 767 Lambin, E.F., Turner, B.L., Geist, H.J., Agbola, S.B., Angelsen, A., Bruce, J.W.,
768 Coomes, O.T., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K.,
769 Imbernon, J., Leemans, R., Li, X.B., Moran, E.F., Mortimore, M.,
770 Ramakrishnan, P.S., Richards, J.F., Skanes, H., Steffen, W., Stone, G.D.,
771 Svedin, U., Veldkamp, T.A., Vogel, C., and Xu, J.C., 2001. The causes of
772 land-use and land-cover change: moving beyond the myths. *Global
773 Environmental Change-Human and Policy Dimensions* 11, 261-269.
- 774 Liu, J.G., Dietz, T., Carpenter, S.R., Alberti, M., Folke, C., Moran, E., Pell, A.N.,
775 Deadman, P., Kratz, T., Lubchenco, J., Ostrom, E., Ouyang, Z., Provencher,
776 W., Redman, C.L., Schneider, S.H., and Taylor, W.W., 2007. Complexity of
777 coupled human and natural systems. *Science* 317, 1513-1516.
- 778 Luus, K.A., Robinson, D.T., and Deadman, P.J., 2011. Representing ecological
779 processes in agent-based models of land use and cover change. *Journal of
780 Land Use Science iFirst* 1–24.
- 781 McLane, A.J., Semeniuk, C., McDermid, G.J., and Marceau, D.J., 2011. The role of
782 agent-based models in wildlife ecology and management. *Ecological
783 Modelling* 222(8), 1544-1556.
- 784 Milner-Gulland, E.J., 2012. Interactions between human behaviour and ecological
785 systems. *Philosophical Transactions of the Royal Society B* 367(1586), 270-
786 278.
- 787 Morris, A.J., Smith, B., Jones, N.E., and Cook, S.K., 2007. Experiment 1.1 –
788 Manipulate within crop agronomy to increase biodiversity: Crop architecture.
789 In *The SAFFIE Project Report HGCA No. 416*, 21-107.
- 790 Murray-Rust, D., Dendoncker, N., Dawson, T., Acosta-Michlik, L., Karali, E.,
791 Guillem, E., and Rounsevell, M., 2011. Conceptualising the analysis of socio-
792 ecological systems through ecosystem services and agent based modelling.
793 *Journal of Land Use Science* 6, 83-99.
- 794 Murray-Rust, D., Rounsevell, M.D.A., Robinson, D.T, Guillem, E.E., and Karali, E.,
795 2014. [An open framework for agent based modelling of agricultural land use
796 change](#). *Environmental Modelling & Software* 61, 19-38.
- 797 Odderskaer, P., Prang, A., Poulsen, J.G., Andersen, P.N., and Elmegaard, N., 1997.
798 Skylark (*Alauda arvensis*) utilisation of micro-habitats in spring barley fields.
799 *Agriculture Ecosystems & Environment* 62, 21-29.

800 Pelosi, C., Goulard, M., and Balent, G., 2010. The spatial scale mismatch between
801 ecological processes and agricultural management: Do difficulties come from
802 underlying theoretical frameworks? *Agriculture Ecosystems & Environment*
803 139, 455-462.

804 Pinheiro, J., Bates. D., DebRoy. S., Sarkar. D., and R Core team., 2009. nlme: Linear
805 and nonlinear mixed effects models. R package version 3.1-92.

806 Polasky, S., Nelson, E., Pennington, D., and Johnson, K.A., 2011. The Impact of
807 Land-Use Change on Ecosystem Services, Biodiversity and Returns to
808 Landowners: A Case Study in the State of Minnesota. *Environmental &*
809 *Resource Economics* 48, 219-242.

810 Poulsen, J.G., Sotherton, N.W., and Aebischer, N.J., 1998. Comparative nesting and
811 feeding ecology of skylarks *Alauda arvensis* on arable farmland in southern
812 England with special reference to set-aside. *Journal of Applied Ecology* 35,
813 131-147.

814 Power, A.G., 2010. Ecosystem services and agriculture: tradeoffs and synergies.
815 *Philosophical Transactions of the Royal Society B-Biological Sciences* 365,
816 2959-2971.

817 Robinson, R.A., 2005. Bird Facts: profiles of birds occurring in Britain & Ireland.
818 BTO Research Report 407. BTO, Thetford, <http://www.bto.org/birdfacts>

819 Robinson, D.T., Murray-Rust, D., Rieser, V., Rounsevell, M.D.A., Karali, E.,
820 Hersperger, A., and Guillem, E.E., 2011. Estimating the impacts of land-use
821 change on human well-being and the provision of ecosystem services.
822 Proceeding of the 18th annual conference of the international association for
823 landscape ecology , 6-8 september 2011, Wolverhampton, UK.

824 SAC, 2000 to 2008. The Farm Management Handbook. SAC, Edinburgh, UK.

825 Sage, R., Cunningham, M., Haughton, A.J., Mallott, M.D., Bohan, D.A., Riche, A.,
826 and Karp, A., 2010. The environmental impacts of biomass crops: use by birds
827 of miscanthus in summer and winter in south western England. *Ibis* 152, 487-
828 499.

829 Schwarz, G., Moxey, A., McCracken, D., Huband, S., and Cummins, R., 2008. An
830 analysis of the potential effectiveness of a Payment-by-Results approach to the
831 delivery of environmental public goods and services supplied by Agri-
832 Environment Schemes. Report to the Land Use Policy Group, UK. Macaulay
833 Institute, Pareto Consulting and Scottish Agricultural College.

834 Scotland's Environment. 2014. Crops and livestock.
835 [http://www.environment.scotland.gov.uk/get-informed/land/crops-and-](http://www.environment.scotland.gov.uk/get-informed/land/crops-and-livestock/)
836 [livestock/](http://www.environment.scotland.gov.uk/get-informed/land/crops-and-livestock/)

837 Scottish Executive. 2003. Guidelines for farmers in Nitrate Vulnerable Zones. Water
838 Environment Unit, Scottish Executive Environment and Rural Affairs
839 Department, Edinburgh.

840 Scottish Government, 2007. Final Results of the 2007 June Agricultural Census.

841 Semere, T. and Slater, F.M., 2007. Ground flora, small mammal and bird species
842 diversity in miscanthus (*Miscanthus x giganteus*) and reed canary-grass
843 (*Phalaris arundinacea*) fields. *Biomass & Bioenergy* 31, 20-29.

844 Setälä, H., Bardgett, R.D., Birkhofer, K., Brady, M., Byrne, L., de Ruiter, P.C., de
845 Vries, F.T., Gardi, C., Hedlund, K., Hemerik, L., Hotes, S., Liiri, M.,
846 Mortimer, S.R., Pavao-Zuckerman, M., Pouyat, R., Tsiafouli, M, and van der
847 Putten, W.H., 2014. Urban and agricultural soils: conflicts and trade-offs in
848 the optimization of ecosystem services. *Urban Ecosystems* 17, 239-253.

849 Settele, J., Carter, T.R., Kuhn, I., Spangenberg, J.H., and Sykes, M.T., 2012.
850 Scenarios as a tool for large-scale ecological research: experiences and legacy
851 of the ALARM project. *Global Ecology and Biogeography* 21, 1-4.

852 Siriwardena, G.M., Baillie, S.R., Buckland, S.T., Fewster, R.M., Marchant, J.H., and
853 Wilson, J.D., 1998. Trends in the abundance of farmland birds: a quantitative
854 comparison of smoothed Common Birds Census indices. *Journal of Applied
855 Ecology* 35, 24-43.

856 Sitch, S., Smith, B., Prentice, I.C., Arneth, A., Bondeau, A., Cramer, W., Kaplan,
857 J.O., Levis, S., Lucht, W., Sykes, M.T., Thonicke, K., and Venevsky, S., 2003.
858 Evaluation of ecosystem dynamics, plant geography and terrestrial carbon
859 cycling in the LPJ dynamic global vegetation model. *Global Change Biology*
860 9, 161-185.

861 SNH. 2005. East Scotland Grassland Scheme.
862 <http://www.snh.org.uk/pdfs/NatCare/GrasslandScheme.pdf>

863 Spangenberg, J.H., Bondeau, A., Carter, T.R., Fronzek, S., Jaeger, J., Jylha, K., Kuhn,
864 I., Omann, I., Paul, A., Reginster, I., Rounsevell, M., Schweiger, O., Stocker,
865 A., Sykes, M.T., and Settele, J., 2012. Scenarios for investigating risks to
866 biodiversity. *Global Ecology and Biogeography* 21, 5-18.

867 Staav R., and Fransson T., 2008. EURING list of longevity records for European
868 birds, http://www.euring.org/data_and_codes/longevity.htm.

869 Sutherland, W.J., and Freckleton, R.P., 2012. Making predictive ecology more
870 relevant to policy makers and practitioners. *Philosophical Transactions of the
871 Royal Society B* 367(1586), 322-330.

872 Swagemakers, P., Wiskerke, H., and van der Ploeg, J.D., 2009. Linking birds, fields
873 and farmers. *Journal of Environmental Management* 90, S185-S192.

874 Topping, C.J., Hansen, T.S., Jensen, T.S., Jepsen, J.U., Nikolajsen, F., and
875 Odderskaer, P., 2003. ALMaSS, an agent-based model for animals in
876 temperate European landscapes. *Ecological Modelling* 167, 65-82.

877 Topping, C.J., Sibly, R.M., Akcakaya, H.R., Smith, G.C., and Crocker, D.R., 2005.
878 Risk assessment of UK skylark populations using life-history and individual-
879 based landscape models. *Ecotoxicology* 14, 925-936.

880 Tranter, R.B., Swinbank, A., Wooldridge, M.J., Costa, L., Knapp, T., Little, G.P.J.,
881 and Sottomayor, M.L., 2007. Implications for food production, land use and
882 rural development of the European Union's Single Farm Payment: Indications
883 from a survey of farmers' intentions in Germany, Portugal and the UK. *Food
884 Policy* 32, 656-671.

885 Valentine, J., Heaton, R., Randerson, P., and Duller, C. J., 2008. The economics of
886 short-rotation coppice willow in the UK. *Proceedings of 16th European
887 Biomass Conference & Exhibition, Valencia, Spain, 2-6 June 2008*, 527-528.

888 Verburg, P.H. and Overmars, K.P., 2009. Combining top-down and bottom-up
889 dynamics in land use modeling: exploring the future of abandoned farmlands
890 in Europe with the Dyna-CLUE model. *Landscape Ecology* 24, 1167-1181.

891 Verhulst, J., Kleijn, D., and Berendse, F., 2007. Direct and indirect effects of the most
892 widely implemented Dutch agri-environment schemes on breeding waders.
893 *Journal of Applied Ecology* 44, 70-80.

894 Vinten, A.J.A., Abel, C., Watson, H., and Taylor, A., 2009. Lunan monitored priority
895 catchment project., *Annual Report*, Macauley Institute, Aberdeen, UK.

896 Wilson, G.A., 1997. Factors influencing farmer participation in the environmentally
897 sensitive areas scheme. *Journal of Environmental Management* 50, 67-93.

898 Wolfenden I.H., and Peach W.J., 2001. Temporal changes in the survival rates of
899 skylarks *Alauda arvensis* breeding in duneland in northwest England, In
900 Donald P.F., Vickery J.A. (eds) *The Ecology and Conservation of Skylarks*
901 *Alauda arvensis*, p 3-9, RSPB, Sandy.

902
903
904
905

906 **List of Figure captions**

907

908 **Figure 1** - Location of the case study, the Lunan catchment, in Scotland, and farm
909 (shaded colours) and parcels boundaries within the catchment (SIACS, 2007). (1590
910 parcels, min=0.03 Ha, max=85.86 Ha)

911 **Figure 2** – Conceptual framework of the coupled ABM/IBM. The dotted lines
912 represent feedbacks that are not implemented in the current version of the model.

913 **Figure 3** – Behavioural rules applied to individual adult skylarks. The rules of If-
914 Then type are in grey (Y: yes, N: no). The “Die” rule is not linked with another
915 behaviour but with life-cycle stages.

916 **Figure 4** – Crop vegetation curves derived from survey data. The darker grey
917 represents the suitable vegetation height for skylark nest establishment (10-120 cm).

918 **Figure 5** - Relationship between number of adult skylarks and a: Bioenergy produced,
919 b: Food energy produced; and between number of fledglings produced and c:
920 Bioenergy (MJ), d: Food energy (MJ).

921 **Figure 6** - Ecosystem services for the three ALARM scenarios, a: Average number of
922 adult skylarks (age>300 days), b: Average number of fledglings produced, c:
923 Bioenergy (MJ), d: Food energy (MJ).

924 **Figure 7** – Proportion of land cultivated by the different land use types considered in
925 the model under the three scenarios, B: BAMBU, G: GRAS, S: SEDG.

926 **Figure 8** - Average number of adult skylarks over the 2008-2050 simulation period in
927 a landscape managed exclusively by multifunctionalist (MULTIFUNCTIONAL),
928 profit-oriented (PROFIT), traditionalist (TRADITIONAL) farmers, and in the
929 landscape managed by the actual population of farmers (ALL).

930

931

932 **List of tables**

933

934 **Table 1** – Parameters and values for the suitability of nest sites. T is the maximum
935 territory density per hectare

936 **Table 2** – Parameters and values of life cycle traits in skylarks used in the model

937 **Table 3** – Energy conversion from food and bioenergy products

938

939

940

941

942

943

944

945

946

947

948

949

950

Table 1 – Parameters and values for the suitability of nest sites. *T* is the maximum number of territories per hectare

Parameters	Value	References
Vegetation height	Min=10cm; Max=120cm	Own field survey ^a
$T_{WinterWheat}$	0.141	Own field survey ^a
$T_{SpringCereal}$	0.135	Own field survey ^a
$T_{WinterBarley,Oat}$	0.077	Own field survey ^a
$T_{OilseedRape}$	0.062	Own field survey ^a
$T_{RootCrops}$	0.091	Own field survey ^a
$T_{Legumes}$	0.173	Own field survey ^a
$T_{GrassMowing}$	0.072	Own field survey ^a
$T_{IntensiveGrazing}$	0.084	Browne et al., 2000
$T_{ExtensiveGrazing}$	0.101	Browne et al., 2000
$T_{RoughGrazing}$	0.059	Browne et al., 2000
$T_{Miscanthus}$	0.030	Sage et al., 2010
T_{Willow}	0.095	Sage et al., 2010
$T_{SetAside}$	0.360	Browne et al., 2000

^a Field survey carried out in the Lunan catchment in 2009; unpublished data.

Table 2 – Parameters and values of life cycle traits in skylarks used in the model

Parameters	Value	References
Age of maturity (days)	300	Delius, 1965
Territory search space	ø 500m	maximum territory size ø 250m, Odderskaer et al., 1997
Number of eggs laid	4	Delius, 1965; Robinson, 2005
Daily probability of egg mortality^a	0.0293 ±0.1%	Chamberlain and Crick, 1999
Daily probability of nestling mortality^a	0.0536 ±0.1%	Chamberlain and Crick, 1999
Daily probability of fledgling mortality^a	0.027 ±0.1%	Poulsen et al., 1998
Daily probability of adult mortality (breeding season)^a	0.00197 ±0.1%	Wolfender and Peach, 2001
Daily probability of adult mortality (winter)^a	0.00275 ±0.1%	Topping et al., 2005
Lifespan (days)	max 3285	Staav and Fransson, 2008
Sex ratio	1:1	Dougall, 1997
Mating to egg laying (days)	5	Wilson et al., 1997
Egg laying interval (days)	1	Delius, 1965
Incubation (days)	11	Wilson et al., 1997
Caring for young (days)	19	Delius, 1965

^a These values are transformed from yearly rate (S) to daily rate (d) using the following equation: $d = 1 - (S^{(1/n)})$, with n the length of a given lifecycle stage (days).

Table 3 – Energy conversion from food and bioenergy products

	Energy (MJ/ton)	Reference
Wheat	13975	FAO ^b
Barley	13891	FAO ^b
Oat	16108	FAO ^b
OSR	20669	FAO ^b
Potatoes	32217	USDA ^c
Turnips	15062	USDA ^c
Carrots	30125	USDA ^c
Peas	33890	USDA ^c
Beans	28033	USDA ^c
Willow^a	17200	Valentine et al., 2008
Miscanthus	17000	Natural England ^d
Beef	6070	USDA ^e

^a net energetic value of wood at 35% moisture. value in MJ/oven dried ton

^b <http://www.fao.org/economic/ess/ess-data/ess-fs/ess-nutritive/en/>

^c <http://www.ars.usda.gov/SP2UserFiles/Place/12354500/Data/SR23/reports/sr23fg11.pdf>

^d http://www.naturalengland.org.uk/Images/miscanthus-guide_tcm6-4263.pdf

^e <http://nutritiondata.self.com/facts/beef-products/3477/2>





Figure 1 - Location of the case study, the Lunan, in Scotland, and farm (shaded colours) and parcels boundaries within the catchment (SIACS, 2007). (1590 parcels, min=0.03 Ha, max=85.86 Ha)




Figure 2 – Conceptual framework of the coupled ABM/IBM. The dotted lines represent feedbacks that are not implemented in the current version of the model.




Figure 3 – Behavioural rules applied to individual adult skylarks. The rules of If-Then type are in grey (Y: yes, N: no). The “Die” rule is not linked with another behaviour but with life-cycle stages.




Figure 4 – Crop vegetation curves derived from survey data. The darker grey represents the suitable vegetation height for skylark nest establishment (10-120 cm)




Figure 5 – Proportion of land cultivated by the different land use types considered in the model under the three scenarios, B: BAMBU, G: GRAS, S: SEDG




Figure 6 - Ecosystem services for the three ALARM scenarios, a: Average number of adult skylarks (age>300 days), b: Average number of fledglings produced, c: Bioenergy (MJ), d: Food energy (MJ)




Figure 7 - Average number of adult skylarks over the 2008-2050 simulation period in a landscape managed exclusively by traditionalist (TRADITIONAL), profit-oriented (PROFIT), multifunctionalist (MULTIFUNCTIONAL) farmers, and in the landscape managed by the actual population of farmers (ALL)




Figure 8 - Relationship between number of adult skylarks and a: Bioenergy produced, b: Food energy produced; and between number of fledglings produced and c: Bioenergy (MJ), d: Food energy (MJ)

