

Ex vivo T cell cytokine expression predicts survival in patients with severe alcoholic hepatitis

Ashwin D. Dhanda^{1,2}, Euan Yates², Lauren P. Schewitz-Bowers³, Philippa J. Lait³, Richard W.J. Lee³, Matthew E. Cramp¹²

¹South West Liver Unit, University Hospitals Plymouth NHS Trust, Plymouth, UK

²Hepatology Research Group, University of Plymouth, Plymouth, UK

³Translational Health Sciences, University of Bristol, Bristol, UK

Corresponding author:

Dr Ashwin Dhanda
South West Liver Unit, Level 7, Derriford Hospital
University Hospitals Plymouth NHS Trust
Plymouth PL6 8DH
United Kingdom
Email: ashwin.dhanda@plymouth.ac.uk
Tel: +44 1752 432723
Fax: +44 1752 517576

Running title: T cell cytokines in alcoholic hepatitis

Author contributions:

Study concept: AD, RL; Performed experimental work: EY, LSB, PL; Statistical analysis: AD, PL; Study supervision: RL, MC; Drafting manuscript: AD, LSB; Finalising manuscript: AD, RL, MC

Abstract

Aim

Alcoholic hepatitis (AH) is an acute inflammatory liver condition with high early mortality. Steroids have proven short-term survival benefit but non-responders have the worst outcomes. There is a clinical need to identify these high-risk individuals at presentation. T cells have been implicated in alcoholic hepatitis and steroid responsiveness. We aimed to measure *ex vivo* T cell cytokine expression as a candidate biomarker of outcome in patients with AH.

Methods

Consecutive patients with AH (bilirubin $>80\mu\text{mol/L}$ and $\text{AST:ALT}>1.5$ in heavy alcohol consumers with discriminant function $[\text{DF}]\geq 32$), were recruited from University Hospitals Plymouth NHS Trust. T cells were obtained and stimulated *ex vivo* before cytokine expression was determined by flow cytometry and protein multiplex analysis.

Results

Twenty-three patients were recruited (10 male; median age 51; baseline DF 67; 30% 90-day mortality). Compared to non-survivors at day 90, T cells from survivors had higher baseline intracellular IL-10:IL-17A ratio (0.43 v 1.20; $p=0.02$). Multiplex protein analysis identified IFN γ and TNF α as independent predictors of 90-day mortality ($p=0.04$ and $p=0.01$ respectively). The ratio of IFN γ /TNF α was predictive of 90-day mortality (1.4 v 0.2; $p=0.03$).

Conclusions

These data demonstrate the potential utility of T cell cytokine release assays performed on pre-treatment blood samples as biomarkers of survival in severe AH. Our key findings were that both the ratio of intracellular IL-10 to IL-17A and the ratio of IFN γ to TNF α in culture supernatants were predictors of 90-day mortality. This offers the promise of developing T cell based diagnostic tools for risk stratification.

Keywords: Alcoholic hepatitis, T cells, cytokines, biomarker

Introduction

Alcoholic hepatitis (AH) is an acute inflammatory condition, which carries a high mortality of 30% within 90 days¹. Steroid treatment is the only therapy with proven short-term survival benefit². However, the worst outcomes are in the 30-40% of patients with a poor response to steroid treatment³, and there is a pressing need to identify these high-risk individuals at presentation.

T cells have long been implicated in the pathogenesis of AH and we have previously reported that suppression of lymphocyte proliferation correlates with clinical outcome following steroid treatment^{4,5}. Given the barriers to the development of this radiation-based assay for clinical application and the precedent of functional cytokine release assays for other clinical applications⁶, we sought to evaluate T cell cytokine expression as a candidate biomarker of AH mortality.

Case report

Study design

The study protocol was approved by the UK's Health Research Authority (reference: 15/LO/1501) and all participants provided written informed consent. Consecutive patients with severe AH, defined as recent onset jaundice with bilirubin > 80 $\mu\text{mol/L}$ and AST:ALT > 1.5 in heavy alcohol consumers (> 60 g or 80 g ethanol / day in females and males respectively) with a discriminant function (DF) score ≥ 32 , were recruited from University Hospitals Plymouth NHS Trust. Patients received standard care including steroid treatment in the absence of active infection, hepatorenal syndrome or gastrointestinal haemorrhage.

T cell isolation, stimulation and statistical analysis

Whole blood samples were taken before steroid treatment was started with CD4⁺ T cells isolated by negative selection (Stemcell Technologies, Cambridge, UK) and cultured for 4 days in supplemented

media with interleukin (IL)-2 (Sigma-Aldrich, Poole, UK) and T cell receptor stimulation (anti-CD3/CD28 microbeads; Thermo Fisher Scientific, Loughborough, UK). Cells cultured without T cell receptor stimulation were included as controls. T cell receptor stimulation with anti-CD3/CD28 microbeads was selected as a standardized method of T cell activation, which has been optimized for use in other conditions⁷. For the final 4 hours, cultures were stimulated with T cell mitogen phorbol 12-myristate 13-acetate (PMA) and ionomycin (Sigma) with golgi export inhibitor (BD Biosciences, Oxford, UK). Cells were fixed, permeabilised and stained with fluorescently labelled antibodies to IL-10, IL-17A and interferon gamma (IFN γ) (Thermo Fisher Scientific), quantified on a BD Accuri flow cytometer (BD Biosciences) and analysed in FlowJo (FlowJo LLC, Ashland, OR, USA). Protein concentration in cell culture supernatants prior to the final 4 hour stimulation was analysed in duplicate for CCL20, GM-CSF, IFN γ , IL-10, IL-12p70, IL-17A, IL-21, IL-23, IL-4, IL-6 and TNF α using a magnetic bead array (R&D Systems, Minneapolis, MN) on a Luminex 200 analyser (Luminex Corp, Austin, TX). Statistical analysis was performed using IBM SPSS version 24 (IBM, Armonk, NY). The primary clinical outcome was death within 90 days of presentation. The data presented are median values and comparisons were made with non-parametric tests.

Findings

23 consecutive patients were recruited between April 2016 and November 2017 (10 male; median age 51, baseline DF 67, model for end-stage liver disease [MELD] score 19). 90-day mortality was 30%. Four patients did not receive steroids because of active infection. Samples were obtained prior to steroid treatment in all cases at a median of 5 days from hospital admission. Median time from admission to steroid treatment was 6 days.

Compared to non-survivors at day 90, CD4⁺ T cells from non-survivors had a higher baseline intracellular IL-10:IL-17A ratio (percent of T cells expressing IL-10: 2.4% [non-survivors] v 1.6% [survivors], p=0.07; IL-17A: 2.1% [non-survivors] v 3.5% [survivors], p=0.08; median IL-10:IL-17A ratio

1.20 [non-survivors] v 0.43 [survivors]; $p=0.02$; **figure 1A**) with an area under the curve of the receiver operating characteristic (AUROC) score of 0.82 (95% confidence interval 0.64-1.00). The proportion of T cells expressing IFN γ was similar between groups (14.2% v 11.9%; $p=0.55$). In comparison, expression of cytokines in control cultures was lower than stimulated conditions: IL-10: 0.2% v 1.8%; IL-17: 2.5% v 3.3%; IFN γ : 6.2% v 13.3% (all $p=0.07$; Wilcoxon signed ranks test).

Multiplex protein analysis performed on 22 of the 23 cases identified IFN γ and TNF α as most differentially expressed between survivors and non-survivors at day 90 (7044 v 2741 pg/ml; $p=0.19$ and 2610 v 4098 pg/ml; $p=0.08$ respectively). Multivariate analysis confirmed that both IFN γ and TNF α were independent predictors of 90-day mortality ($p=0.04$ and $p=0.01$ respectively). The ratio of IFN γ /TNF α was predictive of 90-day mortality (1.4 v 0.2; $p=0.03$; **figure 1B**) with an AUROC score of 0.79 (95% CI 0.58-0.99). The concentration of other mediators tested in the multiplex was similar between survivors and non-survivors. No cytokine concentration correlated with baseline DF or MELD scores. Steroid response, as determined by Lille score with a threshold of 0.45 at day 7 of steroid treatment, was not associated with differences in any cytokine measured by either flow cytometry or protein multiplex. Similarly, the ratios of intracellular IL-10/IL-17A and IFN γ /TNF α were the same in Lille steroid responders and non-responders.

Discussion

These data from a pilot prospective cohort of more than 20 patients demonstrate the potential utility of T cell cytokine release assays performed on pre-treatment blood samples as biomarkers of survival in severe AH. Our key findings were that both the ratio of intracellular IL-10 to IL-17A measured by flow cytometry in CD4⁺ cells and the ratio of IFN γ to TNF α in culture supernatants were predictors of 90-day mortality. These data also demonstrate that flow cytometry and protein

multiplex are complementary techniques that can reveal different alterations in protein expression either in proportion of cells secreting a cytokine or in cytokine concentration, respectively.

The pathogenesis of AH is orchestrated by several immune subsets including neutrophils and monocytes, which are activated by gut derived signals such as endotoxin^{8,9}. However, T cells also play a vital role in responding to antigen presenting cells and propagating inflammation in AH¹⁰. Transcriptome studies have confirmed upregulation of TNF and T cell pathways^{11,12}. In particular, the Th17 (IL-17 expressing T cell) pathway is upregulated with high expression of related chemokines such as CCL20^{12,13}. Furthermore, recent studies have demonstrated that monocytes activated by exposure to gut microbial signals interact with CD4⁺ T cells to alter expression of exhaustion markers and cytokines including IFN γ ¹⁴. These data confirm the relevance of T cells in AH and support the measurement of T cell cytokines as a broad measure of immune response.

These findings offer the promise of developing T cell based diagnostic tools for risk stratification both in the context of routine clinical practice and also to inform the design of smart clinical trials seeking to test new interventions in patients most likely to fail conventional corticosteroid therapy.

Acknowledgements

Grant support:

AD received funding from Plymouth Hospitals Charity. AD and RL receive salary funding from the National Institute for Health Research. L.P.S-B, P.J.L and R.W.J.L are supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology. The views expressed are those of the Author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health.

Disclosures:

L.P.S.-B. and R.W.J.L are named inventors on a US patent application (no. 61/919,404), which incorporates biomarkers to identify patients who will benefit from treatments for inflammatory diseases

References

1. Thursz MR, Richardson P, Allison M, et al. Prednisolone or pentoxifylline for alcoholic hepatitis. *N Engl J Med* 2015;372:1619-1628.
2. Louvet A, Thursz MR, Kim DJ, et al. Corticosteroids Reduce Risk of Death Within 28 Days for Patients With Severe Alcoholic Hepatitis, Compared With Pentoxifylline or Placebo—a Meta-analysis of Individual Data From Controlled Trials. *Gastroenterology* 2018;155:458-468 e458.
3. Louvet A, Naveau S, Abdelnour M, et al. The Lille model: a new tool for therapeutic strategy in patients with severe alcoholic hepatitis treated with steroids. *Hepatology* 2007;45:1348-1354.
4. Dhanda AD, di Mambro AJ, Hunt VL, et al. Long-term outcome in patients with severe alcoholic hepatitis can be reliably determined using an in vitro measure of steroid sensitivity. *Hepatology* 2015;61:1099.
5. di Mambro AJ, Parker R, McCune A, Gordon F, Dayan CM, Collins P. In vitro steroid resistance correlates with outcome in severe alcoholic hepatitis. *Hepatology* 2011;53:1316-1322.
6. Diel R, Goletti D, Ferrara G, et al. Interferon-gamma release assays for the diagnosis of latent *Mycobacterium tuberculosis* infection: a systematic review and meta-analysis. *Eur Respir J* 2011;37:88-99.
7. Lee RW, Schewitz LP, Nicholson LB, Dayan CM, Dick AD. Steroid refractory CD4+ T cells in patients with sight-threatening uveitis. *Invest Ophthalmol Vis Sci* 2009;50:4273-4278.
8. Mookerjee RP, Stadlbauer V, Lidder S, et al. Neutrophil dysfunction in alcoholic hepatitis superimposed on cirrhosis is reversible and predicts the outcome. *Hepatology* 2007;46:831-840.
9. Vergis N, Khamri W, Beale K, et al. Defective monocyte oxidative burst predicts infection in alcoholic hepatitis and is associated with reduced expression of NADPH oxidase. *Gut* 2017;66:519-529.
10. Dhanda AD, Collins PL. Immune dysfunction in acute alcoholic hepatitis. *World J Gastroenterol* 2015;21:11904-11913.
11. Affo S, Dominguez M, Lozano JJ, et al. Transcriptome analysis identifies TNF superfamily receptors as potential therapeutic targets in alcoholic hepatitis. *Gut* 2013;62:452-460.
12. Affo S, Morales-Ibanez O, Rodrigo-Torres D, et al. CCL20 mediates lipopolysaccharide induced liver injury and is a potential driver of inflammation and fibrosis in alcoholic hepatitis. *Gut* 2014.
13. Lemmers A, Moreno C, Gustot T, et al. The interleukin-17 pathway is involved in human alcoholic liver disease. *Hepatology* 2009;49:646-657.
14. Markwick LJ, Riva A, Ryan JM, et al. Blockade of PD1 and TIM3 restores innate and adaptive immunity in patients with acute alcoholic hepatitis. *Gastroenterology* 2015;148:590-602 e510.

Figure 1: (A) Ratio of the percentage of CD4⁺ T cells expressing IL-10:IL-17A measured by flow cytometry (0.4 v 1.2; p=0.02) and (B) ratio of IFN γ :TNF α concentration measured by protein multiplex in patients with AH alive and dead at day 90 (1.4 v 0.2; p=0.03). *p<0.05

