

Regionally-defined niche-breadth of tropical African freshwater plant species predicts their global latitudinal range

Kennedy, M. , Lang, P. , Tapia-Grimaldo, J. , Varandas-Martins, S. , Bruce, A. , Moore, I. , Sichingabula, H. , Bottino, F. and Murphy, K.

Published PDF deposited in [Curve](#) May 2016

Original citation:

Kennedy, M. , Lang, P. , Tapia-Grimaldo, J. , Varandas-Martins, S. , Bruce, A. , Moore, I. , Sichingabula, H. , Bottino, F. and Murphy, K. (2016) 'Regionally-defined niche-breadth of tropical African freshwater plant species predicts their global latitudinal range' In: Geophysical Research Abstracts, 'European Geosciences Union General Assembly 2016'. Held 17-22 April 2016 at Vienna, Austria. , 9372.

URL: <http://meetingorganizer.copernicus.org/EGU2016/posters/20288>

Publisher: European Geosciences Union

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

CURVE is the Institutional Repository for Coventry University



Regionally-defined niche-breadth of tropical African freshwater plant species predicts their global latitudinal range

Michael Kennedy (1,2), Pauline Lang (3,4), Julissa Tapia-Grimaldo (4), Sara Varandas-Martins (4), Allannah Bruce (4), Isabel Moore (4), Henry Sichingabula (5), Flávia Bottino (6), and Kevin Murphy (4)

(1) School of Energy, Construction and Environment, Coventry University, Coventry, CV1 5FB, UK (michael.kennedy@coventry.ac.uk), (2) School of Geosciences, University of Aberdeen, Aberdeen, AB24 5BW, UK, (3) Scottish Environment Protection Agency, 6 Parklands Avenue, Eurocentral, Holytown, North Lanarkshire, ML1 4WQ, UK, (4) University of Glasgow, Glasgow, G12 8QQ, UK, (5) Department of Geography and Environmental Studies, University of Zambia, Lusaka, Zambia, (6) Universidade Federal de São Carlos, Departamento de Hidrobiologia, Rodovia 17 Washington Luís, km 235, CEP: 13565-905, São Carlos, São Paulo State, Brasil

This study tested the hypothesis that the measured niche-breadth of river plant species (macrophytes) occurring within a closely-defined geographical area in southern tropical Africa (Zambia), may predict the larger-scale biogeographical range of these species.

Two measures of niche-breadth were calculated for 44 riverine macrophyte species commonly occurring in Zambia, using an approach based on PCA ordination with bio-physico-chemical ordination input variables: altitude, stream order, stream flow, pH, conductivity, soluble reactive phosphate concentration (SRP), benthic macroinvertebrate Average Score per Taxon (ASPT), and individual abundance of nine benthic macroinvertebrate families showing differing water quality tolerance, as indicated by their Sensitivity Weightings within the Zambian Invertebrate Scoring System (ZISS). Macrophyte large-scale latitudinal range was derived from world geospatial records held by online databases, supplemented by records held by the authors.

The two niche-breadth metrics divided the species into narrow-niche and intermediate/broad-niche categories, showing significant variation in altitude, stream flow, conductivity, SRP and ASPT, but not stream order or pH. There was no evidence to suggest that macrophyte alpha-diversity (as a measure of number of individual niches that may co-exist in a given habitat) showed any significant relationship with individual species niche-breadth. However, macrophyte alpha-diversity was significantly positively correlated with altitude, and significantly negatively related to conductivity, pH, ASPT, SRP, stream flow, and stream order. Narrow-niche macrophyte species included a higher proportion of Afrotropical endemics than did species with broader niche size. There were significant predictive relationships between macrophyte niche breadth and latitudinal range of the target species at global and Afrotropical scales, but not for the Neotropical range of species which occurred in both the Neotropics and Zambia. Rare, narrow-niche species distributional data may account for this.

The study found statistically-significant evidence that macrophyte species niche breadth, measured at sites within a limited latitudinal range of 9.83° in Zambian rivers, is a significant predictor of the latitudinal range size of these plants at global and Afrotropical bioregion scales; but not Neotropical range, for those species which co-occurred both in Zambia and the Neotropics.