

# Experiences of cognitive behavioural therapy formulation in clients with depression

Kahlon, S. , Neal, A. and Patterson, T.

**Author post-print (accepted) deposited in CURVE February 2016**

**Original citation & hyperlink:**

Kahlon, S. , Neal, A. and Patterson, T. (2014) Experiences of cognitive behavioural therapy formulation in clients with depression. *The Cognitive Behaviour Therapist*, volume 7 (e8)  
<http://dx.doi.org/10.1017/S1754470X14000075>

ESSN 1754-470X

DOI 10.1017/S1754470X14000075

**Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.**

**This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.**

## ABSTRACT

**Background:** Whilst clinicians have described the benefits of using formulations within therapy, little is understood about the client's experience of the formulation process. Cognitive - behavioural therapy (CBT) is currently the treatment of choice for many adult mental health difficulties. However, research exploring clients' experiences of CBT formulation is very limited.

**Aims:** The present study set out to explore experiences of CBT formulation among clients with depression.

**Method:** Seven participants were interviewed and the data analysed using Thematic Analysis

**Results:** The analysis identified key themes as: 'feeling trapped or restricted by depression', 'The development of the formulation – from coming to my own conclusions to something the therapist developed', 'From negative towards mixed feelings: emotional reactions to the formulation during the therapeutic process' and 'a new journey: towards making a new sense of oneself'.

**Conclusion:** The results of the study highlight the personal and emotional challenge of the formulation process for clients.

## INTRODUCTION

### **Benefits of formulation for clinicians and clients**

Over the past two decades, formulation has become an integral component of a number of psychological approaches to the treatment of mental health difficulties. Formulation is considered to be the basis for achieving an understanding of the client's difficulties in CBT (Aston, 2009; Eels, 1997; Beck, 1995). The importance of collaborative development of formulation has been emphasised in the extant CBT literature (see Dudley and Kuyken, 2006; Kinderman and Lobban, 2000). The current evidence base appears to focus on the benefits of CBT formulation from clinicians' perspectives, for example, Pain, Chadwick and Abba (2008) state that formulation can aid clinicians' understanding of the client and help clients to comprehend and adapt to the cognitive model. Furthermore, CBT formulation allows for a focus on the 'here and now' which is considered to be particularly helpful to clinicians (Flitcroft, James and Freeston, 2007), as well as informing appropriate interventions (Kuyken, 2006).

In contrast, less is understood regarding the benefits of formulation for clients (Aston, 2009). In the UK, NHS clinicians have a commitment to involving service users throughout the research process (DoH, 2004) as well as a need to utilise evidence-based interventions in the treatment of mental health difficulties. Some interventions, such as manualised CBT, do not use formulation therefore clinicians need to be able to justify the use of formulation.

### **Research on the client's experience of formulation**

There appears to be an absence of the client's voice in the extant empirical literature (Gray and Grant, 2005), as well as a dearth of research conducted into clients' experiences of CBT formulation (Aston, 2009). Interestingly, a search of the literature identified only three empirical papers which investigated clients' experiences of CBT formulation. Firstly, Chadwick, Williams and Mackenzie (2003) examined the impact of case formulation in CBT for psychosis using structured questionnaires. Thirteen participants took part within this quantitative study where no significant findings emerged. Although, participants reported some benefits of formulation, these were not described in any detail and the authors of the study suggest that the benefits of formulation might not have been captured due to limitations within the quantitative methodology employed. Secondly, Pain, Chadwick and Abba (2008) investigated experiences of CBT formulation among individuals with psychosis, using qualitative methodology. They used a sample of fifteen participants, investigating clients' experience using semi-structured interviews. The data was analysed using content analysis through which a number of themes were identified, the most important being *clients experiencing an emotional reaction to receiving the formulation* (within this category, 40% had a negative reaction and 22% a positive reaction). Thirdly, Brown (2008) explored clients' experiences of formulation in a sample of six participants with differing mental health presentations who had received formulations from a variety of psychological orientations (including dynamic/systemic, integrative, CBT/systemic and CAT/integrative). A linear

model of clients' experiences was developed consisting of three stages: *initial doubt*, *assimilation* and *empowerment*. Brown (2008) highlighted an under-representation of negative reactions to formulation which raises the question of whether all aspects of clients' reactions were adequately captured in the study.

Building upon the relatively small body of published research in this area, the present study explores clients' experiences of formulation within a single therapeutic model, CBT, and focusing on a specific clinical population, clients referred for the treatment of depression. Through recruiting a sample of clients with similar psychological difficulties and who have all received formulations using the same therapeutic modality, the present study also attempts to address some of the limitations of sample heterogeneity present in previous research.

## **METHODOLOGY**

### **Design**

Thematic Analysis (Boyatzis, 1998; Braun & Clarke, 2006) was used to inform the data collection and analysis of the present study, which elicits and analyses first-person accounts from clients with depression who offer a unique perspective on CBT formulation. The use of semi-structured interviews allowed participants to provide a rich, and open, description of their experiences, a widely adopted approach in qualitative research (Smith & Eatough, 2007).

## **Participants**

### ***Table 2.1 here***

The sample consisted of seven adults between 19 and 54 years of age, referred to psychological services for treatment of depression and recruited using purposive sampling methods. Table 2.1. summarises participant demographic information whilst presenting the pseudonyms used in the analysis to anonymise participants.

The inclusion criteria required that participants in the present study should be adults of working age (18 - 65), who had been referred to psychological services for treatment of depression, and who had not previously received a formulation. Potential participants were excluded if they had co-morbid severe and enduring psychological difficulties or if they did not have the capacity to consent to the study. To control for participants not having received a formulation before, all participants who had received psychological therapy prior to CBT were excluded from the study.

## **Materials**

A semi-structured interview schedule was developed based on discussion with and feedback from Clinical Psychologists, the first author's own clinical experience of developing formulations collaboratively with service users, and a review of relevant literature. Key themes that the interview covered were:

1. Formulation development.
2. Emotional reactions of the client to receiving the formulation.
3. How the formulation informed the therapeutic process.

4. The impact that the formulation had on the client's difficulties and their day-to-day life.
5. What the client learnt from the formulation.

Additional questions were asked dependent on participants' responses during the interview.

### **Procedure**

Ethical approval for the study was granted by the National Research Ethics Service.

Clinical Psychologists and Psychological Therapists working in primary and secondary care psychological services for adults and using a CBT approach were asked to review their caseloads for clients who met the study criteria. Suitable potential participants who completed and returned the opt-in form were contacted by the first author. Seven clients expressed an interest in the study and subsequently consented to participate in the study. Participants were interviewed by the first author where all interviews were audio recorded.

### **Data Analysis**

The interviews were transcribed verbatim and analysed using Thematic Analysis (Boyatzis, 1998; Braun and Clarke, 2006). This entailed reading and re-reading the first transcript and developing line by line comments and

descriptions on participants' thoughts and experiences, from which points of convergence and similarity were identified. A list of emergent themes was developed from this initial commentary. This process was subsequently repeated for each transcript following which, patterns of themes and frequency of theme occurrence across all the data set were considered in order to develop subordinate themes. Superordinate themes were developed by considering difference and commonality between subordinate themes.

An independent audit was conducted whereby an independent researcher was provide with excerpts of transcripts to develop their own line by line coding and themes. These were then compared to the themes identified by the first author. Discussions with the independent researcher were used to highlight both the similarities and some subtle differences in coding and themes, and this, in addition to research supervision discussions, served as a quality control procedure within the data analysis.

### **Researcher's position**

The data analysis was conducted by the first author who at the point of conducting the research was a trainee clinical psychologist with a keen interest in CBT, having used this model with adults on placement where she had developed CBT formulations with clients. Her own experience of CBT formulation combined with her experience of other theoretical and therapeutic perspectives had resulted in a particular interest in thinking about the formulation process from an eclectic perspective.



The second author is a clinical psychologist who has interests in integrative and CBT formulation, therapy process, and the therapeutic relationship.

The third author is a clinical psychologist and university lecturer, with research interests in older adult clinical psychology, person-centred psychology, third wave cognitive-behavioural therapies and process variables in psychological therapy.

### **Writing up the data**

In writing up the findings of the present study, it was decided to separate the results from the analysis to enable the reader to differentiate between the participants' and the researcher's voice. Thus, the results section provides a descriptive summary of the data found whilst the discussion incorporates the authors' interpretations of this. By separating the analysis in this way readers are invited to consider their own interpretations of the findings.

## RESULTS

Four superordinate themes were drawn from the participants data, these were: 'feeling trapped or restricted by depression', 'the development of the formulation – from coming to my own conclusions to something the therapist developed', 'from negative towards mixed feelings: emotional reactions to the formulation during the therapeutic process' and 'a new journey: towards making a new sense of oneself'. The themes along with their subordinate themes are illustrated in Table 2.2. The extent to which themes and sub-themes represent the dataset is noted in parentheses. Superordinate themes, along with their subordinate themes will be described and illustrated with quotations from participant interviews.

***Table 2.2 here***

### **Feeling trapped or restricted by depression**

Although the first superordinate theme does not directly relate to participants' experience of CBT formulation, it will be briefly explored in order to provide a context to the process participants have gone through. Even though this was not an area explored by the interviewer, all participants discussed their experience of depression prior to therapy.

The data showed that prior to therapy, participants experienced a lack of control over their emotions and thought processes and felt caught up in these. All participants reflected on this time in their life as having been more challenging or negative than their present situation. In particular, participants

## CLIENTS' EXPERIENCE OF FORMULATION

described a sense of feeling stuck within or constricted by depression. One participant made a link between the potential hazards within their external world and feeling trapped but was unable to describe what they were trapped by illustrating the difficulty in describing this complex experience:

'...every situation is a hazard, reflected, negative, of being trapped in something, with difficulties or something'. (Arun)

Another participant described this experience as one of feeling trapped within one's mind and having an inability to take in his entire external world:

'As I say, going along blinkered, mired in your own mind. I think you do feel trapped in your own mind'.  
(Sam)

In the extract below, the experience of being depressed is described as the client feeling tortured as a result of her expectations upon herself and social interactions:

'I think because the main problem I had was that I just used to be tortured by my thoughts, from any little innocent little interaction with anybody, and now it is just, I used to have little noisy voices saying, it sounds schizophrenic but oh for god's sake, kind of why just

giving myself a hard time and having such high expectations of myself'. (Jill)

Understandably, individuals feeling this way had been socially isolating themselves and, prior to therapy, had been struggling to find a way to change their current emotional world alone.

**The development of the formulation – from coming to my own conclusions to something the therapist developed**

***Exploring thoughts, feelings, actions and previous experiences – developing the formulation***

The formulation was developed between the clinician and participant by exploring the individuals' experience through questions and answers for all seven participants. All participants described a process of exploring thought, feelings and personal experiences with their clinician. One participant shared her experience of the development of the formulation through initially discussing her past which was still hard to talk about:

'It was basically developed from where I had to think about when I was younger and what kind of life I'd wanted. Different feelings and thoughts that I'd have, bad feelings in my life, well, to relate back to bad things that have happened in the past - who it was and when it

## CLIENTS' EXPERIENCE OF FORMULATION

was, and what it was, and started from that and then basically carried on'. (Lisa)

Another participant described a further element of the development of a formulation where they and their clinician explored not only feelings, but also the client's actions, suggesting that behaviours as well as thoughts and feelings were a focus during the process of developing the formulation:

'It was developed I think, through questions being asked slowly. First of all I think it was quite distant unrelated questions, and then it started to be a bit more focussed, and questioning why I would do something. I think originally it was on why I would feel bad about something'. (Arun)

A further participant described the formulation process as involving the development of an account of why she had become depressed. Furthermore, the process of developing a formulation is described by this participant as something which was occurring or evolving throughout the course of the therapy:

'The formulation was developed really over the course of the entire 16 weeks. Erm, I think the first couple of times that [THERAPIST] and I met they just explored

how I was feeling, and what had been happening with me, and then just slowly it evolved into a bit of an explanation of why I had reached that point'. (Jill)

***Collaboration, directive, or coming up with it myself: views on who developed the formulation***

Even though all participants appeared to agree that the formulation had emerged as a result of the discourse between client and clinician, their comments varied regarding their views of who had been responsible for developing the formulation. Four participants viewed the process as a collaborative process, one of whom considered the formulation as being something initially developed together with her therapist within the session but then further elaborated out of sessions by herself alone:

`Well it was probably, I probably best say developed together at first but then I, obviously because we met once a week I developed it on my own on the rest of the days`. (Penny)

In contrast, another participant considered it to be more a process of coming up with her own formulation which seemed to have a greater impact than a clinician developed explanation:

'I think, retrospectively, I realised it was more of a validation for me because it was coming from me and I

## CLIENTS' EXPERIENCE OF FORMULATION

was talking through everything myself and coming to my own conclusions. And I think it had more of an impact, and my clinician would affirm what I was saying or nod and I think, because she obviously thought the same things'. (Jill)

The remaining two participants viewed the formulation as having been developed predominantly by the clinician. Whilst one of these participants felt this was due to them needing more prompting during their early therapy sessions as a result of not being ready to commence therapy, another expressed the view that the formulation was not an interactive process but more something that the clinician brought along, which could not be changed, having developed it without him:

'I would think that it was something, when I looked at it, it was something that he developed. It wasn't something so much, it wasn't an interactive thing where we said 'oh okay what about this, what about that?', it almost felt like one day he came up with this thing and I was like looking at it going 'oh okay' rather than 'oh shall we change this?' and 'do you feel like this and like this?' (Arun)

**From negative to mixed feelings: emotional reactions to the formulation during the therapeutic process**

All of the participants experienced a broad range of feelings towards the formulation which appeared to differ during the therapeutic process. With the exception of one individual, feelings towards the formulation appeared more negative earlier on. However, these individuals also expressed more positive feelings later on in the therapeutic process.

***Receiving a formulation is difficult***

All the participants had more negative feelings towards their formulation to begin with. For one participant these feelings were combined with that person's reservations regarding whether therapy would help them, apparently linked to the experience of medication not helping:

'Not sure really, you know just confusion, thinking I hope it's going to work and thinking things like that, and "I hope I'm going to get better from all of this" at the time because tablets haven't been working or anything'.

(Lisa)

In line with Lisa's experience, other participants also found the formulation to be difficult to comprehend with one participant stating that it was 'very technical' and another finding the initial formulation emotionally very difficult. Although this participant found it very difficult to be open or receptive to the content of the formulation, he also recognised that although parts of the formulation were unexpected, other aspects were already familiar to him:



'Oh it was very hard to look at it at the beginning and to think that I had been thinking this way for so long. I wanted to just close my eyes and ignore what was in front of me. Some aspects were completely new to me. But once I started to revisit the formulation I started to realise that not all of the information was new, in fact I already knew some of it. But along the way I had stopped seeing the whole view – like selective vision'.

(Alex)

Another participant described the formulation as serving the role of reminding him of a part of himself which he had forgotten or perhaps lost. Whilst another participant found it more difficult to connect to his formulation, in part, due to him feeling that some aspects of the explanation were not accurate. Finally, Karolina felt that the written formulation, although helpful, seemed easier to talk about and develop, then to put into practice:

'How easy it is actually on the paper (laughter)...I think it does because you know I am a person who can talk and you can say whatever you want, but feeling it and practicing it, going into practise I think it would be much more harder than actually writing on a paper. Erm, but I did find it quite okay with it and useful for me to be honest'. (Karolina)

***'It was something I went away with' – sharing feelings with the clinician***

Four participants felt unable to share their feelings about the formulation with their clinician. One of these was a participant who had only received their formulation that week and therefore whether or not they would share their feelings with their clinician remains unknown. Two participants chose to keep their feelings to themselves. In response to being asked about whether the participant was able to share her feelings with her clinician, here one participant talks about leaving a session without discussing their feelings with their clinician:

'It was something I went away with really and thought about again'. (Lisa)

In contrast to this experience, Jill was able to create a more open dialogue about her explanation of her difficulties with her clinician. She felt that she was able to share her views with her clinician openly:

'Yes there was a lot of debate really actually, there was a lot of discussion and I felt like I could disagree or you know, I would try and take her points on board, erm, I felt like I was able to be very open, you know whether it was something I would consider or just give it a moment

or say "it is not for me", or "I don't really agree that that is the case" '. (Jill)

In part, it seems that Jill's experience was different to the other participants as she developed a reason for her behaviour and why she felt the way she had over the entire course of therapy.

***Revelation, empowerment or relief: from negative towards mixed feelings***

Participant reactions to the formulation appeared to reflect a sense that for some individuals the past was still difficult to talk about, even though a more positive focus for the future had been established. One participant described the process of incorporating the new information received from the formulation as a process which, though revelatory, would take time to integrate having realised they had been missing out on aspects of their life in the past:

'It's a bit of a revelation, once you start to realise that you really have been missing out, and that takes, that does take a long time, it's not a quick process at all I don't think'. (Sam)

One participant described a sense of newly developed awareness of things he had been ignoring. However, it seems that although the formulation helped him to see things he had previously been putting out his mind, though

he recognised the process of being able to be fully present and active in his life would take time. Another participant described feeling hesitant initially due to feelings of fear and doing things wrong whilst later feeling relief:

'Yes, I was a bit wary at first in case it wasn't for me or, in case, or I would do something wrong because that is what I have always been afraid of, but you don't really have anything to be afraid of because it does work and it is more of a relief when you realise that. (Penny)

For Arun, a sense of relief was expressed when thinking about his formulation, linked to an understanding of why he had been feeling the way he had and being able to challenge his previously held perspective on situations:

'so I think it more relief it was an understanding that, you know, that everything isn't so bad in the world but sometimes this thing can, you yourself through how you act and how you are going to act can create, how can I put it? A relief of understanding why you feel that way'.  
(Arun)

**A new journey: towards making sense of oneself**

It would appear that once clients had received their formulation or had discussed it with their clinician and begun to process associated feelings, they also began a journey of making a new sense of themselves. This may possibly be due to the emergence of a new or different self or a different way of responding to making sense of oneself, resulting from changes made during the intervention. With the exception of one participant, this process was apparent for all the participants even though the process of making sense of themselves may have differed. This observation appears to be less relevant for the one participant who had received a formulation very recently and thus her voice is less represented in this section. Through the process of making sense of themselves it seems that participants were also developing a changed perspective on their difficulties.

***re(Finding) me***

It seems that the formulation gave participants an insight into themselves which appeared to be positive for the majority of the participants in the study. In the following extract, one participant shares his experience where, although he at times still felt the same way as he did in the past, he now understands his feelings better. Additionally, recognising his feelings to be normal, which makes him feel more confident and equipped to deal with everyday situations in his life:

‘ I just seem to know myself much better now – I know,  
erm, why I feel the way I do and this has kind of made

## CLIENTS' EXPERIENCE OF FORMULATION

me more self-confident. I still feel the same feelings as before, but now I know that the way I feel is no different to other people. And I know the dark feelings will pass and when I feel that way it's like I now know I'll be able to deal with it no matter what life throws my way..

(Alex)

A second participant described rediscovering his identity, while also being able to think for himself again:

'Erm, day-to-day. I can think clearer. I was always saying 'I'm confused, I can't think straight, I can't work things out, I can't make decisions' and I couldn't. I genuinely thought I couldn't. But I think I'm now starting, you know, at the end of the tunnel sort of thing. I think I'm now starting to be able to think for myself again, which is nice. It's horrible to be, I suppose the best way to describe it is walking around in a fog, and you're lost. You are lost. You don't know which way to go, you don't, you lose your identity even. And to get that back is, it's lovely. To be able to think for yourself again, yeah, that's important'. (Sam)

Interestingly, another participant realised that although the formulation had not given him the answers to making his life easier, it had helped him realise

how he could be contributing to his own mood states. This newly developed understanding had also helped him to begin to look at things more positively which he continues to do in the present:

'I think it certainly helped me, erm, in that it made me realise that in a way I was creating my own moods, through how I perceived everything in life and how I react. It didn't give me, erm, it wasn't a route map to make things easier, because I, but after the realisation that how I could have got into this position, then it certainly helped me to start thinking about things a little bit more positively. I still do it now'. (Arun)

### ***New perspectives and skills***

All the participants described a changed perception of their difficulties. For some participants this came with new skills in managing emotions and thoughts. For example, Penny shared how her difficulties reduced due to a knowing how to manage and prevent low mood :

'My problems kind of shrunk for me, after grasping what to do, everything kind of seemed much better for me in myself and people around me. I started to get less anxious and more, I don't know, kind of bouncy instead because you feel low and you feel high, but I

## CLIENTS' EXPERIENCE OF FORMULATION

suppose you don't shut the lows out – you kind of prevent them instead.' (Penny)

In a similar vein, Lisa described how the formulation helped her to not only make sense of her feelings but how she could make changes to her thinking in order to be less negative:

'Yeah, it made a bit more sense actually, it seemed to make sense of why I'd been feeling the way I'd been feeling, and how I could change things and change the way I think about things to make a difference basically, instead of thinking negative all the time'. (Lisa)

One participant expressed having a greater feeling of control over their feelings and thoughts; describing being able to recognise that it was in fact he who makes the decisions within his life. Another participant realised that he does in fact have a choice by being aware of when he may be getting stuck and how he might respond differently at those times:

'It's made me realise that you don't have to be trapped, there are choices, you've just got to be aware of when you're trapping yourself and what opportunities, what sort of techniques, to try and get out of it'. (Sam)



## CLIENTS' EXPERIENCE OF FORMULATION

Finally another, participant spoke about how their perspective may have been inaccurate and that through the therapeutic process they were able to be more open to consider all the facts around them and influence their perspective:

[therapist] helped show me the whole picture and then I realised I had it all wrong. I was missing important information out and blaming myself for things that were not my fault. I guess, I'd say therapy really helped me to have a more balanced view of things. (Alex)

## **DISCUSSION**

### **Feeling trapped or restricted by depression**

All the participants described their thoughts and feelings prior to coming to therapy. This was unexpected by the interviewer, who had not directly asked questions about this aspect of participants' experience. The pervasive nature of symptoms described by participants appears consistent with descriptions in the current NICE guidelines for depression (NICE, 2010) as well as in a qualitative research paper on experience of depression conducted by Smith and Rhodes (2010).

### **The development of the formulation – from coming to my own conclusions to something the therapist developed**

For most participants, the formulation was developed through the client exploring their thoughts, feelings and life experiences with the clinician. Significant variability was observed, ranging from formulations being client developed with clinicians only providing validation, through to clinicians developing formulations more autonomously and sharing that with the client (participant). Participants appeared to feel understood and also more able to contribute when formulations were developed collaboratively over time. In contrast, the therapist-led formulation appeared to leave participants unable to share their feelings and uncertain of the accuracy of the formulation, suggesting that formulation development requires collaboration and time in order for clients to feel sufficiently comfortable to address perceived inaccuracies within the formulation.

Where the clinician was perceived as working to the client's pace, the participant reported mainly positive feelings towards the formulation process, such as being able to openly share feelings and to question aspects of the formulation which they did not agree with. This experience is consistent with research which suggests that clients seem to have a more positive experience of formulation when the formulation is developed collaboratively (McManus, Peerbhoy, Larkin and Clark, 2010). Furthermore, it highlights the importance of collaborative formulation development in clinical practice.

For one participant, it seemed that the process of receiving their formulation was particularly difficult. This may have been because the participant felt less connected to the formulation as they disagreed with some aspects of it. This may also be related to the fact that the participant had received a relatively brief intervention lasting only five sessions. Therefore, perhaps the participant felt less able to share their feelings with their clinician as they had not developed a 'good enough' therapeutic relationship within this time. Having received the formulation in the penultimate session, the participant felt that the clinician did not allow adequate time to review the original formulation before ending the therapeutic relationship. This experience may have specific significance for short term CBT interventions, and is a reminder for clinicians of the importance of allowing sufficient time both for the development of the therapeutic relationship and for a truly collaborative process of formulation development to occur. .

**From negative to mixed feelings: emotional reactions to the formulation during the therapeutic process**

It seems that participants initially struggled to connect with their formulation for many reasons. For some, it seemed that the formulation itself may have been technically difficult to comprehend due to unfamiliarity to CBT terminology. However, other participants were also debating whether the intervention would actually work for them. One aspect of this may have been the pervasive nature of negative automatic thoughts in depression making it difficult to initially consider change in thinking, feelings, behaviour and physiology. These feelings of initial doubt and questioning whether the intervention will work appear to be in line with Brown's (2008) model of 'client's view of client's experience of formulation'. However, an additional dimension to this experience was identified in the present study, with some participants struggling to relate to memories and feelings linked to difficult past experiences.

It is also clear that clinicians face a challenge at the beginning of therapy in terms of balancing empathy for the client's mixed feelings along with challenging the client within therapy so that they can begin to consider the potential benefits of trying something different. This finding, of initially experiencing negative feelings towards the formulation, is consistent with research by Pain, Chadwick and Abba's (2008), who found that forty per cent of participants reported negative feelings after receiving their formulation. It is noteworthy that they interviewed their participants relatively soon (2-3 weeks) after receiving a formulation and the responses appear to be in line

with the 'initial doubt' period highlighted in Brown's (2008) formulation model. Perhaps a richer picture of clients' experiences would have been gained if the researchers had interviewed their participants again at the end of their intervention. Such research may have captured a broader range of clients' views towards their formulation at different points of optimism or hopelessness.

Participants may have struggled to share their feelings with their clinician because they themselves were still processing how they felt, or perhaps because they did not feel able to do this due to the power differential or lack of trust between client and clinician at the beginning of therapy. This poses a difficulty for clinicians as they may be unsure of which aspects of the formulation their clients disagree with initially. Some participants in the present study expressed that they would not have continued with therapy if they could not agree with their formulation, indicating the need for clinicians to formulate in a sensitive way inviting clients to share their opinions. Clients being unable to connect to their formulation may suggest one possible explanation for early drop-out of CBT interventions

While initial responses to the formulations seemed to be predominantly negative, it seems that after some time processing their formulation and perhaps through having time to make changes in their life through their therapeutic intervention, participants felt more positive, even though the journey towards making changes felt challenging. Participants reported feeling liberated and enlightened which, compared to the entrapment they

felt prior to therapy, appeared to be a significant positive change. These subsequent feelings seem consistent with Townend and Grant's (2008) view that CBT formulation instils hope and optimism in clients.

### **A new journey towards understanding oneself**

It was anticipated that as a result of the formulation and intervention, participants would report an improvement in mood. However, what actually seemed to change was the way participants perceived themselves, their thoughts and feelings. Participants reported rediscovering lost identities and realised that difficult feelings pass. Additionally, participants were able to think more clearly and become more confident and self-accepting. It may be that these positive changes stemmed from having better strategies to manage negative thoughts and feelings, as well as a normalised perspective about their problems. It appears that this new perspective may have been just as important as the reduction of their depressive symptoms, pointing to a need for therapists to use a broader range of outcome measures to capture the effects of psychological interventions.

Along with a changed perspective and tools to manage negative thoughts and feelings, participants appeared to have regained control in their lives which perhaps empowered individuals who had previously felt stuck within depression.

The subordinate themes within this superordinate theme appeared to be closely linked, perhaps because new skills in managing depression are

necessary to the development of a positive perspective of oneself and the external world. Future research could perhaps focus on furthering our understanding the relationship between these factors as they may be key to successful outcomes in therapy.

### **Limitations**

The clinicians of all of the participants' were experienced Clinical Psychologists who primarily, though not necessarily exclusively, used the CBT approach in their therapeutic work. As a result it is unclear whether the findings of this study would have been the same if the clinicians had practiced purely from a CBT perspective. Furthermore, the nature and power of the therapeutic relationship was not considered within this study, for example the interpersonal style of the participants was not established which may have had a significant impact on the therapeutic process.

The length of depressive episode varied for each participant and the research was unable to establish whether duration or chronicity of the depressive episode impacts upon the experience formulation. In light of this limitation to the present study, future quantitative research where such variables can be controlled for is indicated. Additionally, clients may have experienced different levels of formulation and future studies could investigate this more closely by collecting data at multiple time points during therapy.

Finally, another limitation to the study may have been the use of a purposive sampling style. It remains possible that participants who were recruited were either clients who had responded well to a CBT approach or those who were perceived as having formed a good relationship with their clinician. Although some negative experiences were identified in the present study, it may also be that a sampling bias resulted in an over reporting of positive reactions of formulations.

### **Future research**

As a relatively new area for research there are several further aspects of formulation which could usefully be explored in future studies. Firstly, research exploring clients' experiences of formulation with other mental health difficulties other than depression would allow for comparison of findings with those of the present study. Secondly, clients' accounts could be explored at different time points after receiving their formulation, in order to explore similarities and difference across time. Thirdly, it may be helpful for a future study to specifically focus on individuals who have had negative experiences of formulation, in order to help guide clinical practice in this area. Finally, as an evidence base is gradually developed in the area of clients' experiences of formulation, further quantitative research with larger samples and using longitudinal designs is also indicated.



## REFERENCES

- Aston, R. (2009). A literature review exploring the efficacy of case formulations in clinical practice. What are the themes and pertinent issues? *The Cognitive Behaviour Therapist*, 2, 63-74.
- Beck, J. (1995). *Cognitive therapy: Basics and beyond*. New York: The Guildford Press.
- Boyatzis, R. (1998). *Transforming Qualitative Information: Thematic Analysis and Code Development*. London: Sage Publications Ltd.
- Braun, V. & Clarke, V. (2006). Using Thematic Analysis in Psychology. *Qualitative Research in Psychology*, 3, 77 – 101.
- Brocki, J. M., and Wearden, A. J. (2006). A critical evaluation of the use of interpretative phenomenological analysis (IPA) in health psychology. *Psychology and health*, 21, 87-108.
- Brown, H. (2008). *Clients' experience of formulation: The views of clients and therapists - A grounded theory exploration*. Unpublished D Clin Psy, Coventry University.
- Chadwick, P., Williams, C., and Mackenzie, J. (2003). Impact of case formulation in cognitive behaviour therapy for psychosis. *Behaviour Research and Therapy*, 41, 671-680.
- Department of Health. (2004). *The national service framework for mental health – five years on*. London: Department of Health.

- Dudley, R., and Kuyken, W. (2006). Formulation in cognitive-behaviour therapy. In L. Johnstone, and R. Dallos (Eds.). *Formulation in psychology and psychotherapy: Making sense of people's problems*. London: Routledge.
- Eells, T. D. (1997). *Handbook of psychotherapy case formulation*. New York: The Guildford Press.
- Flitcroft, A., James, I. A., Freston, M. (2007). Determining what is important in a good formulation. *Behavioural and Cognitive Psychotherapy*, 35, 325-33.
- Golofshani, N. (2003). Using reliability and validity in qualitative research. *The Qualitative Report*, 8, 597-607.
- Gray, J. C., and Grant, A. (2005). Cognitive behavioural therapy: helping the client find her voice. *Mental Health Practice*, 8, 34-37.
- Kinderman, P., and Lobban, F. (2000). Evolving formulations: sharing complex information with clients. *Behavioural and Cognitive Psychotherapy*, 28, 307-310.
- Kuyken, W. (2006). Evidence-based case formulation: is the emperor clothed? In N. Tarrier. (Eds). *Case Formulation in Cognitive Behaviour Therapy*. London: Routledge.
- Malan, D.H. (1995). *Individual Psychotherapy and the Science of Psychodynamics*. London: Butterworth.

- McManus, F., Peerbhoy, D., Larkin, M., and Clark, D. M. (2010). Learning to change a way of being: An interpretative phenomenological perspective on cognitive therapy for social phobia. *Journal of Anxiety Disorders, 24*, 581-589.
- National Institute for Health and Clinical Excellence. (2010). *Depression – the treatment and management of depression in adults (updated edition)*. Leicester: The British Psychological Society and The Royal College of Psychiatrists.
- Pain, C. M., Chadwick, P., and Abba, N. (2008). Clients' experience of case formulation in cognitive behaviour therapy for psychosis. *British Journal of Clinical Psychology, 47*, 127 -138.
- Smith, J. & Eatough, V. (2007). Interpretive phenomenological analysis. In E. Lyons & A. Coyle (Eds.). *Analysing Qualitative Data in Psychology* (pp. 34 – 50). London: Sage
- Tarrier, N. (2006). An introduction to case formulation and its challenges. In N. Tarrier. (Eds). *Case formulation in cognitive behavior therapy: the treatment of challenging and complex cases*. Hove: Routledge.
- Townend, M., and Grant, A. (2008). Assessment in CBT: the ideographic approach. In A. Grant, M. Townend, Mills, J and Cockx, A. (Eds.). *Assessment and case formulation in cognitive behavioural therapy*. Trowbridge: The Cromwell Press.

**Table 2.1. Participant summary of pseudonyms, age, ethnicity, number of CBT sessions and duration of depressive symptoms prior to CBT.**

<b>Participant pseudonym</b>	<b>Age</b>	<b>Ethnicity</b>	<b>Duration of depressive symptoms prior to CBT</b>
Lisa	28	White British	12 years
Sam	54	White British	5 years
Jill	42	White British	3 years
Arun	32	Asian British	1 year
Karolina	35	White Other	1.5 years
Penny	19	White British	6 years
Alex	22	Asian other	2 years

**Table 2.2. Superordinate and subordinate themes with participant representation in parenthesis**

<b>Superordinate themes</b>	<b>Subordinate themes</b>
Feeling trapped or restricted by depression (7/7)	
The development of the formulation – from coming to my own conclusions to	-Exploring thoughts, feelings, action and previous experiences – developing the formulation (7/7)

CLIENTS' EXPERIENCE OF FORMULATION

<p>something the therapist developed (7/7)</p>	<p>-Collaboration, directive or coming up with it myself: views on who developed the formulation (7/7)</p>
<p>From negative to mixed feelings: emotional reactions to the formulation during the therapeutic process (7/7)</p>	<p>-Receiving a formulation is difficult (7/7)</p> <p>-‘It was something I went away with’ –sharing feelings with the clinician (7/7)</p> <p>-Revelation, empowerment or relief: from negative towards mixed feelings (5/7)</p>
<p>A new journey: towards making a new sense of oneself (7/7)</p>	<p>- (re)Finding me (6/7)</p> <p>-‘New perspectives and skills (7/7) (5/7)</p>