

Carbohydrate Mouth Rinse Improves Morning High-Intensity Exercise Performance

Clarke, N, Hammond, S, Kornilios, E & Mundy, P

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Clarke, N, Hammond, S, Kornilios, E & Mundy, P 2017, 'Carbohydrate Mouth Rinse Improves Morning High-Intensity Exercise Performance' *Journal of Sports Sciences*, vol (in press), pp. (in press)
<https://dx.doi.org/>

DOI

ISSN 0264-0414

ESSN 1466-447X

Publisher: Taylor and Francis

This is an Accepted Manuscript of an article published by Taylor & Francis in Journal of Sports Sciences on 5th June 2017, available

online: <http://www.tandfonline.com/doi/full/10.1080/17461391.2017.1333159>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

Article Title: Carbohydrate Mouth Rinse Improves Morning High-Intensity Exercise Performance.

Running Head: Morning high-intensity exercise and CHO rinsing

Authors: Neil D. Clarke., Scott Hammond, Evangelos Kornilios, and Peter D. Mundy.

Author Affiliation: School of Life Sciences, Faculty of Health and Life Sciences, Coventry University, Coventry, UK.

Contact details for corresponding author: Neil D. Clarke, School of Life Sciences, Faculty of Health Health and Life Sciences, Coventry University, Priory Street, CV1 5FB, UK.

Telephone: (44) 24776 65 8559

Fax: (44) 2476 88 7688

Email: Neil.clarke@coventry.ac.uk

Abstract

Oral carbohydrate rinsing has been demonstrated to provide beneficial effects on exercise performance of durations of up to one hour. The aim of the present study was to investigate the effects of carbohydrate mouth-rinsing on morning high-intensity exercise performance. Following institutional ethical approval and familiarisation, twelve healthy males (mean±SD age: 23±3 years, height: 175.5±7.4 cm, body mass: 75.4±7.5 kg) participated in this study. Countermovement jump (CMJ) height, isometric mid-thigh pull peak force, 10 m sprint time, and bench press and back squat repetitions to failure were assessed following carbohydrate (CHO) and placebo (PLA) rinsing or a control condition (CON). All testing took place at 07:30 following an eleven hour overnight fast. Performance of CMJ height (CHO: 39±7 cm; PLA: 38±7 cm; CON: 36±6 cm; $P=0.003$, $\eta_p^2=0.40$), 10 m sprint time (CHO: 1.78±0.07 s; PLA: 1.81±0.07 s; CON: 1.85±0.05 s; $P=0.001$, $\eta_p^2=0.47$), the number of bench press (CHO: 25±3; PLA: 24±4; CON: 22±4; $P<0.001$, $\eta_p^2=0.55$) and squat (CHO: 31±4; PLA: 29±5; CON: 26±6; $P<0.001$, $\eta_p^2=0.70$) repetitions and mean felt arousal (CHO: 5±1; PLA: 4±0; CON: 4±0; $P=0.009$, $\eta_p^2=0.25$) improved following carbohydrate rinsing. However, isometric mid-thigh pull peak force was unchanged (CHO: 2262±288 N; PLA: 2236±354 N; CON: 2212±321 N; $P=0.368$, $\eta_p^2=0.08$). These results suggest that oral carbohydrate rinsing solution significantly improved the morning performance of CMJ height, 10 m sprint times, bench press and squat repetitions to failure and felt arousal, although peak force during an isometric mid-thigh pull, RPE and heart rate were unaffected.

Keywords: Maltodextrin, Oral receptors, Arousal

Introduction

Oral rinsing of a carbohydrate solution prior to, and during, exercise can improve performance without altering metabolic responses (e.g. Carter, Jeukendrup & Jones, 2004; Kasper et al. 2015; Rollo, Williams, Gant & Nute, 2008; Rollo, Homewood, Williams, Carter, & Goosey-Tolfrey, 2015). The underlying mechanism is believed to relate to the presence of carbohydrate in the mouth inducing increased brain activity within the orbitofrontal cortex (De Pauw et al. 2015). In addition, Gant, Stinear and Byblow (2010) demonstrated that carbohydrate ingestion can immediately affect performance by increasing corticomotor excitability through non-sweet receptors in the oral cavity area and counteract the decreasing motor activity. Similar findings were reported by Chambers, Bridge and Jones (2009), in that, independent of sweetness, carbohydrate can activate brain regions related to reward and motor control, possibly through non-sweet taste receptors found in the mouth.

Several 30-minute to 1-hour time trial studies exist where the effect of carbohydrate mouth-rinsing has been investigated during cycling (Chambers et al. 2009; Lane, Bird, Burke & Hawley, 2013; Pottier, Bouckaert, & Derave, 2010) and running (Clarke, Thomas, Kagka, Ramsbottom, & Delextrat. 2016b; Rollo et al. 2008; Rollo, Williams, & Nevill, 2011). However, most of the literature focuses on endurance-based exercise, so evidence of possible ergogenic benefits on high-intensity and resistance exercise is lacking. Furthermore, those studies investigating the effect of carbohydrate rinsing on high intensity exercise have produced inconsistent results (Beaven, Maulder, Pooley, Kilduff, & Cook, 2013; Bortolotti, Pereira, Oliveira, Cyrino, & Altimari, 2013; Chong, Guelfi, & Fournier, 2011; Dorling and Earnest, 2013; Kasper et al. 2016; Phillips, Findlay, Kavaliuskas, & Grant, 2014; Přibyslavská

et al. 2016). Carbohydrate rinsing has been shown to induce no significant improvements in performance whilst completing one repetition maximum or muscular endurance (Clarke, Kornilios, & Richardson, 2016a; Dunkin and Phillips, 2017; Painelli et al. 2011), but beneficial in attenuating torque reduction during a series of maximal voluntary contractions (Jensen, Stellingwerff, & Klimstra, 2015). Similarly, Rollo et al. (2015) and Dorling and Earnest (2013) demonstrated that sprint performance during the Loughborough Intermittent Shuttle Running Test was not significantly improved following carbohydrate rinsing, whereas Beaven et al. (2013) reported that mean and peak power output were increased during the first of five six-second sprints. However, information relating to other high-intensity activities is limited, although Rollo et al. (2015) speculated that in the presence of a perceived impending increase in fuel supply during carbohydrate rinsing, the balance of excitation and inhibition of the brain's motor cortex may be altered in favour of excitation, allowing athletes to improve their performance. Furthermore, although the effect size of this improvement on performance may be trivial to small, it is likely to be practically significant (Peart, 2016).

One possible explanation for inconsistent findings is the nutritional status of the participants, with participants arriving at the laboratory after an overnight fast (Jensen et al. 2015) or their usual pre-exercise meal (Clarke et al. 2016a). Although the effects of carbohydrate rinsing appear more profound after an overnight fast, there is still evidence to support beneficial effects after the ingestion of a meal (Fares and Kayser, 2011; Jeukendrup, 2013). Furthermore, athletes training in the early morning frequently choose to begin activity without eating, with some athletes preferring to consume water over carbohydrate during the session. Such choices may be explained by the reported gastrointestinal problems and distress reported by participants regarding carbohydrate consumption before and during exercise (de Oliveira, Burini, & Jeukendrup, 2014), which are likely due to stress of the abdominal organs caused by the

carbohydrate and movements during exercise (Van Nieuwenhoven, Brouns, & Kovacs, 2005). Therefore, the aim of the present study was to investigate the effects of carbohydrate mouth-rinsing on morning high-intensity exercise performance.

Methods

Following institutional ethical approval and familiarisation, twelve healthy males (mean \pm SD age: 23 ± 3 years, height: 175.5 ± 7.4 cm, body mass: 75.4 ± 7.5 kg) who trained three to four times a week for between eight and 12 months, and therefore met the intermediate resistance training experience classification as per the National Strength and Conditioning Association (Sheppard and Triplett, 2015) participated in this study. Participants were instructed to avoid caffeine ingestion for a minimum of 12 hours prior to the trials and refrain from strenuous exercise for 24 hours. In addition, a 24-hour dietary record was completed by each participant during the familiarisation session; it was then photocopied and handed back to the participants so that the same diet could be repeated for subsequent trials. All procedures were undertaken in accordance with the Declaration of Helsinki.

A randomised, Latin-square, crossover, placebo-controlled design was employed during this study. Each participant attended the laboratory on four occasions. The first session was to establish one repetition maximum (1-RM) for the squat and bench press, and to allow familiarisation with the exercises. Following a standardised warm-up, squat and bench press 1-RM testing was then conducted following the National Strength and Conditioning Association guidelines (McGuigan, 2015). The 1RM testing began with a warm-up at 50% of their

predicted 1RM. The load was then increased to the predicted 1RM for the first attempt. If that attempt was successful, five minutes of rest were given after which another 1RM was attempted; this sequence was repeated until the 1RM attempt was unsuccessful or the subject refused to continue; the highest load successfully lifted was recorded as the 1RM value. The subsequent three sessions were to complete the exercise battery under each condition and were separated by seven days to allow recovery.

All testing took place at 07:30 after the ingestion 500 mL of water 60 minutes prior to arrival and following an eleven hour overnight fast. On arrival participants completed a warm-up which involved five minutes of sub-maximal cycling on a cycle ergometer at 150 W and then performed no more than two sets of 12 repetitions at a self-selected, light intensity for the squat and bench press. The warm-up weights selected during the initial trial were recorded and repeated prior to subsequent trials. The participants then orally rinsed either 25 mL of carbohydrate or taste-matched-placebo solution for ten seconds before each exercise or no fluid. The participants performed a series of exercise tests and were monitored on the performances of countermovement jump (CMJ) height, isometric mid-thigh pull peak force, 10 m sprint times, and bench press and back squat repetitions to failure at 60% 1-RM. Each exercise, excluding the bench press and squat performance, was performed three times and the best result recorded. Furthermore, there was a five-minute rest period between each exercise.

The participants performed three vertical countermovement jump with the arms held akimbo on a force platform (AMTI AccuPower, Watertown, MA, USA) sampling at 400 Hz. All participants were instructed to jump as high as possible and jump height was calculated using the velocity at take-off (Hatze, 1998) (Technical error of measurement (TEM) = 0.68). The

participants then performed a maximal isometric mid-thigh pull on a portable standing force plate (AMTI AccuPower, Watertown, MA, USA) sampling at 400 Hz. For this test a custom designed rig was used that allowed a fixed bar to be adjusted to meet the required height for each participant with the stationary bar positioned in the middle of their thigh. The participants used lifting straps to improve the grip and were instructed to get in a comfortable and stable position with self-selected knee and hip angles (Comfort, Jones, McMahon, & Newton, 2015). Once in this position participants performed a maximal isometric pull for five seconds. All participants were instructed to pull as hard and fast as possible and there was a 90-second break recovery period between each repetition. Peak force was identified as the highest instantaneous value observed during the trial (TEM = 6.36).

The participants then completed a maximal 10 m sprint from a standing position. The time was measured using an infrared timing system (Smart Speed, Fusion Sports, Australia) and all participants were instructed to run as fast as possible. Each participant performed this task three times with 90 s between each trial and the fastest time recorded (TEM = 0.14). After five minutes, the participants began the 60% to failure bench-press protocol (TEM = 0.79). A five-minute rest period was allowed before participants began the squat protocol (TEM = 1.06). For both exercises a metronome was used to provide a cadence of two seconds for both the eccentric and concentric phases of movement.

The felt arousal scale (Svebak and Murgatroyd, 1985), used to monitor arousal throughout the trials, and rating of perceived exertion (RPE) (Borg, 1973) was recorded immediately after each of the exercises. Heart rate (HR) (Polar Electro Oy, Kempele, Finland) was recorded

immediately before and after administration of the treatment, after the warm up, and following each exercise.

Rinsing protocol

Prior to each exercise either 25 mL of a 6% carbohydrate solution (maltodextrin: My Protein, Manchester, UK) (CHO) or water (PLA) were rinsed around the buccal cavity for ten seconds. Participants then expectorated the solution back into the plastic cup before starting the exercise protocol. All solutions were flavoured with orange (No added sugar orange squash, Sainsbury's, London, UK). During the remaining session, no solution was rinsed (CON).

Statistical analysis

Data are reported as the mean \pm the standard deviation (SD). The Shapiro-Wilk test was used to confirm normal distribution. Furthermore, due to variation between participants and the suggestion that carbohydrate mouth rinsing may be worth investigating on an individual basis (Peat, 2016), individual performance responses are also presented. A one-way analysis of variance (ANOVA) with repeated measures was used to compare all data except for RPE, felt arousal and heart rate, which were analysed with two-way ANOVA with repeated measures. Sphericity was analysed by Mauchly's test of sphericity followed by the Greenhouse-Geisser adjustment where required. Where any differences were identified, 95% confidence intervals and pairwise comparisons with Bonferroni correction were used to show where they lay. All statistical procedures were conducted using IBM SPSS Statistics for Windows, Version 20.0

(Armonk, NY: IBM Corp.). Sample size was calculated using G*Power software (version 3.1.9.2, Franz Faul, Universitat Kiel, Dusseldorf, Germany) for repeated measures ANOVA for detecting a small effect size (0.3) with α as 0.05 and a 1- β error probability of 0.8 revealed that a sample size of 12 participants was required. Furthermore, effect sizes using partial eta squared (η_p^2) and Cohen's d were calculated, which were defined as trivial (0-0.19), small (0.20-0.49), moderate (0.50-0.79) or large (≥ 0.80) (Cohen, 1992).

Results

A small increase in CMJ height during both rinsing conditions was observed ($F_{(2,22)}=7.395$; $P=0.003$, $\eta_p^2=0.40$; Figure 1a). Jump height was significantly higher following CHO (95%CI; 1.2, 6.2; $P=0.008$; $d=0.54$) and PLA (95%CI; 0.5, 5; $P=0.047$; $d=0.36$) compared with CON. Furthermore, only a trivial difference in jump height between CHO and PLA (95%CI; -0.1, 2.7; $P=0.070$; $d=0.18$) was observed. Individual CMJ results (Figure 2a) show that 67% of the participants had a greater CMJ height in the CHO trial. A small decrease in the time to complete the 10 m sprint following CHO and PLA was observed ($F_{(2,22)}=9.683$; $P=0.001$, $\eta_p^2=0.47$; Figure 1b). Sprint time was significantly faster following CHO compared with CON (95%CI; -0.1, -0.3; $P=0.003$; $d=1.14$) and PLA (95%CI; -0.6, -0.01; $P=0.024$; $d=0.45$) trials. Furthermore, sprint time was significantly faster following PLA than CON (95%CI; -0.8, -0.003; $P=0.036$; $d=0.65$). Individual times (Figure 2b) show that 58% of the participants were faster following CHO rinsing.

Only trivial differences in the peak force produced during the isometric mid-thigh pull were observed ($F_{(2,22)}=1.048$; $P=0.368$, $\eta_p^2=0.08$; Figure 1c). There were no significant differences between CHO and CON (95%CI; -26, 125; $P=0.174$; $d=0.16$), CHO and PLA (95%CI; -70, 122; $P=0.564$; $d=0.10$) and PLA and CON (95%CI; -24, 71.4; $P=0.294$; $d=0.08$). A moderate increase in the number of bench press repetitions performed during the rinsing trials compared with CON was observed ($F_{(2,22)}=13.253$; $P<0.001$, $\eta_p^2=0.55$; Figure 1d). The number of bench press repetitions following CHO was significantly greater than PLA (95%CI; 0.1, 3.2; $P=0.039$; $d=0.46$) and CON (95%CI; 2.1, 4.4; $P<0.001$; $d=0.86$). Furthermore, a significant greater number of bench press repetitions following PLA compared with CON was observed (95%CI; 0.2, 3; $P=0.029$; $d=0.39$). Individual bench press repetitions to failure (Figure 2d) show that 75% of the participants performed a greater number of repetitions following CHO. Moderately more squat repetitions were performed following CHO and PLA compared with CON ($F_{(2,22)}=25.729$; $P<0.001$, $\eta_p^2=0.70$; Figure 1e). A significantly higher number of squat repetitions were observed following CHO compared with PLA (95%CI; 3.9, 10.6; $P=0.001$; $d=0.45$) and CON (95%CI; 5.4, 12.2; $P<0.001$; $d=0.97$). In addition, significantly more squats repetitions were performed following PLA compared with CON (95%CI; 0.2, 3; $P=0.029$; $d=0.53$). Individual squat repetitions to failure (Figure 2e) show that 75% of the participants performed a greater number of repetitions following CHO.

A small interaction between condition and time was observed for felt arousal ($F_{(12, 132)}=1.046$; $P<0.001$, $\eta_p^2=0.25$; Table 1). A moderate increase in felt arousal occurred throughout the protocol ($F_{(6,66)}=6.713$; $P<0.001$, $\eta_p^2=0.38$) and a moderate difference between conditions was observed ($F_{(2,22)}=18.295$; $P<0.001$, $\eta_p^2=0.63$). Felt arousal was greater following CHO (95%CI; 0.5, 1.3; $P<0.001$; $d=2.04$) and PLA (95%CI; 0.2, 1.2; $P=0.009$; $d=1.30$) compared

with CON, although no significant difference between CHO and PLA (95%CI; -0.2, 0.6; $P=0.629$; $d=0.46$) was observed. A large increase in RPE (Table 1) was observed during the exercise protocol ($F_{(4,4)}=282.563$; $P<0.001$, $\eta_p^2=0.96$), although only trivial differences were observed between conditions ($F_{(2,22)}=0.840$; $P=0.445$, $\eta_p^2=0.07$). Similarly, a large increase in heart rate (Table 1) occurred during the protocol ($F_{(11,121)}=204.961$; $P<0.001$, $\eta_p^2=0.95$). However, there were no significant differences between conditions ($F_{(2,22)}=1.100$; $P=0.351$, $\eta_p^2=0.01$).

Discussion

The key finding of the present study was that oral carbohydrate rinsing significantly improved the morning performance of CMJ height, 10 m sprint times, bench press and squat repetitions to failure and felt arousal, without changes in perceived exertions or heart rate. Furthermore, small to moderate performance improvements in countermovement jump, 10 m sprint time, and bench and squat repetitions to failure were observed following rinsing with the placebo when compared with the control condition. The results of this study concur with those of Gant et al. (2010), who indicated similar effects following orally rinsing carbohydrate on isometric elbow flexion, Jensen et al (2015) who reported decreased torque attenuation following carbohydrate mouth rinsing in a fatigued state, and Peart (2016), who concluded a trivial to small overall positive effect of CHO mouth-rinsing on performance. Furthermore, it has been demonstrated that CHO rinsing improves peak and mean power output during sprinting (Beaven et al. 2013; Phillips et al. 2014). The proposed mechanism suggests that carbohydrate mouth-rinsing activates regions in the brain related to motor output and pleasure/reward

(Chambers et al. 2009) and increasing arousal. Similarly, De Pauw et al. (2015) reported that the presence of carbohydrate within the mouth activates the reward centres of the brain, due to a direct link between the buccal mucosa and the brain (Nicolazzo, Reed, & Finnin, 2003). Furthermore, Gant et al. (2010) indicated orally rinsing carbohydrate immediately increased the excitability of the corticomotor pathway. Therefore, the presence of carbohydrate during the rinsing process, the balance of excitation and inhibition of the brain's motor cortex may be altered in favour of excitation, allowing athletes to improve their performance.

Peak isometric force was evaluated using isometric mid-thigh pull performance, although no differences between trials were observed. Several previous studies involving high-intensity exercises (Chong et al, 2011), multiple sprints (Dorling and Earnest, 2013), and maximum strength (Painelli et al, 2011) concluded that carbohydrate mouth rinsing provided did not improve performance. In contrast, Gant et al. (2010) reported that isometric contraction force during elbow flexion increased following carbohydrate mouth-rinsing. One potential explanation for the present study demonstrating no improvement during the isometric mid-thigh pull could be that when exercise intensity elicits near maximal effort, despite increases in felt arousal following carbohydrate rinsing, as seen in the present study, it creates a "ceiling effect" which makes any appreciable differences between conditions extremely difficult to distinguish (Beaven et al. 2013, Clarke et al. 2016a). Furthermore, it is possible that the presence of carbohydrate stimulated the reward and/or motivation centres in the brain, but this stimulus was insufficient to affect maximal strength performance (Painelli et al. 2011), as strength-trained individuals usually present little to no neural activation deficits (Ahtiainen and Hakkinen, 2009).

The present study included a fasting period of eleven hours prior to testing consistent with the protocol used by Haase, Cerf-Ducastel, and Murphy (2009) and Turner, Byblow, Stinear, and Gant (2014). However, Beelen et al. (2009) suggested overnight fasting reduces the validity of the findings because in a practical setting athletes typically ingest a high carbohydrate meal two hours prior to competition and in many situations, ingest the carbohydrate source. In contrast, Přibyslavská et al. (2016) reported female soccer players commonly refuse to eat before early morning training or competitive matches and often ingested water rather than carbohydrate. Therefore, the findings of the present study may be of interest to those athletes who commonly train in a fasted state in the early morning. This nutritional intervention could be practically used as a performance enhancement without the gastrointestinal discomfort often associated high-intensity exercise (de Oliveira et al. 2014). Furthermore, Kasper et al, (2016) highlighted the method may also be practical for those athletes that integrate phases of carbohydrate restrictions into their training programmes. Consequently, carbohydrate mouth rinsing has recently been incorporated into the nutritional guidelines for short higher intensity exercise (i.e. <1 hour) where glycogen is not a limiting factor for performance (Jeukendrup, 2013; Stellingwerff and Cox, 2014).

The present study employed a non-rinse control trial that has identified a potential placebo effect. Small to moderate performance improvements in countermovement jump, 10 m sprint time, and bench and squat repetitions to failure were observed after orally rinsing a non-carbohydrate solution. Gam, Guelfi, and Fournier (2013) suggested that a placebo effect of carbohydrate rinsing cannot be excluded. In support of this proposal the present study observed that when compared with the control condition, mouth rinsing with a placebo solution caused a large increase in felt arousal. This occurrence may at least partially explain the improved performance observed in the placebo trial as Kerr (1997) reports that relatively high levels of

felt arousal are a feature of successful performance. One suggestion for this occurrence in the present study is that both solutions were flavoured with artificial sweeteners. However, Chambers et al. (2009) demonstrated that the artificial sweeter placebo caused no effect to areas of the brain such as the anterior cingulate cortex and ventral striatum. Consequently, the effect of the mouth rinse itself may provide ergogenic benefits (Gam et al. 2013). Therefore, due to the potential placebo effect demonstrated in the present study, it would be recommended that future research incorporates a non-rinse control trial.

This study is not without limitations. Despite an inclusion criteria, large standard deviations are evident for some variables. The reason for this is primarily attributed to the variability of athletic standards amongst the participants, which had implications for all recorded measures. Ideally, a more homogeneous population would have been recruited thus avoiding a large range in characteristics and abilities which can result in a greater increase in 'noise' within the data. In addition, it is unknown whether a full familiarization of the 1RM protocols would have impacted on the estimation of the maximal dynamic strength and subsequent weight lifted by the participants during the main trials. However, Comfort and McMahon (2015) demonstrated the back squat to be highly reliable. Finally, it is acknowledged that variable such as CMJ and sprint performance demonstrate within-subject variation. However, TEM was calculated and excluding the bench press and squat performance, all tests was performed three times with the best result recorded.

In conclusion, these results suggest that following an overnight fast, oral rinsing with a carbohydrate solution significantly improved the morning performance of CMJ height, 10 m sprint times, bench press and squat repetitions to failure and felt arousal. However, peak force

during an isometric mid-thigh pull, RPE and heart rate were unaffected, possibly due to the nature of the exercise causing a "ceiling effect". Furthermore, a placebo effect of carbohydrate rinsing cannot be excluded.

References

Ahtiainen, J.P., & Hakkinen, K. (2009). Strength athletes are capable to produce greater muscle activation and neural fatigue during high-intensity resistance exercise than nonathletes. *Journal of Strength and Conditioning Research*, 23, 1129-1134. doi:10.1519/JSC.0b013e3181aa1b72

Beaven, C.M., Maulder, P., Pooley, A., Kilduff, L., & Cook, C. (2013). Effects of caffeine and carbohydrate mouth rinses on repeated sprint performance. *Applied Physiology, Nutrition, and Metabolism*, 38, 633-637. doi: 10.1139/apnm-2012-0333.

Beelen, M., Berghuis, J., Bonaparte, B., Ballak, S.B., Jeukendrup, A.E., & Van Loon, L.J. (2009). Carbohydrate mouth rinsing in the fed state, lack of enhancement of time-trial performance. *International Journal of Sport Nutrition & Exercise Metabolism*, 19, 400-409. doi:10.1123/ijsnem.19.4.400.

Borg, G. (1973). Perceived exertion, A note on history and methods. *Medicine and Science in Sport and Exercise*, 5, 90-93. doi: 10.1249/00005768-197300520-00017.

Bortolotti, H., Pereira, L.A., Oliveira, R.S., Cyrino, E.S., & Altimari, L.R. (2013). Carbohydrate mouth rinse does not improve repeated sprint performance. *Revista Brasileira de Cineantropometria & Desempenho Humano*, *15*, 639-645. doi; 10.5007/1980-0037.2013v15n6p639.

Carter, J.M., Jeukendrup, A.E., & Jones, D.A. (2004). The Effect of Carbohydrate Mouth Rinse on 1-h Cycle Time Trial Performance. *Medicine and Science in Sport and Exercise*, *36*, 2107-2111. doi: 10.1249/01.MSS.0000147585.65709.6F.

Chambers, E.S., Bridge, M.W., & Jones, D.A. (2009). Carbohydrate sensing in the human mouth, effects on exercise performance and brain activity. *Journal of Physiology*, *587*, 1779-1794. doi:10.1113/jphysiol.2008.164285.

Chong, E., Guelfi, K.J., & Fournier, P.A., (2011). Effect of a carbohydrate mouth rinse on maximal sprint performance in competitive male cyclists. *Journal of Science and Medicine in Sport*, *14*, 162-167. doi: 10.1016/j.jsams.2010.08.003.

Clarke, N.D., Kornilios, E., & Richardson, D.L. (2016a) Carbohydrate and Caffeine Mouth Rinses Do Not Affect Maximum Strength and Muscular Endurance Performance. *Journal of Strength and Conditioning Research*, *11*, 108-115. doi: 10.1519/JSC.0000000000000945.

Clarke, N.D., Thomas, J.R., Kagka, M., Ramsbottom, R., & Delextrat, A., (2016b). No Dose-Response Effect of Carbohydrate Mouth Rinse Concentration on 5 Km Running Performance in Recreational Athletes. *Journal of Strength and Conditioning Research*, Epub ahead of print. doi: 10.1519/JSC.0000000000001531.

Cohen, J. (1992). A Power Primer. *Psychological Bulletin*, 112, 155-159. doi: 10.1037/0033-2909.112.1.155.

Comfort, P., & McMahon, J.J. (2015) Reliability of maximal back squat and power clean performances in inexperienced athletes. *Journal of Strength and Conditioning Research*, 29, 3089-3096. doi: 10.1519/JSC.0000000000000815.

Comfort, P., Jones, P.A., McMahon, J.J., & Newton, R. (2015). Effect of knee and trunk angle on kinetic variables during the isometric mid-thigh pull; test-retest reliability. *International Journal of Sports Physiology and Performance*, 10, 58-63. doi: 10.1123/ijsp.2014-0077.

de Oliveira, E.P., Burini, R.C., & Jeukendrup, A., (2014). Gastrointestinal complaints during exercise; prevalence, etiology, and nutritional recommendations. *Sports Medicine*, 44, S79-S85. doi: 10.1007/s40279-014-0153-2.

De Pauw, K., Roelands, B., Knaepen, K., Polfliet, M., Stiens, J., & Meeusen, R. (2015). Effects of caffeine and maltodextrin mouth rinsing on P300, brain imaging and cognitive performance. *Journal of Applied Physiology*, *118*, 776-782. doi, 10.1152/jappphysiol.01050.2014.

Dorling, J.L., & Earnest, C.P. (2013). Effect of carbohydrate mouth rinsing on multiple sprint performance. *Journal of the International Society of Sports Nutrition*, *10*, 41. doi: 10.1186/1550-2783-10-41.

Dunkin, J., & Phillips, S.M. (2017). The effect of a carbohydrate mouth rinse on upper body muscular strength and endurance. *Journal of Strength and Conditioning Research*, Epub ahead of print. doi:10.1519/JSC.0000000000001668

Fares, E.J., & Kayser, B. (2011). Carbohydrate mouth rinse effects on exercise capacity in pre- and postprandial states. *Journal of Nutrition and Metabolism*, 385962. doi:10.1155/2011/385962

Gam, S., Guelfi, K.J., & Fournier, P.A. (2013) The presence of carbohydrate in a mouth rinse solution opposes the detrimental effect of mouth rinsing during cycling time trials. *International Journal of Sport Nutrition & Exercise Metabolism*, *3*, 48–56. doi:10.1123/ijsnem.23.1.48.

Gant, N., Stinear, C., & Byblow, W. (2010). Carbohydrate in the mouth immediately facilitates motor output. *Brain Research*, *1350*, 151-158. doi:10.1016/j.brainres.2010.04.004.

Haase, L., Cerf-Ducastel, B., & Murphy, C., (2009). Cortical activation in response to pure taste stimuli during the physiological states of hunger and satiety. *Neuroimage*, *44*, 1008-1021. doi: 10.1016/j.neuroimage.2008.09.044.

Hatze, H. (1998) Validity and reliability of methods for testing vertical jumping performance. *Journal of Applied Biomechanics*, *14*, 127-140. doi:10.1123/jab.14.2.127

Jensen, M., Stellingwerff, T., & Klimstra, M. (2015). Carbohydrate mouth rinse counters fatigue related strength reduction. *International Journal of Sport Nutrition & Exercise Metabolism*, *25*, 252-261. doi: 10.1123/ijsnem.2014-0061.

Jeukendrup, A.E. (2013). Oral Carbohydrate Rinse, Placebo or Beneficial? *Current Sports Medicine Reports*, *12*, 222-227. doi:10.1016/j.nut.2004.04.017.

Kasper, A.N., Cocking, S., Cockayne, M., Barnard, M., Tench, J., Parker, L., McAndrew, J., Langan-Evans, C., Close, G.L., & Morton, J.P. (2016). Carbohydrate mouth rinse and caffeine improves high-intensity interval running capacity when carbohydrate restricted. *European Journal of Sport Science*, *16*, 560-568. doi, 10.1080/17461391.2015.1041063.

Lane, S.C., Bird, S.R., Burke, L.M., & Hawley, J.A. (2013). Effect of a carbohydrate mouth rinse on simulated cycling time-trial performance commenced in a fed or fasted state. *Applied Physiology, Nutrition, and Metabolism*, 38, 134-139. doi:10.1139/apnm-2012-0300.

Kerr, J.H. (1997). *Motivation and emotion in sport: Reversal theory*. Hove, Psychology Press.

McGuigan, M. (2015). Principles of Test Selection and Administration. In G.G. Haff and N.T. Triplett (Eds.), *Essentials of Strength Training and Conditioning (4th ed.)* (pp.249-258). Champaign, IL, Human Kinetics.

Nicolazzo, J.A., Reed, B.L., & Finnin, B.C. (2003). The effect of various in vitro conditions on the 477 permeability characteristics of the buccal mucosa. *Journal of Pharmaceutical Sciences*, 92, 2399-2410. doi: 10.1002/jps.10505.

Painelli, V.S., Roschel, H., Gualano, B., Del-Favero, S., Benatti, F.B., Ugrinowitsch, C., Tricoli, V., & Lancha, A.H. Jr. (2011). The effect of carbohydrate mouth rinse on maximal strength and strength endurance. *European Journal of Applied Physiology*, 111, 2381-2386. doi: 10.1007/s00421-011-1865-8.

Peart, D. (2016) Quantifying the Effect of Carbohydrate Mouth Rinsing on Exercise Performance. *Journal of Strength and Conditioning Research*, Epub ahead of print. doi: 10.1519/JSC.0000000000001741.

Phillips, S.M., Findlay, S., Kavaliauskas, M., & Grant, M.C., (2014). The influence of serial carbohydrate mouth rinsing on power output during a cycle sprint. *Journal of Sports Science and Medicine*, 13.

Pottier, A., Bouckaert, J., Roels, T., & Derave, W. (2010). Mouth rinse but not ingestion of a carbohydrate solution improves 1-h cycle time trial performance. *Scandinavian Journal of Medicine and Sport in Science*, 20, 105-111. doi: 10.1111/j.1600-0838.2008.00868.x.

Přibyslavská, V., Scudamore, E.M., Johnson, S.L., Green, J.M., Stevenson Wilcoxson, M.C., Lowe, J.B., & O'Neal, E.K., (2016). Influence of carbohydrate mouth rinsing on running and jumping performance during early morning soccer scrimmaging. *European Journal of Sport Science*, 16, 441-447. doi: 10.1080/17461391.2015.1020345.

Rollo, I., Williams, C., & Nevill, M. (2011). Influence of ingesting versus mouth rinsing a carbohydrate solution during a 1-h run. *Medicine and Science in Sport and Exercise*, 43, 468-475. doi:10.1249/MSS.0b013e3181f1cda3.

Rollo, I., Williams, C., Gant, N., & Nute, M. (2008). The Influence of Carbohydrate Mouth Rinse on Self-Selected Speeds during a 30-min Treadmill Run. *International Journal of Sport Nutrition & Exercise Metabolism*, 18, 585-600. doi:10.1123/ijsnem.18.6.585.

Rollo, I., Homewood, G., Williams, C., Carter, J., & Goosey-Tolfrey, V.L. (2015). The Influence of Carbohydrate Mouth Rinse on Self-Selected Intermittent Running Performance. *International Journal of Sport Nutrition & Exercise Metabolism*, 25, 550-558. doi: 10.1123/ijsnem.2015-0001.

Sheppard, J.M. & Triplett, N.T (2015). Program Design for Resistance Training. In G.G. Haff and N.T. Triplett (Eds.), *Essentials of Strength Training and Conditioning (4th ed.)* (pp.249-258). Champaign, IL, Human Kinetics.

Stellingwerff, T., & Cox, G.R., (2014). Systematic Review, Carbohydrate supplementation on exercise performance or capacity of varying durations. *Applied Physiology, Nutrition, and Metabolism*, 39, 998-1011. doi: 10.1139/apnm-2014-0027.

Svebak, S., & Murgatroyd, S. (1985). Metamotivational Dominance, a multimethod validation of reversal theory constructs. *Journal of Personality and Social Psychology*, 48, 107-116. doi:10.1037/0022-3514.48.1.107.

Turner, C.E., Byblow, W.D., Stinear, C.M., & Gant, N., (2014). Carbohydrate in the mouth enhances activation of brain circuitry involved in motor performance and sensory perception. *Appetite*, 80, 212-219. doi: 10.1016/j.appet.2014.05.020.

Van Nieuwenhoven, M.A., Brouns, F., & Kovacs, E.M.R., (2005). The effect of two sports drinks and water on GI complaints and performance during an 18-km run. *International Journal of Sports Medicine*, 26, 281-285. doi: 10.1055/s-2004-820931.

Whitham, M., & McKinney, J. (2007). Effect of a carbohydrate mouthwash on running time-trial performance. *Journal of Sports Sciences*, 25, 1385-1392. doi:10.1080/02640410601113676.




Figure 1: Mean \pm SD (n=12) countermovement jump height (a), 10 m sprint time (b), mid-thigh pull peak force (c), number of bench-press repetitions (d) and number of squat repetitions (e). * Significantly greater than CON. † Significantly greater than PLA. # Significantly faster than PLA. ‡ Significantly faster than CON.




Figure 2: Individual countermovement jump height (a), 10 m sprint time (b), mid-thigh pull peak force (c), number of bench-press repetitions (d) and number of squat repetitions (e).

Table 1: Mean \pm SD (n=12) felt arousal, rating of perceived exertion (RPE) and heart rate throughout the exercise battery. * Significantly greater than CON.

	Pre-Rinse	Post-Rinse	CMJ	MTP	Sprint	Bench	Squat
<i>Arousal</i>							
CHO	3 \pm 1	5 \pm 1*	5 \pm 1*	5 \pm 0*	5 \pm 0*	5 \pm 1*	4 \pm 1
PLA	4 \pm 1	4 \pm 1*	4 \pm 1	5 \pm 1*	5 \pm 1	4 \pm 1	4 \pm 1
CON	3 \pm 1	3 \pm 1	4 \pm 1	4 \pm 1	4 \pm 1	4 \pm 1	3 \pm 1
<i>RPE</i>							
CHO			6 \pm 1	10 \pm 1	10 \pm 1	16 \pm 1	18 \pm 1
PLA			7 \pm 1	10 \pm 0	11 \pm 0	16 \pm 0	19 \pm 1
CON			6 \pm 1	10 \pm 2	11 \pm 3	16 \pm 1	18 \pm 1
<i>Heart Rate</i> (beats \cdot min ⁻¹)							
CHO	77 \pm 7	86 \pm 8	109 \pm 10	124 \pm 16	136 \pm 9	149 \pm 14	176 \pm 8
PLA	80 \pm 11	88 \pm 14	108 \pm 15	120 \pm 16	132 \pm 11	142 \pm 13	173 \pm 8
CON		81 \pm 7	99 \pm 11	117 \pm 13	128 \pm 7	147 \pm 10	171 \pm 11