

Semi-natural habitats support biological control, pollination and soil conservation in Europe: A review

Holland, J, Douma, J, Crowley, L, Kor, L, Stevenson, D & Smith, B

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Holland, J, Douma, J, Crowley, L, Kor, L, Stevenson, D & Smith, B 2017, 'Semi-natural habitats support biological control, pollination and soil conservation in Europe: A review' *Agronomy for Sustainable Development*, vol 37, no. 4, 31

<https://dx.doi.org/10.1007/s13593-017-0434-x>

DOI 10.1007/s13593-017-0434-x

ISSN 1774-0746

ESSN 1773-0155

Publisher: Springer

The final publication is available at Springer via <http://dx.doi.org/10.1007/s13593-017-0434-x>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

**Semi-natural habitats support biological control, pollination and soil conservation in
Europe: A review**

**John M. Holland¹, Jacob C. Douma^{2,3}, Liam Crowley¹, Laura James¹, Laura Kor¹, David
Stevenson¹, Barbara M. Smith^{1,4}**

¹Game and Wildlife Conservation Trust, Fordingbridge, Hampshire, SP6 1EF, UK

*²Wageningen University, Centre for Crop System Analysis, Droevendaalsesteeg 1, 6708PB
Wageningen, The Netherlands*

*³Laboratory of Entomology, Wageningen University, Droevendaalsesteeg 1, 6708PB, Wageningen,
The Netherlands*

*⁴Present address: Centre for Agroecology, Water and Resilience, Coventry University, Ryton Gardens, Wolston
Lane, Coventry, Warwickshire, CV8 3LG, UK*

*Corresponding author: John Holland, Game and Wildlife Conservation Trust, Fordingbridge, SP6
1EF, UK; email jholland@gwct.org.uk

Abstract

Semi-natural habitats are integral to most agricultural areas and have the potential to support ecosystem services, especially biological control and pollination by supplying resources for the invertebrates providing these services and for soil conservation by preventing erosion and run-off. Some habitats are supported through agri-environment scheme funding in the European Union, but their value for ecosystem service delivery has been questioned. An improved understanding of previous research approaches and outcomes will contribute to the development of more sustainable farming systems, improve experimental designs and highlight knowledge gaps especially for funders and researchers. Here we compiled a systematic map to allow for the first time a review of the quantity of evidence collected in Europe that semi-natural habitats support biological control, pollination and soil conservation. A literature search selected 2252 publications, and, following review, 270 met the inclusion criteria and were entered into the database. Most publications were of pest control (143 publications) with less on pollination (78 publications) or soil related aspects (31). For pest control and pollination most publications reported a positive effect of semi-natural habitats. There were weaknesses in the evidence base though because of bias in study location and the crops, whilst metrics (e.g. yield) valued by end users were seldom measured. Hedgerows, woodland and grassland were the most heavily investigated semi-natural habitats and the wider landscape composition was often considered. Study designs varied considerably yet only 24% included controls

or involved manipulation of semi-natural habitats. Service providers were commonly measured and used as a surrogate for ecosystem service delivery. Key messages for policy makers and funders are that they should encourage research that includes more metrics required by end users, be prepared to fund longer-term studies (61% were of only 1-year duration) and investigate the role of soils within semi-natural habitats in delivering ecosystem services.

Keywords

agroecology; ecosystem services; agricultural policy; pollinators; sustainable agriculture; integrated pest management; experimental design; agricultural research

1. Introduction

2. Method

3. Results

3.1 The extent to which the ecosystem services (biocontrol, pollination, soil conservation) in relation to semi-natural habitats been investigated across Europe

3.1.1 Geographic variation of publications

3.2 The types of semi-natural habitats that have been investigated and impact for a range of crops.

3.3 Other factors effecting ecosystem service delivery

3.4 Study setup, sampling methodology, sampled locations and farming systems

3.5 Studies of pest control

3.6 Studies of pollination

3.7 Approaches for investigating service providers

3.8 Outcomes for ecosystem service delivery

4. Conclusions

4.1 Extending the geographic extent of ecosystem services research across Europe

4.2 Ways of improving ecosystem service delivery

4.3 Filling the gaps

4.3.1 Crops

4.3.2 Interactions between ecosystem services

4.3.3 Yield

4.3.4 Pollination

4.3.5 Seldom investigated ecosystem services

4.4 Recommendations for improving ecosystem services delivery

4.5 Recommendations / notes for policy makers

5. Acknowledgements

6. References

7. References of the systematic map

1. Introduction

Throughout the European Union member states are implementing agri-environment schemes through the Common Agricultural Policy that protect or create semi-natural habitats, here defined as “any habitat within or outside of the crop containing a community of non-crop plant species.” Agri-environment schemes typically have multiple objectives that include nature conservation and resource protection (e.g. water quality), yet despite their considerable cost, their benefits have been questioned. This has been highlighted most with respect to conservation, because the target species have not always shown the expected response (Klein et al., 2006). However, semi-natural habitats have the potential to support ecosystem services and since the Millennium Ecosystem Assessment (MA2005) there has been increasing interest in investigating and quantifying these services in agricultural systems (e.g. Vihervaara et al., 2010; van Zanten et al., 2014). In addition, EU member states are obliged to adopt Integrated Pest Management principles as of 2014 under the Sustainable Use Directive (2009/128/EC) and semi-natural habitats can contribute towards this through enhancement of biological control. The other important ecosystem services provided or supported by semi-natural habitats are pollination and soil conservation (Wratten et al., 2012). Semi-natural habitats provide habitats and resources for the agents delivering biological control and pollination. These resources include shelter for aestivation or overwintering, breeding sites, food resources and an appropriate environment protected from agricultural operations (Landis 2000; Holland et al., 2016). Semi-natural habitats themselves can sequester carbon both within the vegetation and soil (Walter et al., 2003), whilst the vegetation’s physical structure can also hinder soil loss through erosion (Lowrance et al., 2002).

The role that semi-natural habitats play in providing ecosystem services has been widely investigated using a range of empirical approaches across the European Union. The types of investigation vary from evaluations of individual habitats (Holland et al., 2016) to landscape scale studies (e.g. Thies & Tschardtke, 1999). How such studies were conducted varied considerably. For pollination, a relatively straightforward ecosystem service to measure, 62 unique metrics were used in 121 publications (Liss

et al., 2013). Such variation makes comparisons difficult between publications and can lead to discrepancies in the interpretation of findings and recommendations (Liss et al., 2013). A similar review of the approaches used to investigate other ecosystem services, is not readily available. Yet, ensuring consistent methodologies are used could ensure that there is no bias in emphasis for policy and may make it possible to assess trade-offs between ecosystem services.



Fig. 1

Grass strip between fields with pan traps for collecting pollinators and example of a flower-rich margin.

Knowing which approaches have been used previously and their sensitivity to measuring ecosystem services has a number of advantages: 1) it can help ensure standardised ones are used in future research, 2) reviewing the success of previous publications across Europe can help identify which semi-natural habitats and cropping systems have already been investigated allowing users to identify research from countries with similar ecological infrastructure and cropping. 3) it can help to highlight knowledge gaps and identify topics for in-depth meta-analysis, 4) such reviews can help determine the most appropriate metrics, scale and experimental design on which to base further investigations and can help researchers and funding bodies to better target their research strategy.

If semi-natural habitats are to be managed optimally to improve ecosystem services provision on farmland across Europe thereby helping justify the public money invested in it through the Common Agricultural Policy, then it is important to understand the current state of knowledge. The main objective in this study was to summarise the quantity of evidence that has been collected in Europe on semi-natural habitats with respect to the key ecosystem services that semi-natural habitats support,

namely biological control, pollination and soil conservation. The secondary objectives were to synthesise this knowledge for use by other researchers, funders and end-users, and to provide a map on this topic that is searchable. Specifically, we asked:

- 1) To what extent have these ecosystem services (biocontrol, pollination, soil conservation) in relation to semi-natural habitats been investigated across Europe's agricultural production areas?
- 2) Which semi-natural habitats types have been investigated and for which ecosystem services?
- 3) What methodologies were used and how did this influence the outcomes?
- 4) What were the outcomes for ecosystem services delivery?

The outcomes can: 1) provide researchers and funding bodies with summary information that will support them in targeting research, 2) enable policy makers to consider the strength of evidence currently available to underpin decision-making, 3) help them identify priorities for research scoping 4) identify consistent benefits can be that can be utilised in the development of more sustainable farming systems.

2. Method

A systematic map approach was chosen as the appropriate method for this review as it allowed the identification and categorisation of available evidence to form a searchable database (Grant & Booth, 2009). Systematic Maps are methodical overviews of the quantity and quality of evidence in relation to a broad (open) question of policy or management relevance. They help to understand the breadth and depth of the evidence available. The process and rigour of the mapping exercise is the same as for systematic review except that they make no attempt to answer a specific question but instead collates all the evidence available on a topic of interest (James et al., 2016). However, the systematic map still allows areas to be identified where there is sufficient evidence for meta-analysis. In addition, it allows it allows the identification of gaps for future research. Systematic maps include all the publications that meet a set of apriori inclusion rules set by the authors thereby effectively summarising the relevant literature. Guidance on the compilation of systematic maps has been published (James et al., 2016) and is available online

http://www.environmentalevidence.org/Instructionsforauthors_maps.html [27/04/2016].

The types of semi-natural habitat that occur and are supported by agri-environment schemes across Europe vary considerably (Keenleyside et al., 2011) although there are some broad categories. Of the 63 types of entry-level options classified from the EU-27 Rural Development Plans, 15 broader categories were identified some of which are semi-natural habitats (grassland, cover in permanent

crops, buffer strips, management for wildlife and land taken out of production), but the list does not encompass all semi-natural habitats commonly occurring on farmland (e.g. woodland, hedgerows) and there are some semi-natural habitats designed specifically for ecosystem services such as pollination (flower-rich areas) and biological control (beetle banks). In addition, semi-natural habitats may also occur within fields such as the cover under perennial crops or established to prevent soil erosion. The final list selected for investigation therefore included some additional semi-natural habitat types (see search terms below).

A literature search was conducted that was comprehensive and precise enough to find as much of the relevant literature as is realistically possible whilst avoiding the capture of too much irrelevant literature. Only peer-reviewed, published papers describing empirical, original research was included (reviews, meta-analyses and landscape modelling were excluded) and searches were carried out using the online database: Web of Knowledge (v.5.10) for the period 1950-2015 (November). A scoping process that included review by the partners of the QuESSA project was used to refine and select the final search terms (Holland et al., 2014). Searches were only conducted in the English language and a wildcard (*) was used to select multiple word endings or plurals. The final search term was: ((woodland OR "field margin" OR "grass margin" OR hedge* OR "unimproved grass*" OR "field boundary" OR "cover crop" OR fallow OR "semi-natural grass*" OR landscape*) AND ("ecosystem service*" OR pollinat* OR "pest control" OR biocontrol OR "biological control" OR "seed predation" OR "soil erosion" OR "soil organic matter") AND (agricultur* OR farm*)). The search was restricted to research conducted in Europe by adding 31 European countries to the address option (see Appendix D). All results from the search results were reviewed at title and abstract level to ensure they met a set of inclusion criteria as follows:

- (i) European: The study must have been entirely carried out in a European country.
- (ii) Agricultural interest: it must not only have a conservation concern, but must also be relevant to farming systems.
- (iii) Semi-natural habitat OR Landscape complexity: the effects of semi-natural habitats OR landscape complexity must be investigated in the study.
- (iv) Invertebrates OR Other service provider: the organisms providing an ecosystem service must be invertebrates OR an ecosystem service must be influenced by semi-natural habitats or landscape complexities (e.g. use of semi-natural habitats to prevent soil erosion).

- (v) Empirical study: The article must report the results of a primary empirical study. Correlative and manipulative studies are included but literature reviews and statistical models are not.
- (vi) Natural populations: papers investigating the effect of introduced pest enemies were not included but enhancement of existing populations was acceptable.

Following initial scoping exercises to test search terms and responses, two reviewers divided the workload to determine whether the papers met inclusion criteria. A search was conducted in October 2014. Quality was controlled through reciprocal sub-sampling between assessors. If one or more of these criteria were not met, the paper was not included in the final systematic map. Articles that passed the inclusion criteria were then read in full, and entered into the database by extraction of the relevant data (Table 1). The search was then repeated in November 2015 adding in the term for landscape and the whole process was repeated, adding any additional literature, which provided a further check on the original search.

Table 1. Fields used to code publications in the database

ID	Numerical code identification code for each entry, assigned by WoK during search
First Author	Surname, Initial. of first author
Title	Full Article Title
Year	Four digit year of publication
Reference	Full reference of article (Harvard)
Ref Type	Journal What type of source did the entry come from
Text read	Full Text How much of the text was read by review author when entering
Linked studies	ID codes of all other entries in review that are linked by same first author and/or same study
Coded	YES Has the article been coded into review
Intervention	What is the independent variable
English	Y/N Is the language of the article English
Countries	Which country/countries was the study conducted in
Length of study in years	How long, in full years, did the study take place for
Study type	Manipulative/ Correlative Was the study manipulative or correlative
Control	Y/N Was there a control
Randomised	Y/N Was randomisation incorporated into the study design
Spatial replicate	Y/N Was there a spatial replicate
Temporal replicate	Y/N Was there a temporal replicate
Study Scale	Farm/Multi-farm Was the study restricted to one farm or did it incorporate multiple farms
Time of year of measurements	Which season(s) was the study conducted in
Farm system	Arable/Orchard/Livestock What type of farming system was being used

Sampling Location	In what specific location(s) within the farm was the study conducted
Reason heterogeneity results	Reasons that led to any heterogeneity in results
Semi-natural habitat types	What types of semi-natural habitats were involved in the study
ES provided by invertebrates?	Y/N Is the ES being provided done so via invertebrates, or from other source (e.g. semi-natural habitats improving soil quality in adjacent fields)
Providers	<i>Only applicable when ES is provided by invertebrates</i> Pest natural enemies/Pollinators/Nutrient cycling What type of ES were the arthropods providing
Sampling method for predation	<i>Only applicable when ES is provided by invertebrates</i> What method was used to sample the providers
Provider Sampling Location	<i>Only applicable when ES is provided by invertebrates</i> Crop/semi-natural habitats Which location were the providers sampled from
Sampling for provider resources	<i>Only applicable when ES is provided by invertebrates</i> What sampling (if any) for provider resources was conducted
ES Type	Regulating/Provisioning/Cultural/Supporting What type of ES is being provided
ES measured: Provisioning	<i>Only applicable when ES type = Provisioning</i> Food/Water/Energy/Pharmaceuticals/Minerals What type of provision is being examined
ES measured: Regulating	<i>Only applicable when ES type = Regulating</i> Pollination/Pest control/Carbon sequestration/Soil resilience/Water purification/Waste decomposition What type of regulation is being examined
ES measured: Cultural	<i>Only applicable when ES type = Cultural</i> Cultural inspiration/Recreational experience/Scientific discovery What type of cultural ES is being examined
ES measured: Supporting	<i>Only applicable when ES type = Supporting</i> Nutrient cycling/Seed dispersal/Soil formation/ Primary production What type of supporting ES is being examined
ES methodology	What was measured in order to quantify ES provided
Effect on yield	Positive/ Negative/No effect/Not measured What effect of the intervention on the yield
% change in yield	The % increase or decrease in yield that was found
Type of semi-natural habitats effecting yield	The type semi-natural habitats with the greatest impact on yield
Effect on Pest Control	Positive/Negative/No effect/Not measured What effect did the intervention have on the level of pest control
% change in pests	The % increase or decrease upon the level of pest control
Type of semi-natural habitats effecting pest control	The type of semi-natural habitats with the greatest effect on pest control
Effect on Pollination	Positive/ Negative/No effect/Not measured What effect did the intervention have pollination
% change in pollination	The % increase or decrease upon the level of pollination
Type of semi-natural habitats effecting pollination	The type of semi-natural habitats with the greatest effect on pollination

Effect on seed predation	Positive/ Negative/No effect/Not measured What effect did the intervention have on the level of seed predation
% change in seed predation	The % increase or decrease upon the level of seed predation
Type of semi-natural habitats effecting seed predation	The type of semi-natural habitats with the greatest effect on seed predation
Effect on soil Erosion	Positive/ Negative/No effect/Not measured What effect did the intervention have on the level of pest control
% change in soil erosion	The % increase or decrease upon the level of soil erosion
Type of semi-natural habitats effecting soil erosion	The type of semi-natural habitats with the greatest effect on soil erosion
Effect on SOM	Positive/ Negative/No effect/Not measured What effect did the intervention have on the level of SOM
Type of semi-natural habitats effecting SOM	The type of semi-natural habitats with the greatest effect on SOM
Extracting data	Easy/Moderate/Difficult What level of difficulty was experienced in extracting data from the publication
Organism investigated	Which organism(s) where the subject of the study
Notes	Any additional relevant notes about the entry

3. Results

The search terms returned 2252 publications and these were reviewed for suitability for inclusion; of these 270 met the inclusion criteria and were entered into the systematic map database. The number of publications investigating the selected ecosystem services covered by the systematic map rose steadily over the last 10 years reaching approximately 30 per year by 2014 (Fig. 2).




Fig. 2

The number of publications in the systematic map per annum (n = 270).

3.1 The extent to which the ecosystem services (biocontrol, pollination, soil conservation) in relation to semi-natural habitats been investigated across Europe

The majority of publications looked at regulating ecosystem services (250), while 28 investigated supporting services, 12 measured provisioning and three cultural services. The most commonly investigated regulating ecosystem services were pest control (55%) and pollination (30%), with the other ecosystems services having 20 (7%) or less publications (Fig. 3). 31 publications investigated soil related ecosystem services such as soil resilience and carbon sequestration. Of supporting services, 26 investigated nutrient cycling or levels of soil organic matter and one publication each on seed dispersal and soil formation. 23 publications investigated two or more ecosystem services, these most often being pest control and pollination.

3.1.1 Geographic variation of publications

Publications originated from 23 countries, but there was geographic bias because most were conducted in 9 countries that had >10 publications. The highest number of publications originated from Germany (64), followed by France, Spain, Sweden and the UK (21-26). (Fig. 4). Only nine publications reported on studies conducted in more than one country. Pest control was the most widely researched ecosystem services, investigated in all but three countries. Pollination was most heavily investigated in Germany, Sweden and UK with each 10 or more publications. Few publications on other ecosystem services were included in the database.




Fig 3

The number of publications for each regulating ecosystem service (n=260).




Fig. 4

The number of publications originating from European countries and the type of ecosystem service investigated.

3.2 The types of semi-natural habitats that have been investigated and impact for a range of crops.

A total of 245 publications specifically investigated a type of semi-natural habitat of which half (137 publications) investigated more than one habitat type. 31% of these publications included hedgerows/field margins, 23% woodland or shrubland and 28% grassland (Fig. 5). Investigations of fallows were reported in 8% of publications whilst the other habitats were addressed by only 12 (2%)

or fewer publications. In these publications, the locations where service providers or ecosystem services were most commonly measured were hedgerows or field margins (44%), woodland/shrubland (19%) and grassland (25%) (Fig. 6). Only areas within fields, orchards, olive groves or vineyards were sampled in 106 publications.




Fig. 5

The number of publications for each type of semi-natural habitat.




Fig. 6

The number of publications for the semi-natural habitats that were sampled (n=225).

Of the 230 publications that looked at ecosystem services within crops, the most studied were wheat (28%) and oilseed rape (17%), followed by barley (10%) and other cereals such as triticale, rye, spelt and oats (10%) (Fig. 7). Perennial, vegetable and root crops were relatively poorly investigated.




Fig. 7

The number of publications for each crop type.

3.3 Other factors effecting ecosystem service delivery

A high proportion of publications (43%) investigated whether the composition of the surrounding landscape effected the selected ecosystem services. These publications used for example, the proportion of arable land, uncropped land or specific habitats to provide an indication of the landscapes complexity. Some other factors that may also effect these ecosystem services were also investigated. These included 41 publications in which farming systems such as conventional and organic or those with different levels of inputs were examined. There were 30 publications in which the type of ground management, fertiliser inputs, grazing intensity or seed mix, was investigated and 18 publications of cover crops. Only two publications examined grazing management.

3.4 Study setup, sampling methodology, sampled locations and farming systems

The experimental design used in each study was extracted. Excluding 12 publications in which long-term data (>8 years) was analysed, 64% were single year studies, 18% of two and 13% of 3-8 years duration. The most common design was one using a non-randomised, correlative approach without a

control (34%), whilst a further 20% used the same design but with a randomised approach. To show effects on the selected ecosystem services the correlative studies usually employed some quality or quantity of semi-natural habitats rather than formal randomised designs with experimental manipulations of semi-natural habitat type or area. Overall only 24% of all publications included any type of control treatment. The majority of publications included spatial or temporal replication (Fig. 8). In manipulative studies that tested interventions through experimental manipulations to create treatments, more studies did not use controls (59 publications) than those that did (44), however, randomisation was more commonly employed when there was a control. The most robust experimental approach using manipulation, randomization and controls was only used in 11% of publications. It was not possible to identify whether the type of experimental design determined the final recommendations because where an effect was reported most were positive regardless of the approach or ecosystem service (pest control 83%, pollination 79%).




Fig. 8

Type of experimental design used to study semi-natural habitats: whether correlative (use of existing landscape elements) or manipulative (habitats created or manipulated), randomized or not, and with or without control treatments.

Overall 73% of publications reported on studies conducted on more than one farm (site), whilst the remainder were of a single study site. Most publications originated from studies conducted on arable farms (81%), 12% were in orchards and 6% on livestock farms; these were the most frequently investigated systems regardless of whether pest control or pollination was investigated (Fig. 9).




Fig. 9

The effect of semi-natural habitats on pest control and pollination separated by the scale of the study and the cropping system.

3.5 Studies of pest control

Of the 143 publications that investigated the ecosystem service “pest control” none included measurements of yield. Of these, 138 publications measured the pests, or levels of predators and parasitism. Of these a remarkably low percentage, 22% reported on actual pest levels, and only 9% of these also measured pest predators or parasitism rates (6%) or both of these (2%) (Fig. 10). The majority of the publications on pest control measured predators (70%) and fewer parasitism (25%).




Fig.10

Number of publications separated by the service providers that were measured.

3.6 Studies of pollination

There were 85 publications that investigated pollinators or pollination and most were conducted in arable crops (Fig. 9). Of these 87% measured the number of pollinators, however, only 24% measured any metric of pollination such as fruit or seed set. The predominant method for assessing pollinators was through transect counts (57%), pan traps (14%) (Fig. 11), flower visiting observations (14%) or netting (10%).

3.7 Approaches for investigating service providers

The most commonly used methods for assessing invertebrates were visual counts of some form either along transect or on the vegetation (34%) and pitfall traps (19%) which collected epigeal invertebrates (Fig. 11). The transect counts and pan traps for most often used for assessments of pollinators, whilst pitfall traps, suction sampling, sweep nets, window traps and sticky traps were used for collecting predators. Considering all sampling methods, it was more common for either the crop or the semi-natural habitats to be sampled than for both, with the exception of sweep netting (Fig. 11). The crop was also more frequently sampled than the semi-natural habitats for the majority of methods, except where the soil was sampled or when transect counts and water traps were used.




Fig. 11

Number of times in which different invertebrate sampling methods were used in the crop, semi-natural habitat (SNH) or both locations.

The most extensively studied ecosystem service providers were invertebrates (217) of which 66% were of pest natural enemies and 36% of pollinators, although only 1% of papers covered both groups and there were only two studies of organisms responsible for nutrient recycling. The botanical resources for pest natural enemies and pollinators were less well investigated. Vegetation surveys were conducted in 13% (35 publications) and 5% (13 publications) conducted assessments of flower abundance or other measures of floral resources.

Of the invertebrates, Hymenoptera were studied in the most publications (123) because this order includes both bees (pollinators) and parasitic wasps (pest regulators) (Fig. 12). Coleoptera (beetles) were investigated in 37 publications. 32 studies were more general in their approach and only assessed functional groups such as predators and aphidophagous predators or total invertebrates.




Fig. 12

The number and percentage of publications for each type of service provider.

3.8 Outcomes for ecosystem service delivery

Of the 24% publications that reported an effect of pest control, 81% of these concluded that there was a positive effect, although of these publications usually the abundance of predators or parasitism (67%) was used as a proxy for pest control and pest levels were only measured in 10%. A negative effect was found in only three publications and all of these measured the abundance of predators or parasitism to obtain the recommendation (Fig. 9). Most positive effects were reported for arable crops, but also for other systems. Only two publications reported on yield with one finding a positive and the other both negative and positive effects of semi-natural habitats on weed levels.

Whether a positive or negative recommendation was provided was unrelated to the scale of the study. 58 publications provided a recommendation as to the best semi-natural habitats for enhancing pest control which included some landscape scale studies. Of these publications 52% recommended field boundary habitats such as hedgerows, hedgebase or field margins, 17% recommended cover crops, leys or fallows to enhance pest control, woodland or forest was recommended in 12% and grassy habitats in 9%. Some publications recommended more than one habitat type.

28 publications reported on whether there was any effect of the interventions on pollination or pollinators and of these 79% reported a positive effect of the landscape, one found a negative response and 5% no effect (Fig. 9). In all but one study, however, pollination itself was not measured but abundance of pollinators and floral resources instead. Those studies found an increase in pollinators or floral resources after interventions and assumed it to be positively related to pollination. Three times as many of the multi-farm scale publications reported a positive effect compared to single farm studies. Of those 28 publications reporting an effect of interventions, six recommended field boundary habitats such as hedgerows, hedge base or field margins and seven mentioned that floral abundance was important irrespective of the specific semi-natural habitats. An increase in the area of semi-natural habitats *per se* was advised in five publications. The negative effect reported occurred in wild cherry trees where increasing vegetation complexity around the trees decreased bee species richness and crop flower visitation rates.

Of the 10 publications investigating soil erosion, 8 reported a positive effect of semi-natural habitats and one no effect. In 17 publications, a positive effect on soil organic matter was found whilst two reported no effect. In the 15 publications that provided a recommendation on ways to increase soil organic matter, seven recommended using grass leys, two using fallows and four using cover crops. Woodland and hedgerows were also found to have high soil organic matter.

4. Conclusions

The systematic map revealed that the majority of the publications found a positive effect of semi-natural habitats on pollination and pest control or their proxies, although there were some weaknesses to the evidence base. The study design and sampling methodology varied largely across studies, which were conducted in a limited number of countries and crops. Yield was not measured in any of the pest control studies nor were metrics such as reduced pesticide use that are valued by end-users. Likewise for pollination, only 24% of publications measured a metric of pollination such as pollen deposition or measures that end-users are interested in such as seed quality, fruit set or quality (Bommarco et al., 2012). To some extent these omissions are because the studies were not designed to measure ecosystem services. Yet they can provide a useful indication of what further information may be gained in future studies.

4.1 Extending the geographic extent of ecosystem services research across Europe

The systematic map revealed that biocontrol, pollination and soil conservation have been relatively poorly investigated in most European countries with only five having more than 20 publications suitable for inclusion. There may be many different reasons for this geographic bias, apart from the

size of the countries research community, funding opportunities or publication in national languages. The great majority of publications were from Germany followed by Spain, France, Sweden and Switzerland, whilst Eastern Europe and the far south of Europe were poorly represented. This may be a reflection on the types of landscapes that occur in Western Europe where there has been a history of smaller farms and fields, surrounded by distinct boundary types or woodland, as opposed to Eastern Europe where large state farms predominated during the 20th Century. Thus, the geographical bias in the evidence-base prevents researchers from generalising effectively across pedo-climatic zones. If further studies were conducted to allow generalisation across pedo-climatic zones this would give insight into the role of semi-natural habitats on pest control and pollination across Europe. For some services it may be prudent to focus on those that are particularly relevant to the pedo-climatic zone, for example, some zones may be more susceptible to soil erosion.

4.2 Ways of improving ecosystem service delivery

The most heavily investigated semi-natural habitats were hedgerows or other field boundary habitats, woodland and grassland as these represent the main areas of non-crop land on farmland. For these semi-natural habitats the abundance of service providers was typically measured. This provided some indication of the resource potential of the habitats, yet if the semi-natural habitat is studied in interaction with the crop and its surrounding landscape, a better understanding of ecosystem services provisioning may be gained. This is because there will be local and landscape influences on ecosystem services provision due to for, example, differences in the mobility of the various service providers (Tscharntke et al., 2005). For less mobile providers it is the interchange between the semi-natural habitats and the adjacent crop that may determine the level of ecosystem services provision (Bianchi et al., 2006). For more mobile service providers such as bumblebees and hoverflies, it is important to consider the surrounding landscapes that may provide additional service providers to those of local semi-natural habitats.

Different experimental approaches are needed to evaluate local and landscape effects. At the local level it is valuable to determine the zone of influence of semi-natural habitats yet such studies are difficult and costly to conduct, relying on extensive sampling networks and service providers that exhibit a degree of gradation from the habitat edge into the crop or across the landscape (for examples see Holland et al., 1999; Holland et al., 2005). That explains why these studies are less frequently conducted and instead spatially explicit individual based models are employed, although these still require realistic data on the service provider's mobility (Bianchi & van der Werf, 2003). Tracking the movement of individuals is sometimes feasible either by marking physically (Holland et al., 2004; Winder et al., 2005), using chemical markers such as Rubidium (Tillman et al., 2007) or for larger insects using a tracking device (Tahir & Brooker, 2011). All of these require considerably more resources. Directional trapping using window, sticky or pitfall traps can also be employed for some

species (Winder et al., 2001; Muirhead-Thomson, 1991) to show movement in or out of habitats. Sampling for ecosystem services providers was conducted in crops adjacent to semi-natural habitats and/or the semi-natural habitat. When appraising the value of semi-natural habitats adjacent to different crops the crop management should also be taken into account because there is some evidence that this can also influence service providers within adjacent semi-natural habitats (Rand et al., 2006).

For landscape scale studies, the influence of landscape composition has been widely investigated and the literature on this periodically reviewed for biocontrol (e.g. Bianchi et al., 2006; Tschardt et al., 2007, 2012; Chaplin-Kramer et al., 2011) and pollination and pollinators (Viana et al., 2012). Meta-analyses were also conducted to examine the effects of landscape and local features on the comparative abundance of pollinators and natural enemies (Shackelford et al., 2013; Rusch et al., 2014) and on levels of pest control (Rusch et al., 2016). Evidence from these studies can be contradictory, but is not surprising given the complexity of the ecosystems and number of interacting factors that eventually determine the level of ecosystem service that is provided. In addition, these studies typically make use of existing landscapes with contrasting levels of landscape composition that may have unforeseen and unaccountable influences on the service provision. For example, the metric of the proportion of non-crop habitats (or cropped area) is commonly used and related to the level ecosystem service provision. Yet the composition of the non-crop areas may differ considerably and some types may be more influential than others. More compelling evidence could be gained from manipulations of existing farms or landscapes with ecosystem services evaluations pre- and post-habitat enhancement. Such designs were only used in 11% of publications. However, we were unable to identify if specific designs led to stronger recommendations because positive effects were predominantly reported. Meta-analyses, more detailed landscape mapping (Garcia-Feced et al., 2015) and use of spatial models (Ekroos et al., 2014) may help in identifying if there are optimal designs for utilising the ecosystem services supported by semi-natural habitats.

4.3 Filling the gaps

4.3.1 Crops

This investigation revealed that arable crops, predominantly cereals and oilseed rape, were the most heavily investigated whilst those for which pests and pollination are more important (orchards and vegetable crops) were less frequently studied. This may be because horticultural crops occupy less land and are economically less important in the countries conducting most of the research. In addition, the development of integrated pest management is more problematic for high value crops with stringent cosmetic quality targets. On the other hand, such a finding may indicate that research is not being driven by end-user requirements.

4.3.2 Interactions between ecosystem services

More than one ecosystem service was rarely investigated in the same publication. It may be that other ecosystem services were investigated but published elsewhere, however, these findings suggest that there could be potential to investigate trade-offs between services and to include disservices. If semi-natural habitats are to be optimally utilised it is essential that trade-offs are understood as they may be beneficial or disadvantageous (Rodriguez et al., 2006).

4.3.3 Yield

The majority of publications examined supporting ecosystem services, notably pest control and/or pollination, which is to be expected as most semi-natural habitats occur outside of the crop and these services rely on mobile agents that will benefit from semi-natural habitat enhancement. Yield was only examined in 13 publications despite its importance. This may be because yield is affected by many other inputs including levels of agrochemical inputs, crop variety, water availability, soil type and environmental conditions that may have a much larger impact than that of the service. Moreover, such factors may vary considerably between plots or fields that were chosen to provide a particular habitat and not to standardise management inputs for example, therefore making it difficult to ascertain any difference in yield attributable to a supporting service alone. Nevertheless, many of these other variables can be accounted for using an appropriate design. However, the majority of studies were of one-year duration even though many factors that influence the level of service provision are likely to vary considerably between years, such as the weather, levels of invertebrate service providers, disease and pest pressure. More long-term experiments are therefore needed.

Impacts on pests or yield are the most compelling evidence for farmers and are needed if wider adoption of semi-natural habitats for pest control is to be advocated. Of the publications that investigated insect pest control the emphasis was on measurement of service providers and only 22% measured pest levels. Yield was not measured for pest control studies yet 57 publications made recommendations on the value of semi-natural habitats for pest control. Most predation publications focussed on insect pests and only five publications were included in the map that investigated seed predation in relation to semi-natural habitats, despite the agronomic importance of weeds. The proportion of sales accounted for by insecticides was less than 5% in most EU countries whereas sales of herbicides were usually over 50% and reached 80% in some countries (Eurostat, 2012). These findings and the EU policies on Sustainable Use of Pesticides indicate that further research is needed, incorporating measurements of impact on either yield or suitable surrogates are needed if farmers are to move towards IPM utilising conservation biocontrol.

4.3.4 Pollination

Pollination was investigated in 78 publications of which only six measured pollen transfer or pollinator foraging activity on flowers, with a further 14 measuring seed or fruit set. Most measured pollinators as a surrogate for pollination. There is consequently a knowledge gap for measurement of

pollination and its effect on yield with respect to the value of semi-natural habitats. With respect to the study of pollination itself there is also a need for the development and use of standardised methods as recommended by Liss et al. (2013).

The majority of publications of pest control and pollination focussed on measurements of the service providers whereas few also measured factors that determine their abundance, such as the type and quality of the vegetation in the semi-natural habitats. As a consequence, identification of plant species or traits that are particularly beneficial are seldom made (Lavorel et al., 2011), nor are recommendations on how to create or improve semi-natural habitats.

4.3.5 Seldom investigated ecosystem services

Few other ecosystem services have been investigated in relation to semi-natural habitats, but these included levels of soil organic matter or carbon storage that reflect carbon sequestration, soil erosion that impacts on water quality, nutrient levels as a measure of nutrient cycling and the aesthetic value of semi-natural habitats. There was evidence that semi-natural habitats stored carbon and helped prevent soil erosion. The role of soil organisms in pest control, carbon sequestration, nutrient cycling and erosion was seldom investigated. All of these areas require further investigation.

4.4 Recommendations for improving ecosystem services delivery

Agriculture has been long reliant on artificial agrochemical inputs and farmers may be reluctant to adopt alternative pest control strategies whilst such inputs remain relatively cheap and reliable. As a consequence, there has not been much pressure to develop alternative techniques and it is only recently with changes in legislation and the advent of more widespread insecticide resistance that alternatives are being investigated. Yet, the most compelling evidence on the benefits of semi-natural habitats will come from measures on yield, pesticide use, level of pollination and/or seed quality.

Despite this, it was apparent from this review that many of the publications focussed on one aspect of ecosystem services, such as abundance of service providers rather than the ecosystem services delivery. This was in part because the studies had a different focus, such as nature conservation, rather than ecosystem services delivery. However, even where an ecosystem service was the focus there was often a tendency to focus on the service providers and typically those that are easiest to sample and identify, are most prolific or are typically used as indicators rather than focussing on end-user requirements. By measuring the ecosystem service itself (yield, pest control, pesticide use, or level of pollination, seed quality) more persuasive evidence may be gained. In addition, there is a need to harmonise data collection with standard methodologies so that studies can be compared and integrated. This will include selecting metrics that are biologically meaningful and developing indices that can advise us when, for example, sufficient service providers are present. The development of rapid assessment methods would also help reduce the sampling effort per unit resource.

If more resources become available or if sampling becomes more efficient, we recommend to focus on:

- 1) Measuring the ecosystem services, including year-to-year variability, rather than only service providers to generate recommendations for end-users.
- 2) Investigations that include more than one ecosystem services including disservices so that trade-offs can be identified.
- 3) Extending the geographical scope of studies to Eastern Europe because at present the majority are from Western Europe.
- 4) The impact of semi-natural habitats in preventing the movement of soil and water. This was identified as a knowledge gap. In addition the potential of semi-natural habitats to sequester carbon was rarely measured, although there was evidence that soil organic matter and thereby carbon storage is higher than in the surrounding fields (Walter et al., 2003).
- 5) Study the effectiveness of semi-natural habitats in relation to the (trait) composition of crops and non-crop habitats in the landscape (e.g. Lavorel et al., 2011).
- 6) Developing proxies that are easier to measure yet still give insight if semi-natural habitats are advantageous (see discussion earlier) and need to be stimulated through policy.

4.5 Recommendations / notes for policy makers

Knowledge exchange for policy making should be a two-way process, policy makers need reliable advice that is evidence based and the policy makers should work with researchers and practitioners to develop research priorities.

This work suggests that policy makers could call for:

- 1) Estimation of how effective semi-natural habitats are at providing ecosystem services such as food production, pest control, pollination and soil preservation. For farmers, some estimate of the reliability and risk associated with more reliance on ecosystem services would also be needed if they are to be encouraged to adopt more sustainable farming practices reliant on ecosystem services (Duru et al., 2015). The predominance of short term studies of 1-3 years' duration does little to help meet this requirement. Research funders need to recognise that ecosystems change slowly and that longer-term support is needed.
- 2) Some semi-natural habitats in EU member states are supported through agri-environment schemes funding and therefore economic evaluations of the costs and benefits of semi-natural habitats to ecosystem services would help in the justification for this spend. Other habitats that are not supported yet may be important need to be identified and measures put in place to encourage their retention and improvement where necessary.

Caveat: the authors acknowledge that although the search terms used were devised to identify all relevant publications, they were not comprehensive and therefore may not cover the entire record. Moreover, only those papers that met the inclusion criteria, which was subjective to some extent, were included in the systematic map and other publications exist that also investigated aspects of these ecosystem services.

5. Acknowledgements

The review formed part of the QuESSA project that received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement No 311879. JCD was supported by an NWO Earth and Life Sciences (NWO-ALW) VENI grant project 863.14.018.

6. References

- Bianchi FJJA, Booij CJH, Tscharntke T (2006) Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proc R Soc Biol Sci Ser B* 273:1715-1727. doi:10.1098/rspb.2006.3530
- Bianchi FJJA, van der Werf W (2003) The effect of the area and configuration of hibernation sites on the control of aphids by *Coccinella septempunctata* (Coleoptera : Coccinellidae) in agricultural landscapes: A simulation study. *Environ Entomol* 32:1290-1304. doi:10.1603/0046-225X-32.6.1290
- Bommarco R, Marini L, Vaissiere BE (2012) Insect pollination enhances seed yield, quality, and market value in oilseed rape. *Oecologia (Berl)* 169:1025-1032. doi:10.1007/s00442-012-2271-6
- Chaplin-Kramer R, O'Rourke ME, Blitzer EJ, Kremen C (2011) A meta-analysis of crop pest and natural enemy response to landscape complexity. *Ecol Lett* 14:922-932. doi:10.1111/j.1461-0248.2011.01642
- Duru M, Therond O, Martin G, Martin-Clouaire R, Magne M-A, Justes E, Journet E-P, Aubertot J-N, Savary S, Bergez J-E, Sarthou JP (2015) How to implement biodiversity-based agriculture to enhance ecosystem services: a review. *Agron Sustain Dev* 35:1259-1281. doi:10.1007/s13593-015-0306-1
- Ekroos J, Olsson O, Rundlöf M, Wätzold F, Smith HG (2014) Optimizing agri-environment schemes for biodiversity, ecosystem services or both? *Biol Conserv* 172:65-71. doi:10.1016/j.biocon.2014.02.013
- García-Feced C, Weissteiner CJ, Baraldi A, Paracchini ML, Maes J, Zulian G, Kempen M, Elbersen B, Pérez-Soba M (2015) Semi-natural vegetation in agricultural land: European map and links to ecosystem service supply. *Agron Sustain Dev* 35:273-283. doi:10.1007/s13593-014-0238-1

- Grant MJ, Booth A (2009) A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J* 26:91-108. doi:10.1111/j.1471-1842.2009.00848
- Holland J, Begbie M, Birkett T, Southway S, Thoms S, Alexander C, Thomas C (2004) The spatial dynamics and movement of *Pterostichus melanarius* and *P. madidus* (Carabidae) between and within arable fields in the UK. *Int J Ecol Environ Sci* 30:35-50.
- Holland JM, Bianchi FJJA, Entling MH, Moonen AC, Smith BM, Jeanneret P (2016) Structure, function and management of semi-natural habitats for conservation biological control: A review of European studies. *Pest Manag Sci* 72:1638-1651. doi: 10.1002/ps.4318
- Holland JM, Jeanneret P, Herzog F, Moonen A-C, Rossing W, van der Werf W, Kiss J, van Helden M, Paracchini ML, Cresswell J, Pointereau P, Heijne B, Veromann E, Antichi D, Entling M, Balázs B (2014) The QuESSA Project: Quantification of Ecological Services for sustainable agriculture. *Landscape Management for Functional Biodiversity International Organisation for Biological Control/West Palaearctic Regional Section Bulletin* 100:55-58.
- Holland JM, Perry JN, Winder L (1999) The within-field spatial and temporal distribution of arthropods in winter wheat. *Bull Entomol Res* 89:499-513. doi:10.1017/S0007485399000656
- Holland JM, Thomas C, Birkett T, Southway S, Oaten H (2005) Farm-scale spatiotemporal dynamics of predatory beetles in arable crops. *J Appl Ecol* 42:1140-1152. doi:10.1111/j.1365-2664.2005.01083
- James KL, Randall NP, Haddaway NR (2016) A methodology for systematic mapping in environmental sciences. *Environ Evidence* 5. doi:10.1186/s13750-016-0059-6
- Keenleyside C, Allen B, Hart K, Menaude H, Stefanova V, Prazan J, Herzon I, Clement T, Povellato A, Maciejczak M, Boatman N (2011) Delivering environmental benefits through entry level agri-environment schemes in the EU. Report Prepared for DG Environment, Project ENV.B.1/ETU/2010/0035. Institute for European Environmental Policy, London
- Kleijn D, Baquero RA, Clough Y, Diaz M, De Esteban J, Fernandez F, Gabriel D, Herzog F, Holzschuh A, Johl R, Knop E, Kruess A, Marshall EJP, Steffan-Dewenter I, Tschamtkke T, Verhulst J, West TM, Yela JL (2006) Mixed biodiversity benefits of agri-environment schemes in five European countries. *Ecol Lett* 9:243-254. doi:10.1111/j.1461-0248.2005.00869
- Lavorel S, Grigulis K, Lamarque P, Colace M-P, Garden D, Girel J, Pellet G, Douzet R (2011) Using plant functional traits to understand the landscape distribution of multiple ecosystem services. *J Ecol* 99:135-147. doi:10.1111/j.1365-2745.2010.01753
- Liss KN, Mitchell MGE, MacDonald GK, Mahajan SL, Méthot J, Jacob AL, Maguire DY, Metson GS, Ziter C, Dancose K, Martins K, Terrado M, Bennett EM (2013) Variability in ecosystem service measurement: a pollination service case study. *Front Ecol Environ* 11:414-422. doi:10.1890/120189

- Lowrance R, Dabney S, Schultz R (2002) Improving water and soil quality with conservation buffers. *J Soil Water Conserv* 57:36A-43A.
- Muirhead-Thomson RC (1991) Flight Traps and Interceptor Traps. In: Muirhead-Thomson RC (ed) *Trap Responses of Flying Insects: The Influence of Trap Design on Capture Efficiency* Academic Press, London, pp 152-179.
- Rand TA, Tylianakis JM, Tscharntke T (2006) Spillover edge effects: the dispersal of agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecol Lett* 9:603-614. doi:10.1111/j.1461-0248.2006.00911
- Rodriguez E, Fernandez-Anero FJ, Ruiz P, Campos M (2006) Soil arthropod abundance under conventional and no tillage in a Mediterranean climate. *Soil Tillage Res* 85:229-233. doi:10.1016/j.still.2004.12.010
- Rusch A, Birkhofer K, Bommarco R, Smith HG, Ekbom B (2014) Management intensity at field and landscape levels affects the structure of generalist predator communities. *Oecologia* 175:971-983. doi:10.1007/s00442-014-2949
- Rusch A, Chaplin-Kramer R, Gardiner MM, Hawro V, Holland J, Landis D, Thies C, Tscharntke T, Weisser WW, Winqvist C, Woltz M, Bommarco R (2016) Agricultural landscape simplification reduces natural pest control: A quantitative synthesis. *Agr Ecosyst Environ* 221:198-204. doi:10.1007/s10980-016-0390
- Shackelford G, Steward PR, Benton TG, Kunin WE, Potts SG, Biesmeijer JC, Sait SM (2013) Comparison of pollinators and natural enemies: a meta-analysis of landscape and local effects on abundance and richness in crops. *Biol Rev* 88:1002-1021. doi:10.1111/bry.12040
- Tahir N, Brooker G (2011) Recent developments and recommendations for improving harmonic radar tracking systems. In: *Proceedings of the 5th European Conference on Antennas and Propagation (EUCAP)*, 11-15 April 2011, pp 1531-1535.
- Thies C, Tscharntke T (1999) Landscape structure and biological control in agroecosystems. *Science* 285:893-895. doi:10.1126/science.285.5429.893
- Tillman PG, Prasifka JR, Heinz KM (2007) Rubidium marking to detect dispersal of pest and predator from corn into sorghum and cotton in Georgia. *J Entomol Sci* 42:383-391.
- Tscharntke T, Bommarco R, Clough Y, Crist TO, Kleijn D, Rand TA, Tylianakis JM, van Nouhuys S, Vidal S (2007) Conservation biological control and enemy diversity on a landscape scale. *Biol Control* 43:294-309. doi:10.1016/j.biocontrol.2007.08.006
- Tscharntke T, Rand TA, Bianchi FJJA (2005) The landscape context of trophic interactions: insect spillover across the crop-noncrop interface. *Ann Zool Fenn* 42:421-432.
- Tscharntke T, Tylianakis JM, Rand TA, Didham RK, Fahrig L, Peter B, Bengtsson J, Clough Y, Crist TO, Dormann CF, Ewers RM, Freund J, Holt RD, Holzschuh A, Klein AM, Kleijn D, Kremen C, Landis DA, Laurance W, Lindenmayer D, Scherber C, Sodhi N, Steffan-Dewenter I, Thies C, van der Putten WH, Westphal C (2012) Landscape moderation of biodiversity

- patterns and processes - eight hypotheses. *Biol Rev (Camb)* 87:661-685. doi:10.1111/j.1469-185X.2011.00216
- van Zanten BT, Verburg PH, Espinosa M, Gomez-y-Paloma S, Galimberti G, Kantelhardt J, Kapfer M, Lefebvre M, Manrique R, Piorr A, Raggi M, Schaller L, Targetti S, Zasada I, Viaggi D (2014) European agricultural landscapes, common agricultural policy and ecosystem services: a review. *Agron Sustain Dev* 34:309-325. doi:10.1007/s13593-013-0183-4
- Viana BF, Boscolo D, Neto E, Lopes L, Lopes A, Ferreira PA, Pigozzo C, Primo L (2012) How well do we understand landscape effects on pollinators and pollination services? *J Pollinat Ecol* 7:31-42
- Vihervaara P, Rönkä M, Walls M (2010) Trends in Ecosystem Service Research: Early Steps and Current Drivers. *Ambio* 39:314-324. doi:10.1007/s13280-010-0048
- Walter C, Merot P, Layer B, Dutin G (2003) The effect of hedgerows on soil organic carbon storage in hillslopes. *Soil Use Manage* 19:201-207. doi:10.1079/sum2002190
- Winder L, Alexander CJ, Holland JM, Symondson WO, Perry JN, Woolley C (2005) Predatory activity and spatial pattern: the response of generalist carabids to their aphid prey. *J Anim Ecol* 74:443-454. doi:10.1111/j.1365-2656.2005.00939
- Winder L, Holland JM, Perry JN, Woolley C, Alexander CJ (2001) The use of barrier-connected pitfall trapping for sampling predatory beetles and spiders. *Entomol Exp Appl* 98:249-258. doi:10.1046/j.1570-7458.2001.00781
- Wratten SD, Gillespie M, Decourtye A, Mader E, Desneux N (2012) Pollinator habitat enhancement: Benefits to other ecosystem services. *Agr Ecosyst Environ* 159:112-122. doi:10.1016/j.agee.2012.06.020

7. References of the systematic map

- Aguilar-Fenollosa, E., Ibáñez-Gual, M. V., Pascual-Ruiz, S., Hurtado, M., Jacas, J. A. (2011) Effect of ground-cover management on spider mites and their phytoseiid natural enemies in clementine mandarin Orchards (I): Bottom-up regulation. *Biological Control*, 59, 158-170.
- Aguilar-Fenollosa, E., Ibáñez-Gual, M. V., Pascual-Ruiz, S., Hurtado, M., Jacas, J. A. (2011) Effect of ground-cover management on spider mites and their phytoseiid natural enemies in clementine mandarin Orchards (II): Top-down regulation. *Biological Control*, 2011, 59, 171-179.
- Aguilar-Fenollosa, E., Pascual-Ruiz, S., Hurtado, M. A., Jacas, J. A. (2011) Efficacy and economics of ground cover management as a conservation biological control strategy against *Tetranychus urticae* in clementine mandarin orchards. *Crop Protection*, 30, 1328-1333.

- Ailincăi, C., Jităreanu, G., Bucur, D., Ailincăi, D., Mercuș, A. D. (2011) Evolution of some chemical properties of soil under influence of soil erosion and different Cropping systems. *Cercetari Agronomice in Moldova*, Vol. XLIV , No. 4 (148), 5-18.
- Al Hassan, D., Georgelin, E., Delattre, T., Burel, F., Plantegenest, M., Kindlmann, P., Butet, A. (2013) Does the presence of grassy strips and landscape grain affect the spatial distribution of aphids and their carabid predators? *Agricultural and Forest Entomology*, 15, 24-33.
- Alanen, E. L., Hyvönen, T., Lindgren, S., Härmä, O., Kuussaari, M. (2011) Differential responses of bumblebees and diurnal Lepidoptera to vegetation succession in long-term set-aside. *Journal of Applied Ecology*, 48, 1251-1259.
- Albrecht, M., Duelli, P., Müller, C., Kleijn, D., Schmid, B. (2007) The Swiss agri-environment scheme enhances pollinator diversity and plant reproductive success in nearby intensively managed farmland. *Journal of Applied Ecology*, 44, 813-822.
- Albrecht, M., Duelli, P., Schmid, B., Müller, C. B. (2007) Interaction diversity within quantified insect food webs in restored and adjacent intensively managed meadows. *Journal of Animal Ecology*, 76, 1015-1025.
- Alcántara, C., Pujadas, A., Saavedra, M. (2011) Management of cruciferous cover crops by mowing for soil and water conservation in southern Spain. *Agricultural Water Management*, 98, 1071-1080.
- Alebeek, F., Kamstra, J-H., Kruistum, G., Visser, A. (2006) Improving natural pest suppression in arable farming: field margins and the importance of ground dwelling predators. *Landscape Management For Functional Biodiversity, IOBC wprs Bulletin*, 29, 137.
- Alhmedi, A., Haubruge, E., Francis, F. (2011) Effect of stinging nettle habitats on aphidophagous predators and Hymenoptera in wheat and green pea fields with special attention to the invader *Harmonia*. *Entomological Science*, 12, 349-358.
- Alhmedi, A., Haubruge, E., D'Hoedt, S., Francis, F. (2011) Quantitative food webs of herbivore and related beneficial community in non-crop and crop habitats. *Biological Control*, 58, 103-112.
- Alomar, Ò., Goula, M., Albajes, R. (2002) Colonisation of tomato fields by predatory mirid bugs (Hemiptera: Heteroptera) in Northern Spain. *Agriculture, Ecosystems & Environment*, 89, 105-115.
- Ameixa, O. L. G. A., Kindlmann, P. A. V. E. L. (2008) Agricultural policy-induced landscape changes: effects on carabid abundance and their biocontrol potential. *European Journal of Entomology*, 105, 467-476.
- Andersen, A. (1997) Densities of overwintering Coleoptera and staphylinids (Col., Carabidae and Staphylinidae) in cereal and grass fields and their boundaries. *Journal of Applied Entomology*, 121, 77-80.

- Anderson, A., Carnus, T., Helden, A. J., Sheridan, H., Purvis, G. (2013) The influence of conservation Field margins in intensively managed grazing land on communities of five arthropod trophic groups. *Insect Conservation and Diversity*, 6, 201-211.
- Andersson, G. K., Ekroos, J., Stjernman, M., Rundlöf, M., Smith, H. G. (2014) Effects of farming intensity, crop rotation and landscape heterogeneity on field bean pollination. *Agriculture, Ecosystems & Environment*, 184, 145-148.
- Anjum-Zubair, M., Schmidt-Entling, M. H., Querner, P., Frank, T. (2010) Influence of within-field position and adjoining habitat on carabid beetle assemblages in winter wheat. *Agricultural and Forest Entomology*, 12, 301-306.
- Antil, R. S., Gerzabek, M. H., Haberhauer, G., Eder, G. (2005) Long-term effects of cropped vs. fallow and fertilizer amendments on soil organic matter - I. Organic Carbon. *Journal of Plant Nutrition And Soil*, 168, 108-116.
- Anti, R. S., Gerzabek, M. H., Haberhauer, G., Eder, G. (2005) Long-term effects of cropped vs. fallow and fertilizer amendments on soil organic matter II. Nitrogen. *Journal of Plant Nutrition And Soil Science*, 168, 212-218.
- Anyszka, Z., Dobrzanski, A. (2006) Impact of cover crops and herbicides usage on weed infestation, growth and yield of transplanted leek. *Journal of Plant Diseases and Protection*, 20, 733-738.
- Bailey, D., Schmidt-Entling, M. H., Eberhart, P., Herrmann, J. D., Hofer, G., Kormann, U., Herzog, F. (2010) Effects of habitat amount and isolation on biodiversity in fragmented traditional orchards. *Journal of Applied Ecology*, 47, 1003-1013.
- Bailey, S., Requier, F., Nusillard, B., Roberts, S. P., Potts, S. G., Bouget, C. (2014) Distance from forest edge affects bee pollinators in oilseed rape fields. *Ecology and Evolution*, 4, 370-380.
- Baraibar, B., Westerman, P. R., Carrión, E., Recasens, J. (2009) Effects of tillage and irrigation in cereal fields on weed seed removal by seed predators. *Journal of Applied Ecology*, 46, 380-387.
- Bartomeus, I., Potts, S. G., Steffan-Dewenter, I., Vaissiere, B. E., Woyciechowski, M., Krewenka, K. M., ... Bommarco, R. (2014) Contribution of insect pollinators to crop yield and quality varies with agricultural intensification. *Peer J*, 2, e328.
- Bartomeus, I., Vila, M., Steffan-Dewenter, I. (2010) Combined effects of *Impatiens glandulifera* invasion and landscape structure on native plant pollination. *Journal of Ecology*, 98, 440-450.
- Basedow, T. (1998) The species composition and frequency of Araneae (Araneae) in fields of winter wheat grown under different conditions in Germany. *Journal of Applied Entomology*, 122, 585-590.
- Batáry, P., Sutcliffe, L., Dormann, C. F., Tschardtke, T. (2013) Organic farming favours insect-pollinated over non-insect pollinated forbs in meadows and wheat Fields. *PloS One*, 8, e54818.

- Bennewicz, J. (2011) Aphidivorous Diptera (Diptera: Syrphidae) at field boundaries and woodland edges in an agricultural landscape. *Polish Journal of Entomology*, 80, 129-149.
- Berg, A., Ahrne, K., Ockinger, E., Svensson, R., Soderstrom, B. (2011) Butterfly distribution and abundance is affected by variation in the Swedish forest-farmland landscape. *Biological Conservation*, 144, 2819-2831.
- Bianchi, F. J. J. A., Goedhart, P. W., Baveco, J. M. (2008) Enhanced pest control in cabbage crops near forest in The Netherlands. *Landscape Ecology*, 23, 595-602.
- Bianchi, F. J. J. A., Van Wingerden, W. K. R. E., Griffioen, A. J., Van Der Veen, M., Van Der Straten, M. J. J., Wegman, R. M. A., Meeuwsen, H. A. M. (2005) Landscape factors affecting the control of *Mamestra brassicae* by natural enemies in Brussel sprout. *Agriculture, Ecosystems & Environment*, 107, 145-150.
- Birge, T., Fred, M. (2011) New ideas for old landscapes: using a social-ecological approach for conservation of traditional rural biotopes-a case study from Finland. *European Countryside*, 3, 51-67.
- Boccaccio, L., Petacchi, R. (2009) Landscape effects on the complex of *Bactrocera oleae* Hymenoptera and implications for conservation biological control. *BioControl*, 54, 607-616.
- Bodner, G., Himmelbauer, M., Loiskandl, W., Kaul, H. P. (2010) Improved evaluation of cover crop species by growth and root factors. *Agronomy for Sustainable Development*, 30, 455-464.
- Bodner, G., Loiskandl, W., Buchan, G., Kaul, H. P. (2008) Natural and management-induced dynamics of hydraulic conductivity along a cover-cropped field slope. *Geoderma*, 146, 317-325.
- Boller, E. F., Remund, U., Candolfi, M. P. (1988) Hedges as potential sources of *Typhlodromus pyri*, the most important predatory mite in vineyards of Northern Switzerland. *Entomophaga*, 33, 249-255.
- Bommarco, R., Marini, L., Vaissière, B. E. (2012) Insect pollination enhances seed yield, quality, and market value in oilseed rape. *Oecologia*, 169, 1025-1032.
- Bonte, D., Lanckacker, K., Wiersma, E., Lens, L. (2008) Web building flexibility of an orb-web spider in a heterogeneous agricultural landscape. *Ecography*, 31, 646-653.
- Borin, M., Passoni, M., Thiene, M., Tempesta, T. (2010) Multiple functions of buffer strips in farming areas. *European Journal of Agronomy*, 32, 103-111.
- Boulal, H., Gómez-Macpherson, H. (2010) Dynamics of soil organic carbon in an innovative irrigated permanent bed system on sloping land in southern Spain. *Agriculture, Ecosystems & Environment*, 139, 284-292.
- Breitbach, N., Tillmann, S., Schleuning, M., Grünewald, C., Laube, I., Steffan-Dewenter, I., Böhning-Gaese, K., (2012) Influence of habitat complexity and landscape configuration on pollination and seed-dispersal interactions of wild cherry trees. *Oecologia*, 168, 425-437.

- Brittain, C., Bommarco, R., Vighi, M., Settele, J., Potts, S. G. (2010) Organic farming in isolated landscapes does not benefit flower-visiting insects and pollination. *Biological Conservation*, 143, 1860-1867.
- Burgio, G., Ferrari, R., Pozzati, M., Boriani, L. (2004) The role of ecological compensation areas on predator populations: an analysis on biodiversity and phenology of Coleoptera (Coleoptera) on non-crop plants within hedgerows in Northern Italy. *Bulletin of Insectology*, 57, 1-10.
- Buri, P., Humbert, J. Y., Arlettaz, R. (2014) Promoting pollinating insects in intensive agricultural matrices: field-scale experimental manipulation of hay-meadow mowing regimes and its effects on Hymenoptera. *PloS One*, 9, e85635.
- Caballero-López, B., Bommarco, R., Blanco-Moreno, J. M., Sans, F. X., Pujade-Villar, J., Rundlöf, M., Smith, H. G. (2012) Aphids and their natural enemies are differently affected by habitat features at local and landscape scales. *Biological Control*, 63, 222-229.
- Cammeraat, E. L. (2004) Scale dependent thresholds in hydrological and erosion response of a semi-arid catchment in southeast Spain. *Agriculture, Ecosystems & Environment*, 104, 317-332.
- Campiglia, E., Mancinelli, R., Radicetti, E., Caporali, F. (2010) Effect of Cover crops and mulches on weed control and nitrogen fertilization in tomato (*Lycopersicon esculentum* Mill.). *Crop Protection*, 29, 354-363.
- Carré, G., Roche, P., Chifflet, R., Morison, N., Bommarco, R., Harrison-Cripps, J., ... Vaissière, B. E. (2009) Landscape context and habitat type as drivers of bee diversity in European annual Crops. *Agriculture, Ecosystems & Environment*, 133, 40-47.
- Carvell, C., Meek, W. R., Pywell, R. F., Goulson, D., Nowakowski, M. (2007) Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *Journal of Applied Ecology*, 44, 29-40.
- Carvell, C., Osborne, J. L., Bourke, A. F. G., Freeman, S. N., Pywell, R. F., Heard, M. S. (2011) Bumble bee species' responses to a targeted conservation measure depend on landscape context and habitat quality. *Ecological Applications*, 21, 1760-1771.
- Casalí, J., Gastesi, R., Álvarez-Mozos, J., De Santisteban, L. M., Lersundi, J., Giménez, R., ... Donézar, M. (2008) Runoff, erosion, and water quality of agricultural watersheds in central Navarre (Spain). *Agricultural Water management*, 95, 1111-1128.
- Castro, J., Fernandez-Ondoño, E., Rodríguez, C., Lallena, A. M., Sierra, M., Aguilar, J. (2008) Effects of different olive-grove management systems on the organic carbon and nitrogen content of the soil in Jaen (Spain). *Soil & Tillage Research*, 98, 56-67.
- Chifflet, R., Klein, E. K., Lavigne, C., Le Feon, V., Ricroch, A. E., Lecomte, J., Vaissiere, B. E. (2011) Spatial scale of insect-mediated pollen dispersal in oilseed rape in an open agricultural landscape. *Journal of Applied Ecology*, 48, 689-696.

- Clough, Y., Kruess, A., Tschardtke, T. (2007) Local and landscape factors in differently managed arable fields affect the insect herbivore community of a non-crop plant species. *Journal of Applied Ecology*, 44, 22-28.
- Clough, Y., Kruess, A., Kleijn, D., Tschardtke, T. (2005) Spider diversity in cereal fields: comparing factors at local, landscape and regional scales. *Journal of Biogeography*, 32, 2007-2014.
- Constantin, J., Mary, B., Laurent, F., Aubrion, G., Fontaine, A., Kerveillant, P., Beudoin, N. (2010) Effects of catch crops, no till and reduced nitrogen fertilization on nitrogen leaching and balance in three long-term experiments. *Agriculture, Ecosystems & Environment*, 135, 268-278.
- Cotes, B., Campos, M., Pascual, F., Ruano, F. (2010) The ladybeetle community (Coleoptera: Coleoptera) in southern olive agroecosystems of Spain. *Environmental Entomology*, 39, 79-87.
- Cranmer, L., McCollin, D., Ollerton, J. (2012) Landscape structure influences pollinator movements and directly affects plant reproductive success. *Oikos*, 121, 562-568.
- Daedlow, D., Sommer, T., Westermann, P. R. (2012) Weed seed predation in organic and conventional cereal fields. *Julius-Kühn-Archiv*, 434, 265.
- Dalin, P. (2006) Habitat difference in abundance of willow leaf beetle *Phratora vulgatissima* (Coleoptera: Chrysomelidae): Plant quality or natural enemies? *Bulletin of Entomological Research*, 96, 629-635.
- Dauber, J., Biesmeijer, J. C., Gabriel, D., Kunin, W. E., Lamborn, E., Meyer, B., ... Petanidou, T. (2010) Effects of patch size and density on flower visitation and seed set of wild plants: a Pan-European approach. *Journal of Ecology*, 98, 188-196.
- De Cauwer, B. (2005) Biodiversity and agro-ecology in field margins. PhD Thesis, Faculty of Bioscience Engineering, Universiteit Gent.
- Debras, J. F., SeNoussi, R., Rieux, R., Buisson, E., Dutoit, T. (2008) Spatial distribution of an arthropod community in a pear orchard (southern France) - identification of a hedge effect. *Agriculture, Ecosystems & Environment*, 127, 166-176.
- Debras, J., Senoussi, R., Dutoit, T. (2010) Hedgerow effects on the distribution of beneficial arthropods in a pear orchard in Southern France. *Revue Internationale D'écologie Méditerranéenne International Journal of Mediterranean Ecology*, 37, 75-82.
- Dennis, P., and Fry, G. L. A. (1992) Field margins - can they enhance natural enemy population-densities and general arthropod diversity on farmland. *Agriculture, Ecosystems and Environment*, 40, 95-115.
- Denys, C., Tschardtke, T. (2002) Plant-insect communities and predator-prey ratios in field margin strips, adjacent crop fields, and fallows. *Oecologia*, 130, 315-324.
- Díaz, M. (1992) Spatial and temporal patterns of granivorous ant seed predation in patchy cereal crop areas of central Spain. *Oecologia*, 91, 561-568.

- Diehl, E., Mader, V. L., Wolters, V., Birkhofer, K. (2013) Management intensity and vegetation complexity affect web-building Araneae and their prey. *Oecologia*, 173, 579-589.
- Diekötter, T., Kadoya, T., Peter, F., Wolters, V., Jauker, F. (2010) Oilseed rape crops distort plant-pollinator interactions. *Journal of Applied Ecology*, 47, 209-214.
- Diekötter, T., Walther-Hellwig, K., Conradi, M., Suter, M., Frankl, R. (2006) Effects of landscape elements on the distribution of the rare bumblebee species *Bombus muscorum* in an agricultural landscape. *Biodiversity and Conservation*, 15, 57-68.
- Ditner, N., Balmer, O., Beck, J., Blick, T., Nagel, P., Luka, H. (2013) Effects of experimentally planting non-crop flowers into cabbage fields on the abundance and diversity of predators. *Biodiversity and Conservation*, 22, 1049-1061.
- Drapela, T., Frank, T., Heer, X., Moser, D., Zaller, J. G. (2011) Landscape structure affects activity density, body size and fecundity of *Pardosa wolf* Araneae (Araneae: Lycosidae) in winter oilseed rape. *European Journal of Entomology*, 108, 609-614.
- Drapela, T., Moser, D., Zaller, J. G., Frank, T. (2008) Spider assemblages in winter oilseed rape affected by landscape and site factors. *Ecography*, 31, 254-262.
- Duso, C., Malagnini, V., Paganelli, A., Aldegheri, L., Bottini, M., Otto, S. (2004) Pollen availability and abundance of predatory phytoseiid mites on natural and secondary Hedgerows. *BioControl*, 49, 397-415.
- Ekroos, J., Piha, M., Tiainen, J. (2008) Role of organic and conventional field boundaries on boreal bumblebees and Lepidoptera. *Agriculture, Ecosystems and Environment*, 124, 155-159.
- Evans, D. M., Pocock, M. J., Brooks, J., Memmott, J. (2011) Seeds in farmland food-webs: resource importance, distribution and the impacts of farm management. *Biological Conservation*, 144, 2941-2950.
- Fabian, Y., Sandau, N., Bruggisser, O. T., Aebi, A., Kehrli, P., Rohr, R. P., ... Bersier, L. F. (2013) The importance of landscape and spatial structure for Hymenopteran-based food webs in an agro-ecosystem. *Journal of Animal Ecology*, 82, 1203-1214.
- Fabian, Y., Sandau, N., Bruggisser, O. T., Aebi, A., Kehrli, P., Rohr, R. P., ... Bersier, L. F. (2014) plant diversity in a nutshell: testing for small-scale effects on trap nesting wild Hymenoptera and wasps. *Ecosphere*, 5, 1-18.
- Farwig, N., BaiLey, D., Bochud, E., Herrmann, J. D., Kindler, E., Reusser, N., ... Schmidt-Entling, M. H. (2009) Isolation from forest reduces pollination, seed predation and insect scavenging in Swiss farmland. *Landscape Ecology*, 24, 919-927.
- Felipe Lucia, M., Comín, F. A., Bennett, E. M. (2014) Interactions among ecosystem services across land uses in a floodplain agroecosystem. *Ecology and Society*, 19, 20.
- Fischer, C., Thies, C., Tschardtke, T. (2011) Mixed effects of landscape complexity and farming practice on weed seed removal. *Perspectives in Plant Ecology, Evolution and Systematics*, 13, 297-303.

- Fliszkiewicz, M., Kuśnierczak, A., Szymaś, B. (2012) The accompanying fauna of solitary bee *Osmia bicornis* (L.) Syn. *Osmia rufa* (L.) nests settled in different biotopes. *Journal of Apicultural Science*, 56, 51-58.
- Follain, S., Walter, C., Legout, A., Lemercier, B., Dutin, G. (2007) Induced effects of hedgerow networks on soil organic carbon storage within an agricultural landscape. *Geoderma*, 142, 80-95.
- Fontana, V., Radtke, A., Walde, J., Tasser, E., Wilhalm, T., Zerbe, S., Tappeiner, U. (2014) What plant traits tell us: Consequences of land-use change of a traditional agro-forest system on biodiversity and ecosystem service provision. *Agriculture, Ecosystems & Environment*, 186, 44-53.
- Fournier, E., Loreau, M. (1999) Effects of newly planted hedges on ground-beetle diversity (Coleoptera, Carabidae) in an agricultural landscape. *Ecography*, 22, 87-97.
- Frank, T. (1999) Density of adult Diptera (Dipt., Syrphidae) in sown weed strips acid adjacent fields. *Journal of Applied Entomology*, 123, 351-355.
- Franzen, M., Nilsson, S. G. (2008) How can we preserve and restore species richness of pollinating insects on agricultural land? *Ecography*, 31, 698-708.
- Gabarra, R., Alomar, Ò., Castañé, C., Goula, M., Albajes, R. (2004) Movement of greenhouse whitefly and its predators between in-and outside of Mediterranean greenhouses. *Agriculture, Ecosystems & Environment*, 102, 341-348.
- Gabriel, D., Tschardtke, T. (2007) Insect pollinated plants benefit from organic farming. *Agriculture, Ecosystems & Environment*, 118, 43-48.
- Gagic, V., Hänke, S., Thies, C., Tschardtke, T. (2014) Community variability in aphid Hymenoptera versus predators in response to agricultural intensification. *Insect Conservation and Diversity*, 7, 103-112.
- Gagic, V., Hänke, S., Thies, C., Scherber, C., Tomanović, Ž., Tschardtke, T. (2012) Agricultural intensification and cereal aphid–parasitoid–hyperparasitoid food webs: network complexity, temporal variability and parasitism rates. *Oecologia*, 170, 1099-1109.
- Gagic, V., Tschardtke, T., Dormann, C. F., Gruber, B., Wilstermann, A., Thies, C. (2011) Food web structure and biocontrol in a four-trophic level system across a landscape complexity gradient. *Proceedings of the Royal Society B: Biological Sciences*, rspb20102645.
- Geiger, F., Bianchi, F. J. J. A., Wäckers, F. L. (2005) Winter ecology of the cabbage aphid *Brevicoryne brassicae* (L.)(Homo., Aphididae) and its parasitoid *Diaeretiella rapae* (McIntosh)(Hym., Braconidae: Aphidiidae). *Journal of Applied Entomology*, 129, 563-566.
- Geiger, F., Wäckers, F. L., Bianchi, F. J. J. A. (2009) Hibernation of predatory arthropods in semi-natural habitats. *BioControl*, 54, 529-535.
- Geslin, B., Gauzens, B., Thébault, E., Dajoz, I. (2013) Plant pollinator networks along a gradient of urbanisation. *PloS One*, 8, e63421.

- Gladbach, D. J., Holzschuh, A., Scherber, C., Thies, C., Dormann, C. F., Tschamntke, T. (2011) Crop-noncrop spillover: Arable fields affect trophic interactions on wild plants in surrounding habitats. *Oecologia*, 166, 433-441.
- Glendell, M., Granger, S. J., Bol, R., Brazier, R. E. (2014) Quantifying the spatial variability of soil physical and chemical properties in relation to mitigation of diffuse water pollution. *Geoderma*, 214, 25-41.
- Goidts, E., van Wesemael, B. A. S. Van Oost, K. (2009) Driving forces of soil organic carbon evolution at the landscape and regional scale using data from a stratified soil monitoring. *Global Change Biology*, 15, 2981-3000.
- Goller, E., Nunnenmacher, L., Goldbach, H. E. (1997) Faba beans as a cover crop in organically grown hops: influence on aphids and aphid antagonists. *Entomological Research in Organic Agriculture*, 15, 279-284.
- Gómez, J. A., Llewellyn, C., Basch, G., Sutton, P. B., Dyson, J. S. Jones, C. A. (2011) The effects of cover crops and conventional tillage on soil and runoff loss in vineyards and olive groves in several Mediterranean countries. *Soil Use and Management*, 27, 502-514.
- Good, J. A., Giller, P. S. (1991) The effect of cereal and grass management on staphylinid (Coleoptera) assemblages in south-west Ireland. *Journal of Applied Ecology*, 28, 810-826.
- Haenke, S., Kovács-Hostyánszki, A., Fründ, J., Batáry, P., Jauker, B., Tschamntke, T., Holzschuh, A. (2014) Landscape configuration of crops and hedgerows drives local syrphid fly abundance. *Journal of Applied Ecology*, 51, 505-513.
- Haenke, S., Scheid, B., Schaefer, M., Tschamntke, T., Thies, C. (2009) Increasing syrphid fly diversity and density in sown flower strips within simple vs. complex landscapes. *Journal of Applied Ecology*, 46, 1106-1114.
- Haschek, C., Drapela, T., Schuller, N., Fiedler, K., Frank, T. (2012) Carabid beetle condition, reproduction and density in winter oilseed rape affected by field and landscape parameters. *Journal of Applied Entomology*, 136, 665-674.
- Helenius, J. K., Holopainen, J. K., Huusela-Veistola, E. (2008) Ground beetle (Coleoptera, Carabidae) diversity in Finnish arable land. *Agricultural and Food Science*, 10, 261-276.
- Henriksen, C. I., Langer, V. (2013) Road verges and winter wheat fields as resources for wild Hymenoptera in agricultural landscapes. *Agriculture, Ecosystems & Environment*, 173, 66-71.
- Henry, M., Fröchen, M., Maillet-Mezeray, J., Breyné, E., Allier, F., Odoux, J. F., Decourtye, A. (2012) Spatial autocorrelation in honeybee foraging activity reveals optimal focus scale for predicting agro-environmental scheme efficiency. *Ecological Modelling*, 225, 103-114.
- Hirsch, M., Pfaff, S., Wolters, V. (2003) The influence of matrix type on flower visitors of *Centaurea jacea* L. *Agriculture, Ecosystems & Environment*, 98, 331-337

- Holland, J. M., Birkett, T., Southway, S. (2009) Contrasting the farm-scale spatio-temporal dynamics of boundary and field overwintering predatory Coleoptera in arable crops. *Biocontrol*, 54, 19-33.
- Holland, J. M., Oaten, H., Southway, S., Moreby, S. (2008) The effectiveness of field margin enhancement for cereal aphid control by different natural enemy guilds. *Biological Control*, 47, 71-76.
- Holzschuh, A., Dormann, C. F., Tschardtke, T., Steffan-Dewenter, I. (2013) Mass-flowering crops enhance wild bee abundance. *Oecologia*, 172, 477-484.
- Holzschuh, A., Dormann, C. F., Tschardtke, T., Steffan-Dewenter, I. (2011) Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the Royal Society B: Biological Sciences*, rspb20110268.
- Holzschuh, A., Dudenhöffer, J. H., Tschardtke, T. (2012) Landscapes with wild bee habitats enhance pollination, fruit set and yield of sweet cherry. *Biological Conservation*, 153, 101-107.
- Holzschuh, A., Steffan-Dewenter, I., Tschardtke, T. (2009) Grass strip corridors in agricultural landscapes enhance nest-site colonization by solitary wasps. *Ecological Applications*, 19, 123-132.
- Holzschuh, A., Steffan-Dewenter, I., Tschardtke, T. (2010) How do landscape composition and configuration, organic farming and fallow strips affect the diversity of Hymenoptera, wasps and their Hymenoptera? *Journal of Animal Ecology*, 79, 491-500.
- Holzschuh, A., Steffan-Dewenter, I., Kleijn, D., Tschardtke, T. (2007) Diversity of flower-visiting Hymenoptera in cereal fields: effects of farming system, landscape composition and regional context. *Journal of Applied Ecology*, 44, 41-49.
- Hradetzky, R., Kromp, B. (1997) Spatial distribution of flying insects in an organic rye field and an adjacent hedge and forest edge. *Entomological Research in Organic Agriculture*, 15, 353-357.
- Iekarska-Boniecka, H., Wilkaniec, B., Dolanska-Niedbala, E. (2008) Parasitic wasps of the Pimplinae subfamily (Hymenoptera, Ichneumonidae) of agricultural landscape refugium habitats in central Wielkopolska. *Acta Oecologica*, 7, 23-30.
- Inclán, D. J., Cerretti, P., Marini, L. (2014) Interactive effects of area and connectivity on the diversity of tachinid Hymenoptera in highly fragmented landscapes. *Landscape Ecology*, 29, 879-889.
- Jakobsson, A., Ågren, J. (2014) Distance to semi-natural grassland influences seed production of insect-pollinated herbs. *Oecologia*, 175, 199-208.
- Jankauskas, B., Jankauskiene, G., Fullen, M. A. (2004) Erosion-preventive crop rotations and water erosion rates on undulating slopes in Lithuania. *Canadian Journal of Soil science*, 84, 177-186.
- Jauker, F., Diekotter, T., Schwarzbach, F., Wolters, V. (2009) Pollinator dispersal in an agricultural matrix: opposing responses of wild Hymenoptera and Diptera to landscape structure and distance from main habitat. *Landscape Ecology*, 24, 547-555.

- Jonason, D., Smith, H. G., Bengtsson, J., Birkhofer, K. (2013) Landscape simplification promotes weed seed predation by carabid Coleoptera (Coleoptera: Carabidae). *Landscape Ecology*, 28, 487-494.
- Josso, C., Le Ralec, A., Raymond, L., Saulais, J., Baudry, J., Poinot, D., Cortesero, A. M. (2013) Effects of field and landscape variables on crop colonization and biological control of the cabbage root fly *delia radicum*. *Landscape Ecology*, 28, 1697-1715.
- Kells, A. R., Holland, J. M., Goulson, D. (2001) The value of uncropped field margins for foraging bumblebees. *Journal of Insect Conservation*, 5, 283-291.
- Kiss, J., Kádár, F., Kozma, E., Tóth, I. (1993) Importance of various habitats in agricultural landscape related to integrated pest management: a preliminary study. *Landscape and Urban Planning*, 27, 191-198.
- Klaus, V. H., Kleinebecker, T., Prati, D., Gossner, M. M., Alt, F., Boch, S., ... Hölzel, N. (2013) Does organic grassland farming benefit plant and arthropod diversity at the expense of yield and soil fertility? *Agriculture, Ecosystems & Environment*, 177, 1-9.
- Koerner, W., Dupouey, J. L., Dambrine, E., Benoit, M. (1997) Influence of past land use on the vegetation and soils of present day forest in the Vosges mountains, France. *Journal of Ecology*, 85, 351-358.
- Kohler, F., Verhulst, J., KNop, E., Herzog, F., Kleijn, D. (2007) Indirect effects of grassland extensification schemes on pollinators in two contrasting European countries. *Biological Conservation*, 135, 302-307.
- Kohler, F., Verhulst, J., Van Klink, R., Kleijn, D. (2008) At what spatial scale do high-quality habitats enhance the diversity of forbs and pollinators in intensively farmed landscapes? *Journal of Applied Ecology*, 45, 753-762.
- Korpela, E. L., Hyvönen, T., Lindgren, S., Kuussaari, M. (2013) Can pollination services, species diversity and conservation be simultaneously promoted by sown wildflower strips on farmland? *Agriculture, Ecosystems & Environment*, 179, 18-24.
- Kosmas, C., Danalatos, N., Cammeraat, L. H., Chabart, M., Diamantopoulos, J., Farand, R., ... Vacca, A. (1997) The effect of land use on runoff and soil erosion rates under Mediterranean conditions. *Catena*, 29, 45-59.
- Kovács-Hostyánszki, A., Batáry, P., Báldi, A. (2011) Local and landscape effects on bee communities of Hungarian winter cereal fields. *Agricultural and Forest Entomology*, 13, 59-66.
- Kovács-Hostyánszki, A., Haenke, S., Batáry, P., Jauker, B., Báldi, A., Tschardtke, T., Holzschuh, A. (2013) Contrasting effects of mass-flowering crops on bee pollination of hedge plants at different spatial and temporal scales. *Ecological Applications*, 23, 1938-1946.
- Krause, B., Culmsee, H. (2013) The significance of habitat continuity and current management on the compositional and functional diversity of grasslands in the uplands of Lower Saxony, Germany. *Flora-Morphology, Distribution, Functional Ecology of Plants*, 208, 299-311.

- Krauss, J., Gallenberger, I., Steffan-Dewenter, I. (2011) Decreased functional diversity and biological pest control in conventional compared to organic crop fields. *Plos One*, 6, e19502.
- Krewenka, K. M., Holzschuh, A., Tschardtke, T., Dormann, C. F. (2011) Landscape elements as potential barriers and corridors for Hymenoptera, wasps and Hymenoptera. *Biological Conservation*, 144, 1816-1825.
- Kromp, B., Steinberger, K-H., 1992. Grassy field margins and arthropod diversity - a case-study on ground Coleoptera and Araneae in Eastern Austria (Coleoptera, Carabidae, Arachnida, Aranei, Opiliones). *Agriculture, Ecosystems & Environment*, 40, 71-93.
- Kruess, A., Tschardtke, T. (1994) Habitat fragmentation, species loss, and biological control. *Science (Washington)*, 264, 1581-1584.
- Kruess, A. (2003) Effects of landscape structure and habitat type on a plant-herbivore-parasitoid community. *Ecography*, 26, 283-290.
- Kuussaari, M., Hyvönen, T., Härmä, O. (2011) Pollinator insects benefit from rotational fallows. *Agriculture, Ecosystems and Environment*, 143, 28-36.
- Lagerlof, J., Wallin, H. (1993). The abundance of arthropods along two field margins with different types of vegetation composition - an experimental-study. *Agriculture, Ecosystems & Environment*, 4, 141-154.
- Lagerlöf, J., Starkb, J., Svensson, B. (1992) Margins of agricultural fields as habitats for pollinating insects. *Agriculture, Ecosystems & Environment*, 40, 117-124.
- Langer, V. (2001). The potential of leys and short rotation coppice hedges as reservoirs for Hymenoptera of cereal aphids in organic agriculture. *Agriculture, Ecosystems & Environment*, 87, 81-92.
- Lanzoni, A., Masetti, A., Plankesteiner, D., and Burgio, G. (2003) Role of field margin habitat and annual flowering plant mixture on parasitization of economic agromyzid pests. *Landscape Management For Functional Biodiversity, IOBC wprs Bulletin*, 26, 95-100.
- Lautenbach, S., Kugel, C., Lausch, A., Seppelt, R. (2011) Analysis of historic changes in regional ecosystem service provisioning using land use data. *Ecological Indicators*, 11, 676-687.
- Le Féon, V., Burel, F., Chifflet, R., Henry, M., Ricroch, A., Vaissière, B. E., Baudry, J. (2013) Solitary bee abundance and species richness in dynamic agricultural landscapes. *Agriculture, Ecosystems & Environment*, 166, 94-101.
- Le Féon, V., Schermann-Legionnet, A., Delettre, Y., Aviron, S., Billeter, R., Bugter, R., ... Burel, F. (2010) Intensification of agriculture, landscape composition and wild bee communities: a large scale study in four European countries. *Agriculture, Ecosystems & Environment*, 137, 143-150.
- Lemke, A., Poehling, H. M. (2002) Sown weed strips in cereal fields: overwintering site and "source" habitat for *Oedothorax apicatus* (Blackwall) and *Erigone atra* (Blackwall)(Araneae: Erigonidae). *Agriculture, Ecosystems & Environment*, 90, 67-80.

- Lye, G., Park, K., Osborne, J., Holland, J., Goulson, D. (2009) Assessing the value of Rural Stewardship schemes for providing foraging resources and nesting habitat for bumblebee queens (Hymenoptera: Apidae). *Biological Conservation*, 142, 2023-2032.
- Maalouly, M., Franck, P., Bouvier, J-C., Toubon, J-F., Lavigne, C. (2013) Codling moth parasitism is affected by semi-natural habitats and agricultural practices at orchard and landscape levels. *Agriculture, Ecosystems & Environment*, 169, 33-42.
- MacFadyen, S., Gibson, R., Raso, L., Sint, D., Traugott, M., Memmott, J. (2009) Parasitoid control of aphids in organic and conventional farming systems. *Agriculture, Ecosystems & Environment*, 133, 14-18.
- MacLeod, A. (1999) Attraction and retention of *Episyrphus balteatus* DeGeer (Diptera: Syrphidae) at an arable field margin with rich and poor floral resources. *Agriculture, Ecosystems & Environment*, 73, 237-244.
- Rundlöf, M., Persson, A.S., Smith, H.G., Bommarco, R. (2014) Late-season mass-flowering red clover increases bumble bee queen and male densities. *Biological Conservation*, 172, 138-145.
- Mänd, M., Mänd, R., Williams, I. H. (2002) Bumblebees in the agricultural landscape of Estonia. *Agriculture, Ecosystems & Environment*, 89, 69-76.
- Marchi, C., Andersen, L. W., Loeschke, V. (2013) Effects of land management strategies on the dispersal pattern of a beneficial arthropod. *PloS One*, 8, e66208.
- Menta, C., Leoni, A., Gardi, C., Conti, F. D. (2011) Are grasslands important habitats for soil microarthropod conservation? *Biodiversity and Conservation*, 20, 1073-1087.
- Meyer, B., Jauker, F., Steffan-Dewenter, I. (2009) Contrasting resource-dependent responses of hoverfly richness and density to landscape structure. *Basic And Applied Ecology*, 10, 178-186.
- Miñarro, M., Prida, E. (2013) Hedgerows surrounding organic apple orchards in North-west Spain: potential to conserve beneficial insects. *Agricultural and Forest Entomology*. 15, 382–390.
- Molinillo, M., Lasanta, T., García-Ruiz, J. M. (1997) Managing mountainous degraded landscapes after farmland abandonment in the Central Spanish Pyrenees. *Environmental Management*, 21, 587-598.
- Monteiro, L. B., Lavigne, C., Ricci, B., Franck, P., Toubon, J-F., Sauphanor, B. (2013) Predation of codling moth eggs is affected by pest management practices at orchard and landscape levels. *Agriculture, Ecosystems and Environment*, 166, 86-93.
- Moonen, A. C., Castro-Rodas, N., Bàrberi, P., Petacchi, R. (2006) Field margin structure and vegetation composition effects on beneficial insect diversity at farm scale: a case study on an organic farm near Pisa (Italy). *Landscape Management for Functional Biodiversity, IOBC wprs Bulletin*, 29, 77-80.

- Moreno, B., Garcia-Rodriguez, S., Canizares, R., Castro, J., Benitez, E. (2009) Rainfed olive farming in south-eastern Spain: long-term effect of soil management on biological indicators of soil quality. *Agriculture, Ecosystems & Environment*, 131, 333-339.
- Moreno, G., Obrador, J. J., Garcia, A. (2007) Impact of evergreen oaks on soil fertility and crop production in intercropped dehesas. *Agriculture, Ecosystems and Environment*, 119, 270-280.
- Morlat, R., Jacquet, A. (2003) Grapevine root system and Soil characteristics in a vineyard maintained long-term with or without interrow sward. *American Journal of Enology and Viticulture*, 54, 1-7.
- Moschini, V., Migliorini, P., Sacchetti, P., Casella, G., Vazzana, C. (2012) Presence of aphid predators in common wheat (*Triticum aestivum* L.) in organic and conventional agroecosystems of Tuscany. *Mediterranean Journal of Economics, Agriculture and Environment*, 11, 58-61.
- Murray, T. E., Fitzpatrick, U., Byrne, A., Fealy, R., Brown, M. J., Paxton, R. J. (2012) Local-scale factors structure wild bee communities in protected areas. *Journal of Applied Ecology*, 49, 998-1008.
- Nielsen, A., Steffan-Dewenter, I., Westphal, C., Messinger, O., Potts, S. G., Roberts, S. P., ... Petanidou, T. (2011) Assessing bee species richness in two Mediterranean communities: importance of habitat type and sampling techniques. *Ecological Research*, 26, 969-983.
- Novara, A., Gristina, L., SaladiNo, S. S., Santoro, A., Cerda, A. (2011) Soil erosion assessment on tillage and alternative soil managements in a Sicilian vineyard. *Soil and Tillage Research*, 117, 140-147.
- Öberg, S. (2009) Influence of landscape structure and farming practice on body condition and fecundity of wolf Araneae. *Basic and Applied Ecology*, 10, 614-621.
- Oberg, S. (2007) Diversity of Araneae after spring sowing - influence of farming system and habitat type. *Journal of Applied Entomology*, 131, 524-531.
- Ockinger, E., Smith, H. G. (2007) Semi-natural grasslands as population sources for pollinating insects in agricultural landscapes. *Journal of Applied Ecology*, 44, 50-59.
- Osborne, J. L., Martin, A. P., Shortall, C. R., Todd, A. D., Goulson, D., Knight, M. E., ... Sanderson, R. A. (2008) Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. *Journal of Applied Ecology*, 45, 784-792.
- Östman, Ö., Ekbom, B., Bengtsson, J. (2001) Landscape heterogeneity and farming practice influence biological control. *Basic and Applied Ecology*, 2, 365-371.
- Östman, Ö., Ekbom, B., Bengtsson, J., Weibull, A. C. (2001) Landscape complexity and farming practice influence the condition of polyphagous carabid Coleoptera. *Ecological Applications*, 11, 480-488.
- Paoletti, M. G. (1988) Soil Invertebrates in cultivated and uncultivated soils in north eastern Italy. *Estratto da redia*, 71, 501-563

- Paoletta, M. G., Boscoloa, P., Sommaggioa, D. (1997) Beneficial insects in fields surrounded by hedgerows in north eastern Italy. *Biological Agriculture & Horticulture*, 15, 310-323.
- Paredes, D., Cayuela, L., Campos, M. (2013) Synergistic effects of ground cover and adjacent vegetation on natural enemies of olive insect pests. *Agriculture, Ecosystems & Environment*, 173, 72-80.
- Parmentier, L., Meeus, I., Cheroutre, L., Mommaerts, V., Louwye, S., Smagghe, G. (2014) Commercial bumblebee hives to assess an anthropogenic environment for pollinator support: a case study in the region of Ghent (Belgium). *Environmental Monitoring and Assessment*, 186, 2357-2367.
- Patzold, S., Klein, C., Brummer, G. W. (2007) Run-off transport of herbicides during natural and simulated rainfall and its reduction by vegetated filter strips. *Soil Use and Management*, 23, 294-305.
- Peregrina, F., Pérez-Álvarez, E. P., Colina, M., García-Escudero, E. (2012) Cover crops and tillage influence soil organic matter and nitrogen availability in a semi-arid vineyard. *Archives of Agronomy and Soil Science*, 58(sup1), SS95-SS102.
- Persson, A. S., Smith, H. G. (2013) Seasonal persistence of bumblebee populations is affected by landscape context. *Agriculture, Ecosystems & Environment*, 165, 201-209.
- Péter, G., Kádár, F., Kiss, J., Tóth, F. (2001) Role of Field margin in the winter phenophase of carabid Coleoptera (Coleoptera: Carabidae) in winter wheat field. *Integrated Control in Cereal Crops, IOBC wprs Bulletin*, 24, 91-94
- Pfiffner, L., Luka, H. (2000) Overwintering of arthropods in soils of arable fields and adjacent semi-natural habitats. *Agriculture, Ecosystems and Environment*, 78, 215-222.
- Piekarska-Boniecka, H., Siatkowski, I., Ratynska, H. (2010) The influence of agricultural landscape structure on the flora and communities of Hymenoptera in statistical terms. *Polish Journal of Entomology*, 79, 353-365.
- Plećaš, M., Gagić, V., Janković, M., Petrović-Obradović, O., Kavallieratos, N. G., TomaNović, Ž., ... Ćetković, A. (2014) Landscape composition and configuration influence cereal aphid–parasitoid–hyperparasitoid interactions and biological control differentially across years. *Agriculture, Ecosystems & Environment*, 183, 1-10.
- Pollard, K. A., Holland, J. M. (2006) Arthropods within the woody element of hedgerows and their distribution pattern. *Agricultural and Forest Entomology*, 8, 203-211.
- Pommeresche, R., Bakken, A. K., Korsaeath, A. (2013) Abundance and diversity of Araneae in barley and young leys. *The Journal of Arachnology*, 41, 168-175.
- Pons, X., Lumbierres, B., Comas, J., Madeira, F., Starý, P. (2013) Effects of surrounding landscape on parasitism of alfalfa aphids in an IPM Crop system in Northern Catalonia. *BioControl*, 58, 733-744.

- Ponti, L., Ricci, C., Torricelli, R. (2003) The ecological role of hedges on population dynamics of *Anagrus* (Hymenoptera: Mymaridae) in vineyards of Central Italy. *Landscape Management for Functional Biodiversity, IOBC wprs Bulletin*, 26, 117-122.
- Ponti, L., Ricci, C., Veronesi, F., Torricelli, R. (2005) Natural hedges as an element of functional biodiversity in agroecosystems: the case of a Central Italy vineyard. *Bulletin of Insectology*, 58, 19-23.
- Potts, S. G., Woodcock, B. A., Roberts, S. P. M., Tscheulin, T., Pilgrim, E. S., Brown, V. K., Tallowin, J. R. (2009) Enhancing pollinator biodiversity in intensive grasslands. *Journal of Applied Ecology*, 46, 369-379.
- Pywell, R. F., James, K. L., Herbert, I., Meek, W. R., Carvell, C., Bell, D., Sparks, T. H. (2005) Determinants of overwintering habitat quality for Coleoptera and Araneae on arable farmland. *Biological Conservation*, 123, 79-90.
- Rand, T. A., Tschardtke, T. (2007) Contrasting effects of natural habitat loss on generalist and specialist aphid natural enemies. *Oikos*, 116, 1353-1362.
- Rand, T. A., van Veen, F. J., Tschardtke, T. (2012) Landscape complexity differentially benefits generalized fourth, over specialized third, trophic level natural enemies. *Ecography*, 35, 97-104.
- Raymond, L., Sarthou, J. P., Plantegenest, M., Gauffre, B., Ladet, S., Vialatte, A. (2014) Immature Diptera overwinter in cultivated fields and may significantly control aphid populations in autumn. *Agriculture, Ecosystems & Environment*, 185, 99-105.
- Reyniers, M., Maertens, K., Vrindts, E., De Baerdemaeker, J. (2006) Yield variability related to landscape properties of a loamy soil in central Belgium. *Soil and Tillage Research*, 88, 262-273.
- Ricci, B., Franck, P., Bouvier, J. C., Casado, D., Lavigne, C. (2011) Effects of hedgerow characteristics on intra-orchard distribution of larval codling moth. *Agriculture, Ecosystems and Environment*, 140, 395-400.
- Riedinger, V., Renner, M., Rundlöf, M., Steffan-Dewenter, I., Holzschuh, A. (2014) Early mass-flowering crops mitigate pollinator dilution in late-flowering crops. *Landscape Ecology*, 29, 425-435.
- Rieux, R., Simon, S., Defrance, H. (1999) Role of hedgerows and ground cover management on arthropod populations in pear orchards. *Agriculture, Ecosystems and Environment*, 73, 119-127.
- Roschewitz, I., Hücker, M., Tschardtke, T., Thies, C. (2005) The influence of landscape context and farming practices on parasitism of cereal aphids. *Agriculture, Ecosystems & Environment*, 108, 218-227.

- Roume, A., Deconchat, M., Raison, L., Balent, G., Ouin, A. (2011) Edge effects on ground Coleoptera at the woodlot-field interface are short-range and asymmetrical. *Agricultural and Forest Entomology*, 13, 395-403.
- Rundlöf, M., Nilsson, H., Smith, H. G. (2008) Interacting effects of farming practice and landscape context on bumble bees. *Biological Conservation*, 141, 417-426.
- Rusch, A., Birkhofer, K., Bommarco, R., Smith, H. G., Ekbom, B. (2014) Management intensity at field and landscape levels affects the structure of generalist predator communities. *Oecologia*, 1-13.
- Rusch, A., Bommarco, R., Jonsson, M., Smith, H. G., Ekbom, B. (2013) Flow and stability of natural pest control services depend on complexity and crop rotation at the landscape scale. *Journal of Applied Ecology*, 50, 345-354.
- Rusch, A., Valantin-Morison, M., Roger-Estrade, J., Sarthou, J. P. (2012) Local and landscape determinants of pollen beetle abundance in overwintering habitats. *Agricultural and Forest Entomology*, 14, 37-47.
- Rusch, A., Valantin-Morison, M., Roger-Estrade, J., Sarthou, J. P. (2012) Using landscape indicators to predict high pest infestations and successful natural pest control at the regional scale. *Landscape and Urban Planning*, 105, 62-73.
- Rusch, A., Valantin-Morison, M., Sarthou, J. P., Roger-Estrade, J. (2011) Multi-scale effects of landscape complexity and crop management on pollen beetle parasitism rate. *Landscape Ecology*, 26, 473-486.
- Samnegård, U., Persson, A. S., Smith, H. G. (2011) Gardens benefit bees and enhance pollination in intensively managed farmland. *Biological Conservation*, 144, 2602-2606.
- Samu, F., Beleznai, O., Tholt, G. (2013) A potential spider natural enemy against virus vector leafhoppers in agricultural mosaic landscapes—corroborating ecological and behavioral evidence. *Biological Control*, 67, 390-396.
- Sárospataki, M., Báldi, A., Batáry, P., Józán, Z., Erdős, S., Rédei, T. (2009) Factors affecting the structure of bee assemblages in extensively and intensively grazed grasslands in Hungary. *Community Ecology*, 10, 182-188.
- Saska, P., Vodde, M., Heijerman, T., Westerman, P., van der Werf, W. (2007) The significance of a grassy field boundary for the spatial distribution of Coleoptera within two cereal fields. *Agriculture, Ecosystems and Environment*, 122, 427-434.
- Scheid, B. E., Thies, C., Tschardtke, T. (2011) Enhancing rape pollen beetle parasitism within sown flower fields along a landscape complexity gradient. *Agricultural and Forest Entomology*, 13, 173-179.
- Schmidt, M. H., Tschardtke, T. (2005) The role of perennial habitats for central European farmland spiders. *Agriculture, Ecosystems and Environment*, 105, 235-242.

- Schmidt, M. H., Tschardtke, T. (2005) Landscape context of sheetweb spider (Araneae: Linyphiidae) abundance in cereal fields. *Journal of Biogeography*, 32, 467-473.
- Schmidt, M. H., Roschewitz, I., Thies, C., Tschardtke, T. (2005) Differential effects of landscape and management on diversity and density of ground-dwelling farmland spiders. *Journal of Applied Ecology*, 42, 281-287.
- Schneider, G., Krauss, J., Steffan-Dewenter, I. (2013) Predation rates on semi-natural grasslands depend on adjacent habitat type. *Basic and Applied Ecology*, 14, 614-621.
- Schroeter, L., Irmeler, U. (2013) Organic cultivation reduces barrier effect of arable fields on species diversity. *Agriculture, Ecosystems and Environment*, 164, 176-180.
- Schüepp, C., Herzog, F., Entling, M. H. (2014) Disentangling multiple drivers of pollination in a landscape-scale experiment. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20132667.
- Schüepp, C., Uzman, D., Herzog, F., Entling, M. H. (2014) Habitat isolation affects plant–herbivore–enemy interactions on cherry trees. *Biological Control*, 71, 56-64.
- Schulp, C. J., Veldkamp, A. (2008) Long-term landscape–land use interactions as explaining factor for soil organic matter variability in Dutch agricultural landscapes. *Geoderma*, 146, 457-465.
- Sciarretta, A., Trematerra, P. (2011) Spatio-temporal distribution of *Ceratitis capitata* population in a heterogeneous landscape in Central Italy. *Journal of Applied Entomology*, 135, 241-251.
- Scohier, A., Ouin, A., Farruggia, A., Dumont, B. (2013) Is there a benefit of excluding sheep from pastures at flowering peak on flower-visiting insect diversity? *Journal of Insect Conservation*, 17, 287-294.
- Scutareanu, P., Lingeman, R., Drukker, B., Sabelis, M. W. (1999) Cross-correlation analysis of fluctuations in local populations of pear psyllids and anthocorid bugs. *Ecological Entomology*, 24, 354-362.
- Sheridan, H., Finn, J. A., O'Donovan, G. (2009) Botanical rejuvenation of field margins and benefits for invertebrate fauna on a drystock farm in County Longford. *Biology and Environment- Proceedings of The Royal Irish Academy*, 109B(2), 95-106.
- Shvaleyeva, A., e Silva, F. C., Costa, J. M., Correia, A., Anderson, M., Lobo-do-Vale, R., ... Cruz, C. (2014) Comparison of methane, nitrous oxide fluxes and CO₂ respiration rates from a Mediterranean cork oak ecosystem and improved pasture. *Plant and Soil*, 374, 883-898.
- Silva, E. B., Franco, J. C. Vasconcelos, T., Branco, M. (2010) Effect of ground cover vegetation on the abundance and diversity of beneficial arthropods in citrus orchards. *Bulletin of Entomological Research*, 100, 489-499.
- Simón, N., Montes, F., Díaz-Pinés, E., Benavides, R., Roig, S., Rubio, A. (2013) Spatial distribution of the soil organic carbon pool in a Holm oak dehesa in Spain. *Plant and Soil*, 366, 537-549.

- Sjödin, N. E., Bengtsson, J., Ekbom, B. (2008) The influence of grazing intensity and landscape composition on the diversity and abundance of flower-visiting insects. *Journal of Applied Ecology*, 45, 763-772.
- Skórka, P., Lenda, M. (2010) Abandoned fields as refuges for Lepidoptera in the agricultural landscapes of Eastern Europe. In: *Insect Habitats: Characteristics, Diversity and Management*, Eds. Edina L. Harris and Newell E. Davies. Nova Science Publishers, New York, pp. 83-103.
- Smith, J., Potts, S., Eggleton, P. (2008) The value of sown grass margins for enhancing soil macrofaunal biodiversity in arable systems. *Agriculture, Ecosystems and Environment*, 127, 119-125.
- Smits, N., Dupraz, C., Dufour, L. (2012) Unexpected lack of influence of tree rows on the dynamics of wheat aphids and their natural enemies in a temperate agroforestry system. *Agroforestry Systems*, 85, 153-164.
- Sonneveld, M. P. W., Van Den Akker, J. J. H. (2011) Quantification of C and N stocks in grassland topsoils in a dutch region dominated by livestock farming. *The Journal of Agricultural Science*, 149, 63-71.
- Starý, P., Havelka, J. (2008) Fauna and associations of aphid parasitoids in an up-dated farmland area (Czech Republic). *Bulletin of Insectology*, 61, 251-276.
- Steckel, J., Westphal, C., Peters, M. K., Bellach, M., Rothenwoehrer, C., Erasmí, S., ... Steffan-Dewenter, I. (2014) Landscape composition and configuration differently affect trap-nesting Hymenoptera, wasps and their antagonists. *Biological Conservation*, 172, 56-64.
- Steffan-Dewenter, I., Leschke, K. (2003) Effects of habitat management on vegetation and above-ground nesting Hymenoptera and wasps of orchard meadows in Central Europe. *Biodiversity & Conservation*, 12, 1953-1968.
- Steffan-Dewenter, I., Tschardtke, T. (1999) Effects of habitat isolation on pollinator communities and seed set. *Oecologia*, 121, 432-440.
- Steffan-Dewenter, I., Munzenberg, U., Burger, C., Thies, C., Tschardtke, T. (2002) Scale-dependent effects of landscape context on three pollinator guilds. *Ecology*, 83, 1421-1432.
- Świtoniak, M. (2014) Use of soil profile truncation to estimate influence of accelerated erosion on soil cover transformation in young morainic landscapes, North-Eastern Poland. *Catena*, 116, 173-184.
- Tatzber, M., Stemmer, M., Spiegel, H., Katzlberger, C., Landstetter, C., Haberhauer, G., Gerzabek, M. H. (2012) ¹⁴C-labeled organic amendments: Characterization in different particle size fractions and humic acids in a long-term field experiment. *Geoderma*, 177-178, 39-48.
- Thies, C., Tschardtke, T. (1999) Landscape structure and biological control in agroecosystems. *Science*, 285, 893-895.

- Thies, C., Haenke, S., Scherber, C., Bengtsson, J., Bommarco, R., Clement, L. W., ... Tscharrntke, T. (2011) The relationship between agricultural intensification and biological control: experimental tests across Europe. *Ecological Applications*, 21, 2187-2196.
- Thies, C., Roschewitz, I., Tscharrntke, T. (2005) The landscape context of cereal aphid–parasitoid interactions. *Proceedings of the Royal Society B: Biological Sciences*, 272, 203-210.
- Thies, C., Steffan-Dewenter, I., Tscharrntke, T. (2003) Effects of landscape context on herbivory and parasitism at different spatial scales. *Oikos*, 101, 18-25.
- Thies, C., Steffan-Dewenter, I., Tscharrntke, T. (2008) Interannual landscape changes influence plant–herbivore–parasitoid interactions. *Agriculture, Ecosystems & Environment*, 125, 266-268.
- Tkaczuk C., Krzyczkowski T., Wegensteiner R.. (2012) The occurrence of entomopathogenic fungi in soils from mid-field woodlots and adjacent small-scale arable fields. *Acta Mycol.* 47, 191–202
- Trichard, A., Alignier, A., Biju-Duval, L., Petit, S. (2013) The relative effects of local management and landscape context on weed seed predation and carabid functional groups. *Basic and Applied Ecology*, 14, 235-245.
- Tscharrntke, T., Steffan-Dewenter, I., Kruess, A., Thies, C. (2002) Contribution of small habitat fragments to conservation of insect communities of grassland-cropland landscapes. *Ecological Applications*, 12, 354-363.
- Tscheulin, T., Neokosmidis, L., Petanidou, T., Settele, J. (2011) Influence of landscape context on the abundance and diversity of Hymenoptera in Mediterranean olive groves. *Bulletin of entomological research*, 101, 557-564.
- van Berkel, D. B., Verburg, P. H. (2014) Spatial quantification and valuation of cultural ecosystem services in an agricultural landscape. *Ecological Indicators*, 37, 163-174.
- Van Geert, A., Van Rossum, F., Triest, L. (2010) Do linear landscape elements in farmland act as biological corridors for pollen dispersal?. *Journal of Ecology*, 98, 178-187.
- Verboven, H. A., Uyttenbroeck, R., Brys, R., Hermy, M. (2014) Different responses of Hymenoptera and Diptera to land use in an urban–rural gradient show the importance of the nature of the rural land use. *Landscape and Urban Planning*, 126, 31-41.
- Veres, A., Tóth, F., Kiss, J., Fetykó, K., Orosz, S., Lavigne, C., ... Bohan, D. (2012) Spatio-temporal dynamics of *Orius* spp. (Heteroptera: Anthocoridae) abundance in the agricultural landscape. *Agriculture, Ecosystems & Environment*, 162, 45-51.
- Vidal, S. (1997) Factors influencing the population dynamics of *Brevicoryne brassicae* in undersown brussels sprouts. *Entomological Research in Organic Agriculture*, 15, 285-295.
- Vollhardt, I. M., Tscharrntke, T., Wäckers, F. L., Bianchi, F. J., Thies, C. (2008) Diversity of cereal aphid Hymenoptera in simple and complex landscapes. *Agriculture, Ecosystems & Environment*, 126, 289-292.

- Weihrauch, F. (2008) Overwintering of common green lacewings in hibernation shelters in the Hallertau hop growing area. *Bulletin of Insectology*, 61, 67-71.
- Westphal, C., Steffan-Dewenter, I., Tschamtkke, T. (2003) Mass flowering crops enhance pollinator densities at a landscape scale. *Ecology Letters*, 6, 961-965.
- Westphal, C., Steffan-Dewenter, I., Tschamtkke, T. (2009) Mass flowering oilseed rape improves early colony growth but not sexual reproduction of bumble bees. *Journal of Applied Ecology*, 46, 187-193.
- Wiesmeier, M., Spörlein, P., Geuß, U., Hangen, E., Haug, S., Reischl, A., ... Kögel-Knabner, I. (2012) Soil organic carbon stocks in southeast Germany (Bavaria) as affected by land use, soil type and sampling depth. *Global Change Biology*, 18, 2233-2245.
- Williams, A., Hedlund, K. (2013) Indicators of soil ecosystem services in conventional and organic arable fields along a gradient of landscape heterogeneity in southern Sweden. *Applied Soil Ecology*, 65, 1-7.
- Williams, A., Hedlund, K. (2014) Indicators and trade-offs of ecosystem services in agricultural soils along a landscape heterogeneity gradient. *Applied Soil Ecology*, 77, 1-8.
- Winkler, K., Wäckers, F. L., Termorshuizen, A. J., van Lenteren, J. C. (2010) Assessing risks and benefits of floral supplements in conservation biological control. *BioControl*, 55, 719-727.
- Winqvist, C., Bengtsson, J., Aavik, T., Berendse, F., Clement, L. W., Eggers, S., ... Bommarco, R. (2011) Mixed effects of organic farming and landscape complexity on farmland biodiversity and biological control potential across Europe. *Journal of Applied Ecology*, 48, 570-579.
- Wissuwa, J., Salamon, J. A., Frank, T. (2012) Effects of habitat age and plant species on predatory mites (Acari, Mesostigmata) in grassy arable fallows in eastern Austria. *Soil Biology and Biochemistry*, 50, 96-107.
- Woodcock, B. A., Potts, S. G., Tschulin, T., Pilgrim, E., Ramsey, A. J., Harrison-Cripps, J., Brown, V. K., Tallwin, J. R. (2009) Responses of invertebrate trophic level, feeding guild and body size to the management of improved grassland field margins. *Journal of Applied Ecology*, 46, 920-929.
- Yang, Z., Singh, B. R., Sitaula, B. K. (2004) Fractions of organic carbon in soils under different crop rotations, cover crops and fertilization practices. *Nutrient Cycling in Agroecosystems*, 70, 161-166.
- Zádorová, T., Žižala, D., Penížek, V., Čejková, Š. (2014) Relating extent of colluvial soils to topographic derivatives and soil variables in a Luvisol sub-catchment, Central Bohemia, Czech Republic. *Soil and Water Research*, 9, 47-57.
- Zaller, J. G., Moser, D., Drapela, T., Schmöger, C., Frank, T. (2008) Insect pests in winter oilseed rape affected by field and landscape characteristics. *Basic and Applied Ecology*, 9, 682-690.

Zaller, J. G., Moser, D., Drapela, T., Schmöger, C., Frank, T. (2009) Parasitism of stem weevils and pollen beetle in winter oilseed rape is differentially affected by crop management and landscape characteristics. *BioControl*, 54, 505-514.