

# Histopathology, vitellogenin and chemical body burden in mosquitofish (*Gambusia holbrooki*) sampled from six river sites receiving a gradient of stressors

Scott, P.D., Coleman, H.M., Khan, S., Lim, R., McDonald, J.A., Mondon, J., Neale, P.A., Prochazka, E., Tremblay, L.A., Warne, M.S., Leusch, F.D.L.

Author post-print (accepted) deposited by Coventry University's Repository

## Original citation & hyperlink:

Scott, P.D., Coleman, H.M., Khan, S., Lim, R., McDonald, J.A., Mondon, J., Neale, P.A., Prochazka, E., Tremblay, L.A., Warne, M.S., Leusch, F.D.L. (2017) Histopathology, vitellogenin and chemical body burden in mosquitofish (*Gambusia holbrooki*) sampled from six river sites receiving a gradient of stressors. *Science of the Total Environment*, (in press)

<https://dx.doi.org/10.1016/j.scitotenv.2017.10.148>

DOI 10.1016/j.scitotenv.2017.10.148

ISSN 0048-9697

ESSN 1879-1026

Publisher: Elsevier

**NOTICE: this is the author's version of a work that was accepted for publication in *Science of the Total Environment*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Science of the Total Environment*, in press, (2017)]**  
DOI: 10.1016/j.scitotenv.2017.10.148

© 2017, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively

**from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.**

**This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.**

1 Histopathology, vitellogenin and chemical body burden in mosquitofish (*Gambusia*  
2 *holbrooki*) sampled from six river sites receiving a gradient of stressors

3

4 Philip D. Scott <sup>a</sup>, Heather M. Coleman <sup>b,1</sup>, Stuart Khan <sup>b</sup>, Richard Lim <sup>c</sup>, James A.  
5 McDonald <sup>b</sup>, Julie Mondon <sup>d</sup>, Peta A. Neale <sup>a</sup>, Erik Prochazka <sup>a</sup>, Louis A. Tremblay <sup>e</sup>,  
6 Michael St. J. Warne <sup>a,f,2</sup>, and Frederic D.L. Leusch <sup>a,\*</sup>

7

8 <sup>a</sup> Australian Rivers Institute, Griffith School of Environment, Griffith University,  
9 Queensland, 4222, Australia

10 <sup>b</sup> School of Civil & Environmental Engineering, University of New South Wales,  
11 New South Wales, 2052, Australia

12 <sup>c</sup> School of Life Sciences, University of Technology Sydney, PO Box 123, Broadway,  
13 New South Wales, 2007, Australia

14 <sup>d</sup> Centre for Integrative Ecology, Deakin University, Warrnambool, Victoria, 3280,  
15 Australia

16 <sup>e</sup> Cawthron Institute, 98 Halifax St. East, Nelson 7042, New Zealand, and School of  
17 Biological Sciences, University of Auckland, PO Box 92019, Auckland, 1142, New  
18 Zealand

19 <sup>f</sup> Water Quality and Investigations, Department of Science, Information Technology  
20 and Innovation, Queensland Government, Queensland, 4001, Australia

21

---

<sup>1</sup> Present address: School of Pharmacy and Pharmaceutical Science, Ulster University,  
Coleraine, BT52 1SA, Northern Ireland

<sup>2</sup> Present address: Centre for Agroecology, Water and Resilience, Coventry  
University, West Midlands, United Kingdom

22 \* Corresponding Author.

23 e: [f.leusch@griffith.edu.au](mailto:f.leusch@griffith.edu.au)

24 p: +61 7 5552 7832

25 Mailing address:

26 A/Prof Frederic Leusch

27 Australian Rivers Institute, Griffith School of Environment

28 Griffith University, Gold Coast Campus

29 Southport, Queensland, 4222

30 Australia

31

32 **Abstract**

33 There are over 40,000 chemical compounds registered for use in Australia, and only a  
34 handful are monitored in the aquatic receiving environments. Their effects on fish  
35 species in Australia are largely unknown. Mosquitofish (*Gambusia holbrooki*) were  
36 sampled from six river sites in Southeast Queensland identified as at risk from a range  
37 of pollutants. The sites selected were downstream of a wastewater treatment plant  
38 discharge, a landfill, two agricultural areas, and two sites in undeveloped reaches  
39 within or downstream of protected lands (national parks). Vitellogenin analysis,  
40 histopathology of liver, kidney and gonads, morphology of the gonopodium, and  
41 chemical body burden were measured to characterize fish health. Concentrations of  
42 trace organic contaminants (TrOCs) in water were analyzed by *in vitro* bioassays and  
43 chemical analysis. Estrogenic, anti-estrogenic, anti-androgenic, progestagenic and  
44 anti-progestagenic activities and TrOCs were detected in multiple water samples.  
45 Several active pharmaceutical ingredients (APIs), industrial compounds, pesticides  
46 and other endocrine active compounds were detected in fish carcasses at all sites,

47 ranging from <4 – 4700 ng/g wet weight, including the two undeveloped sites. While  
48 vitellogenin protein was slightly increased in fish from two of the six sites, the  
49 presence of micropollutants did not cause overt sexual endocrine disruption in  
50 mosquitofish (*i.e.*, no abnormal gonads or gonopodia). A correlation between lipid  
51 accumulation in the liver with total body burden warrants further investigation to  
52 determine if exposure to low concentrations of TrOCs can affect fish health and  
53 increase stress on organs such as the liver and kidneys via other mechanisms,  
54 including disruption of non-sexual endocrine axes involved in lipid regulation and  
55 metabolism.

56

## 57 **Keywords**

58 Australia; Endocrine disruption; GeneBLAzer; micropollutant; pharmaceutical and  
59 personal care products; QuEChERS.

60

## 61 **1. Introduction**

62 There are many tens of thousands of chemicals in current use, with over 100,000  
63 chemical substances deemed to be in commercial use in the European Union  
64 (European Chemicals Agency 2017), more than 84,000 in the United States (U.S.  
65 Environmental Protection Agency 2017) and over 40,000 in Australia (Australian  
66 Department of Health 2017). Without even taking into account the large number of  
67 potential environmental transformation products, an optimistic estimate would  
68 suggest that fewer than 1% of these compounds have been monitored in wastewater  
69 treatment plant (WWTP) effluents and rivers worldwide to date (Bradley et al. 2017,  
70 Kolpin et al. 2002, Leusch et al. 2014b, Loos et al. 2009, Schäfer et al. 2011, Scott et  
71 al. 2014a, b, Tousova et al. 2017).

72 There is both a paucity of monitoring data and a lack of information regarding the  
73 toxic effects that these compounds may have on biota in the receiving environments,  
74 particularly in Australia (Woods and Kumar 2011). There is some evidence indicating  
75 that trace organic contaminants (TrOCs) and endocrine active compounds (EACs)  
76 from anthropogenic sources may negatively impact fish in Australian rivers (Table 1);  
77 however, the occurrence of endocrine disruption is not always consistent.

78 Mosquitofish (*Gambusia holbrooki*) is a wide-spread invasive species in Australia  
79 that has been particularly useful in studies on endocrine disruption (Table 1) due to a  
80 high level of sexual dimorphism, the presence of secondary sexual characteristics that  
81 can be affected by exposure to EACs and a well understood endocrinology (Leusch et  
82 al. 2006, Rawson et al. 2010). In particular, development of a specialized anal fin in  
83 males, the gonopodium, used for sexual reproduction is under hormonal control and  
84 can be reduced upon exposure to estrogenic EACs (Doyle and Lim 2002) or elongated  
85 upon exposure to androgenic EACs (Angus et al. 2001). Three studies have reported  
86 significantly reduced gonopodial length (an androgen-mediated secondary sex  
87 characteristic) in male mosquitofish at sites impacted by point (WWTP effluent) and  
88 non-point (agricultural and residential) sources (Table 1), indicating exposure to  
89 estrogenic or anti-androgenic EACs. At the same time, no significant effects were  
90 reported in mosquitofish sampled at different sites impacted by WWTP effluent,  
91 industrial contamination, and residential effluents (Table 1).

92 To help determine the significance of endocrine disruption in Australian rivers, this  
93 study prioritized six sites in Southeast Queensland for *in situ* sampling based on  
94 recent monitoring and *in vivo* exposure data (Scott et al. 2014a, b, Scott et al. 2017).  
95 The sites were chosen to represent a selection of point (*e.g.*, WWTP, landfill) and  
96 non-point sources (*e.g.*, agricultural activities) of potential EAC contamination.

97 Mosquitofish and grab water samples were collected from four impacted sites and two  
98 sites in undeveloped reaches of the catchment within or downstream of protected  
99 lands (national parks) to assess endocrine disruption using the following techniques:  
100 1) histopathology of gonads, liver and kidneys, 2) measurement of gonopodium  
101 length and vitellogenin (Vtg) concentration in adult males, 3) chemical analysis of  
102 TrOC and EAC body burden in males and females, and 4) chemical and *in vitro*  
103 analysis of grab water samples.

104 **Table 1.** Summary of *in situ* studies in Australia assessing endocrine disruption in aquatic environments. WWTP = Wastewater Treatment Plant.

Endpoint(s)	Species	Result(s)	Influence(s)	Environment	Reference	
Aromatase mRNA	<i>Lates calcarifer</i>	Slight increase	Agricultural	River/estuary	Kroon et al. (2015)	
Gonadal histology	<i>Carassius auratus</i>	No effect	Urban and WWTP	River	Kellar et al. (2014)	
	<i>Crocodylus johnstoni</i>	No effect	Agricultural	River	Yoshikane et al. (2006)	
	<i>Cyprinus carpio</i>	No effect	WWTP effluent	River	Hassell et al. (2016)	
	<i>Gambusia holbrooki</i>	No effect	Urban and WWTP	River	Kellar et al. (2014)	
	<i>Gambusia holbrooki</i>	No effect	WWTP effluent	River	Leusch et al. (2006)	
	<i>Melanotaenia fluviatilis</i>	Suppression of spermatogenesis	100% WWTP effluent	Mobile lab	Vajda et al. (2015)	
	<i>Rutilus rutilus</i>	No effect	WWTP effluent	River	Hassell et al. (2016)	
	<i>Saccostrea glomerata</i>	No effect	WWTP effluent	Marine	Andrew-Priestley et al. (2012)	
Gonopodium morphology	<i>Saccostrea glomerata</i>	No effect	WWTP effluent	River	Anderson et al. (2010)	
	<i>Gambusia holbrooki</i>	No effect	Agricultural (rural)	River	Chinathamby et al. (2013)	
	<i>Gambusia holbrooki</i>	No effect	Residential/industrial	River/estuary	Rawson et al. (2009)	
	<i>Gambusia holbrooki</i>	No effect	Urban	River	Chinathamby et al. (2013)	
	<i>Gambusia holbrooki</i>	No effect	Urban and WWTP	River	Kellar et al. (2014)	
	<i>Gambusia holbrooki</i>	No effect	WWTP effluent	River	Leusch et al. (2014a)	
	<i>Gambusia holbrooki</i>	No effect	WWTP effluent	River	Chinathamby et al. (2013)	
	<i>Gambusia holbrooki</i>	Significantly reduced length	Agricultural	Lake	Game et al. (2006)	
	<i>Gambusia holbrooki</i>	Significantly reduced length	Agricultural / residential	Lake	Game et al. (2006)	
	<i>Gambusia holbrooki</i>	Significantly reduced length	WWTP effluent	River	Batty and Lim (1999)	
	<i>Gambusia holbrooki</i>	Significantly reduced length	WWTP effluent	River	Doyle et al. (2003)	
	<i>Gambusia holbrooki</i>	Slightly increased length	WWTP effluent	River	Leusch et al. (2006)	
	Hormones (plasma)	<i>Crocodylus johnstoni</i>	No effect	Agricultural	River	Yoshikane et al. (2006)
	Morphometrics	<i>Gambusia holbrooki</i>	Slightly reduced testes weight	WWTP effluent	River	Doyle et al. (2003)
Phenotype	<i>Morula granulata</i>	Significant increase in imposex	Industrial (shipping)	Marine	Reitsema and Spickett (1999)	
	<i>Morula marginalba</i>	Significant increase in imposex	Industrial (shipping)	Marine	Andersen (2004)	
Reproductive output	<i>Gambusia holbrooki</i>	No effect on spermatozeugmata	WWTP effluent	River	Batty and Lim (1999)	
Sex ratio	<i>Gambusia holbrooki</i>	Decrease in mature males	WWTP effluent	River	Rawson et al. (2008)	
Skeletal morphology	<i>Gambusia holbrooki</i>	No effect	WWTP effluent	River	Rawson et al. (2008)	
Vitellogenin (plasma)	<i>Carassius auratus</i>	No effect	Urban and WWTP	River	Kellar et al. (2014)	
	<i>Cyprinus carpio</i>	No effect	WWTP effluent	River	Hassell et al. (2016)	
	<i>Gambusia holbrooki</i>	No effect	WWTP effluent	River	Leusch et al. (2014a)	
	<i>Rutilus rutilus</i>	No effect	WWTP effluent	River	Hassell et al. (2016)	



Vitellogenin (protein/mRNA)	<i>Tetractenos glaber</i>	Significantly increased	Agricultural / residential	River/estuary	Booth and Skene (2006)
	<i>Tetractenos glaber</i>	Significantly increased	WWTP effluent	River/estuary	Booth and Skene (2006)
	<i>Melanotaenia fluviatilis</i>	No effect	100% WWTP effluent	Mobile lab	Vajda et al. (2015)
Vitellogenin mRNA	<i>Lates calcarifer</i>	Significantly increased	Agricultural	River/estuary	Kroon et al. (2015)
	<i>Plectropomus sp.</i>	Significantly increased	Agricultural	Coastal lagoons	Kroon et al. (2015)
	<i>Saccostrea glomerate</i>	Significantly increased	WWTP effluent	Marine	Andrew-Priestley et al. (2012)
	<i>Saccostrea glomerate</i>	Significantly increased	WWTP effluent	River	Anderson et al. (2010)

---

105 **2. Experimental section**

106 *2.1. Site identification*

107 Historical chemical and *in vitro* (endocrine activity) monitoring data and catchment  
108 analysis (Scott et al. 2014a, b) were used to identify four sites in Southeast  
109 Queensland (sites labelled AGR1, AGR2, WWEF, LNDF) with some of the highest  
110 chemical concentrations and estrogenic activity from a previous year-long survey  
111 (2011-2012 data in Table 5), and one site with low anthropogenic contaminants in an  
112 undeveloped part of the catchment as a comparison site (UND1) Catchment analysis  
113 was used to identify another undeveloped comparison site (UND2), which had not  
114 previously been monitored. Basic physico-chemical measurements and more details  
115 about the sites are provided in the supplementary material (SI Table S1 and kml  
116 geolocation file). Site designation was assigned based on the most significant land-use  
117 in the catchment at each sampling location: AGR1 and AGR2 were located within  
118 areas of agricultural land-use, WWEF was influenced primarily by wastewater  
119 effluent, and LNDF was a short distance downstream of a landfill site. Sites UND1  
120 and UND2 were located in undeveloped areas within or downstream of national parks  
121 that experience some recreational use, but otherwise no known anthropogenic  
122 pollution source.

123

124 *2.2. Sampling*

125 Mosquitofish were sampled between April and May 2013 using an electro-fishing unit  
126 set to 100 Hz, 20% duty cycle and 225 V, until a sufficient number of fish were  
127 obtained for analysis, or until fishing effort dropped below one fish per 10 min  
128 (whichever came first) at all six sites. A total of 190 fish were captured for this study,  
129 ranging from 14 (AGR2) to 47 (WWEF) with a median of 29.5 across all sites (Table

130 2). All animal handling was conducted with respect and in accordance with  
131 Department of Science, Information Technology and Innovation (DSITI) animal  
132 ethics permit. Fish were euthanized on site in 80 mg/L Aqui-S anesthetic (Lower  
133 Hutt, New Zealand). Wet weight, standard length and gonopodium length (from the  
134 body following the curve of the gonopodium to the tip) were then immediately  
135 recorded to the nearest tenth of a mg or mm, respectively (Table 2). Condition factor  
136 (K) was calculated using the following equation:  $K=10^5 \times W/L^3$ , where W is the wet  
137 weight (in g) and L is the standard length (in mm). The gonopodium index was  
138 calculated as gonopodium length (mm) divided by standard length (mm).

139 Fish carcasses were split into three roughly equivalent groups by sex for each one of  
140 the three separate analyses: Vtg, body burden and histology. As a limited number of  
141 fish were captured at site AGR2, biochemical analyses were prioritized and fish were  
142 split into two groups only for Vtg and body burden analysis. Male and female whole  
143 fish were then placed in 50 mL of Davidson's Fixative (pure ethanol, 10% neutral  
144 buffered formalin, glacial acetic acid and deionized water at a ratio of 3:2:1:3) for  
145 histological analysis. Fish were transferred to 50 mL pure ethanol after 48 h in  
146 Davidson's Fixative until slide preparation (Section 2.3). Fish selected for whole body  
147 Vtg analysis (Section 2.4) or body burden analysis (Section 2.5) were wrapped in  
148 aluminium foil and frozen in liquid nitrogen, and then stored at -80°C. A 2-L water  
149 grab sample was also collected when fish were sampled at each site, adjusted to pH 2  
150 using 12 M HCl, stored at 4°C and extracted within 24 h for chemical and *in vitro*  
151 analyses (Section 2.6).

152

### 153 2.3. Histopathology

#### 154 2.3.1. Slide preparation

155 Mosquitofish were removed from their ethanol solution, decapitated with a sharp  
156 blade, sliced along the sagittal plane (longitudinally) to fully expose the internal  
157 cavity and immediately fixed in 100% ethanol, then triple rinsed in 100% ethanol for  
158 24 h, cleared in histolene for 4 h and then impregnated with paraffin for 3 h. Wax  
159 impregnated sections were then transferred to disposable molds and embedded in  
160 paraffin. Tissue was sectioned to 5  $\mu\text{m}$  thickness using a microtome (Micron HS  
161 355S) and stained with standard hematoxylin and eosin (H&E).

162

### 163 2.3.2. Pathology analysis

164 Gonads, liver and kidneys were all visually inspected for pathology under a  
165 microscope (Nikon Eclipse 80i, 4 $\times$ , 10 $\times$ , and 20 $\times$  magnification; Nikon, Sydney,  
166 NSW). The classification of “healthy” testes was based on the presence of  
167 spermatocytes in all stages of development (SI Fig. S1A), while ovaries were  
168 considered to be of good reproductive health if they consisted of several stages of  
169 follicular development (SI Fig. S1B), as proposed by Hou et al. (2011). Livers were  
170 assessed for fat storage (“lacy liver” or “fatty liver”), granulomas,  
171 liquification/haemorrhage, and degenerative fatty necrosis. Kidneys were examined  
172 for liquification/haemorrhage and inclusions. Liquification was defined as liquid  
173 produced through cellular degeneration and haemorrhage was defined as blood  
174 breaching the circulatory system and spreading into tissues (liver and/or kidneys).  
175 Inclusions were defined as the separation of kidney tissues from the renal corpuscles,  
176 beyond that of the Bowman’s space (space between parietal and visceral layers of the  
177 Bowman’s capsule; Genten et al. 2009). Liver and kidney pathology was described  
178 quantitatively by calculating the percentage of affected area relative to total area of  
179 the visible organ. Preliminary analysis determined significant effect variation

180 resulting from using smaller cross-sections/higher magnification, and sections were  
181 therefore selected based on the largest cross-section of targeted organ available in  
182 order to obtain the most representative section and minimize variation. With the  
183 exception of liver fat storage, all tissue areas were calculated using Nikon NIS-  
184 Elements BR software (Tokyo, Japan) (SI Fig. S2A). Liver fat storage area was  
185 determined using a combination of Adobe Photoshop CS 6 (California, USA) and FIJI  
186 (Image J version 1.48o for OSX, National Institutes of Health, USA). Colour images  
187 (SI Fig. S2B) were first converted to binary images in Photoshop CS 6 using the  
188 threshold function (SI Fig. S2C), followed by area analysis using the threshold and  
189 analyse particle functions in FIJI.

190

#### 191 *2.4. Vitellogenin analysis by LC-MS/MS*

192 Vitellogenin (Vtg) protein content was determined by liquid chromatography as  
193 described in Scott et al. (2017). In brief, frozen whole fish were homogenized on ice  
194 using a tissue homogenizer in 1:4 (mass/volume ratio) buffer of 3 mM Tris and 0.1  
195  $\mu$ M phenylmethylsulfonyl fluoride (PMSF). Thereafter the homogenate was  
196 centrifuged at 12000 g for 98 min at 4°C and the supernatant was stored at -80°C until  
197 analysis. The homogenate protein content was  $10.5 \pm 2.9$  mg/mL, as quantified using  
198 the Bradford method (Bradford, 1976).

199 Tryptic digestions were conducted using the In-Solution Tryptic Digestion and  
200 Guanidination Kit (Thermo Fisher Scientific, Victoria, Australia) following the  
201 manufacturer's protocol. Briefly, 1  $\mu$ L of sample was added to ammonium  
202 bicarbonate digestion buffer and dithiothreitol reducing buffer, along with *Gallus*  
203 *gallus* lysozyme, which was spiked into each sample at a final concentration of 16.1  
204  $\mu$ g/mL as a loading control. Samples were incubated at 95°C for 5 min, after which

205 iodoacetamide alkylation buffer was added and the samples were incubated at room  
206 temperature for 20 min in the dark. After incubation, 1  $\mu$ L of activated trypsin was  
207 added to each sample, giving a final concentration of 3.2  $\mu$ g/mL. The sample was  
208 incubated at 37°C for 3 h, then at 30°C overnight.

209 All chromatographic separations were performed using a 5  $\mu$ L injection volume onto  
210 an Agilent 1290 HPLC, fitted with an Phenomenex Aeris C8 column (column  
211 dimensions 2.1  $\times$  100 mm with 1.8  $\mu$ M particle size, 100 Å pore size), and high  
212 resolution mass spectral data were acquired on an Agilent 6530 QTOF using an ESI  
213 source fitted with Agilent Jetstream technology, as described in Scott et al. (2017).

214

## 215 *2.5. Body burden analysis*

### 216 *2.5.1. Extraction of organic compounds from whole fish*

217 A “quick, easy, cheap, effective, rugged and safe” (QuEChERS) method was applied  
218 to extract organic compounds from whole male and female mosquitofish for chemical  
219 analysis. Originally developed for pesticide residue analysis in vegetable produce  
220 (Anastassiades et al. 2003), QuEChERS has recently been validated for extraction of  
221 TrOCs in fish (Lopes et al. 2012, Munaretto et al. 2013, Norli et al. 2011). Frozen fish  
222 were sectioned finely (using a razor blade) and placed into 1 mL deionized water with  
223 an internal standard consisting of isotope-labelled analytes (details in SI Table S2).

224 The tissue was homogenized for 30 s using an Ultra-Turrax (IKA, Malaysia) at 4000  
225 rpm. The fish homogenate was transferred to a 50 mL centrifuge tube with 8 mL of  
226 1% glacial acetic acid in acetonitrile and the contents of one QuEChERS extraction  
227 packet (6 mg MgSO<sub>4</sub>, 1.5 mg sodium acetate; Agilent, Victoria, Australia). The tube  
228 was mixed at 400 rpm with a platform mixer (Ratek, Victoria, Australia) for 1 min  
229 and then centrifuged at 3000 g for 5 min (Hercules Multifuge X3R, Thermo

230 Scientific, Victoria, Australia). The supernatant was transferred to a pre-made 15 mL  
231 dispersive solid phase extraction (SPE) centrifuge tube (50 mg primary secondary  
232 amine, 50 mg graphitized carbon black, 150 mg MgSO<sub>4</sub>; Agilent, Victoria, Australia),  
233 mixed vigorously again for 30 s and centrifuged again at 3000 g for 5 min. The  
234 supernatant was collected and evaporated under a gentle nitrogen stream and  
235 reconstituted into the equivalent volume of methanol (in  $\mu\text{L}$ ) corresponding to the  
236 original wet weight of the fish (in mg) (*e.g.* extract from a fish weighing 150 mg was  
237 reconstituted into 150  $\mu\text{L}$  methanol).

238

#### 239 *2.5.2. Chemical analysis of body residue*

240 Fish whole body homogenates were analyzed for 38 compounds including 1 industrial  
241 compound, 1 personal care product, 19 active pharmaceutical ingredients (APIs), 5  
242 pesticides, 10 steroids and 2 synthetic hormones (Table 4). Chemical analysis was  
243 performed using LC-MS/MS and GC-MS/MS following previously detailed methods  
244 (Scott et al. 2014a, b, Trinh et al. 2011, Vanderford and Snyder 2006). Concentrations  
245 were corrected to account for any losses during extraction by adding an internal  
246 standard prior to extraction (see SI Table S2).

247

#### 248 *2.6. Analysis of water samples*

##### 249 *2.6.1. Solid phase extraction and chemical analysis*

250 Grab water samples obtained concurrently with fish sampling were concentrated using  
251 SPE (Oasis HLB SPE cartridges; 500 mg sorbent, 6 cc; Waters, New South Wales,  
252 Australia) for chemical analysis and *in vitro* bioassays. The SPE was performed as  
253 previously described in Scott et al. (2014b). Chemical analysis of water extracts was  
254 performed using LC-MS/MS and GC-MS/MS as described in Section 2.5.2 except for

255 ethinylestradiol (EE2), which was measured by a commercially available enzyme-  
256 linked immunosorbent assay (ELISA; Takiwa Chemical Industries, Japan) as detailed  
257 in (Scott et al. 2014a).

258

#### 259 *2.6.2. Bioassay of water samples*

260 Endocrine activity was measured using three CellSensor GeneBLAzer assays  
261 (Invitrogen, ThermoFisher Scientific, New South Wales, Australia) to test for  
262 estrogenic (ER $\alpha$ ), androgenic (AR) and progestagenic (PR) receptor induction  
263 (Wilkinson et al. 2008), in both agonist and antagonist modes. The assays were  
264 performed as previously described in Escher et al. (2014).

265

#### 266 *2.7. Statistical analysis*

267 Chemical occurrence and morphometric data were not normally distributed, and thus  
268 non-parametric Kruskal-Wallis test followed by Dunn's multiple comparison test  
269 were used to determine significant differences ( $\alpha = 0.05$ ) between sample sites. All  
270 statistics were performed using IBM SPSS Statistics 21 (New York, USA). Analysis  
271 of Vtg LC-MS data was performed as previously described in Scott et al. (2017). As  
272 there is no pure Vtg protein standard for mosquitofish, whole body Vtg protein  
273 concentration was expressed as fold Vtg expression compared to unexposed  
274 laboratory reference males as described in Scott et al. (2017).



275 **Table 2.** Sample sizes, sex ratio and morphological measurements for mosquitofish (*Gambusia holbrooki*) collected at six sites in Southeast  
 276 Queensland. Mass, standard length and gonopodium index data are presented as average  $\pm$  standard error of the mean.

Site label	AGR1	AGR2	WWEF	LNDF	UND1	UND2
Downstream of ...	Agricultural	Agricultural	WWTP	Landfill	Undeveloped	Undeveloped
<b>Sample size for</b>						
Histology (n) <sup>1</sup>	9 (2/7)	0 (0/0)	15 (9/6)	7 (5/2)	6 (2/4)	22 (11/11)
Vitellogenin (n) <sup>2</sup>	6	5	11	10	5	10
QuEChERS (n)	13 (5/8)	9 (4/5)	21 (9/12)	14 (7/7)	13 (3/10)	14 (6/8)
<b>Male (n)</b>	<b>13</b>	<b>9</b>	<b>29</b>	<b>22</b>	<b>10</b>	<b>27</b>
Mass (mg)	198.6 $\pm$ 11.1	208.0 $\pm$ 9.1	164.5 $\pm$ 6.7	296.9 $\pm$ 84.2	177.0 $\pm$ 20.6	173.6 $\pm$ 8.1
Standard length (mm)	23.9 $\pm$ 0.4	24.1 $\pm$ 0.6	24.0 $\pm$ 0.3	24.8 $\pm$ 0.4	22.9 $\pm$ 1.0	22.7 $\pm$ 0.4
Condition factor (K) <sup>3</sup>	1.45 $\pm$ 0.03	1.48 $\pm$ 0.05	1.34 $\pm$ 0.02	1.94 $\pm$ 0.58	1.48 $\pm$ 0.04	1.53 $\pm$ 0.09
Gonopodium index <sup>4</sup>	0.31 $\pm$ 0.01	0.32 $\pm$ 0.01	0.31 $\pm$ 0.01	0.32 $\pm$ 0.01	0.30 $\pm$ 0.02	0.32 $\pm$ 0.01
<b>Female (n)</b>	<b>15</b>	<b>5</b>	<b>18</b>	<b>9</b>	<b>14</b>	<b>19</b>
Mass (mg)	340.3 $\pm$ 29.7 ab	321.2 $\pm$ 39.0 ab	189.4 $\pm$ 13.1 b	569.0 $\pm$ 63.9 a	311.6 $\pm$ 56.2 b	277.1 $\pm$ 35.0 b
Standard length (mm)	27.5 $\pm$ 0.7 ab	27.5 $\pm$ 1.3 ab	23.5 $\pm$ 0.5 b	32.9 $\pm$ 1.0 a	26.2 $\pm$ 1.6 b	24.9 $\pm$ 1.2 b
Condition factor (K) <sup>3</sup>	1.58 $\pm$ 0.05	1.53 $\pm$ 0.07	1.42 $\pm$ 0.02	1.56 $\pm$ 0.08	1.53 $\pm$ 0.04	1.83 $\pm$ 0.22

277 <sup>1</sup> total n (male n/ female n); <sup>2</sup> male fish only; <sup>3</sup> Condition factor (K) calculated as  $K = 100,000 \times W / L^3$ , where W is the wet weight (in g) and L  
 278 is the standard length (in mm); <sup>4</sup> Gonopodium index = gonopodium length (in mm) / standard length (in mm); Abbreviations: "WWTP" =  
 279 Wastewater treatment plant. Different letters indicate statistically different groups for each measure (Kruskal-Wallis ANOVA on ranks).

### 280 3. Results and discussion

#### 281 3.1. Morphological measurements and histopathology

282 A total of 190 mosquitofish were sampled, ranging from 14 (site AGR2) to 47 (site WWEF).  
283 Female fish were larger and longer downstream of the landfill site (LNDF) compared to those  
284 at the undeveloped (UNDF1 and UNDF2) site and downstream of the wastewater discharge  
285 (WWEF) (Kruskall-Wallis ANOVA on rank,  $p < 0.05$ ), but all female fish had a similar  
286 condition factor ( $p = 0.148$ ; Table 2). There were no significant differences in morphological  
287 measures for male fish (Kruskall-Wallis ANOVA on ranks,  $p > 0.05$ ; Table 2). Mosquitofish  
288 from each site (with the exception of site AGR2, due to the small sample size) were used for  
289 histological analysis of gonads, liver and kidney tissues (Table 3). There was no evidence of  
290 ovotestis in any of the specimens. Furthermore, there was no significant difference ( $p > 0.05$ )  
291 in gonopodium elongation (calculated as gonopodial index) between male fish from the  
292 different sites (Table 2).

293 Percent liver fat was highly variable with mosquitofish from undeveloped sites (UND1 and  
294 UND2) exhibiting anywhere between 1 and 55% liver fat (relative to total liver area), and  
295 livers from mosquitofish from impacted sites (AGR1, WWEF and LNDF) exhibiting between  
296 7 and 72% fat. The only statistically significant result was a higher liver fat content at the  
297 wastewater effluent downstream site (WWEF) compared to that at sites AGR1 and UND2  
298 (Kruskal-Wallis;  $p = 0.020$  and  $p < 0.001$ , respectively). Haemorrhage and liquification in liver  
299 tissue was minimal (average of  $4.3 \pm 0.9\%$ ), and there were no significant differences  
300 between fish from the various sites ( $p > 0.05$ ).

301 The average area affected by haemorrhages and liquification in kidney tissue of fish from all  
302 sites was  $23.0 \pm 2.4\%$ . At impacted sites, averages varied from 15.0% (site AGR1) to 23.1%  
303 (site WWEF), while those at the two undeveloped sites displayed higher affected areas ( $32.9$   
304  $\pm 11.1\%$  and  $24.9 \pm 3.4\%$  for UND1 and UND2, respectively); however, they were not

305 significantly different (Kruskal-Wallis;  $p>0.05$ ). Mosquitofish sampled at UND1 and UND2  
306 exhibited less kidney tissue damage from inclusions ( $1.3 \pm 0.7\%$  and  $1.8 \pm 0.5\%$ ,  
307 respectively) compared to that in fish from the other sites (3.1% to 5.8%), but again the data  
308 were quite variable and there were no statistically significant differences between fish from  
309 the various sites (Kruskal-Wallis;  $p>0.05$ ). Based on histological analysis (Table 3),  
310 mosquitofish from impacted sites (sites AGR1, WWEF and LNDF) were not dissimilar to  
311 fish from undeveloped locations (UND1 and UND2), suggesting that there were no gross  
312 adverse effects on this fish species at the sites monitored in this study.

313 The histological analysis of the gonads did not produce any evidence of endocrine disruption  
314 in mosquitofish. All gonads inspected were healthy and there was no evidence of ovotestis  
315 tissue. This is consistent with previous Australian studies, which have usually found no  
316 evidence of ovotestis tissue in fish exposed to treated municipal sewage effluent (Table 1).

317

318 **Table 3.** Analysis of histopathology in mosquitofish gonads, livers and kidneys from three impacted and two undeveloped sites. Data (average  $\pm$   
319 SEM) are expressed as percent area affected relative to total organ area. Histopathology was not carried out at site AGR2 because the few fish  
320 collected (Table 2) were only sufficient for biomarker and chemical analysis.

Pathology	Impacted sites			Undeveloped sites	
	AGR1	WWEF	LNDF	UND1	UND2
<b><i>Gonads</i></b>					
n (male/female)	5 (2/3)	12 (9/3)	4 (2/2)	4 (2/2)	21 (11/8)
Condition	Healthy	Healthy	Healthy	Healthy	Healthy
<b><i>Liver</i></b>					
n (male/female)	8 (2/6)	15 (9/6)	7 (5/2)	6 (2/4)	19 (10/9)
Fatty (lacy) % *	25.6 $\pm$ 4.8 bc	50.0 $\pm$ 3.8 ab	36.0 $\pm$ 6.9 abc	38.7 $\pm$ 2.8 abc	20.3 $\pm$ 3.7 bc
Haem/Liq %	6.4 $\pm$ 3.6	2.9 $\pm$ 0.6	4.1 $\pm$ 1.7	4.5 $\pm$ 1.7	4.7 $\pm$ 1.9
<b><i>Kidney</i></b>					
n (male/female)	8 (2/6)	14 (8/6)	5 (4/1)	6 (2/4)	17 (9/8)
Haem/Liq %	15.0 $\pm$ 4.7	23.1 $\pm$ 5.2	17.9 $\pm$ 2.7	32.9 $\pm$ 11.1	24.9 $\pm$ 3.4
Inclusion %	3.1 $\pm$ 1.1	4.2 $\pm$ 0.8	5.8 $\pm$ 1.5	1.3 $\pm$ 0.7	1.8 $\pm$ 0.5


321 "Haem/Liq" = percentage of area of kidney affected by haemorrhage and liquification. \* Letters indicate statistically different groups (Kruskall-  
322 Wallis,  $p < 0.05$ ).

323

324 3.2. Biomarker analysis

325 Low concentrations of Vtg protein were detected in all male fish (Fig. 1A). This was not  
326 unexpected, as trace concentrations of Vtg in males are not uncommon due to low levels of  
327 circulating natural estrogens in the blood (Bowman et al. 2000). Fish from UND1 had the  
328 lowest Vtg protein, while fish from AGR1 and the other undeveloped site (UND2) had  
329 significantly elevated Vtg protein compared to those from unexposed laboratory reference  
330 male fish (NC; Mann-Whitney test,  $p=0.041$  and  $p=0.044$ , respectively). It is worth noting  
331 that chemical analysis likewise suggests that site UND2 may not be as “pristine” as expected,  
332 with EE2 detected at 0.07 ng/L (Table 5) in a water grab sample and one fish from the site  
333 with 25 ng/g EE2 (Table 4). Vtg protein levels in fish from all other sites were not  
334 significantly different from unexposed laboratory reference males ( $p>0.05$ ). Fish from the  
335 AGR1 site had the highest Vtg protein concentration (up to 2.97-fold the concentration of  
336 unexposed laboratory reference male fish). With the exception of site UND2, there was good  
337 agreement between Vtg protein level and estrogenic activity in the water (determined by *in*  
338 *vitro* bioassay;  $p = 0.032$ ,  $R^2 = 0.828$ , Fig. 1B and Table 5). The discrepancy between  
339 estrogenic activity (from a snapshot grab sample) and Vtg protein level (the result of long-  
340 term exposure to estrogenic stimulation) suggests prior but intermittent estrogenic exposure  
341 at this site.

342



343

344 **Fig. 1.** Vitellogenin (Vtg) protein levels in male mosquitofish sampled at four impacted and  
 345 two undeveloped sites in Queensland, Australia. A) Vtg protein was significantly induced in  
 346 mosquitofish sampled at the first agricultural site (AGR1) and the second undeveloped site  
 347 (UND2) compared to unexposed laboratory reference male (NC), but not other sites. \*  
 348 indicates statistically significant difference from male negative control (Mann-Whitney test,  
 349  $p < 0.05$ ). B) *In vitro* estrogenic activity was correlated with Vtg protein levels in most  
 350 samples, except for UND2.

351

### 352 3.3. Body burden

353 Limits of quantification (LOQ) for the 38 analyzed compounds ranged from 4 to 61 ng/g wet  
354 weight (ww; Table 4), in the same range reported in other studies (Munaretto et al. 2013,  
355 Togunde et al. 2012). The majority of fish tissues analyzed (52/87, or 60%) contained at least  
356 one TrOC, with a maximum of eight (out of 38) compounds in one mosquitofish from site  
357 WWEF, which was also the site with the highest liver fat content (Table 3). Fish from site  
358 WWEF had the highest detection frequency of synthetic compounds in fish homogenates (*i.e.*  
359 number of fish with at least one quantifiable TrOC other than the natural hormones; 81%)  
360 and the highest average number of compounds per fish ( $2.6 \pm 0.4$ ). The analytical method  
361 used here was developed specifically for wastewater-derived TrOCs (Vanderford and Snyder  
362 2006), and thus this result is not unexpected. This was followed by UND1 (77%;  $2.5 \pm 0.7$ ),  
363 AGR2 (56%,  $1.2 \pm 0.4$ ), UND2 (43%;  $1.4 \pm 0.6$ ), LNDF (21%;  $1.1 \pm 1.4$ ) and AGR1 (15%;  
364  $0.4 \pm 0.3$ ). In terms of chemical complexity, site WWEF again had the highest total number  
365 of different TrOC detected in fish homogenates at 49% (16/38), followed by UND2 at 34%  
366 (13/38), LNDF and UND1 at 29% (11/38), AGR2 at 21% (8/38) and AGR1 at 11% (4/38).

367 The industrial compound tris(2-chloroethyl) phosphate (TCEP) was detected in 14% of fish  
368 carcasses (12 of 84) and had the highest overall concentration (4703 ng/g ww in a fish from  
369 site WWEF). It was detected in fish at site UND1 (46%; maximum of 1191 ng/g ww) and  
370 fish at site WWEF (29%) only. While TCEP was not detected in grab water samples from site  
371 UND1 at the time of sampling (Table 5; Apr/May 2013), it had been detected in the grab  
372 water samples at that site on previous occasions (Table 5; 2011-2012). The result illustrates  
373 that even “remote” sites can be intermittently contaminated by human TrOCs in developed  
374 regions of the world.

375 Active pharmaceutical ingredients (APIs) were detected in 40% of all fish samples, with a  
376 maximum of five APIs detected in one fish (at site LNDF). Clozapine, an antipsychotic drug,  
377 was the most commonly detected API with an average detection frequency of 17% and a  
378 maximum concentration of 155 ng/g ww. It was detected most frequently in fish downstream  
379 of the wastewater discharge (WWEF, 43%). Other commonly detected APIs include the anti-  
380 histamine hydroxyzine and the anxiety medication meprobamate (13 and 11% of all fish  
381 carcasses, respectively). Omeprazole, a proton pump inhibitor, was detected at the highest  
382 concentrations (1017 ng/g ww) in one fish from site WWEF, but was otherwise only detected  
383 in a few samples (5% of total fish carcasses; Table 4). Fluoxetine, a selective serotonin  
384 reuptake inhibitor, was detected at a maximum concentration of 240 ng/g ww at WWEF  
385 (Table 4). Fluoxetine was identified in white sucker (*Catostomus commersonii*) liver at a  
386 maximum concentration of 80 ng/g ww in a US study (Ramirez et al. 2009).

387 The personal care product caffeine was detected in 10% of fish carcasses, with a maximum  
388 concentration of 74 ng/g wet weight. It was not detected in fish from site AGR1 or AGR2  
389 (Table 4).

390 At least one of five pesticides analyzed (atrazine, chlorpyrifos, diazinon, linuron and  
391 simazine) was detected in 26% of mosquitofish samples. Chlorpyrifos, one of the most  
392 widely used insecticides in Australia (ATSE 2002), was the most commonly detected  
393 compound overall and quantified in 19% (16 of 84) of carcasses. It was detected only in fish  
394 from site WWEF (48%; 10 of 21) and UND1 (46%; 6 of 13). The herbicide results confirm  
395 that urban wastewater treatment plant contribute to a great extent to herbicide pollution of  
396 surface water (Nitschke and Schüssler 1998).

397 Nine natural and three synthetic steroid hormones were measured in whole body homogenate  
398 with LOQs ranging from 4-61 ng/g ww (Table 4). Natural hormones were detected in 30%




399 (25 of 87) of mosquitofish sampled. The only hormones not detected were  $17\alpha$ -estradiol,  
400 estriol and etiocholanolone. The natural estrogen estrone was the most frequently detected  
401 hormone (13%; 11 of 87), with a maximum concentration of 105 ng/g. The androgen  
402 androstenedione was detected in 19% of mosquitofish from WWEF, with maximum  
403 concentration of 88 ng/g ww. Androsterone was detected twice at concentrations exceeding  
404 1000 ng/g wet weight. Testosterone and dihydrotestosterone were also detected at their  
405 maximum concentrations (357 and 119 ng/g wet weight, respectively) in fish from site  
406 WWEF. Mosquitofish from WWEF and AGR2 had the highest number of different hormones  
407 detected (5 of 12), and concentrations were typically highest in WWEF. Only one of the three  
408 synthetic hormones measured was detected: the synthetic hormone  $17\alpha$ -ethinylestradiol  
409 (EE2) was detected in two fish (sites AGR2 and UND2, at 37 and 25 ng/g ww, respectively).  
410 Site UND2 has no known WWTP influence and is in a rural area so the presence of this  
411 compound may be due to defective septic systems, leaking or overflowing sewers or  
412 recreational activities.

413 It is difficult to relate any of the body burden concentrations to possible adverse effects, as  
414 most studies to that end measure water concentration, not body burden. To put body burden  
415 concentrations for APIs in relative context, an approach could be to compare body residues to  
416 the human daily therapeutic dose, assuming a 70-kg adult. While APIs do not necessarily  
417 produce similar effects in humans and aquatic wildlife (Rand-Weaver et al. 2013), this  
418 comparison is presented here as a means to put aquatic animal exposure in context with  
419 intentional human therapeutic dosing. If this is done, only two APIs occur within the range of  
420 human daily therapeutic dose. At 1017 ng/g ww, omeprazole was present at  $2\times$  the equivalent  
421 human daily therapeutic dose to treat ulcers (40 mg/day converts to 571 ng/g for a 70-kg  
422 adult). Fluoxetine was detected at 240 ng/g ww, comparable to the equivalent daily human  
423 therapeutic dose to treat depression (20 mg/day converts to 286 ng/g for a 70-kg adult). Other

424 detected APIs, such as dilantin, hydroxyzine, meprobamate and metformin, were present at  
425 concentrations that were one to three orders of magnitude lower than the equivalent human  
426 therapeutic dose in a 70-kg adult.

427 Depending on octanol-water partition coefficients, the concentrations of compounds that  
428 were found in fish tissue and not in water samples may decrease without sustained exposure.  
429 For example, carbamazepine was detected in liver and muscle tissue of bluntnose minnows  
430 after the first day of a 28 d exposure to 298  $\mu\text{g/L}$ , but decreased within a day of depuration  
431 and had returned to a baseline concentration after 14 d depuration (Garcia et al. 2012).

432 Some concentrations of compounds present in the tissue samples were very high (*e.g.*, TCEP,  
433 omeprazole, chlorpyrifos, androsterone, dihydrotestosterone, fluoxetine; Table 4). Although  
434 histological analysis did not identify endocrine disruption of sexual axes as a problem,  
435 increased body burden of these TrOCs may induce organism stress in other ways and on  
436 other endocrine functions (*e.g.*, glucocorticoid). For instance, WWEF was the most polluted  
437 site (the fish had higher liver fat content and greater body burden, and the water was  
438 chemically more complex compared to that of other sites, Table 5). This could indicate a  
439 correlation between water chemistry, chemical body burden and ultimately organism stress  
440 such as alteration in lipid metabolism (Fig. 2). While fish showed no observable effects of  
441 sexual endocrine disruption, further studies should investigate whether TrOCs in their  
442 environment are inducing stress on exposed organisms in different ways, such as via other  
443 modes of action, oxidative stress, inflammation, etc.



444

445 **Fig. 2.** Correlation between body burden as the sum of all trace organic contaminants  
 446 (TrOCs) detected in fish carcasses vs. lacy liver in mosquitofish captured from 6 sites in  
 447 Southeast Queensland.

448

449 **Table 4.** Body burden analysis (ng/g wet weight) of mosquitofish from six rivers across South East Queensland, Australia.

	Impacted sites (ng/g wet weight)									Undeveloped sites (ng/g wet weight)									
	LOQ (ng/g)	AGR1 (n = 13)			AGR2 (n = 9)			WWEF (n = 21)			LNDF (n = 14)			UND1 (n = 13)			UND2 (n = 14)		
		DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max
<i>Industrial compound</i>																			
TCEP	38	0%	<38	<38	0%	<38	<38	<b>29%</b>	<b>653</b>	<b>4703</b>	0%	<38	<38	<b>46%</b>	<b>1148</b>	<b>1191</b>	0%	<38	<38
<i>Pharmaceutical ingredients</i>																			
Amtriptyline	19	0%	<19	<19	<b>11%</b>	<b>13</b>	<b>63</b>	0%	<19	<19	0%	<19	<19	0%	<19	<19	<b>7%</b>	<19	<b>88</b>
Atenolol	19	<b>8%</b>	<19	<b>150</b>	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Carbamazepine	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Clozapine	19	0%	<19	<19	0%	<19	<19	<b>43%</b>	<b>102</b>	<b>155</b>	<b>7%</b>	<19	<b>33</b>	<b>31%</b>	<b>67</b>	<b>113</b>	0%	<19	<19
Diazepam	19	<b>8%</b>	<19	<b>27</b>	0%	<19	<19	0%	<19	<19	<b>7%</b>	<19	<b>32</b>	0%	<19	<19	0%	<19	<19
Dilantin	19	0%	<19	<19	0%	<19	<19	<b>10%</b>	<19	<b>141</b>	0%	<19	<19	0%	<19	<19	<b>7%</b>	<19	<b>118</b>
Enalapril	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Fluoxetine	19	0%	<19	<19	0%	<19	<19	<b>5%</b>	<19	<b>240</b>	0%	<19	<19	0%	<19	<19	0%	<19	<19
Hydroxyzine	19	0%	<19	<19	<b>33%</b>	<b>53</b>	<b>56</b>	<b>14%</b>	<b>41</b>	<b>64</b>	<b>7%</b>	<19	<b>46</b>	<b>23%</b>	<b>73</b>	<b>121</b>	<b>7%</b>	<19	<b>81</b>
Meprobamate	19	<b>8%</b>	<19	<b>92</b>	<b>11%</b>	<19	<b>68</b>	<b>14%</b>	<b>70</b>	<b>164</b>	0%	<19	<19	<b>8%</b>	<19	<b>155</b>	<b>21%</b>	<b>89</b>	<b>133</b>
Metformin	38	0%	<38	<38	0%	<38	<38	0%	<38	<38	<b>14%</b>	<b>85</b>	<b>214</b>	0%	<38	<38	<b>7%</b>	<19	<b>74</b>
Paracetamol	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Primidone	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Risperidone	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Omeprazole	19	0%	<19	<19	0%	<19	<19	<b>5%</b>	<19	<b>1017</b>	<b>7%</b>	<19	<b>38</b>	<b>8%</b>	<19	<b>95</b>	<b>7%</b>	<19	<b>97</b>
Sulfamethoxazole	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	<b>7%</b>	<19	<b>26</b>	<b>8%</b>	<19	<b>49</b>	0%	<19	<19
Triamterene	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	<b>14%</b>	<19	<b>38</b>
Trimethoprim	19	0%	<19	<19	0%	<19	<19	<b>5%</b>	<19	<b>33</b>	<b>14%</b>	<19	<b>32</b>	0%	<19	<19	0%	<19	<19
Verapamil	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
$\Sigma$ pharmaceuticals	--	<b>15%</b>	<b>73</b>	<b>177</b>	<b>44%</b>	<b>71</b>	<b>131</b>	<b>71%</b>	<b>249</b>	<b>1017</b>	<b>21%</b>	<b>140</b>	<b>292</b>	<b>46%</b>	<b>208</b>	<b>321</b>	<b>36%</b>	<b>209</b>	<b>250</b>
<i>Personal care product</i>																			
Caffeine	38	0%	<38	<38	0%	<38	<38	<b>14%</b>	<b>52</b>	<b>74</b>	<b>14%</b>	<38	<b>56</b>	<b>15%</b>	<38	<b>54</b>	0%	<38	<38
<i>Pesticides</i>																			
Atrazine	19	0%	<19	<19	0%	<19	<19	<b>14%</b>	<b>33</b>	<b>183</b>	0%	<19	<19	<b>15%</b>	<b>43</b>	<b>72</b>	<b>14%</b>	<b>42</b>	<b>82</b>

	Impacted sites (ng/g wet weight)									Undeveloped sites (ng/g wet weight)									
	LOQ	AGR1 (n = 13)			AGR2 (n = 9)			WWEF (n = 21)			LNDF (n = 14)			UND1 (n = 13)			UND2 (n = 14)		
	(ng/g)	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max	DF (%)	90th	Max
Chlorpyrifos	38	0%	<38	<38	0%	<38	<38	<b>48%</b>	<b>357</b>	<b>1557</b>	0%	<38	<38	<b>46%</b>	<b>533</b>	<b>713</b>	0%	<38	<38
Diazinon	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Linuron	19	0%	<19	<19	0%	<19	<19	<b>10%</b>	<19	<b>78</b>	<b>14%</b>	<b>35</b>	<b>54</b>	0%	<19	<19	0%	<19	<19
Simazine	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
<i>Σ pesticides</i>	--	<i>0%</i>	--	--	<i>0%</i>	--	--	<b>52%</b>	<b>364</b>	<b>1740</b>	<b>14%</b>	<b>35</b>	<b>65</b>	<b>54%</b>	<b>533</b>	<b>785</b>	<b>14%</b>	<b>42</b>	<b>82</b>
<i>Natural hormones</i>																			
Androstenedione	19	0%	<19	<19	0%	<19	<19	<b>19%</b>	<b>63</b>	<b>88</b>	0%	<19	<19	0%	<19	<19	0%	<19	<19
Androsterone	4	0%	<4	<4	<b>22%</b>	<b>341</b>	<b>1012</b>	<b>5%</b>	<4	<b>1251</b>	0%	<4	<4	0%	<4	<4	0%	<4	<4
Dihydrotestosterone	61	0%	<61	<61	<b>11%</b>	<61	<b>134</b>	<b>5%</b>	<61	<b>357</b>	0%	<61	<61	0%	<61	<61	<b>7%</b>	<61	<b>82</b>
17 $\alpha$ -Estradiol	4	0%	<4	<4	0%	<4	<4	0%	<4	<4	0%	<4	<4	0%	<4	<4	0%	<4	<4
17 $\beta$ -Estradiol	4	0%	<4	<4	<b>11%</b>	<b>11</b>	<b>53</b>	0%	<4	<4	0%	<4	<4	0%	<4	<4	<b>7%</b>	<4	<b>54</b>
Estriol	12	0%	<12	<12	0%	<12	<12	0%	<12	<12	0%	<12	<12	0%	<12	<12	0%	<12	<12
Estrone	4	<b>15%</b>	<b>27</b>	<b>42</b>	<b>11%</b>	<b>13</b>	<b>63</b>	<b>5%</b>	<4	<b>104</b>	0%	<4	<4	<b>31%</b>	<b>91</b>	<b>105</b>	<b>21%</b>	<b>42</b>	<b>148</b>
Etiocholanolone	23	0%	<23	<23	0%	<23	<23	0%	<23	<23	0%	<23	<23	0%	<23	<23	0%	<23	<23
Testosterone	19	0%	<19	<19	0%	<19	<19	<b>14%</b>	<b>58</b>	<b>119</b>	<b>7%</b>	<19	<b>63</b>	<b>23%</b>	<b>64</b>	<b>85</b>	0%	<19	<19
<i>Σ natural hormones</i>	--	<b>15%</b>	<b>27</b>	<b>42</b>	<b>44%</b>	<b>384</b>	<b>1012</b>	<b>33%</b>	<b>104</b>	<b>1608</b>	<b>7%</b>	--	<b>63</b>	<b>54%</b>	<b>92</b>	<b>105</b>	<b>29%</b>	<b>90</b>	<b>148</b>
<i>Synthetic hormones</i>																			
17 $\alpha$ -Ethinylestradiol	4	0%	<4	<4	<b>11%</b>	<b>7</b>	<b>37</b>	0%	<4	<4	0%	<4	<4	0%	<4	<4	<b>7%</b>	<4	<b>25</b>
Levonorgestrel	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
Mestranol	19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19	0%	<19	<19
<i>Σ synthetic hormones</i>		<i>0%</i>	--	--	<b>11%</b>	<b>7</b>	<b>37</b>	<i>0%</i>	--	--	<i>0%</i>	--	--	<i>0%</i>	--	--	<b>7%</b>	--	<b>25</b>
<i>ΣΣ synthetic TrOCs <sup>(1)</sup></i>	--	<b>15%</b>	<b>73</b>	<b>177</b>	<b>56%</b>	<b>71</b>	<b>131</b>	<b>81%</b>	<b>1123</b>	<b>6761</b>	<b>21%</b>	<b>225</b>	<b>384</b>	<b>77%</b>	<b>1837</b>	<b>2341</b>	<b>43%</b>	<b>237</b>	<b>281</b>

450 DF = detection frequency; 90<sup>th</sup> = 90<sup>th</sup> percentile concentration; Max = maximum concentration; LOQ = Limit of quantification. See Table 2 for

451 sample size. <sup>(1)</sup>  $\Sigma\Sigma$  synthetic TrOCs was calculated as the sum of the concentration of all chemicals except the natural hormones.

452 3.4. Water analysis

453 3.4.1. Chemistry

454 Site WWEF had the most chemically complex water sample with 12% of TrOCs (6/51)  
455 detected (Table 5). Three compounds were detected at sites AGR1 and UND2, one  
456 compound detected at sites LNDF and UND1, while no TrOCs were detected in the sample  
457 from AGR2. Historically, site WWEF has had more chemically complex water (using this  
458 analytical method) compared to the other sites, with 37% of compounds detected in at least  
459 one grab sample over a 12-month monitoring period at that site, compared with 22% for  
460 AGR1, 16% for AGR2, 14% for UND1 and 8% for LNDF (2011-2012, Table 5).

461 All compounds detected at site WWEF were APIs (clozapine, gemfibrozil, paracetamol, and  
462 salicylic acid) or personal care products (caffeine and triclosan), with no known estrogenic  
463 properties. The synthetic hormone EE2 was detected at two sites (LNDF and UND2) in  
464 Apr/May 2013 during fish sampling. The sample from site LNDF had a concentration of 0.11  
465 ng/L, slightly above the predicted no-effect concentration (PNEC) of 0.1 ng/L for 17 $\alpha$ -EE2  
466 proposed by Caldwell et al. (2012), while that from UND2 had a concentration of 0.07 ng/L.  
467 Further monitoring is required to determine the temporal variation and persistence of EE2 at  
468 these sites.

469 **Table 5.** Chemical and *in vitro* monitoring data of 38 trace organic pollutants (TrOCs) from water extracts from the present study (Apr and May  
470 2013) and historical data (May 2011 – Feb 2012) adapted from Scott et al. (2014a, 2014b). All values are in ng/L, except where indicated for  
471 bioanalytical equivalent concentrations.

Compound	AGR1		AGR2		WWEF		LNDF		UND1		UND2
	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	Apr 2013	Apr 2013
<i>Industrial compounds</i>											
Bisphenol A	<b>16 - 82</b>	NA	<b>&lt;10 - 106</b>	NA	<b>15 - 22</b>	NA	<b>12 - 25</b>	NA	<b>12 - 50</b>	NA	NA
TCEP	<b>&lt;10 - 15</b>	<10	<b>&lt;10 - 15</b>	<10	<b>&lt;10 - 11</b>	<10	<10	<10	<b>&lt;10 - 17</b>	<10	<10
4-t-Octylphenol	<10	<20	<10	<20	<10	<20	<10	<20	<10	<20	<20
<i>Pharmaceutical ingredients</i>											
Amtriptyline	<10	<10	<10	<10	<b>&lt;10 - 15</b>	<10	<10	<10	<10	<10	<10
Atenolol	<5	<5	<5	<5	<b>&lt;5 - 9</b>	<5	<5	<5	<5	<5	<5
Carbamazepine	<b>&lt;5 - 7</b>	<5	<5	<5	<b>&lt;5 - 166</b>	<b>41</b>	<5	<5	<5	<5	<5
Clozapine	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Diazepam	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Dilantin	<5	<5	<5	<5	<b>&lt;5 - 22</b>	<5	<5	<5	<5	<5	<5
Enalapril	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
Fluoxetine	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Gemfibrozil	<b>&lt;5 - 5</b>	<5	<5	<5	<b>&lt;5 - 95</b>	<b>11</b>	<5	<5	<5	<5	<5
Hydroxyzine	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
Ibuprofen	<b>&lt;5 - 10</b>	<5	<5	<5	<b>&lt;5 - 44</b>	<5	<5	<5	<5	<5	<5
Ketoprofen	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
Meprobamate	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10
Metformin	NA	<5	NA	<5	NA	<5	NA	<5	NA	<5	<5
Naproxen	<b>&lt;5 - 6</b>	<5	<5	<5	<b>&lt;5 - 15</b>	<5	<5	<5	<5	<5	<5
Omeprazole	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Paracetamol	<b>&lt;5 - 5</b>	<b>128</b>	<b>&lt;5 - 314</b>	<5	<b>7 - 28</b>	<b>460</b>	<5	<5	<b>&lt;5 - 8</b>	<5	<5
Primidone	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Risperidone	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Salicylic acid	<b>&lt;20 - 91</b>	<b>20</b>	<b>&lt;20 - 92</b>	<20	<b>&lt;20 - 88</b>	<b>75</b>	<b>&lt;20 - 97</b>	<20	<b>&lt;20 - 46</b>	<b>29</b>	<b>23</b>
Simvastatin	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5

Compound	AGR1		AGR2		WWEF		LNDF		UND1		UND2
	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	Apr 2013	Apr 2013
Simvastatin-hydroxyacid	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Sulfamethoxazole	<5	<5	<5	<5	<5 - 5	<5	<5	<5	<5	<5	<5
Triamterene	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Trimethoprim	<5	<5	<5	<5	<5 - 25	<5	<5	<5	<5	<5	<5
Verapamil	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
<i>Personal care products</i>											
Caffeine	<10 - 186	15	<10 - 234	<10	20 - 285	333	<10 - 142	<10	<10 - 29	<10	11
Propylparaben	<10 - 34	<10	<10 - 33	<10	<10	<10	<10	<10	<10 - 20	<10	<10
Triclocarban	<10	<10	<10	<10	<10 - 55	<10	<10	<10	<10	<10	<10
Triclosan	<10	<10	<10	<10	<10 - 43	7	<10	<10	<10	<10	<10
<i>Pesticides</i>											
Atrazine	<5	<5	<5 - 8	<5	<5	<5	<5	<5	<5	<5	<5
Chlorpyrifos *	<5	<10	<5	<10	<5	<10	<5	<10	<5	<10	<10
Diazinon *	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Linuron	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
2-Phenylphenol	<10	<10	<10	<10	<10 - 20	<10	<10	<10	<10 - 59	<10	<10
Simazine	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
<i>Hormones</i>											
Androstenedione	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
Androsterone	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Dihydrotestosterone	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16
17 $\alpha$ -Estradiol	<1	<1	<1	<1	<1 - 4	<1	<1	<1	<1	<1	<1
17 $\beta$ -Estradiol	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Estriol	<3	<5	<3	<5	<3	<5	<3	<5	<3	<5	<5
Estrone	<1 - 3	<1	<1 - 2	<1	<1 - 10	<1	<1 - 1	<1	<1 - 2	<1	<1
Etiocholanolone	<6	<6	<6	<6	<6	<6	<6	<6	<6	<6	<6
Mestranol	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Testosterone	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
<i>Synthetic hormones</i>											
17 $\alpha$ -Ethinylestradiol	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.11	<0.05	<0.05	0.07
Levonorgestrel	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5
<i>In vitro endocrine activity</i>											



Compound	AGR1		AGR2		WWEF		LNDF		UND1		UND2
	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	May 2013	2011-2012	Apr 2013	Apr 2013
Estrogenic (EEQ, ng/L)	<0.1 - 0.52	1.18	<0.1	0.78	<0.1 - 1.16	0.89	<0.1 - 0.28	0.1	<0.1	0.04	<0.02
Antiandrogenic (FluEQ, µg/L)	<20	2.44	<20	<1	<20	<1	<20	1.46	<20	<1	2.74
Androgenic (DHTEQ, ng/L)	<7	<9	<7	<9	<7	<9	<7	<9	<7	<9	<9
Antiandrogenic (FluEQ, µg/L)	<60	96	<60	80	<60	90	<60	83	<60	90	73
Progestagenic (LevoEQ, ng/L)	<5	0.09	<5	<0.06	<5	<0.06	<5	0.14	<5	<0.06	0.11
Antiprogestagenic (MifEQ, ng/L)	<8	4.2	<8	<1.8	<8	2.2	<8	3.5	<8	1.8	2.1

472 \* Only measured in one sampling event of four (May 2011 – Feb 2012). NA = Not analyzed; TCEP = Tris(2-chloroethyl) phosphate.

473 Bioanalytical equivalents are: EEQ = 17β-estradiol equivalents; TMXEQ = tamoxifen equivalents; DHTEQ = dihydrotestosterone equivalents;

474 FluEQ = flutamide equivalents; LevoEQ = levonorgestrel equivalents; MifEQ = mifepristone equivalents.

### 475 3.4.2. *In vitro* activity

476 Five of six sites had detectable estrogenic activity, with values ranging from 0.04 to 1.18  
477 ng/L EEQ (Table 5). Site UND2 was the only site without detectable estrogenic activity. *In*  
478 *vitro*-specific safe EEQs (EEQ-SSE) between 0.5 and 2 ng/L EEQ for short-term and  
479 between 0.1-0.4 ng/L EEQ for longer-term exposures have been proposed by Jarošová et al.  
480 (2014). Estrogenic activity in samples from AGR1, AGR2 and WWEF (0.78 to 1.18 ng/L  
481 EEQ) exceeded the most conservative EEQ-SSE of 0.1 ng/L EEQ, while samples from  
482 LNDF, UND1 and UND2 did not (Table 5). The least conservative short-term exposure  
483 EEQ-SSE of 2 ng/L EEQ was not exceeded in any of the samples. While there was a  
484 correlation between *in vitro* estrogenic activity and Vtg protein levels (with higher Vtg levels  
485 at higher EEQ; Fig. 1B), Vtg protein levels in fish at most sites were not significantly  
486 elevated compared to unexposed laboratory reference males, except for fish at AGR1 and  
487 UND2 (Fig. 1A). A recent study suggests that mosquitofish may not be as sensitive as other  
488 native species, such as rainbowfish for example (Scott et al. 2017), and further studies with  
489 more endocrine-sensitive and/or sedentary fish or invertebrate species would help determine  
490 whether these levels of estrogenic activity constitute a risk of endocrine disruption in other  
491 organisms in chronic exposure conditions.

492 Slight anti-estrogenic activity was detected at sites AGR1, LNDF and UND1, with a  
493 maximum concentration of 2.44 µg/L tamoxifen equivalents (TMXEQ). Anti-estrogenic  
494 activity has not previously been reported at these sites in the 2011 to 2012 study; however,  
495 our LOQ was more sensitive in the current analysis (1 µg/L compared to 5 µg/L TMXEQ;  
496 Scott et al. (2014a)). A similar inconstant picture is apparent in the literature, with Leusch et  
497 al. (2014b) reporting anti-estrogenic activity in two of nine WWTP effluents examined (up to  
498 4.4 µg/L TMXEQ), while Roberts et al. (2015) did not detect anti-estrogenic activity in  
499 wastewater of a large Australian WWTP. Quantification of antagonistic activity, while

500 technically possible (Neale and Leusch 2015), is difficult to accurately perform *in vitro* due  
501 to the presence of competing agonist and the possible interference from natural organic  
502 matter (Neale et al. 2015).

503 Androgenic activity was not detected in any samples, but anti-androgenic activity was  
504 detected at all six sites ranging from 73 to 96 µg/L flutamide equivalents (FluEQ). Previous  
505 studies of Australian WWTP effluent (Leusch et al. 2014b, Roberts et al. 2015) have  
506 generally not detected anti-androgenic activity, but this could be due to the high LOQ in  
507 those other studies (*e.g.*, 250 µg/L FluEQ compared to 25 µg/L FluEQ in the present study).

508 Progestagenic activity was detected at three sites, up to a maximum of 0.14 ng/L  
509 levonorgestrel equivalents (LevoEQ) at LNDF (Table 5). Progestagenic activity has  
510 previously been reported in Dutch sewage effluent at a concentration up to 2.2 µg/L LevoEQ  
511 (Van der Linden et al. 2008) and in Australian WWTP effluent up to 5.4 ng/L LevoEQ, and  
512 was hypothesized to be associated with human APIs (Leusch et al. 2014b). Progestagenic  
513 activity was not detected at any river sites in a previous Australian national survey (73 sites;  
514 Scott et al. (2014a)), although the LOQ in that study was much higher than that of the present  
515 study (5 vs. 0.06 ng/L LevoEQ, respectively). The potent synthetic progestin levonorgestrel,  
516 often used in combination with the synthetic estrogen EE2 in birth control pills, is currently  
517 difficult to measure by chemical analysis, with LOQs comparable to the 5 ng/L achieved in  
518 the current study. A recent study has calculated a predicted concentration in wastewater  
519 ranging from 0.2 to 0.6 ng/L (King et al. 2016). If present at these concentrations,  
520 levonorgestrel would likely explain a significant portion of the progestagenic activity  
521 detected here. A provisional PNEC of 0.1 ng/L has been derived for levonorgestrel (King et  
522 al. 2016). The progestagenic activity at sites LNDF and UND2 was slightly above this  
523 concentration (0.14 and 0.11 ng/L LevoEQ, respectively), indicating a potential risk if all the  
524 activity is caused by levonorgestrel. Improvements in chemical analysis methods and

525 refinements of the provisional PNEC value are necessary to more firmly quantify the risk that  
526 this potent progestin poses to the receiving environment. Anti-progestagenic activity was  
527 detected at most sites and ranged from <1.8 to 4.2 µg/L mifepristone equivalents (MifEQ;  
528 Table 5). The maximum concentration in the present study was much lower than that in a  
529 recent Australian study, which reported anti-progestagenic activity in 16% of Australian  
530 rivers sampled (73 in total) at concentrations as high as 32 µg/L MifEQ (Scott et al. 2014a).  
531 The concentrations in the current study were also lower than those measured in Chinese  
532 WWTP effluent (29 µg/L MifEQ measured with a yeast based bioassay) (Li et al. 2011). The  
533 compounds responsible for the anti-progestagenic activity measured in the current study are  
534 unidentified, although nonylphenol, which has been shown to significantly inhibit the binding  
535 of progesterone to the human progesterone receptor in a yeast-based bioassay (Viswanath et  
536 al. 2008), is a potential suspect. Unfortunately, due to analytical complications regarding the  
537 quantification of nonylphenol in the environmental samples (as detailed in Scott et al.  
538 (2014a)), nonylphenol was not analyzed in this study.

539

### 540 3.5. Conclusions

541 This study found no overt evidence of endocrine disruption of sexual axes: there was no  
542 evidence of abnormal secondary sexual characteristic (gonopodium) or gonadal development  
543 (including incidence of intersex) in mosquitofish from any of the sites sampled. *In vitro*  
544 bioassays however indicated slight estrogenic and anti-androgenic activity at most sites, and  
545 Vtg protein (a sensitive biomarker of exposure to estrogenic EACs) was elevated at two sites  
546 (AGR1 and UND2). This suggests that while fish at the sites samples are exposed to low  
547 concentrations of EACs, these concentrations are too low to produce significant organism-  
548 level disruption, in agreement with recent suggestions that endocrine disruption in Australian  
549 freshwaters is unlikely to be widespread (Hassell et al. 2016, Vajda et al. 2015).

550 Several TrOCs were detected in fish carcasses, confirming that fish are exposed to and ingest  
551 a wide range of TrOCs; however only a few TrOCs were detected in grab water, the  
552 discrepancy likely illustrating the high variability of concentrations of these TrOCs over time.  
553 TrOCs were detected at the two undeveloped sites, suggesting that even areas relatively  
554 removed from populated areas may still exhibit the chemical traces of human activity.  
555 Concentrations of TrOCs and EACs in water samples were typically not cause for concern,  
556 with one exception at site LNDF where EE2 was detected slightly above the PNEC of 0.1  
557 ng/L. (Anti)estrogenic, anti-androgenic, and (anti)progestagenic activities were all quantified  
558 in water samples from at least three (and up to five of six) sites. Estrogenicity ranged from  
559 0.1 – 1.18 ng/L EEQ, in excess of the *in vitro*-specific safe estrogenicity (EEQ-SSE) value of  
560 0.1 ng/L for chronic exposure proposed in Jarošová et al. (2014) at all sites except the two  
561 undeveloped sites (Table 5). In a prior study, WWEF had the highest estrogenicity (1.16 ng/L  
562 EEQ) compared to 18 other Queensland sites, and the fifth highest estrogenicity out of 73  
563 sites across mainland Australia (Scott et al. 2014a).

564 While the results of this study indicate a low risk of disruption of sexual endocrine systems in  
565 fish, chemical body burdens were correlated with lipid accumulation in liver (hepatic  
566 steatosis), which may indicate that other effects, including hormonal regulation of lipid  
567 synthesis and storage and other subtle mechanisms of toxicity may be of concern downstream  
568 of wastewater discharges and dense human activity. It should be noted that the low sample  
569 size at some sites, particularly when split across three analyses (body burden, Vtg and  
570 histology), meant limited statistical power to identify subtle differences across sites for some  
571 endpoints, and these results should be treated with caution.

572

### 573 **Acknowledgements**

574 The authors wish to thank P. Corbett, S. Knight, B. Matthews, D. Moffat, M. Mulbjerg, C.

575 Peitsch, J. van de Merwe, K. Watson and R. Woods for support in sampling and laboratory  
576 analysis, and M. Bartkow, J. Blackbeard, S. Blockwell, H. Chapman, H. Nice, and V.  
577 Pettigrove for their input in planning and implementation of this project. This study was  
578 funded by the Australian Research Council (ARC Linkage scheme LP100100163) in  
579 collaboration with Water Research Australia, Sydney Water, Seqwater and Melbourne Water,  
580 and supported in-kind by the Queensland Department of Science, Information Technology  
581 and Innovation (DSITI) and the Western Australia Department of Water. P.S. was supported  
582 with an Australian Postgraduate Award (Industry) scholarship and Water Research Australia  
583 PhD top-up scholarship.

584

## 585 **References**

- 586 Anastassiades, M., Lehotay, S.J., Stajnbaher, D. and Schenck, F.J. (2003) Fast and easy  
587 multiresidue method employing acetonitrile extraction/partitioning and "dispersive solid-  
588 phase extraction" for the determination of pesticide residues in produce. *Journal of AOAC*  
589 *International* 86(2), 412-431.
- 590 Andersen, L. (2004) Imposex: A biological effect of TBT contamination in Port Curtis,  
591 Queensland. *Australasian Journal of Ecotoxicology* 10, 105-113.
- 592 Anderson, K., Burnell, F., Roiko, A., Andrew, M., O'Connor, W. and Elizur, A. (2010)  
593 Development of a method for identifying elevated vitellogenin gene expression in the  
594 Sydney Rock Oyster (*Saccostrea glomerata*) as an indicator of endocrine disruption on the  
595 Sunshine Coast. *Ecological Management & Restoration* 11(2), 143-146.
- 596 Andrew-Priestley, M.N., O'Connor, W.A., Dunstan, R.H., Van Zwieten, L., Tyler, T.,  
597 Kumar, A. and MacFarlane, G.R. (2012) Estrogen mediated effects in the Sydney rock  
598 oyster, *Saccostrea glomerata*, following field exposures to sewage effluent containing  
599 estrogenic compounds and activity. *Aquatic Toxicology* 120, 99-108.

600 Angus, R., McNatt, H., Howell, W. and Peoples, S. (2001) Gonopodium development in  
601 normal male and 11-ketotestosterone-treated female mosquitofish (*Gambusia affinis*): a  
602 quantitative study using computer image analysis. *General and Comparative*  
603 *Endocrinology* 123(2), 222-234.

604 ATSE (2002) Pesticide use in Australia, p. 310pp., Australian Academy of Technological  
605 Sciences and Engineering (ATSE), Parkville, Vic, Australia.

606 Australian Department of Health (2017) Australian Inventory of Chemical Substances  
607 (AICS). Retrieved 6 July 2017 from <https://www.nicnas.gov.au/chemical-inventory-AICS>.

608 Batty, J. and Lim, R. (1999) Morphological and reproductive characteristics of male  
609 mosquitofish (*Gambusia affinis holbrooki*) inhabiting sewage-contaminated waters in New  
610 South Wales, Australia. *Archives of Environmental Contamination and Toxicology* 36(3),  
611 301-307.

612 Booth, D.J. and Skene, C.D. (2006) Rapid assessment of endocrine disruption: Vitellogenin  
613 (Vtg) expression in male estuarine toadfish. *Australasian Journal of Ecotoxicology* 12, 3-  
614 8.

615 Bowman, C.J., Kroll, K.J., Hemmer, M.J., Folmar, L.C. and Denslow, N.D. (2000) Estrogen-  
616 Induced Vitellogenin mRNA and Protein in Sheepshead Minnow (*Cyprinodon variegatus*).  
617 *General and Comparative Endocrinology* 120(3), 300-313.

618 Bradley, P.M., Journey, C.A., Romanok, K.M., Barber, L.B., Buxton, H.T., Foreman, W.T.,  
619 Furlong, E.T., Glassmeyer, S.T., Hladik, M.L., Iwanowicz, L.R., Jones, D.K., Kolpin,  
620 D.W., Kuivila, K.M., Loftin, K.A., Mills, M.A., Meyer, M.T., Orlando, J.L., Reilly, T.J.,  
621 Smalling, K.L. and Villeneuve, D.L. (2017) Expanded Target-Chemical Analysis Reveals  
622 Extensive Mixed-Organic-Contaminant Exposure in U.S. Streams. *Environmental Science*  
623 *& Technology* 51(9), 4792-4802.

624 Caldwell, D.J., Mastrocco, F., Anderson, P.D., Länge, R. and Sumpter, J.P. (2012) Predicted-  
625 no-effect concentrations for the steroid estrogens estrone, 17 $\beta$ -estradiol, estriol, and 17 $\alpha$ -  
626 ethinylestradiol. *Environmental Toxicology and Chemistry* 31(6), 1396-1406.

627 Chinathamby, K., Allinson, M., Shiraishi, F., Lopata, A.L., Nugegoda, D., Pettigrove, V. and  
628 Allinson, G. (2013) Screening for potential effects of endocrine-disrupting chemicals in  
629 peri-urban creeks and rivers in Melbourne, Australia using mosquitofish and recombinant  
630 receptor-reporter gene assays. *Environmental Science and Pollution Research* 20(3),  
631 1831-1841.

632 Doyle, C.J., Barker, J., Lim, R.P. and Gray, L.E. (2003) Reproductive morphology of male  
633 mosquitofish (*Gambusia holbrooki*) inhabiting sewage-contaminated waters in the South  
634 Creek catchment of the Hawkesbury-Nepean river, International Water Association &  
635 Australian Water Association, Sydney, NSW, Australia.

636 Doyle, C.J. and Lim, R.P. (2002) The effect of 17 beta-estradiol on the gonopodial  
637 development and sexual activity of *Gambusia holbrooki*. *Environmental Toxicology and*  
638 *Chemistry* 21(12), 2719-2724.

639 Escher, B.I., Allinson, M., Altenburger, R., Bain, P.A., Balaguer, P., Busch, W., Crago, J.,  
640 Denslow, N.D., Dopp, E., Hilscherova, K., Humpage, A.R., Kumar, A., Grimaldi, M.,  
641 Jayasinghe, B.S., Jarosova, B., Jia, A., Makarov, S., Maruya, K.A., Medvedev, A.,  
642 Mehinto, A.C., Mendez, J.E., Poulsen, A., Prochazka, E., Richard, J., Schifferli, A.,  
643 Schlenk, D., Scholz, S., Shiraishi, F., Snyder, S., Su, G., Tang, J.Y., van der Burg, B., van  
644 der Linden, S.C., Werner, I., Westerheide, S.D., Wong, C.K., Yang, M., Yeung, B.H.,  
645 Zhang, X. and Leusch, F.D. (2014) Benchmarking organic micropollutants in wastewater,  
646 recycled water and drinking water with in vitro bioassays. *Environmental Science &*  
647 *Technology* 48(3), 1940-1956.



648 European Chemicals Agency (2017) EC Inventory. Retrieved 6 July 2017 from  
649 <https://echa.europa.eu/information-on-chemicals/ec-inventory>.

650 Game, C., Gagnon, M.M., Webb, D. and Lim, R. (2006) Endocrine disruption in male  
651 mosquitofish (*Gambusia holbrooki*) inhabiting wetlands in Western Australia.  
652 *Ecotoxicology* 15(8), 665-672.

653 Garcia, S.N., Foster, M., Constantine, L.A. and Huggett, D.B. (2012) Field and laboratory  
654 fish tissue accumulation of the anti-convulsant drug carbamazepine. *Ecotoxicology and*  
655 *Environmental Safety* 84, 207-211.

656 Genten, F., Terwinghe, E. and Danguy, A. (2009) Atlas of fish histology. Science Publishers,  
657 New Hampshire, USA.

658 Hassell, K., Pettigrove, V., Beresford, N., Jobling, S. and Kumar, A. (2016) No evidence of  
659 exposure to environmental estrogens in two feral fish species sampled from the Yarra  
660 River, Australia: A comparison with Northern Hemisphere studies. *Ecotoxicology and*  
661 *Environmental Safety* 131, 104-117.

662 Hou, L., Xie, Y., Ying, G. and Fang, Z. (2011) Developmental and reproductive  
663 characteristics of western mosquitofish (*Gambusia affinis*) exposed to paper mill effluent  
664 in the Dengcun River, Sihui, South China. *Aquat Toxicol* 103(3-4), 140-149.

665 Jarošová, B., Bláha, L., Giesy, J.P. and Hilscherová, K. (2014) What level of estrogenic  
666 activity determined by in vitro assays in municipal waste waters can be considered as  
667 safe? *Environment International* 64, 98-109.

668 Kellar, C.R., Hassell, K.L., Long, S.M., Myers, J.H., Golding, L., Rose, G., Kumar, A.,  
669 Hoffmann, A.A. and Pettigrove, V. (2014) Ecological evidence links adverse biological  
670 effects to pesticide and metal contamination in an urban Australian watershed. *Journal of*  
671 *Applied Ecology* 51(2), 426-439.

672 King, O.C., van de Merwe, J.P., McDonald, J.A. and Leusch, F.D.L. (2016) Concentrations  
673 of levonorgestrel and ethinylestradiol in wastewater effluents: Is the progestin also cause  
674 for concern? *Environmental Toxicology and Chemistry* 35(6), 1378-1385.

675 Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B. and  
676 Buxton, H.T. (2002) Pharmaceuticals, hormones, and other organic wastewater  
677 contaminants in U.S. streams, 1999-2000: a national reconnaissance. *Environmental*  
678 *Science and Technology* 36(6), 1202-1211.

679 Kroon, F.J., Hook, S.E., Jones, D., Metcalfe, S., Henderson, B., Smith, R., Warne, M.S.J.,  
680 Turner, R.D., McKeown, A. and Westcott, D.A. (2015) Altered transcription levels of  
681 endocrine associated genes in two fisheries species collected from the Great Barrier Reef  
682 catchment and lagoon. *Marine Environmental Research* 104, 51-61.

683 Leusch, F.D., Khan, S.J., Gagnon, M.M., Quayle, P., Trinh, T., Coleman, H., Rawson, C.,  
684 Chapman, H.F., Blair, P., Nice, H. and Reitsema, T. (2014a) Assessment of wastewater  
685 and recycled water quality: a comparison of lines of evidence from in vitro, in vivo and  
686 chemical analyses. *Water Research* 50, 420-431.

687 Leusch, F.D., Khan, S.J., Laingam, S., Prochazka, E., Froscio, S., Trinh, T., Chapman, H.F.  
688 and Humpage, A. (2014b) Assessment of the application of bioanalytical tools as  
689 surrogate measure of chemical contaminants in recycled water. *Water Research* 49, 300-  
690 315.

691 Leusch, F.D.L., Chapman, H.F., Kay, G.W., Gooneratne, S.R. and Tremblay, L.A. (2006)  
692 Anal fin morphology and gonadal histopathology in mosquitofish (*Gambusia holbrooki*)  
693 exposed to treated municipal sewage effluent. *Archives of Environmental Contamination*  
694 *and Toxicology* 50, 562-574.

695 Li, J., Chen, M., Wang, Z., Ma, M. and Peng, X. (2011) Analysis of Environmental  
696 Endocrine Disrupting Activities in Wastewater Treatment Plant Effluents Using

697 Recombinant Yeast Assays Incorporated with Exogenous Metabolic Activation System.  
698 *Biomedical and Environmental Sciences* 24(2), 132-139.

699 Loos, R., Gawlik, B.M., Locoro, G., Rimaviciute, E., Contini, S. and Bidoglio, G. (2009) EU-  
700 wide survey of polar organic persistent pollutants in European river waters. *Environmental*  
701 *Pollution* 157(2), 561-568.

702 Lopes, R.P., Reyes, R.C., Romero-Gonzalez, R., Vidal, J.L. and Frenich, A.G. (2012)  
703 Multiresidue determination of veterinary drugs in aquaculture fish samples by ultra high  
704 performance liquid chromatography coupled to tandem mass spectrometry. *Journal of*  
705 *Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 895-896,  
706 39-47.

707 Munaretto, J.S., Ferronato, G., Ribeiro, L.C., Martins, M.L., Adaime, M.B. and Zanella, R.  
708 (2013) Development of a multiresidue method for the determination of endocrine  
709 disrupters in fish fillet using gas chromatography-triple quadrupole tandem mass  
710 spectrometry. *Talanta* 116, 827-834.

711 Neale, P.A., Escher, B.I. and Leusch, F.D.L. (2015) Understanding the implications of  
712 dissolved organic carbon when assessing antagonism in vitro: An example with an  
713 estrogen receptor assay. *Chemosphere* 135, 341-346.

714 Neale, P.A. and Leusch, F.D.L. (2015) Considerations when assessing antagonism in vitro:  
715 Why standardizing the agonist concentration matters. *Chemosphere* 135, 20-23.

716 Nitschke, L. and Schüssler, W. (1998) Surface water pollution by herbicides from effluents of  
717 waste water treatment plants. *Chemosphere* 36(1), 35-41.

718 Norli, H.R., Christiansen, A. and Deribe, E. (2011) Application of QuEChERS method for  
719 extraction of selected persistent organic pollutants in fish tissue and analysis by gas  
720 chromatography mass spectrometry. *Journal of Chromatography A* 1218(41), 7234-7241.

721 Ramirez, A.J., Brain, R.A., Usenko, S., Mottaleb, M.A., O'Donnell, J.G., Stahl, L.L., Wathen,  
722 J.B., Snyder, B.D., Pitt, J.L., Perez-Hurtado, P., Dobbins, L.L., Brooks, B.W. and  
723 Chambliss, C.K. (2009) Occurrence of pharmaceuticals and personal care products in fish:  
724 results of a national pilot study in the United States. *Environmental Toxicology and*  
725 *Chemistry* 28(12), 2587-2597.

726 Rand-Weaver, M., Margiotta-Casaluci, L., Patel, A., Panter, G.H., Owen, S.F. and Sumpter,  
727 J.P. (2013) The read-across hypothesis and environmental risk assessment of  
728 pharmaceuticals. *Environmental Science & Technology* 47(20), 11384-11395.

729 Rawson, C.A., Lim, R.P. and Warne, M.S.J. (2008) Skeletal morphology and maturation of  
730 male *Gambusia holbrooki* exposed to sewage treatment plant effluent. *Ecotoxicology and*  
731 *Environmental Safety* 70(3), 453-461.

732 Rawson, C.A., Tremblay, L.A., Warne, M.S.J., Ying, G.G., Kookana, R., Laginestra, E.,  
733 Chapman, J.C. and Lim, R.P. (2009) Bioactivity of POPs and their effects in mosquitofish  
734 in Sydney Olympic Park, Australia. *Science of the Total Environment* 407(12), 3721-3730.

735 Rawson, C.A., Webb, D. and Gagnon, M.M. (2010) Investigations of endocrine disruption  
736 using *Gambusia* species - A review. *Australasian Journal of Ecotoxicology* 16, 53-87.

737 Reitsem, T.J. and Spickett, J.T. (1999) Imposex in *Morula Granulata* as bioindicator of  
738 tributyltin (TBT) contamination in the Dampier Archipelago, Western Australia. *Marine*  
739 *Pollution Bulletin* 39(1), 280-284.

740 Roberts, J., Bain, P.A., Kumar, A., Hepplewhite, C., Ellis, D.J., Christy, A.G. and Beavis,  
741 S.G. (2015) Tracking multiple modes of endocrine activity in Australia's largest inland  
742 sewage treatment plant and effluent- receiving environment using a panel of in vitro  
743 bioassays. *Environmental Toxicology and Chemistry* 34(10), 2271-2281.

744 Schäfer, R.B., Pettigrove, V., Rose, G., Allinson, G., Wightwick, A., von der Ohe, P.C.,  
745 Shimeta, J., Kühne, R. and Kefford, B.J. (2011) Effects of pesticides monitored with three

746 sampling methods in 24 sites on macroinvertebrates and microorganisms. *Environmental*  
747 *Science & Technology* 45(4), 1665-1672.

748 Scott, P.D., Bartkow, M., Blockwell, S.J., Coleman, H.M., Khan, S.J., Lim, R., McDonald,  
749 J.A., Nice, H., Nugegoda, D., Pettigrove, V., Tremblay, L.A., Warne, M.S. and Leusch,  
750 F.D. (2014a) An assessment of endocrine activity in Australian rivers using chemical and  
751 in vitro analyses. *Environmental Science and Pollution Research International* 21(22),  
752 12951-12967.

753 Scott, P.D., Bartkow, M., Blockwell, S.J., Coleman, H.M., Khan, S.J., Lim, R., McDonald,  
754 J.A., Nice, H., Nugegoda, D., Pettigrove, V., Tremblay, L.A., Warne, M.S. and Leusch,  
755 F.D. (2014b) A national survey of trace organic contaminants in Australian rivers. *Journal*  
756 *of Environmental Quality* 43(5), 1702-1712.

757 Scott, P.D., Coleman, H.M., Colville, A., Lim, R., Matthews, B., McDonald, J.A., Miranda,  
758 A., Neale, P.A., Nugegoda, D., Tremblay, L.A. and Leusch, F.D.L. (2017) Assessing the  
759 potential for trace organic contaminants commonly found in Australian rivers to induce  
760 vitellogenin in the native rainbowfish (*Melanotaenia fluviatilis*) and the introduced  
761 mosquitofish (*Gambusia holbrooki*). *Aquatic Toxicology* 185, 105-120.

762 Togunde, O.P., Oakes, K.D., Servos, M.R. and Pawliszyn, J. (2012) Optimization of solid  
763 phase microextraction for non-lethal in vivo determination of selected pharmaceuticals in  
764 fish muscle using liquid chromatography-mass spectrometry. *Journal of Chromatography*  
765 *A* 1261, 99-106.

766 Tousova, Z., Oswald, P., Slobodnik, J., Blaha, L., Muz, M., Hu, M., Brack, W., Krauss, M.,  
767 Di Paolo, C., Tarcai, Z., Seiler, T.-B., Hollert, H., Koprivica, S., Ahel, M., Schollée, J.E.,  
768 Hollender, J., Suter, M.J.F., Hidasi, A.O., Schirmer, K., Sonavane, M., Ait-Aissa, S.,  
769 Creusot, N., Brion, F., Froment, J., Almeida, A.C., Thomas, K., Tollefsen, K.E., Tufi, S.,  
770 Ouyang, X., Leonards, P., Lamoree, M., Torrens, V.O., Kolkman, A., Schriks, M.,

771 Spirhanzlova, P., Tindall, A. and Schulze, T. (2017) European demonstration program on  
772 the effect-based and chemical identification and monitoring of organic pollutants in  
773 European surface waters. *Science of the Total Environment* 601–602, 1849-1868.

774 Trinh, T., Harden, N.B., Coleman, H.M. and Khan, S.J. (2011) Simultaneous determination  
775 of estrogenic and androgenic hormones in water by isotope dilution gas chromatography–  
776 tandem mass spectrometry. *Journal of Chromatography A* 1218(12), 1668-1676.

777 U.S. Environmental Protection Agency (2017) Toxic Substance Control Act (TSCA)  
778 Chemical Substance Inventory. Retrieved 6 July 2017 from [https://www.epa.gov/tsca-](https://www.epa.gov/tsca-inventory)  
779 [inventory](https://www.epa.gov/tsca-inventory).

780 Vajda, A.M., Kumar, A., Woods, M., Williams, M., Doan, H., Tolsher, P., Kookana, R.S. and  
781 Barber, L.B. (2015) Integrated assessment of wastewater treatment plant effluent  
782 estrogenicity in the Upper Murray River, Australia, using the native Murray rainbowfish  
783 (*Melanotaenia fluviatilis*). *Environmental Toxicology and Chemistry* 34(5), 1078-1087.

784 Van der Linden, S.C., Heringa, M.B., Man, H.Y., Sonneveld, E., Puijker, L.M., Brouwer, A.  
785 and Van der Burg, B. (2008) Detection of multiple hormonal activities in wastewater  
786 effluents and surface water, using a panel of steroid receptor CALUX bioassays.  
787 *Environmental Science & Technology* 42(15), 5814-5820.

788 Vanderford, B.J. and Snyder, S.A. (2006) Analysis of pharmaceuticals in water by isotope  
789 dilution liquid chromatography/tandem mass spectrometry. *Environmental Science &*  
790 *Technology* 40(23), 7312-7320.

791 Viswanath, G., Halder, S., Divya, G., Majumder, C.B. and Roy, P. (2008) Detection of  
792 potential (anti)progestagenic endocrine disruptors using a recombinant human  
793 progesterone receptor binding and transactivation assay. *Molecular and Cellular*  
794 *Endocrinology* 295(1-2), 1-9.

795 Wilkinson, J.M., Hayes, S., Thompson, D., Whitney, P. and Bi, K. (2008) Compound  
796 profiling using a panel of steroid hormone receptor cell-based assays. *Journal of*  
797 *Biomolecular Screening* 13(8), 755-765.

798 Woods, M. and Kumar, A. (2011) Vitellogenin induction by 17 $\beta$ -estradiol and 17 $\alpha$ -  
799 ethynylestradiol in male Murray rainbowfish (*Melanotaenia fluviatilis*). *Environmental*  
800 *Toxicology and Chemistry* 30(11), 2620-2627.

801 Yoshikane, M., Kay, W.R., Shibata, Y., Inoue, M., Yanai, T., Kamata, R., Edmonds, J.S. and  
802 Morita, M. (2006) Very high concentrations of DDE and toxaphene residues in crocodiles  
803 from the Ord River, Western Australia: an investigation into possible endocrine disruption.  
804 *Journal of Environmental Monitoring* 8(6), 649-661.

805