

On the critical exponent α of the 5D random-field Ising model

Fytas, N., Martin-Mayor, V., Parisi, G., Picco, M. & Surlas, N.

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Fytas, N, Martin-Mayor, V, Parisi, G, Picco, M & Surlas, N 2019, 'On the critical exponent α of the 5D random-field Ising model' Journal of Statistical Mechanics: Theory and Experiment, vol. 2019, no. 9, 093203.

<https://dx.doi.org/10.1088/1742-5468/ab3987>

DOI 10.1088/1742-5468/ab3987

ESSN 742-5468

Publisher: IOP Publishing:

This is an author-created, un-copyedited version of an article accepted for publication/published in Journal of Statistical Mechanics: Theory and Experiment IOP Publishing Ltd is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at <https://dx.doi.org/10.1088/1742-5468/ab3987>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

On the critical exponent α of the 5D random-field Ising model

Nikolaos G. Fytas¹, Víctor Martín-Mayor^{2,3}, Giorgio Parisi⁴,
Marco Picco⁵, and Nicolas Sourlas⁶

¹Applied Mathematics Research Centre, Coventry University, Coventry CV1 5FB, United Kingdom

²Departamento de Física Teórica I, Universidad Complutense, 28040 Madrid, Spain

³Instituto de Biocomputación y Física de Sistemas Complejos (BIFI), 50009 Zaragoza, Spain

⁴Dipartimento di Fisica, Sapienza Università di Roma, P.le Aldo Moro 2, 00185 Rome, Italy and INFN, Sezione di Roma I, NANONTEC – CNR, P.le A. Moro 2, 00185 Rome, Italy

⁵Laboratoire de Physique Théorique et Hautes Energies, UMR7589, Sorbonne Université et CNRS, 4 Place Jussieu, 75252 Paris Cedex 05, France

⁶Laboratoire de Physique Théorique de l'École Normale Supérieure (Unité Mixte de Recherche du CNRS et de l'École Normale Supérieure, associée à l'Université Pierre et Marie Curie, PARIS VI) 24 rue Lhomond, 75231 Paris CEDEX 05, France

Abstract. We present a complementary estimation of the critical exponent α of the specific heat of the 5D random-field Ising model from zero-temperature numerical simulations. Our result $\alpha = 0.12(2)$ is consistent with the estimation coming from the modified hyperscaling relation and provides additional evidence in favor of the recently proposed restoration of dimensional reduction in the random-field Ising model at $D = 5$.

PACS numbers: 705.50.+q, 75.10.Hk, 64.60.Cn, 75.10.Nr

Submitted to: *Journal of Statistical Mechanics*

The random-field Ising model (RFIM) is one of the archetypal disordered systems [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15], extensively studied due to its theoretical interest, as well as its close connection with experiments in condensed-matter physics [15, 16, 17, 18, 19]. Its beauty stems from the combination of random fields and the standard Ising model that creates rich and complicated physical phenomena, responsible for a great volume of research over the last 40 years and more. It is well established that the physically relevant dimensions of the RFIM lay between $2 < D < 6$, where $D_l = 2$ and $D_u = 6$ are the lower and upper critical dimensions of the model, respectively. For $D \geq D_u$ one expects the standard mean-field behavior [1, 8, 9, 10, 20, 21], whereas exactly at $D = D_u$ the notoriously obscuring logarithmic corrections appear [22].

In the last few years, the development of a powerful panoply of simulation and statistical analysis methods [23] have set the basis for a fresh revision of the problem. In fact, some of the main controversies have been resolved, the most notable being the illustration of critical universality in terms of different random-field distributions [24, 25, 26] and the restoration of supersymmetry and dimensional reduction at $D = 5$ [27, 28, 29] (see also references [30, 31, 32, 33] for additional evidence in this respect).

In particular, the large-scale numerical simulations of the 5D RFIM reported in reference [27] have provided high-accuracy estimates for the spectrum of critical exponents and for several universal ratios (see Table III in reference [27]), with one missing element: that of the direct computation of the critical exponent α of the specific heat. Let us point out that the specific heat of the RFIM is of experimental interest [18] and that the value of α has severe implications for the validity of the fundamental scaling relations, and in particular for the Rushbrooke relation, $\alpha + 2\beta + \gamma = 2$, that has been the most controversial of all [34, 35, 36, 37, 38]. Therefore a strong command on this aspect of the model's critical behavior is necessary. In the current work we fill this gap by performing additional simulations and scaling analysis that allow us to directly compute α for the 5D RFIM and to therefore present a complete picture of the scaling behavior of the specific heat. Our final estimate, $\alpha = 0.12(2)$, agrees well with that of the 3D Ising universality class, 0.110087(12) [39], and therefore constitutes additional evidence in favor of our recently proposed restoration of dimensional reduction at $D = 5$ [27, 28, 29].

The RFIM Hamiltonian is

$$\mathcal{H} = -J \sum_{\langle xy \rangle} S_x S_y - \sum_x h_x S_x, \quad (1)$$

with the spins $S_x = \pm 1$ occupying the nodes of a hyper-cubic lattice in space dimension D with nearest-neighbor ferromagnetic interactions J and h_x independent random magnetic fields with zero mean and dispersion σ . Here we consider the Hamiltonian (1) on a $D = 5$ hyper-cubic lattice with periodic boundary conditions and energy units $J = 1$. Our random fields h_x follow either a Gaussian (\mathcal{P}_G), or a Poissonian (\mathcal{P}_P)




Figure 1. Connected correlation length in units of the system size L versus σ for the 5D Gaussian RFIM and a system of linear size $L = 10$. Four distinct simulation sets are shown, corresponding to different simulation values, $\sigma^{(s)}$, and different sets of random-field realizations. The inset illustrates the reweighting error-evolution for the fourth simulation set with $\sigma = 6.02395$ and $N_{\text{samples}} = 10^6$.

distribution of the form

$$\mathcal{P}_G(h, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{h^2}{2\sigma^2}}; \quad \mathcal{P}_P(h, \sigma) = \frac{1}{2|\sigma|} e^{-\frac{|h|}{\sigma}}, \quad (2)$$

where $-\infty < h < \infty$ and σ the disorder-strength control parameter.

As it is well-established, in order to describe the critical behavior of the model one needs two correlation functions, namely the connected and disconnected propagators, $C_{xy}^{(\text{con})}$ and $C_{xy}^{(\text{dis})}$:

$$C_{xy}^{(\text{con})} \equiv \frac{\partial \overline{\langle S_x \rangle}}{\partial h_y}; \quad C_{xy}^{(\text{dis})} \equiv \overline{\langle S_x \rangle \langle S_y \rangle}, \quad (3)$$

where the $\langle \dots \rangle$ are thermal mean values as computed for a given realization, a sample, of the random fields $\{h_x\}$. Over-line refers to the average over the samples. Following the prescription of reference [23], for each of these two propagators we scrutinize the second-moment correlation lengths, denoted as $\xi^{(\text{con})}$ and $\xi^{(\text{dis})}$, respectively.

Our numerical simulations for the 5D RFIM are described in reference [27]. We therefore outline here the very necessary details. We simulated lattice sizes from $L_{\min} = 4$ to $L_{\max} = 28$. For each pair of (L, σ) values we generated ground states for 10^7 samples – for the additional simulations at the most accurate determinations of the critical points shown below in figures 4 and 5, 10^6 samples were generated – exceeding previous relevant studies [22] by a factor of 10^3 on average. The calculation of the ground states of the RFIM was based on the well-established mapping [34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55] to




Figure 2. Connected correlation length in units of the system size L versus σ for the 5D Gaussian (main panel) and Poissonian (inset) RFIM. An illustrative example of the three lattice-size sequence $(L, 2L, 4L) = (5, 10, 20)$ used in the application of the modified quotients method is shown [see equations (5) and (6)]. Data taken from reference [27].

the maximum-flow problem [56, 57, 58]. We used our own C version of the push-relabel algorithm of Tarjan and Goldberg [59], involving some technical modifications proposed by Middleton and collaborators for further efficiency [46, 47]. Suitable generalized fluctuation-dissipation formulas and reweighting extrapolations have facilitated our analysis, as exemplified in reference [23]. A comparative illustration in favor of the numerical accuracy of our scheme is shown in figure 1 for the universal ratio $\xi^{(\text{con})}/L$ of an $L = 10$ Gaussian RFIM and four different simulation sets, as outlined in the panel.

The specific heat of the RFIM can be estimated using ground-state calculations in two complementary frameworks, both based on the analysis of singularities of the bond-energy density E_J [60]. This bond-energy density is the first derivative $\partial E/\partial J$ of the ground-state energy with respect to the random-field strength σ [34, 35]. The derivative of the sample averaged quantity \overline{E}_J with respect to σ then gives the second derivative with respect to σ of the total energy and thus the sample-averaged specific heat C . The singularities in C can also be studied by computing the singular part of \overline{E}_J , as \overline{E}_J is just the integral of C with respect to σ . Thus, one may estimate α by studying the behavior of \overline{E}_J at $\sigma = \sigma_c$ [34], via the scaling form

$$\overline{E}_J(L, \sigma_c) = E_{J, \infty} + bL^{(\alpha-1)/\nu}(1 + b'L^{-\omega}), \quad (4)$$

where $E_{J, \infty}$, b , and b' are non-universal constants, and ω is the universal corrections-to-scaling exponent.

Of course, the use of equation (4) for the application of standard finite-size scaling methods requires an *a priori* knowledge of the *exact* value of the critical random-field




Figure 3. Effective exponent ratio $(\alpha - 1)/\nu$ versus $L^{-\omega}$ for all random-field distributions and crossing points considered in this work. Note the notation $Z^{(x)}$, where Z stands for the distribution – G for Gaussian and P for Poissonian – and the superscript x for the connected (con) or disconnected (dis) type of the universal ratio $\xi^{(x)}/L$, used for the application of the quotients method [see equations (5) and (6)].

strength σ_c [see also the analysis below in figures 4 and 5]. Although we currently have at hand such high-accuracy estimates of the critical fields for both types of the random-field distributions under study [27], we start our analysis with an alternative to this approach. In particular, we implement a three lattice-size variant of the original quotients method [61], also known as phenomenological renormalization [62, 63, 64] that has been described in detail in reference [26] and already successfully applied to the $D = 3$ [23] and $D = 4$ [25] models. The main idea in this perspective, given that $\alpha - 1 < 0$, is the elimination of the non-divergent background term $E_{J,\infty}$ in equation (4) by considering three lattice sizes in the following sequence: $(L_1, L_2, L_3) = (L, 2L, 4L)$ [see figure 2 for an instructive illustration of the three-lattice variant of the quotients method based on the crossings of $\xi^{(con)}/L$]. Taking the quotient of the differences at the crossings of the pairs $(L, 2L)$ and $(2L, 4L)$

$$\hat{Q}_O = \frac{(\overline{E}_{J,4L} - \overline{E}_{J,2L})|_{(\xi_{4L}/\xi_{2L})=2}}{(\overline{E}_{J,2L} - \overline{E}_{J,L})|_{(\xi_{2L}/\xi_L)=2}}, \quad (5)$$

one obtains the following scaling formula for the bond-energy density [26]

$$\hat{Q}_{\overline{E}_J}^{(cross)} = 2^{(\alpha-1)/\nu} + \mathcal{O}(L^{-\omega}). \quad (6)$$

Our results for the effective exponent ratio $(\alpha - 1)/\nu$ as a function of $L^{-\omega}$ – where ω is set to the 3D Ising value 0.82966 [39] – are shown in figure 3. The dashed line marks the estimate $(\alpha - 1)/\nu = -1.412\ 625\ 34\dots$ of the 3D Ising universality class, where we have used the values $\alpha = 0.110087(12)$ and $\nu = 0.629971(4)$ [39]. A few comments are in




Figure 4. Finite-size scaling behavior of the bond-energy density at the critical random-field strength $\sigma_c(G)$ of the 5D Gaussian RFIM. The lines are fittings of the form (4) with different ω values, as indicated in the panel.

order: (i) Clearly, there exist large corrections to scaling for the sequence of smaller sizes (2, 4, 8) and (3, 6, 12) that obscure the application of any finite-size scaling approach. (ii) The remaining data points [(4, 8, 16), (5, 10, 20), and (6, 12, 24)] do not allow for a safe extrapolation of the ratio $(\alpha - 1)/\nu$ to $L \rightarrow \infty$, although the general trend of the data appears to be on the right track and, in fact, joint polynomial fits with a shared constant term do approach the value $-1.45(6)$ but with a rather bad fitting quality. (iii) Larger system sizes would be needed to clarify this point, but are unfortunately out of reach with our current resources.

Guided by these qualitative results of the phenomenological-renormalization approach, we have performed, at a second stage, additional simulations at the critical points $\sigma_c(G) = 6.02395$ and $\sigma_c(P) = 5.59038$ of the Gaussian and Poissonian models, respectively [27]. In figures 4 and 5 we report on the finite-size scaling behavior of the bond-energy density at these critical points for the whole spectrum of system sizes studied, alongside with the resulting estimates for the ratio $(\alpha - 1)/\nu$. In both panels the solid lines are fits of the form (4), where the different colors correspond to different fixed values of ω . Black curves correspond to the value 0.82966 of the 3D Ising universality class [39], whereas red curves to the value 0.66 estimated in reference [27]. The fitting quality, measured in terms of χ^2/dof , where dof measures the number of degrees of freedom, and the minimum system size, L_{\min} , used in the fits are as follows: $\chi^2/\text{dof} = 1.8/3$, $L_{\min} = 8$ for the Gaussian model (figure 4) and $\chi^2/\text{dof} = 6.1/4$, $L_{\min} = 6$ for the Poissonian model (figure 5). Note that there was practically no variation in the fitting quality moving from $\omega = 0.82966$ down to 0.66 [65]. Using now the estimate $\nu = 0.629971(4)$ for the critical exponent of the correlation length,




Figure 5. Finite-size scaling behavior of the bond-energy density at the critical random-field strength $\sigma_c(P)$ of the 5D Poissonian RFIM. The lines are fittings of the form (4) with different ω values, as indicated in the panel.

simple algebra and error propagation produces values for α within the range $0.10 - 0.13$. Taking an average over the values of α obtained from the black curves with $\omega = 0.82966$, we give our final estimate for the critical exponent α to be

$$\alpha = 0.12(2). \quad (7)$$

This is compatible to the value $0.12(5)$ obtained in reference [27] via the modified hyperscaling relation $\alpha = 2 - \nu(D - 2 + \bar{\eta} - \eta)$, where η and $\bar{\eta}$ are the corresponding anomalous dimensions of the connected and disconnected correlation functions [see equation (3)] and also agrees nicely with the 3D Ising universality benchmark $\alpha = 0.110087(12)$ [39].

As an additional consistency check of our results shown in figures 4 and 5, we depict in figure 6 the scaling behavior of the specific heat C , obtained from the derivative of the bond-energy density with respect to the random-field strength σ , at the critical point. Note that the horizontal axis has been rescaled to $L^{\alpha/\nu}$ (remember that as in the standard case $C \sim L^{\alpha/\nu}$), and α/ν has been set to the value $0.174749\dots$ via $\alpha = 0.110087$ and $\nu = 0.629971$ of the 3D Ising universality class [39]. As expected the data become rather noisy with increasing system size, forcing us to exclude from our fittings the larger system sizes $L = 20$ and $L = 24$, where statistical errors are larger than 30%. Although we illustrate for the benefit of the reader data for the complete spectrum of system sizes studied, the solid lines are simple linear fits within the range $L = 4 - 16$ with a very good fitting quality indeed: $\chi^2/\text{dof} = 4.16/6$ and $2.03/6$ for the Gaussian and Poissonian models, respectively.

To summarize, using extensive numerical simulations at zero temperature we




Figure 6. Scaling behavior of the specific heat C for both models considered in this work, as indicated in the panel. For a detailed discussion on the scaling laws and the fitting tests refer to the main text.

provided a high-precision estimate of the specific-heat’s critical exponent of the 5D RFIM. Our final result $\alpha = 0.12(2)$ is fully consistent with the estimation coming from the modified hyperscaling relation given in reference [27], and also supports the recent results of reference [29] for the restoration of supersymmetry and dimensional reduction in the RFIM at $D = 5$. We close this contribution with figure 7 and an overview of the critical exponent α of the RFIM at all physically relevant dimensions. Two sets of data points are shown, as outlined in the caption, corroborated by a graphical validation of the Rushbrooke relation in the corresponding inset. Whilst the collative results of figure 7 are reassuring and settle down previous controversies in the random-field problem originating from defective estimations of the critical exponent α , for reasons of clarity we would like to point out that the large error at $D = 3$ stems from the joint fits of $[(\alpha - 1)/\nu]^{(\text{eff})}$ performed over several random-field distributions (including the double Gaussian distribution) and the large scaling corrections via $\omega(D = 3) = 0.52$ [23, 24] – for further details and graphical explanations on this aspect we refer the interested reader to figures 6 and 7 of reference [23].

Acknowledgments

We acknowledge partial financial support from Ministerio de Economía, Industria y Competitividad (MINECO, Spain) through Grant No. FIS2015-65078-C2, and from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (Grant No. 694925).




Figure 7. Critical exponent α of the specific heat of the RFIM as a function of the spatial dimension D . Two sets of data points are shown: Estimates from direct computation (open squares) and from the modified hyperscaling relation (filled circles) via previously obtained results for the exponents η , $\bar{\eta}$, and ν . The dashed line marks the 3D Ising value 0.110087(12) [39]. The filled star signals the mean-field (MF) value $\alpha = 0$, expected to hold at $D \geq 6$. **Inset:** Verification of the Rushbrooke scaling relation. For the estimation of the magnetic critical exponents β and γ we have used the standard relations $\beta = \nu(D - 4 + \bar{\eta})/2$ and $\gamma = \nu(2 - \eta)$. The dashed line is located exactly at the value 2. Data taken from this work and from references [23, 24, 25, 26, 27].

References

- [1] Imry Y and Ma S-K 1975 *Phys. Rev. Lett.* **35** 1399
- [2] Aharony A, Imry Y, and Ma S-K 1976 *Phys. Rev. Lett.* **37** 1364
- [3] Young A P 1977 *J. Phys. Condens. Matter* **10** L257
- [4] Fishman S and Aharony A 1979 *J. Phys. C: Solid State Phys.* **12** L729
- [5] Parisi G and Sourlas N 1979 *Phys. Rev. Lett.* **43** 744
- [6] Cardy J L 1984 *Phys. Rev. B* **29** 505
- [7] Imbrie J Z 1984 *Phys. Rev. Lett.* **53** 1747
- [8] Villain J 1984 *Phys. Rev. Lett.* **52** 1543; Villain J 1985 *J. Physique* **46** 1843
- [9] Bray A J and Moore M A 1985 *J. Phys. C: Solid State Phys.* **18** L927
- [10] Fisher D S 1986 *Phys. Rev. Lett.* **56** 416
- [11] Schwartz M and Soffer A 1985 *Phys. Rev. Lett.* **55** 2499 ; Schwartz M and Soffer A 1986 *Phys. Rev. B* **33** 2059 ; Schwartz M 1985 *J. Phys. Condens. Matter* **18** 135 ; Schwartz M, Gofman M, and Nattermann T 1991 *Physica A* **178** 6 ; Schwartz M 1994 *Europhys. Lett.* **15** 777
- [12] Gofman M, Adler J, Aharony A, Harris A B, and Schwartz M 1993 *Phys. Rev. Lett.* **71** 1569
- [13] Esser J and Nowak U 1997 *Phys. Rev. B* **55** 5866
- [14] Barber W C and Belanger D P 2001 *J. Magn. Magn. Mater.* **226** 545
- [15] See, e.g., the articles by Belanger D P and Nattermann T 1998 *Spin Glasses and Random Fields* ed A P Young (Singapore: World Scientific)
- [16] Belanger D P and Young A P 1991 *J. Magn. Magn. Mater.* **100** 272

- [17] Rieger H 1995 *Annual Reviews of Computational Physics II* ed D Stauffer (Singapore: World Scientific) pp 295-341
- [18] Belanger D P, King A R, Jaccarino V, and Cardy J L 1983 *Phys. Rev. B* **28** 2522 ; Belanger D P and Slanič Z 1998 *J. Magn. Magn. Mater.* **186** 65
- [19] Vink R L C, Binder K, and Löwen H 2006 *Phys. Rev. Lett.* **97** 230603
- [20] Berker A N and McKay S R 1986 *Phys. Rev. B* **33**, 4712
- [21] Bricmont J and Kupiainen A 1987 *Phys. Rev. Lett.* **59** 1829
- [22] Ahrens B and Hartmann A K 2011 *Phys. Rev. B* **83** 014205
- [23] Fytas N G and Martín-Mayor V 2016 *Phys. Rev. E* **93**, 063308
- [24] Fytas N G and Martín-Mayor V 2013 *Phys. Rev. Lett.* **110** 227201
- [25] Fytas N G, Martín-Mayor V, Picco M, and Sourlas N 2016 *Phys. Rev. Lett.* **116** 227201
- [26] Fytas N G, Martín-Mayor V, Picco M, and Sourlas N *J. Stat. Mech.* (2017) 033302
- [27] Fytas N G, Martín-Mayor V, Picco M, and Sourlas N 2017 *Phys. Rev. E* **95** 042117
- [28] Fytas N G, Martín-Mayor V, Picco M, and Sourlas N 2018 *J. Stat. Phys.* **172** 665
- [29] Fytas N G, Martín-Mayor V, Parisi G, Picco M, and Sourlas N 2019 *Phys. Rev. Lett.* **122** 240603
- [30] Tissier M and Tarjus G 2011 *Phys. Rev. Lett.* **107** 041601
- [31] Tissier M and Tarjus G 2012 *Phys. Rev. B* **85** 104203
- [32] Tarjus G, Balog I, and Tissier M 2013 *Europhys. Lett.* **103** 61001
- [33] Hikami S, arXiv:1801.09052
- [34] Middleton A A and Fisher D S 2002 *Phys. Rev. B* **65** 134411
- [35] Hartmann A K and Young A P 2001 *Phys. Rev. B* **64** 214419
- [36] Theodorakis P E, Georgiou I, and Fytas N G 2013 *Phys. Rev. E* **87** 032119
- [37] Fytas N G, Theodorakis P E, Georgiou I, and Lelidis I 2013 *Eur. Phys. J. B* **86** 268
- [38] Nowak U, Usadel K D, and Esser J 1998 *Physica A* **250**, 1
- [39] Kos F, Poland D, Simmons-Duffin D, and Vichi A 2016 *J. High Energy Phys.* **08** 036
- [40] Ogielski A T 1986 *Phys. Rev. Lett.* **57** 1251
- [41] Hartmann A K and Usadel K D 1995 *Physica A* **214** 141; Hartmann A K 1998 *Physica A* **248** 1
- [42] Bastea S and Duxbury P M 1998 *Phys. Rev. E* **58** 4261 ; Bastea S 1998 *Phys. Rev. E* **58** 7978 ; Bastea S and Duxbury P M 1999 *Phys. Rev. E* **60** 4941
- [43] Hartmann A K and Nowak U 1999 *Eur. Phys. J. B* **7** 105
- [44] Hartmann A K 2002 *Phys. Rev. B* **65** 174427
- [45] Seppälä E T and Alava M J 2001 *Phys. Rev. E* **63** 066109 ; Seppälä E T, Alava M J, and Duxbury P M 2001 *Phys. Rev. E* **63** 066110 ; Seppälä E T, Pulkkinen A M, and Alava M J 2002 *Phys. Rev. B* **66** 144403
- [46] Middleton A A 2002 *Phys. Rev. Lett.* **88** 017202
- [47] Middleton A A arXiv:cond-mat/0208182 ; Meinke J H and Middleton A A arXiv:cond-mat/0502471 ; Hambrick D C, Meinke J H, and Middleton A A arXiv:cond-mat/0501269
- [48] Dukovski I and Machta J 2003 *Phys. Rev. B* **67** 014413
- [49] Alava M J, Duxbury P M, Moukarzel C F, and Rieger H 2001 *Phase Transitions and Critical Phenomena*, vol 18, ed C Domb and J L Lebowitz (San Diego, CA: Academic)
- [50] Zumsande M, Alava M J, and Hartmann A K 2008 *J. Stat. Mech.: Theory Exp.* P02012
- [51] Shrivastav G P, Krishnamoorthy S, Banerjee V, and Puri S 2011 *Europhys. Lett.* **96** 36003
- [52] Stevenson J D and Weigel M 2011 *Europhys. Lett.* **95** 40001
- [53] Fytas N G, Theodorakis P E, and Georgiou I 2012 *Eur. Phys. J. B* **85** 349
- [54] Hartmann A K and Rieger H 2004 *Optimization Algorithms in Physics* (Berlin: Wiley)
- [55] Hartmann A K and Weigt M 2005 *Phase Transitions in Combinatorial Optimization Problems* (Berlin: Wiley)
- [56] Anglès d'Auriac J-C, Preissmann M, and Rammal R 1985 *J. Phys. Lett.* **46** L173
- [57] Cormen T H, Leiserson C E, and Rivest R L 1990 *Introduction To Algorithms*, (Cambridge: MA MIT Press)
- [58] Papadimitriou C H 1994 *Computational Complexity* (Reading: Addison-Wesley)

- [59] Goldberg A V and Tarjan R E 1988 *J. Assoc. Comput. Mach.* **35** 921 ; Cherkassky B V and Goldberg A V 1997 *Algorithmica* **19** 390
- [60] Holm C and Janke W 1997 *Phys. Rev. Lett.* **78** 2265
- [61] The general approach in the quotients method is to compare observables computed in pair of lattices $(L, 2L)$. We start imposing scale-invariance by seeking the L -dependent critical point: the value of σ such that $\xi_{2L}/\xi_L = 2$, i.e., the crossing point for ξ_L/L [see also figure 2 for the case where $\xi = \xi^{(\text{con})}$]. For dimensionful quantities O , scaling in the thermodynamic limit as $\xi^{x_O/\nu}$, we consider the quotient $Q_O = O_{2L}/O_L$ at the crossing. Thus, we have $Q_O^{(\text{cross})} = 2^{x_O/\nu} + \mathcal{O}(L^{-\omega})$, where x_O/ν , and the scaling-corrections exponent ω are universal
- [62] Ballesteros H G, Fernández L A, Martín-Mayor V, and Muñoz Sudupe A 1996 *Phys. Lett. B* **378** 207
- [63] Amit D J and Martín-Mayor V 2005 *Field Theory, the Renormalization Group and Critical Phenomena*, 3rd edn (Singapore: World Scientific)
- [64] Nightingale M 1976 *Physica A* **83** 561
- [65] See also the relevant statistical tests with respect to the value of ω for the other thermodynamic observables in reference [27]