

# A systematic review on workplace interventions to manage chronic musculoskeletal disorders

Skamagki, G., King, A., Duncan, M. & Wahlin, C.

Author post-print (accepted) deposited by Coventry University's Repository

## Original citation & hyperlink:

Skamagki, G, King, A, Duncan, M & Wahlin, C 2018, 'A systematic review on workplace interventions to manage chronic musculoskeletal disorders' *Physiotherapy Research International*, vol. 23, no. 4, e1738.

<https://dx.doi.org/10.1002/pri.1738>

DOI 10.1002/pri.1738

ISSN 1358-2267

ESSN 1471-2865

Publisher: Wiley

**This is the peer reviewed version of the following article: Skamagki, G, King, A, Duncan, M & Wahlin, C 2018, 'A systematic review on workplace interventions to manage chronic musculoskeletal disorders' *Physiotherapy Research International*, vol 23, no. 4, (e1738), which has been published in final form at**

<https://dx.doi.org/10.1002/pri.1738>

**This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.**

**Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.**

**This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.**

1 A systematic review on workplace interventions to manage chronic musculoskeletal  
2 disorders

3 **ABSTRACT**

4 Background and Purpose: A review to investigate whether there are effective workplace  
5 interventions that manage chronic musculoskeletal disorders.

6 Methods: The literature search included published articles between 2008 and  
7 2017. The databases used in this search were MEDLINE, Scopus, CINAHL, AMED,  
8 PsycINFO, Academic Search Complete, Cochrane, and PEDro. A limited search on  
9 websites for relevant grey literature was also conducted.

10 Results: The review included 12 studies that investigated effectiveness of a specific  
11 strength exercise programme or interventions provided by health professionals at the  
12 workplace when compared with controls or interventions not at the workplace. Seven  
13 studies were classified as high quality (>85% of criteria met) and five studies were  
14 classified as acceptable. Studies were heterogeneous preventing a meta-analysis. No  
15 intervention was clearly superior to another.

16 Discussion: There was some consistency in the results of the selected studies, suggesting  
17 that workplace interventions such as high-intensity strength exercises and/or  
18 integrated health care can decrease pain and symptoms for employees who experience  
19 long-term musculoskeletal disorders. However, the current research is limited.

20 Key Words: Chronic Musculoskeletal Disorders, Management, Systematic Review,  
21 Workplace

22

## 23 **INTRODUCTION**

24 A healthy work environment influences the physical, mental and socioeconomic  
25 behaviours of its employees (Waddell & Burton, 2006) and can promote the well-being of  
26 their families and communities. It can also increase productivity, and reduce absenteeism or  
27 presenteeism (the practice of coming to work with an injury or medical condition) (Johns,  
28 2009; Tehrani, Humpage, Willmott, & Haslam, 2007). The focus of this review is the  
29 workplace, as the place for providing management and treatment for employees who have  
30 long-term musculoskeletal disorders.

31 Musculoskeletal disorders (MSDs) cover a heterogeneous range of health conditions  
32 such as low back pain and upper or lower limb injuries, which have a big impact on  
33 productivity (Buckley, 2015; Walker-Bone & Linaker, 2016). Long-term musculoskeletal  
34 disorders have an even greater impact on people's lives as they are a source of long-term  
35 pain and increase the number of lost working days (Arthritis Research UK, 2014; Arthritis  
36 Research UK, 2017; McGee, Bevan, & Quadrello, 2011). 'Long-term musculoskeletal  
37 disorders' are those that do not resolve and have a long-term or progressive course  
38 (Goodwin & Naylor, 2010). 'Chronic' is defined in this paper as conditions that have lasted  
39 for over three months. The World Health Organization (WHO) has highlighted that long-  
40 term and chronic conditions require continuous management over many years or decades  
41 (World Health Organization, 2002). The morbidity cost is notable as stretched health care

42 services around the world face further financial pressures due to increasing numbers of  
43 people affected by chronic MSDs (MacKenzie and de Melo-Martin 2015). In addition, the  
44 aging workforce in Europe will mean increasing numbers of these people in the workforce,  
45 with implications for health care.

46 Worldwide, a variety of models and recommendations have been suggested to shift  
47 the need for healthcare and sick leave from the healthcare system to the employer  
48 (McGillivray, 2005; NICE, 2015; Wynne-Jones, Mallen, Mottram, Main, & Dunn, 2009). Some  
49 of these models have been tried without success: for example in the UK, workplace  
50 capability assessments were unsuccessful (Safety and Health Practitioner, 2016). But lack of  
51 effectiveness may have been due to employer and employee ignorance of their roles in  
52 managing those chronic conditions.

53 The WHO (WHO, 2016) has identified three main categories of health interventions  
54 that can be used to manage the risk of MSDs at the workplace. These categories relate to  
55 prevention, return to work, and long-term management, and can include specific services,  
56 actions or products developed and implemented to change or improve health, behaviors  
57 and awareness. A variety of Cochrane systematic reviews have summarised scientific  
58 evidence about the effectiveness of workplace interventions for the first two categories,  
59 prevention and return to work (Aas, Tuntland, Ka, Røe, & Labriola, 2009, 2011; Mulimani et  
60 al., 2014; Parry, Coenen, O'Sullivan, Maher, & Straker, 2017; Rla, Cumpston, Peeters, & Sa,  
61 2013; Shrestha, Ijaz, Kt, Kumar, & Cp, 2015). This study focuses on the third category  
62 (Proper et al., 2003) which includes management at the workplace of individuals with  
63 existing conditions. The aim of this systematic review was to identify the workplace

64 management strategies for individuals with existing long-term musculoskeletal disorders  
65 and to highlight whether these interventions are effective.

## 66 **MAIN TEXT**

### 67 **Methods**

#### 68 **Search strategy**

69 This review used methods from traditional systematic review approaches (Cochrane  
70 Handbook for Systematic Reviews of Interventions) for the literature search phase, and then  
71 assessed, analysed and synthesised the relevant data (Higgins & Green, 2011). The PICO  
72 approach was used to structure the research question (Table 1) and identify the inclusion  
73 and exclusion criteria (Stern, Jordan, & McArthur, 2014).

74 The literature search included articles that were published between 2008 and 2017.  
75 The strategy searched MEDLINE, CINAHL, AMED, Cochrane, PsycINFO, Academic Search  
76 Complete and PEDro (Appendix 1). A limited search for Grey literature examined relevant  
77 websites including the Institute for Work and Health, the Return to Work Knowledge, the  
78 Institution of Occupational Safety and Health, and the European Agency for Safety and  
79 Health at Work. Search strategies used Boolean operators (AND/OR/NOT), Subject  
80 Headings, alternative spellings, acronyms, and wild cards. In addition, Scopus was used to  
81 perform post-publication citation searching on identified articles.

#### 82 **Selection of studies**

83

##### 84 *Eligibility criteria*

85

##### 86 **Inclusion criteria**

87           The primary criterion was the testing of effectiveness of workplace interventions to  
88 manage employees with long-term multi-joint conditions and chronic musculoskeletal  
89 disorders (12 weeks or more). Participants' age was between 18-68 years (common working  
90 age range) and both males and females were included. Interventions included strategies or  
91 specific activities that were conducted individually or in groups to manage chronic MSDs.  
92 The period searched was from 2008 to the present, since scoping searches indicated that  
93 earlier studies were of a very low quality (Aas et al., 2009, 2011; Hoe, Urquhart, Kelsall, &  
94 Sim, 2012) and focused on prevention and return to work rather than management.

#### 95 **Exclusion criteria**

96           Workplace interventions focusing purely on prevention and return-to-work  
97 strategies were not included in this review. This review excluded studies including people  
98 with acute MSDs or other serious pathologies (Blangsted, Søgaard, Hansen, Hannerz, &  
99 Sjøgaard, 2008), and those which did not aim to compare the effectiveness of the  
100 interventions used in the workplace arena. In addition, guidelines, policies and other  
101 recommendations were also excluded. The inclusion and exclusion criteria used in this  
102 review are summarised in Table 2.

#### 103 **Outcome**

104           The review's outcomes of interest are symptom modification, pain severity,  
105 presenteeism, and sickness absence at individual, worksite and service level, reflecting the  
106 ICF focus on function and disability (WHO, 2001). Some outcomes can be only measured  
107 subjectively (e.g. pain or presenteeism), so it is important to analyse other outcomes like  
108 sickness absence that can be observed objectively.

109 **Data collection**

110           The titles and abstracts of all identified studies were collected and duplicates were  
111 removed before study selection. Data from the relevant studies were extracted  
112 independently by two reviewers; characteristics of studies were collected including study  
113 design, country where intervention was implemented, participant details, type of  
114 intervention, outcome measures and results.

115 **Risk of bias assessment**

116           Many critical appraisal systems and tools are available and can be used to  
117 assess the rigour of the design, the strength of the resulting evidence and the  
118 implementation of the identified studies. However, disagreement between researchers is  
119 common, since differences in intention, components, construction and psychometric  
120 properties of published critical appraisal tools for research reports have been identified  
121 (Katrak, Bialocerkowski, Massy-Westropp, Kumar, & Grimmer, 2004). Since there is no “gold  
122 standard” critical appraisal tool (Katrak et al., 2004), a systematic and transparent approach  
123 was used to assess both internal and external validity of the studies, identify their relevance  
124 to practice, prevent errors, and facilitate judgments (Figure 1). A recent review of the  
125 grading systems produced by medical specialties (Baker, Young, Potter, & Madan, 2010),  
126 highlighted that the Scottish Intercollegiate Guidelines Network (SIGN) can be selected and  
127 used for RCTs as it is an established and validated tool. The SIGN tool (checklist and an  
128 explanation sheet) was selected for this review. The overall assessment of the strength of  
129 the evidence within each paper was based on grading criteria of “(+) acceptable”, “(++) high  
130 quality”, “(-) low quality” or “(0) un-acceptable/reject”.

131

132           **RESULTS**

133   **Selection of studies**

134           Studies selected were published between 2008 and 2017. One of the advantages of  
135 reviewing studies conducted after 2008 was the higher quality of the RCTs identified. The  
136 search identified 257 references, 21 references in AMED, 108 in Academic Search Complete,  
137 36 in MEDLINE, 29 in CINAHL, 18 references in PsycINFO, 10 in COCHRANE, 17 references in  
138 Scopus, and 18 references in PEDro. After removing duplicates, 159 references remained  
139 (Figure 1). The titles and abstracts were reviewed and, when needed, the full-text articles  
140 were read. The full text of 29 articles was obtained but only nine were included in the  
141 review, as none of the others met the inclusion and exclusion criteria. Hand-searching the  
142 reference lists identified nine more studies that were also assessed; however only three of  
143 them were included in the final review. In summary, 12 articles were included in the review  
144 and consensus on the final results was achieved by a second researcher (AK) who reviewed  
145 and replicated the search strategy identifying the same results.

146   **Study characteristics**

147

148           Of the 12 selected studies one study was conducted in the USA, 8 in Denmark, one in  
149 Finland and two in the Netherlands. All studies followed a randomised or a cluster  
150 randomised controlled trial design, and ethical approval was granted from local ethics  
151 committees. There were no differences within studies in the baseline characteristics of  
152 groups of participants (except in Zebis et al. 2011). Detailed inclusion and exclusion criteria



153 were outlined to ensure patient safety and homogeneity. The characteristics of studies for  
154 this review are presented in Table 3.

### 155 **Quality appraisal**

156 The quality of evidence for each outcome was assessed using the SIGN tool for the  
157 appraisal of RCTs. Seven studies were classified as (++) high quality (>85% of criteria met)  
158 and 5 studies were classified as (+) acceptable. Overall, the studies were of a very good  
159 quality (table 4), minimising the risk of bias for the 'true' effect of the interventions.  
160 Randomisation was achieved with either preratification, labelled paper and selection from  
161 an opaque plastic or with random computer-generated numbers. Participants were  
162 randomly allocated into clusters with the use of a computer-generated random numbers  
163 table and only one study used a coin toss (Zebis et al., 2011). All the authors conducted a  
164 power analysis identifying the appropriate sample size that would detect a 15% or a 10%  
165 change for the selected outcome. However, in one study the drop-out rates reached almost  
166 40% leading to limited interpretation of findings (Hutting et al., 2015). The primary outcome  
167 measures were clearly stated in the studies. Patient outcomes were analysed per the group  
168 to which they were originally allocated, but in one study (Jay et al., 2011) analysis was based  
169 solely on participants who completed the trial. Lastly, statistical analysis was clearly  
170 explained, and appropriate values were given in most of the studies in both texts and tables.  
171 Some of the studies only provided results on histograms making it difficult to identify the  
172 true values (Blangsted et al., 2008; Lambeek et al., 2010). Other the studies identified more  
173 outcomes such as job satisfaction rates, psychological well-being, which are not included in  
174 this review. The quality appraisal of the studies is presented in Table 4.

### 175 **Outcome measures**

176 The outcome measures identified and reviewed for this study were pain and function  
177 (Numeric Pain Rating Scales, Revised Arthritis Impact Measurement Scales),  
178 absenteeism/sick leave days (Work Ability Index Score, DASH work module), Health status  
179 (DASH general module) and presenteeism (Stanford Presenteeism Scale). These are reliable,  
180 validated and responsive instruments that can be used in an occupational health care  
181 setting (Meenan, Mason, Anderson, Guccione, & Kazis, 1992; Roy et al., 2011; Tuomi,  
182 Ilmarinen, Jahkola, Katajarinne, & Tulkki, 1998; Williamson & Hoggart, 2005).

183 The interventions and the outcomes of the studies are presented in Table 5. Some  
184 studies were explicitly interested in the workplace venue: for example, Jakobsen (2015)  
185 compared strength training at the workplace with physical exercise in the home. Other  
186 studies were looking at the workplace primarily as the venue for a form of intervention such  
187 as strength training to be compared with another intervention: for example, Andersen's  
188 group (2012) looked at three different exercise regimes all provided at the workplace, but  
189 also included a no physical training control group. In this study, between groups  
190 comparisons with the control group would have been useful, but these were not available.  
191 Because of the nature of the study design it would be difficult to draw any conclusions  
192 about the benefits of the workplace as a venue over any other venues for interventions.

### 193 **Effectiveness of the interventions**

#### 194 *Effect of different physical exercise interventions at the workplace*

195

196 Two studies, (L. L. Andersen et al., 2008, 2010) investigated the effect of different  
197 physical exercise interventions on musculoskeletal pain in all regions of the body and their  
198 association with specifically the neck and the shoulder. As an example, in one of these  
199 studies (L. L. Andersen et al., 2010), 549 office workers were allocated to 3 separate groups;

200 a specific resistance training group (dumbbell exercises of front raise, lateral raise, reverse  
201 flies, shrugs and wrist extension), an all-around exercise group and a reference intervention  
202 group. The results demonstrated that pain for the strength training group decreased with a  
203 statistically significant difference for neck pain ( $p < 0.01-0.05$ ). The authors of these studies  
204 conducted another randomised controlled trial (Andersen et al., 2012) to measure the  
205 effects of strength training in three different regimes (the first group trained for 1 hour per  
206 week, the second group trained 20 minutes three times a week and the 3rd group trained 7  
207 minutes nine times a week). The results demonstrated reduction ( $p < 0.005$ ) of neck and  
208 shoulder pain in office workers for the weekly one-hour program.

209 A study by the same team (Zebis et al., 2011) evaluated the effect of a strength  
210 training intervention at the workplace on non-specific neck and shoulder pain among  
211 industrial workers, highlighting a reduction of pain in the intervention group. However,  
212 despite randomisation, baseline differences between groups were found for pain intensity  
213 which may have affected the outcome of this study. Another study investigated a different  
214 strength exercise training program for the management of chronic musculoskeletal pain at  
215 the workplace (Jay et al., 2011) and showed that progressive kettlebell training 3 times per  
216 week can reduce the pain intensity of neck and shoulder ( $p < 0.02$ ) and the pain intensity of  
217 the lower back ( $p < 0.05$ ). In addition, more studies from Denmark (Blangsted et al., 2008)  
218 demonstrated the reduction in intensity ( $p < 0.0318$ ) and duration of the pain ( $p < 0.0565$ ) of a  
219 resistance training group and an all-around physical exercise group compared to a reference  
220 group (general health-promoting activities not including physical activity). However, no  
221 significant changes were identified between the different active interventions (e.g. Nordic  
222 walking and running, step count).

223

224 *Effect of physical exercise interventions at the workplace compared to other interventions*

225

226 Jakobsen et al., (2015) investigated the effectiveness of a workplace versus a home-based  
227 exercise program for chronic musculoskeletal neck and back pain conditions. The 200  
228 participants were allocated into two groups and were encouraged to perform a  
229 strengthening exercise program (TheraBand, kettlebells) at the workplace for 10 weeks  
230 whereas the control group performed physical exercises at their houses following  
231 instructions and recommendations from illustrated posters. Although results showed a  
232 significant decrease in pain for both groups ( $p < 0.0001$ ), the workplace chronic MSD group  
233 experienced higher reduction of pain compared with the control group ( $p = 0.003$ ). Baldwin  
234 et al., (2012) compared the use of a self-management manual at home with the use of the  
235 same self-management manual at the workplace in combination with an individual  
236 ergonomic intervention. Employees with rheumatoid arthritis (RA) and osteoarthritis (OA)  
237 followed an intervention that consisted of workstation equipment modifications, person-  
238 specific exercises, postural control or lifestyle changes given by an occupational therapist  
239 trained in ergonomics. The results demonstrated only a within-group statistically significant  
240 improvement in physical functioning and pain for the workplace treatment group after a 12  
241 month ( $p < 0.04$ ) and 24 months ( $p < 0.01$ ). The results however could have been affected  
242 by the heterogeneity in pain intensity and the varying severity of RA and OA at the  
243 beginning of the study.

244

245 *Effect of usual care /ergonomics at the workplace compared to other interventions*

246

247           Hutting et al., (2015) compared a self-management program with a usual care group  
248 at the workplace and identified significant differences in work-status ( $p=0.04$ ) measured by  
249 the Disabilities of the Arm, Shoulder and Hand (DASH) questionnaire. However, no  
250 significant differences emerged from the study for pain intensity and functional status.

251 Sundstrup et al., (2014), compared a strength training program to an ergonomic training and  
252 education program among slaughterhouse workers with chronic musculoskeletal pain.

253 Similarly, no significant differences in pain and function were identified between or within  
254 the two groups during the 10 weeks of testing. Interestingly, the overall score of the Work  
255 Ability Index in the ergonomic group got worse after the intervention ( $p = 0.012$ ) but the  
256 authors have challenged this conclusion as the ergonomic program was based on worksite  
257 analysis and a health and safety systems developed by managers rather than health  
258 professionals with specific knowledge and training in occupational health. In a different  
259 study a physiotherapist assessed the effect of an ergonomic intervention on pain and  
260 sickness absence caused by upper-extremity musculoskeletal disorders (Shiri et al., 2011).

261 There was a decrease in pain intensity ( $p<0.05$ ) in the first two weeks but no significant  
262 differences at the end of a yearly follow up. Unfortunately, this study experienced a lot of  
263 drop-outs and loss of participants at follow-up which could have affected the results. The  
264 use of specific health professionals in this study is echoed by Lambeek et al., (2010) that  
265 assessed the effectiveness of integrated care with usual care at the workplace for  
266 employees with chronic low back pain. All the workplace interventions were provided by  
267 health care professionals, such as a clinical occupational physician, a manual therapist, an  
268 occupational therapist and a physiotherapist. Although pain and functional status improved  
269 in both two groups, the integrated care group demonstrated statistically significant  
270 improvement ( $p<0.001$ ) regarding the functional status.

271 **DISCUSSION**

272           The current review gathered and synthesised updated evidence from the scientific  
273 literature to identify the workplace management strategies for individuals with existing  
274 chronic musculoskeletal disorders, and investigated their effectiveness. Studies included in  
275 this review were assessed for bias and were also rated for their quality. Twelve studies were  
276 categorised with high or acceptable quality and they were selected for the final review. The  
277 RCTs included were highly heterogeneous: they varied in the type of interventions, type of  
278 jobs and outcome measures. The conclusion of this systematic review is that the use of  
279 physical activity and/or the integrated health care at the workplace can decrease pain and  
280 symptoms for employees who experience chronic musculoskeletal disorders. Findings of  
281 these studies highlighted that the type of the exercise program used, the way of delivery  
282 and the regime may affect the outcome. An example providing supervised exercise and  
283 supplementary manuals for self-management, telephone calls for reinforcement and face-  
284 to-face instructions with other supplements showed a positive influence on levels of pain,  
285 function, motivation and lifestyle changes. The use of a specific strength exercise program  
286 appeared to have better effects on pain and functional activity in comparison to other types  
287 of exercises, but all the exercise programs at the workplace showed within-group  
288 improvements.

289           A few systematic reviews (Aas et al., 2011; Hoe et al., 2012; Mischke et al.,  
290 2013; Mulimani et al., 2014) have assessed the effects of workplace ergonomic training  
291 interventions or exercise interventions, but focus only on the prevention of MSK conditions.  
292 Similarly, peer-reviewed literature (Hoe et al., 2012; Menta et al., 2015; Nastasia, Coutu, &  
293 Tcaciuc, 2014) regarding workplace prevention of upper limb musculoskeletal disorders

294 described a variety of interventions of which only a few showed effectiveness (e.g.  
295 resistance training, stretching or forearm support). These results were inconclusive due to  
296 the inclusion of low quality RCTs, poor internal validity and lack of generalisability to the  
297 wider population.

298 Levels of evidence for specific ergonomic interventions emerged also from another  
299 systematic review (Leyshon, 2010) for office workers with musculoskeletal disorders.

300 There was also poor evidence to suggest that self-management programmes are  
301 effective in improving pain and managing MSDs at the workplace, while in some studies the  
302 improvement rate dropped after a year (Blangsted et al., 2008; Hutting et al., 2015; Jay et  
303 al., 2011). On the other hand, the review found positive changes in pain perception and  
304 intensity in response to strength training. However, other type of interventions that could  
305 affect pain were not identified in the literature. As an example, cognitive behavioural  
306 therapy has not been evaluated in a lot of RCTs and results from some moderate quality  
307 studies do not show effectiveness when CBT is applied alone (Basler, Bertalanffy, Quint,  
308 Wilke, & Wolf, 2007; Jørgensen, Faber, Hansen, Holtermann, & Sjøgaard, 2011).  
309 Nevertheless, the present review identified a number of studies that recorded  
310 improvements in pain levels and functional status following a structured and well-delivered  
311 exercise programme at the workplace among employees with chronic musculoskeletal pain.

312 Another important finding from this review was the significant improvement in  
313 functional status and the decrease in pain with the use of a workplace integrated care  
314 program by an allied health professional (e.g. physiotherapist, occupational therapist with  
315 ergonomic training). Our review concluded that the use of private medical insurance with  
316 direct access or other health care services at the workplace (e.g. physiotherapy services) can

317 have a positive effect in managing long-term MSDs (e.g. Lambeek et al., 2010, Shiri et al.,  
318 2011) but further research is necessary to investigate the success of those programs in the  
319 health care environment of different countries. In some countries like the Netherlands the  
320 implementation of a workplace program would not be difficult as the costs of workplace  
321 interventions are covered by the patient's health insurance. In other countries  
322 implementation could be more problematic without financial support by the government or  
323 employers.

324           Additionally, healthcare professionals, like physiotherapists, are able to provide a  
325 well-structured exercise program as part of their role. It is well recognised that a  
326 physiotherapist could be suitably equipped to manage chronic conditions and help  
327 employees to remain healthy at work (Johnston & Shaw, 2013). But, there is as yet no  
328 evidence to show the effectiveness of physiotherapy at the workplace. The grey literature  
329 has identified some one-off successes in individual workplaces, but it is unknown if all  
330 branches of the same company follow the same protocol, if there are long-term results of  
331 the interventions or if these workplaces are still providing the service.

332           Four studies in the review used self-management strategies either as the primary  
333 intervention (Hutting et al., 2015) or as a control group (L. L. Andersen et al., 2008, 2010;  
334 Baldwin et al., 2012). Self-management programmes can include leaflets and manuals, e-  
335 learning modules to prepare people to manage their health conditions or change their  
336 lifestyle. There were no significant differences in any of the selected outcome between the  
337 groups but a small improvement was found within the self-management group. Although  
338 self-management strategies are cost effective (Haas et al., 2005), there is still poor evidence



339 on the effectiveness of these programs for people with chronic MSDs (Nolte & Osborne,  
340 2013).

341 Sick leave was measured in some of the studies included (Baldwin et al., 2012; Shiri  
342 et al., 2011; Sundstrup et al., 2016) but there were no significant differences after the  
343 completion or at follow-up. One possible explanation would be that the intensity or  
344 frequency of the interventions did not meet the level that would result in a positive effect  
345 on reducing sick leave. Another explanation could be that the population size was not big  
346 enough for a change or the fact that pain level in these studies was also very different in the  
347 beginning of each experiment. One study has shown that workers with higher aerobic  
348 capacity had a higher Work Ability Index (WAI) score ( $p < 0.004$ ) and thereby a decreased risk  
349 of having a sick leave episode (Strijk et al., 2011). However, this was an observational study  
350 based on the fact that high levels of aerobic capacity are associated with a reduced  
351 incidence of chronic diseases and therefore might be associated with reduced sick leave  
352 (Kellett, Kellett, & Nordholm, 1991; Macedo, Oakley, Panayi, & Kirkham, 2009). On the other  
353 hand, one study (Sundstrup et al., 2014) found an important deterioration of the  
354 employees' Work Ability Index score results following ergonomic interventions at the  
355 workplace implemented by employers/managers and not by health professionals. Their  
356 results question the role of employers and line managers in this process. Similarly,  
357 presenteeism was measured (Hutting et al., 2015) only in one study without showing  
358 important improvements in the decrease of this phenomenon.

359 Recent research has focused on the effectiveness of interventions in community and  
360 workplace settings to reduce sick leave and job loss among workers with musculoskeletal  
361 disorders (Palmer et al., 2012). The current study has separated the workplace interventions

362 found at individual, worksite and service level from workplace ergonomic interventions  
363 and/or psychosocial risk assessments, control of the workplace risks, ergonomic changes to  
364 the work environment and advice offered by employers. The results of this systematic  
365 review agree with the conclusions of previous systematic reviews (Maher, 2000; Palmer et  
366 al., 2015; Rw, Tuntland, Ka, Røe, & Labriola, 2010) and suggest that a physical activity  
367 program and/or integrated care at the workplace can be effective in the management of  
368 chronic MSK disorders. In addition, the studies in this review showed also clinically and  
369 significantly important differences in favour of some secondary outcomes for the workplace  
370 groups such as well-being, job satisfaction, desire to exercise, energy for family and friends,  
371 motivation to eat better and socializing more with their colleagues.

## 372 **STUDY STRENGTHS AND LIMITATIONS**

373

374 A rigorous systematic search of the literature from 2008 to March 2017 was used to  
375 examine study design, biases, outcome measures and methods of analysis. Strengths of this  
376 review comprise the inclusion of high quality RCTs that investigated workplace interventions  
377 for the management of chronic musculoskeletal disorders. Also, the review excluded studies  
378 before 2008 as previous systematic reviews showed that RCTs from the past decade cannot  
379 be used as supportive evidence due to low quality and poor external validity for their results  
380 to be generalised to the wider population. The likelihood of publication bias was not  
381 assessed but several relevant peer-reviewed studies that reported no effects for important  
382 outcomes were also included in this review. The association of pain with other factors (e.g.  
383 environmental, social, personal, psychological) could have influenced the results of some  
384 studies about the change of the pain levels. Lastly, a meta-analysis was not performed  
385 because the studies demonstrated such heterogeneity: some characteristics like pain

386 intensity, pain duration, occupation or education at the entry level (Baldwin et al., 2012;  
387 Jakobsen et al., 2015; Jay et al., 2014) were so lacking in comparability that such an analysis  
388 would have been meaningless.

389

## 390 **CONCLUSION**

391

392 There was some consistency in the results of the selected studies, suggesting that high  
393 intensity strength exercises and/or integrated health care at the workplace may decrease  
394 pain and symptoms for employees who experience chronic musculoskeletal disorders.  
395 Exercise interventions reported in this review included specific muscle strengthening,  
396 kettlebell training, stretching, and all-round- exercises. Clearly, there are other types of  
397 exercises, such as stabilization exercises, proprioceptive re-education and coordination (e.g.  
398 Tai-Chi, yoga), which might be beneficial for chronic musculoskeletal pain but their  
399 effectiveness at the workplace has not been evaluated. In addition, none of the studies  
400 included psychologically-informed therapy/interventions (e.g. Cognitive behaviour therapy,  
401 motivational interviewing etc.) although the link between mental health, stress, anxiety and  
402 MSDs is now recognised (Magnavita, Elovainio, de Nardis, Heponiemi, & Bergamaschi,  
403 2011). None of the studies in this review identified significant results for sick leave,  
404 presenteeism rates and the use of a self-management programme alone, showing again the  
405 consistency of the findings. There is need for more research since the included studies  
406 showed variety in methodology, intervention, and population, and were conducted in a  
407 variety of countries with different health systems (it is not clear if all employees have access  
408 to the same systems of support at the workplace). This can limit the generalisability of the

409 results to countries like the UK where health care is usually provided outside the workplace.  
410 Lastly, further research needs to consider the study design carefully due to the complexity  
411 of the work environment and the biopsychosocial framework for health. The results of this  
412 literature review suggest the implementation of a multi-component workplace intervention  
413 for the management of long-term MSDs. However, it is crucial to look at this complex topic  
414 with an all-inclusive approach considering the differences within the workforce as this will  
415 benefit both the stakeholders and the providers.

416

417

418

419

420

421

422

423

424

425

426

427

428

430 **REFERENCES**

431 Aas, R., Tuntland, H., Ka, H., Røe, C., & Labriola, M. (2009). Workplace interventions for low-  
432 back pain in workers (Protocol). *Cochrane Database of Systematic Reviews*, (4), 1–14.

433 <https://doi.org/10.1002/14651858.CD008160.pub2>. [www.cochranelibrary.com](http://www.cochranelibrary.com)

434 Aas, R., Tuntland, H., Ka, H., Røe, C., & Labriola, M. (2011). Workplace interventions for neck  
435 pain in workers. *Cochrane Database of Systematic Reviews*, (4), 1–91.

436 <https://doi.org/10.1002/14651858.CD008160.pub>

437 Andersen, C. H., Andersen, L. L., Gram, B., Pedersen, M. T., Mortensen, O. S., Zebis, M. K., &  
438 Sjøgaard, G. (2012). Influence of frequency and duration of strength training for

439 effective management of neck and shoulder pain: a randomised controlled trial. *British*  
440 *Journal of Sports Medicine*, 46(14), 1004–1010. <https://doi.org/10.1136/bjsports-2011->

441 090813

442 Andersen, L. L., Christensen, K. B., Holtermann, A., Poulsen, O. M., Sjøgaard, G., Pedersen,  
443 M. T., & Hansen, E. A. (2010). Effect of physical exercise interventions on

444 musculoskeletal pain in all body regions among office workers: a one-year randomized  
445 controlled trial. *Manual Therapy*, 15(1), 100–104.

446 Andersen, L. L., Kjaer, M., Sogaard, K., Hansen, L., Kryger, A. I., & Sjøgaard, G. (2008). Effect  
447 of two contrasting types of physical exercise on chronic neck muscle pain. *Arthritis and*

448 *Rheumatism*, 59(1), 84–91. <https://doi.org/10.1002/art.23256>

449 Arthritis Research UK. (2014). *Care planning and musculoskeletal health*.

450 Arthritis Research UK. (2017). *State of Musculoskeletal Health 2017*. Retrieved from  
451 file:///Users/mariellabaxter/Downloads/State-of-musculoskeletal-health-2017.PDF

452 Baker, A., Young, K., Potter, J., & Madan, I. (2010). A review of grading systems for evidence-  
453 based guidelines produced by medical specialties. *Clinical Medicine, Journal of the*  
454 *Royal College of Physicians of London*, 10(4), 358–363.  
455 <https://doi.org/10.7861/clinmedicine.10-4-358>

456 Baldwin, D., Johnstone, B., Ge, B., Hewett, J., Smith, M., & Sharp, G. (2012). Randomized  
457 prospective study of a work place ergonomic intervention for individuals with  
458 rheumatoid arthritis and osteoarthritis. *Arthritis Care & Research*, 64(10), 1527–1535.  
459 <https://doi.org/10.1002/acr.21699>

460 Blangsted, A. K., Søgaard, K., Hansen, E. A., Hannerz, H., & Sjøgaard, G. (2008). One-year  
461 randomized controlled trial with different physical-activity programs to reduce  
462 musculoskeletal symptoms in the neck and shoulders among office workers.  
463 *Scandinavian Journal of Work, Environment & Health*, 34(1), 55–65.  
464 <https://doi.org/10.5271/sjweh.1192>

465 Buckley, P. (2015). Work-related Musculoskeletal Disorder ( WRMSDs ) Statistics , Great  
466 Britain , 2015. *Www.Hse.Gov.Uk/Statistics*, 1–20.

467 Goodwin, N., & Naylor, C. (2010). Managing people with long-term conditions. An Inquiry  
468 into the Quality of General Practice in England long-term conditions. *Inquiry*, 98.  
469 Retrieved from <http://www.kingsfund.org.uk/document.rm?id=8757>

470 Haas, M., Group, E., Muench, J., Kraemer, D., Brummel-Smith, K., Sharma, R., ...  
471 Fairweather, A. (2005). Chronic disease self-management program for low back pain in

472 the elderly. *Journal of Manipulative and Physiological Therapeutics*, 28(4), 228–237.  
473 <https://doi.org/10.1016/j.jmpt.2005.03.010>

474 Higgins, J., & Green, S. (2011). *Cochrane Handbook for Systematic Reviews of Interventions*  
475 Version 5.1.0. Retrieved January 4, 2017, from [www.handbook.cochrane.org](http://www.handbook.cochrane.org).

476 Hoe, V. C. W., Urquhart, D. M., Kelsall, H. L., & Sim, M. R. (2012). Ergonomic design and  
477 training for preventing work-related musculoskeletal disorders of the upper limb and  
478 neck in adults. *Cochrane Database Syst Rev*, 8(8), CD008570.  
479 <https://doi.org/10.1002/14651858.CD008570.pub2>

480 Hutting, N., Staal, J. B., Engels, J. A., Heerkens, Y. F., Detaille, S., & der Sanden, M. W. G.  
481 (2015). Effect evaluation of a self-management programme for employees with  
482 complaints of the arm, neck or shoulder: a randomised controlled trial. *Occupational*  
483 *and Environmental Medicine*, 72(12), 852–861. [https://doi.org/10.1136/oemed-2015-](https://doi.org/10.1136/oemed-2015-103089)  
484 103089

485 Jakobsen, M. D., Sundstrup, E., Brandt, M., Jay, K., Aagaard, P., & Andersen, L. L. (2015).  
486 Effect of workplace- versus home-based physical exercise on musculoskeletal pain  
487 among healthcare workers: a cluster randomized controlled trial. *Scandinavian Journal*  
488 *of Work, Environment & Health*, 41(2), 153–163. <https://doi.org/10.5271/sjweh.3479>

489 Jay, K., Brandt, M., Sundstrup, E., Schraefel, M. C. M., Jakobsen, M. D., Sjøgaard, G., &  
490 Andersen, L. L. (2014). Effect of individually tailored biopsychosocial workplace  
491 interventions on chronic musculoskeletal pain, stress and work ability among  
492 laboratory technicians: randomized controlled trial protocol. *BMC Musculoskeletal*  
493 *Disorders*, 15(8), 444. <https://doi.org/10.1186/1471-2474-15-444>

- 494 Jay, K., Frisch, D., Hansen, K., Zebis, M. K., Andersen, C. H., Mortensen, O. S., & Andersen, L.  
495 L. (2011). Kettlebell training for musculoskeletal and cardiovascular health: a  
496 randomized controlled trial. *Scandinavian Journal of Work, Environment & Health*,  
497 37(3), 196–203. <https://doi.org/10.5271/sjweh.3136>
- 498 Johns, G. (2009). Presenteeism in the workplace: A review and research agenda. *Journal of*  
499 *Organizational Behavior*, 31(4), 519–542. <https://doi.org/10.1002/job.630>
- 500 Johnston, V., & Shaw, W. S. (2013). Helping workers help themselves: empowering  
501 physiotherapy clients to manage musculoskeletal problems at work. *Physical Therapy*  
502 *Reviews*, 18(5), 373–378. <https://doi.org/10.1179/1743288X13Y.0000000087>
- 503 Katrak, P., Bialocerkowski, A. E., Massy-Westropp, N., Kumar, S., & Grimmer, K. A. (2004). A  
504 systematic review of the content of critical appraisal tools. *BMC Medical Research*  
505 *Methodology*, 4(1), 22. <https://doi.org/10.1186/1471-2288-4-22>
- 506 Kellett, K. M., Kellett, D. A., & Nordholm, L. A. (1991). Effects of an Exercise Program on Sick  
507 Leave Due to Back Pain. *Physical Therapy*, 71(4), 283–291. Retrieved from  
508 <http://ptjournal.apta.org/content/71/4/283.abstract>
- 509 Lambeek, L. C., Bosmans, J. E., Van Royen, B. J., Van Tulder, M. W., Van Mechelen, W., &  
510 Anema, J. R. (2010). Effect of integrated care for sick listed patients with chronic low  
511 back pain: economic evaluation alongside a randomised controlled trial. *British Medical*  
512 *Journal*, 341(nov30 1), c6414–c6414. <https://doi.org/10.1136/bmj.c6414>
- 513 Leyshon, R. et al. (2010). Ergonomic Interventions for Office Workers with Musculoskeletal  
514 Disorders: A Systematic Review. *Work*, 35(3), 335–348.
- 515 Macedo, A. M., Oakley, S. P., Panayi, G. S., & Kirkham, B. W. (2009). Functional and work



516 outcomes improve in patients with rheumatoid arthritis who receive targeted,  
517 comprehensive occupational therapy. *Arthritis Care and Research*, 61(11), 1522–1530.  
518 <https://doi.org/10.1002/art.24563>

519 MacKenzie, C. R., & de Melo-Martin, I. (2015). Ethical considerations in chronic  
520 musculoskeletal disease. *Current Reviews in Musculoskeletal Medicine*, 8(2), 128–133.  
521 <https://doi.org/10.1007/s12178-015-9271-1>

522 Magnavita, N., Elovainio, M., de Nardis, I., Heponiemi, T., & Bergamaschi, A. (2011).  
523 Environmental discomfort and musculoskeletal disorders. *Occupational Medicine*,  
524 61(3), 196–201. <https://doi.org/10.1093/occmed/kqr024>

525 Maher, G. C. (2000). A systematic review of workplace interventions to prevent low back  
526 pain. *Australian Journal of Physiotherapy*, 46(4), 259–269.

527 McGee, R., Bevan, S., & Quadrello, T. (2011). *Fit For Work? Musculoskeletal Disorders and*  
528 *the Canadian Labour Market. The Work Foundation*. Retrieved from  
529 [http://www.conferenceboard.ca/Libraries/NETWORK\\_PUBLIC/CCDPM\\_report2\\_jul2011.sflb](http://www.conferenceboard.ca/Libraries/NETWORK_PUBLIC/CCDPM_report2_jul2011.sflb)  
530 1.sflb

531 McGillivray, D. (2005). Fitter, happier, more productive: Governing working bodies through  
532 wellness. *Culture and Organization*, 11(2), 125–138.  
533 <https://doi.org/10.1080/14759550500091036>

534 Meenan, R. F., Mason, J. H., Anderson, J. J., Guccione, A. A., & Kazis, L. E. (1992). AIMS2. The  
535 Content and Properties of a Revised and Expanded Arthritis Impact Measurement  
536 Scales Health Status Questionnaire. *Arthritis & Rheumatism*, 35(1), 1–10.  
537 <https://doi.org/10.1002/art.1780350102>

- 538 Menta, R., Randhawa, K., Côté, P., Wong, J. J., Yu, H., Sutton, D., ... Taylor-Vaisey, A. (2015).  
539 The effectiveness of exercise for the management of musculoskeletal disorders and  
540 injuries of the elbow, forearm, wrist, and hand: A systematic review by the Ontario  
541 Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Journal of*  
542 *Manipulative and Physiological Therapeutics*, 38(7), 507–520.  
543 <https://doi.org/10.1016/j.jmpt.2015.06.002>
- 544 Mischke, C., Verbeek, J. H., Job, J., Morata, T. C., Alvesalo-Kuusi, A., Neuvonen, K., ... Pedlow,  
545 R. I. (2013). Occupational safety and health enforcement tools for preventing  
546 occupational diseases and injuries. *Cochrane Database of Systematic Reviews*, 8(8),  
547 CD010183. <https://doi.org/10.1002/14651858.CD010183.pub2>
- 548 Mulimani, P., Hoe, V. C. W., Hayes, M. J., Idiculla, J. J., Abas, A. B. L., & Karanth, L. (2014).  
549 Ergonomic interventions for preventing musculoskeletal disorders in dental care  
550 practitioners. *The Cochrane Library*, (8).  
551 <https://doi.org/10.1002/14651858.CD011261.www.cochranelibrary.com>
- 552 Nastasia, I., Coutu, M., & Tcaciuc, R. (2014). Topics and trends in research on non-clinical  
553 interventions aimed at preventing prolonged work disability in workers compensated  
554 for work-related musculoskeletal disorders (WRMSDs): a systematic, comprehensive  
555 literature review. *Disability Rehabilitation*, 36, 1841–1856.
- 556 NICE. (2015). Workplace health: management practices. Nice Guideline 13., (March).  
557 Retrieved from [https://www.nice.org.uk/guidance/ng13/resources/workplace-health-](https://www.nice.org.uk/guidance/ng13/resources/workplace-health-management-practices-1837269751237)  
558 [management-practices-1837269751237](https://www.nice.org.uk/guidance/ng13/resources/workplace-health-management-practices-1837269751237)
- 559 Nolte, S., & Osborne, R. H. (2013). A systematic review of outcomes of chronic disease self-

560 management interventions. *Quality of Life Research*, 22(7), 1805–1816.  
561 <https://doi.org/10.1007/s11136-012-0302-8>

562 Palmer, K. T., Harris, E. C., Linaker, C., Barker, M., Lawrence, W., Cooper, C., & Coggon, D.  
563 (2012). Effectiveness of community- and workplace-based interventions to manage  
564 musculoskeletal-related sickness absence and job loss: A systematic review.  
565 *Rheumatology*, 51(2), 230–242. <https://doi.org/10.1093/rheumatology/ker086>

566 Parry, S. P., Coenen, P., O’Sullivan, P. B., Maher, C. G., & Straker, L. M. (2017). Workplace  
567 interventions for increasing standing or walking for decreasing musculoskeletal  
568 symptoms in sedentary workers. In S. P. Parry (Ed.), *Cochrane Database of Systematic*  
569 *Reviews*. Chichester, UK: John Wiley & Sons, Ltd.  
570 <https://doi.org/10.1002/14651858.CD012487>

571 Proper, K. I., Koning, M., van der Beek, A., Hildebrandt, V., Bosscher, RJ, & van Mechelen, W.  
572 (2003). The effectiveness of worksite physical activity programs on physical activity,  
573 physical fitness, and health. *Clinical Journal of Sport Medicine*, 13(2), 106–117 12p.  
574 <https://doi.org/10.1097/00042752-200303000-00008>

575 Rla, F., Cumpston, M., Peeters, A., & Sa, C. (2013). Workplace pedometer interventions for  
576 increasing physical activity ( Review ) SUMMARY OF FINDINGS FOR THE MAIN  
577 COMPARISON, (4).  
578 <https://doi.org/10.1002/14651858.CD009209.pub2.www.cochranelibrary.com>

579 Roy, J.-S., MacDermid, J. C., Amick III, B. C., Shannon, H. S., McMurtry, R., Roth, J. H., ...  
580 Beaton, D. (2011). Validity and Responsiveness of Presenteeism Scales in Chronic Work-  
581 Related Upper-Extremity Disorders. *Physical Therapy*, 91(2), 254–266.

582 <https://doi.org/http://dx.doi.org/10.1111/j.1467-9280.2007.01910.x>

583 Rw, A., Tuntland, H., Ka, H., Røe, C., & Labriola, M. (2010). Workplace interventions for low-  
584 back pain in workers ( Protocol ). *Cochrane Database Syst Rev*, (4).  
585 <https://doi.org/10.1002/14651858.CD008160.pub2.www.cochranelibrary.com>

586 Safety and Health Practitioner. (2016). Dame Carol Black: “The workplace capability  
587 assessment has not been a success.” Retrieved from  
588 <http://www.shponline.co.uk/interview-with-dame-carol-black/>

589 Shiri, R., Martimo, K. P., Miranda, H., Ketola, R., Kaila-Kangas, L., Liira, H., ... Viikari-Juntura,  
590 E. (2011). The effect of workplace intervention on pain and sickness absence caused by  
591 upper-extremity musculoskeletal disorders. *Scandinavian Journal of Work, Environment*  
592 *and Health*, 37(2), 120–128. <https://doi.org/10.5271/sjweh.3141>

593 Shrestha, N., Ijaz, S., Kt, K., Kumar, S., & Cp, N. (2015). Workplace interventions for reducing  
594 sitting at work ( Review ) SUMMARY OF FINDINGS FOR THE MAIN COMPARISON. *The*  
595 *Cochrane Library*, (1).  
596 <https://doi.org/10.1002/14651858.CD010912.pub3.www.cochranelibrary.com>

597 Stern, C., Jordan, Z., & McArthur, A. (2014). Developing the review question and inclusion  
598 criteria. *The American Journal of Nursing*, 114(4), 53–6.  
599 <https://doi.org/10.1097/01.NAJ.0000445689.67800.86>

600 Strijk, J. E., Proper, K. I., van Stralen, M. M., Wijngaard, P., van Mechelen, W., & van der  
601 Beek, A. J. (2011). The role of work ability in the relationship between aerobic capacity  
602 and sick leave: a mediation analysis. *Occupational and Environmental Medicine*, 68(10),  
603 753–758. <https://doi.org/10.1136/oem.2010.057646>

604 Sundstrup, E., Jakobsen, M. D., Brandt, M., Jay, K., Aagaard, P., & Andersen, L. L. (2016).  
605 Strength Training Improves Fatigue Resistance and Self-Rated Health in Workers with  
606 Chronic Pain: A Randomized Controlled Trial. *BioMed Research International*, 2016, 1–  
607 11. <https://doi.org/10.1155/2016/4137918>

608 Sundstrup, E., Jakobsen, M. D., Brandt, M., Jay, K., Persson, R., Aagaard, P., & Andersen, L. L.  
609 (2014). Workplace strength training prevents deterioration of work ability among  
610 workers with chronic pain and work disability: a randomized controlled trial.  
611 *Scandinavian Journal of Work, Environment & Health*, 40(3), 244–51.  
612 <https://doi.org/10.5271/sjweh.3419>

613 Tehrani, N., Humpage, S., Willmott, B., & Haslam, I. (2007). *What's happening with well-*  
614 *being at work ? Chartered Institute of Personnel and Development.*  
615 <https://doi.org/10.1037/a0013504>

616 Tuomi, K., Ilmarinen, J., Jahkola, A., Katajarinne, L., & Tulkki, A. (1998). Work ability index.  
617 *Finnish Institute of Occupational Health.*

618 Waddell, G., & Burton, A. K. (2006). *Is Work Good for your health and well-being?* London:  
619 The Stationery Office. Retrieved from [http://iedereen-aandeslag.nl/wp-](http://iedereen-aandeslag.nl/wp-content/uploads/2016/07/hwwb-is-work-good-for-you.pdf)  
620 [content/uploads/2016/07/hwwb-is-work-good-for-you.pdf](http://iedereen-aandeslag.nl/wp-content/uploads/2016/07/hwwb-is-work-good-for-you.pdf)

621 Walker-Bone, K., & Linaker, C. (2016). Prediction of prognosis for people off sick with upper  
622 extremity musculoskeletal disorders. *Occupational and Environmental Medicine*, 73(0),  
623 805–806. <https://doi.org/10.1136/oemed-2016-103884>

624 WHO. (2001). The International Classification of Functioning, Disability and Health. *World*  
625 *Health Organization*, 18, 237. <https://doi.org/10.1097/01.pep.0000245823.21888.71>

626 WHO. (2016). Workplace health promotion. Retrieved September 23, 2016, from  
627 [http://www.who.int/occupational\\_health/topics/workplace/en/](http://www.who.int/occupational_health/topics/workplace/en/)

628 Williamson, A., & Hoggart, B. (2005). Pain: A review of three commonly used pain rating  
629 scales. *Journal of Clinical Nursing, 14*(7), 798–804. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2702.2005.01121.x)  
630 [2702.2005.01121.x](https://doi.org/10.1111/j.1365-2702.2005.01121.x)

631 World Health Organization. (2002). *Innovative Care for Chronic Conditions: building blocks  
632 for action. Noncommunicable Diseases and Mental Health World Health Organization.*  
633 Retrieved from <http://www.who.int/chp/knowledge/publications/icccglobalreport.pdf>

634 Wynne-Jones, G., Mallen, C. D., Mottram, S., Main, C. J., & Dunn, K. M. (2009). Identification  
635 of UK sickness certification rates, standardised for age and sex. *British Journal of  
636 General Practice, 59*(564), 510–516. <https://doi.org/10.3399/bjgp09X453431>

637 Zebis, M. K., Andersen, L. L., Pedersen, M. T. M. M. T., Mortensen, P., Andersen, C. H.,  
638 Pedersen, M. T. M. M. T., ... Sjøgaard, G. (2011). Implementation of neck/shoulder  
639 exercises for pain relief among industrial workers: a randomized controlled trial. *BMC  
640 Musculoskeletal Disorders, 12*(1), 205. <https://doi.org/10.1186/1471-2474-12-205>

641

642

643

644

645

646

647

648

649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662

663 **Tables**

664

665 **Table 1:** PICO approach

Population/problem	Employees with chronic/ long-term MSDs
Intervention	Workplace strategies/interventions to manage MSDs
Comparison	Any or none
Outcome	Pain severity, work status, symptoms, presenteeism and sickness absence

666  
667

668 **Table 2:** Inclusion and exclusion criteria

<b>Participant inclusion criteria</b>	<b>Participant exclusion criteria</b>
Working age male and female adults (18 to 68 years)	Specific pathological conditions (e.g. tumours, infections, fractures)
All sectors and types of jobs	Hypertension or cardiovascular diseases, symptomatic disc prolapses or severe disorders of the cervical spine, postoperative conditions in the neck and shoulder region, history of severe trauma, and pregnancy.
Workers with reported long-term musculoskeletal disorders / chronic MSK conditions (12 weeks or more) at any area of the body	Acute MSK disorders
Group-based and individual interventions conducted at the workplace	Guidelines, policies, recommendations
Interventions focused on management of chronic MSK conditions	Interventions focused on prevention and return to work
RCT design or cluster RCT design	Surveys and qualitative studies

669

670

671

672

673

674

675

676

677

678

679

680



681 Table 3: Study Characteristics

First Author, year	Country	Study design	Sample size	Age in years (mean)	Diagnosis	Occupation	Intervention Provider, Frequency, duration, length	Main Outcomes
Andersen, et al. 2008	Denmark	RCT	N=48 baseline N=48 follow-up	43,6	Neck muscle pain	7 different workplaces	<b>Provider</b> Experienced instructors  <b>Frequency</b> 3 times/week  <b>Duration</b> 20 min  <b>Length of observation</b> 10 weeks	<b>Pain intensity</b>  (in the trapezius muscle 0-100)
Andersen, et al. 2012	Denmark	RCT	N= 449 baseline N=280 follow-up	46	Neck and shoulder pain	Office workers	<b>Frequency/duration Intervention group 1:</b> 1 hour, once a week  <b>Frequency/duration Intervention group 2:</b> 20 mins three times a week  <b>Frequency/duration Intervention group 3:</b> 7 mins nine times a week  <b>Length of observation</b> 20-week intervention	<b>Pain intensity</b>  Neck and shoulders  <b>Health Status</b> (DASH, 1-25)

Andersen, et al. 2010	Denmark	Cluster RCT	N=222 baseline N=173 follow-up	46.5	Musculoskeletal pain symptoms in all regions of the body	Office workers from 12 geographically different units	<b>Provider</b> Experienced instructors  <b>Frequency</b> 3 times/week  <b>Duration</b> 20 min  <b>Length of observation</b> 10 weeks	<b>Pain intensity (0-9)</b>
Baldwin et al. 2012	USA	RCT	N=89 baseline N=75 follow-up	50.54	Rheumatoid Arthritis and Osteoarthritis	Office workers, health care practitioners, business and financial operations, manual workers, other categories	<b>Provider</b> occupational therapist (ergonomist)  <b>Duration</b> 2x 2.5 hours ergonomic sessions  <b>Length of observation</b> 12 and 24 months	<b>Functional status</b> AIMS2 physical component score (0-10 range)  <b>Pain</b> AIMS2 symptom component score (0-10 range)
Blangsted et al. 2008	Denmark	RCT	N= 616 baseline N=440 follow up	45.15	Chronic musculoskeletal symptoms in neck and shoulders	Office workers	<b>Intervention group 1:</b> <b>Frequency</b> 3 sessions per week <b>Duration</b> 20 min	<b>Pain intensity (0-9)</b> <b>Pain duration (days)</b> <b>Work ability Index(7-49)</b>

							<b>Intervention group 2</b> <b>Frequency</b> Visits from instructors 1 to 4 times per month  <b>Length of observation</b> 12 months	
Hutting et al. 2015	Netherlands	RCT	N= 129 baseline N=88 follow up	46.33	Non-specific complaints of the arm, neck or shoulder	Participants from different organisations	<b>Provider:</b> Physical therapist  <b>Frequency</b> 6 weekly sessions  <b>Duration</b> 2.5h  <b>Length of observation</b> 3, 6 and 12 months	<b>Health Status</b> (DASH general module) <b>Work Status</b> (DASH work module) <b>Absenteeism</b> (Days) <b>Pain</b> (NPRS)
Jakobsen et al. 2015	Denmark	RCT	N=200 baseline N=184 follow up N chronic workers in follow up=97	42.5	Musculoskeletal pain	Healthcare workers With acute pain and with chronic pain	<b>Provider</b> Training instructor  <b>Frequency</b> 5 times x 10 min/week  <b>Duration</b> 45-50min total  <b>Length of observation</b> 10 weeks	<b>Pain</b> (0-10)

Jay et al. 2011	Denmark	RCT	N=40 baseline N=33 follow-up	43.5	Neck/shoulder and low-back pain	Laboratory technicians	<p><b>Provider</b> Experienced kettlebell instructor</p> <p><b>Frequency</b> 3 days /week</p> <p><b>Duration</b> 20min sessions</p> <p><b>Length of observation</b> 8-week follow-up</p>	<p><b>Pain intensity of the neck/shoulder (0-10)</b></p> <p><b>Pain intensity of low back (0-10)</b></p>
Lambeek et al. 2010	Netherlands	RCT	N=134 baseline N=126 follow-up	46.15	Chronic low back pain	Any full time or part time paid work	<p><b>Intervention group Provider:</b> -employer, clinicians and OT ergonomists (multilevel focus)</p> <p><b>Control group Provider:</b> -medical specialist, occupational physician, general practitioner, and/or allied health professionals</p> <p><b>Length of observation</b> 3-6-12 months of follow-up</p>	<p><b>Neck Pain (0-10)</b></p> <p><b>Functional status</b> (Roland disability questionnaire, 0-24)</p> <p><b>Sick leave (Days)</b></p>

Shiri et al. 2011	Finland	RCT	N=222 baseline N=173 follow-up	45.2	Upper-extremity musculoskeletal disorders	Healthcare workers, clerical workers and warehouse workers	<b>Provider:</b> occupational therapist or physiotherapist  <b>Length of observation</b> 8-12-52 weeks	<b>Neck Pain (0-10)</b>
Sundstrup et al. 2014	Denmark	RCT	N=66 baseline N=66 follow- up	45.5	Upper-limb chronic pain	Slaughterhouse workers	<b>Provider</b> skilled instructor  <b>Frequency</b> 3 sessions/week  <b>Duration</b> 10 min/session  <b>Length of observation</b> 10 weeks follow up	<b>Work ability index (WAI)</b> (7-49)  Item 5: Sick leave (1-5)
Zebis et al., 2011	Denmark	RCT	N=537 baseline N=448 follow-up	41	Non-specific neck and shoulder pain	Industrial workers	<b>Provider</b> Educated supervisors on the manual  <b>Frequency</b> 3 sessions/week  <b>Duration</b> 20 min per session  <b>Length of observation</b> 20-week period	<b>Neck pain intensity (0-9)</b>  <b>Right shoulder pain intensity (0-9)</b>  <b>Left shoulder pain intensity (0-9)</b>

Table 4: Quality Appraisal using SIGN appraisal tool for RCTs

Checklist for RCTs: SIGN items	Appropriate and clearly focused question	Randomised allocation	Adequate concealment method is used	Blind treatment allocation	Treatment and control groups are similar at the start of the trial.	The only difference between groups is the treatment under investigation	All relevant outcomes are measured in a standard, valid and reliable way.	Drop-out rates	Intention to treat analysis	Results are comparable for all sites.	How well was the study done to minimise bias?	Are the results of this study directly applicable to the patient group targeted by this review?	The overall effect is due to the study
Authors													
Andersen et al. 2008	Yes	Yes	Yes	Yes	Yes	Can't say	Yes	<20 %	Yes	Yes	++	Yes	Yes
Andersen et al. 2010	Yes	Yes	Yes	Yes	Yes	Can't say	Yes	<20 %	Yes	Yes	++	Yes	Yes
Andersen et al. 2012	Yes	Yes	Yes	Yes	Yes	Can't say	Yes	<20 %	Yes	Yes	++	Yes	Yes
Baldwin et al. 2012	Yes	Can't say	Can't say	Yes	No	No	Yes	15.7 0%	Yes	Yes	+	Yes	Can't say
Blangsted et al. 2008	Yes	Yes	Yes	No	Yes	Yes	Yes	19.8 0%	Can't say	Yes	++	Yes	Yes

Checklist for RCTs: SIGN items	Appropriate and clearly focused question	Randomised allocation	Adequate concealment method is used	Blind treatment allocation	Treatment and control groups are similar at the start of the trial.	The only difference between groups is the treatment under investigation	All relevant outcomes are measured in a standard, valid and reliable way.	Drop-out rates	Intention to treat analysis	Results are comparable for all sites.	How well was the study done to minimise bias?	Are the results of this study directly applicable to the patient group targeted by this review?	The overall effect is due to the study
Authors													
Hutting et al. 2015	Yes	Yes	Yes	No	Yes	Yes	Yes	40%	Yes	Yes	++	Yes	Yes
Jakobsen et al. 2015	Yes	Yes	Yes	Yes	Yes	No	Yes	<20 %	Yes	Yes	++	Yes	Yes
Jay et al. 2011	Yes	Yes	Yes	No	Yes	Yes	Yes	17.5 %	No	N/A	+	Yes	Can't say
Lambeek et al. 2010	Yes	Yes	Yes	No	Yes	Yes	Yes	13%	Yes	Yes	++	Yes	Yes
Shiri et al. 2011	Yes	Yes	Yes	Can't say	Yes	No	Yes	<20 %	Can't say	Yes	+	Yes	Yes
Sundstrup et al. 2014	Yes	Can't say	Yes	No	Yes	Yes	Yes	8%	Yes	N/A	+	Yes	Yes

Zebis et al. 2011	Yes	Yes	Yes	Can't say	Yes	No	Yes	15%	Yes	Can't say	+	Yes	Yes
----------------------	-----	-----	-----	--------------	-----	----	-----	-----	-----	--------------	---	-----	-----

685



Author, Year	Intervention	Results			
		Intervention group 1 (SST)	Intervention group 2 (GFT)	Control group	Between groups comparison
Andersen et al. 2008	<b>Intervention group 1</b> Specific strength training (SST)				
	<b>Intervention group 2</b> General fitness training (GFT)				
	<b>Control group</b> Health counselling				
	<b>General pain</b> (in Trapezius muscle 0-100)				
	<b>10-weeks</b>	<b>10-weeks</b>	<b>10-weeks</b>		<b>No between groups comparisons</b>
	$\Delta=-12$ Rate of decrease =1.03 ( $\pm 0.30$ ) $p < 0.0001^{**}$	$\Delta=-6$	$\Delta=-1$		
<b>Worst pain</b> (in Trapezius muscle 0-100)					
<b>10-weeks</b>	<b>10-weeks</b>	<b>10-weeks</b>		<b>No between groups comparisons</b>	
$\Delta=-25$ Rate of decrease=-0.58 ( $\pm 0.22$ ) $p < 0.0001^*$	$\Delta=-11$	$\Delta=-5$			
<b>Acute pain</b> (in Trapezius muscle 0-100)					
<b>10-weeks</b>	<b>10-weeks</b>	<b>10-weeks</b>		<b>No between groups comparisons</b>	
Rate of decrease= 4.8 $p < 0.05^*$  The acute adverse effect lasted 2 hours	Rate of decrease = 5.3 $p < 0.01^{**}$				

Andersen et al. 2010	<b>Intervention group 1</b> Specific resistance training (SRT)	<b>Intervention group 1</b>	<b>Intervention group 2</b>	<b>Control group</b>	<b>Between groups comparison</b>	
	<b>Neck pain (0-9)</b>					
	<b>Intervention group 2</b> All-round physical exercise (APE)	$\Delta=-0.73\pm0.36$ $p<0.05^*$	$\Delta=-0.91\pm0.31$ $p<0.01^{**}$	$\Delta=0.40\pm0.32$ $P>0.05$	Intervention groups 1 and 2 vs Control	
	<b>Control group</b> Reference group (REF): Encouragement and advice					
Andersen et al. 2012	<b>Intervention group 1</b> Specific strength training 1 hour, once a week	<b>Intervention group 1</b>	<b>Intervention group 2</b>	<b>Intervention group 3</b>	<b>Between groups comparison</b>	
	<b>Neck pain (0-10)</b>					
	<b>Intervention group 2</b> Specific strength training 20 min three times a week	<b>20 weeks</b> $\Delta=-0.74$ $p<0.01^{**}$	<b>20 weeks</b> 0.78 $p<0.01^{**}$	<b>20 weeks</b> 0.71	<b>No between groups comparisons</b>	
	<b>Intervention group 3</b> Specific strength training 7 min nine times a week	<b>Right shoulder pain (0-10)</b>				
		$\Delta=-0.94$ $p<0.01^{**}$	$\Delta=-0.61$	$\Delta=-0.83$ $p<0.01^{**}$	<b>No between groups comparisons</b>	
	<b>Control group</b> No physical training	<b>Left shoulder pain (0-10)</b>				
	$\Delta=-0.69$ $p<0.01^{**}$	$\Delta=-0.32$	$\Delta=-0.62$	<b>No between groups comparisons</b>		

		Health Status (DASH, 1-25)				
		$\Delta=-6$ $p<0.01^{**}$	$\Delta=-9$ $p<0.05^*$	$\Delta=-2$	No between groups comparisons	
Baldwin et al. 2012	<b>Intervention group</b>	<b>Intervention group</b>		<b>Control group</b>		
		<b>Functional status (AIMS2 physical component 0-10)</b>				
	1 session of workplace ergonomic assessment and intervention (ergonomic, exercises, workstation equipment modifications and a self-management manual)	<b>12 months</b> $\Delta=-0.24 (\pm 0.94)$ $p < 0.04^*$	<b>12 months</b> $\Delta=-0.09 (\pm 0.66)$ $p < 0.26$	<b>12 months</b> 1.63 ( $\pm 1.27$ ) intervention 1.26 ( $\pm 1.23$ ) control $p = 0.45$		
	1 follow up session	<b>24 months</b> $\Delta=-0.29 (\pm 0.80)$ $p < 0.01^{**}$	<b>24 months</b> $\Delta=-0.12 (\pm 0.82)$ $p < 0.25$	<b>24 months</b> 1.58 ( $\pm 1.09$ ) intervention 1.23 ( $\pm 1.18$ ) control $p = 0.76$		
	Follow-up phone call after a month to determine if modifications to the work plan were desired	<b>Pain (AIMS2 symptom component 0-10)</b>				
	A resource manual with guides for self-management of arthritis and possible ergonomic interventions in the work setting	<b>12 months</b> $\Delta=-1.27 (\pm 2.00)$ $p < 0.01^{**}$	<b>12 months</b> $\Delta=-0.61 (\pm 1.93)$ $p < 0.07$	<b>12 months</b> 4.60 ( $\pm 2.44$ ) intervention 4.16 ( $\pm 2.37$ ) control $p = 0.58$		
<b>Control group</b>	<b>24 months</b> $\Delta=-1.25 (\pm 2.16)$ $p < 0.01^{**}$	<b>24 months</b> $\Delta=-0.29 (\pm 1.94)$ $p < 0.34$	<b>24 months</b> 4.62 ( $\pm 2.22$ ) intervention 4.48 ( $\pm 2.31$ ) control $p = 0.42$			
	Written educational materials (same resource manual that was provided for the intervention group)					

687

<b>Intervention group 1</b>	<b>Between groups comparisons only</b>
-----------------------------	--

Blangsted et al. 2008	Specific resistance training	No specific numerical values provided for changes in each group: Values were presented only on a histogram		
	<p><b>Intervention group 2</b> All-round physical exercise daily to increase physical activity both at the worksite and during leisure time</p> <p><b>Control group</b> Education on general health-promoting activities</p>	<b>12 months</b>		
		<p><b>Comparison of both intervention (group 1 and 2) vs Control group</b>  <b>Pain intensity</b> (p=0.0318) * <i>in favour of the activity interventions</i>  <b>Pain duration</b> (p=0.0565)</p> <p><b>Work ability</b> (p = 0.3073)</p> <p><b>Comparison of intervention group 1 vs intervention group 2</b></p> <p><b>Pain intensity</b> (p=0.5327)  <b>Pain duration</b> (p=0.4046)</p> <p><b>Work ability</b> (p = 0.3073)</p>		
Hutting et al. 2015	<p><b>Intervention group</b></p> <p>Moderated self-management interventions at the workplace within group sessions</p> <p>E-module on Health (available for 12 months)</p>	<b>Self-management group (SU)</b>	<b>Usual care group (UCG)</b>	<b>Between groups comparison</b>
		<b>Health status (DASH general module 0-5)</b>		
		<b>12 months</b> Δ=-7.96	<b>12 months</b> Δ=-7.22	<b>12 months</b> -0.73 p < 0.10

	<b>Control group</b> Usual care and information available within the organisation or outside the organisation.			
		<b>Work Status (DASH work module 0-5)</b>		
		<b>12 months</b> $\Delta=-0.27$	<b>12 months</b> $\Delta=-1.63$	<b>12 months</b> $p=0.04^*$ <i>in favour of the self-management group</i>
		<b>Absenteeism (days)</b>		
		<b>12 months</b> $\Delta=-0.27$	<b>12 months</b> $\Delta=-1.63$	<b>12 months</b> 4.19 $p=0.29$
		<b>Pain the last week (NPRS)</b>		
	<b>12 months</b> $\Delta=-0.61$	<b>12 months</b> $\Delta=-1.2$	<b>12 months</b> -0.63 $p=0.47$	
Jakobsen et al. 2015	<b>Intervention group</b> Strength training at the workplace Ergonomic training and education	<b>Intervention group</b>	<b>Control group</b>	<b>Between groups comparison</b>
	<b>Control group</b> Physical exercise intervention at home with the help of posters and instructions Ergonomic training and education	<b>Average Pain (0-10) (Only the results of the chronic pain groups are reported)</b>		
	<b>10 Weeks</b> $\Delta=-1.7$ $p < 0.0001^{**}$	<b>10 Weeks</b> $\Delta=-0.8$ $p < 0.0001^{**}$	<b>10 Weeks</b> -1.0 $p < 0.0003^{**}$ <i>in favour of the intervention group</i>	
Jay et al. 2011	<b>Intervention group</b>	<b>Intervention group</b>	<b>Control group</b>	<b>Between groups comparison</b>
		<b>Pain intensity of the neck/shoulder (0-10)</b>		

	Progressive worksite intervention using Kettlebell training	<b>8-weeks</b> $\Delta=-1.7$	<b>8-weeks</b> $\Delta=0.3$	<b>8-weeks</b> -2.1 <i>p=0.02* in favour of the intervention group</i>
	<b>Control group</b>	<b>Pain intensity of the low back (0-10)</b>		
	Recommendations	<b>8-weeks</b> $\Delta=-1.6$	<b>8-weeks</b> $\Delta=-0.2$	<b>8-weeks</b> -1.4 <i>p=0.05* in favour of the intervention group</i>
Lambeek et al. 2010	<b>Intervention group</b> Integrated care	<b>Intervention group</b>	<b>Control group</b>	<b>Between groups comparison</b>
	<b>Control group</b> Usual care	<b>Neck pain (0-10)</b>		
		<b>3 months</b> $\Delta=-1.11 (\pm 0.39)$	<b>3 months</b> $\Delta=-1.59 (\pm 0.38)$	<b>3 months</b> -0.99 <i>p &lt; 0.08</i>
		<b>12 months</b> $\Delta=-1.64 (\pm 0.35)$	<b>12 months</b> $\Delta=-1.85 (\pm 0.36)$	<b>12 months</b> $\Delta=-0.21$ <i>p &lt; 0.67</i>
		<b>Functional Status (Roland disability questionnaire 0-24)</b>		
	<b>3 months</b> $\Delta=-3.76 (\pm 0.86)$	<b>3 months</b> $\Delta=-3.82 (\pm 0.85)$	<b>3 months</b> $\Delta=-0.11$ <i>p &lt; 0.93</i>	
	<b>12 months</b> $\Delta=-7.16 (\pm 0.71)$	<b>12 months</b> $\Delta=-4.43 (\pm 0.72)$	<b>12 months</b> $\Delta=-2.86$ <i>p &lt; 0.001** in favour of the intervention group</i>	

		Sick leave (days)		
		<b>3 months</b> 88 days	<b>3 months</b> 208 days	<b>3 months</b> p=0.003** <i>in favour of the intervention group</i>
		<b>12 months</b> 82 days	<b>12 months</b> 175 days	<b>12 months</b> p=0.003** <i>in favour of the intervention group</i>
Shiri et al. 2011	<b>Intervention group</b> Workplace assessment by an occupational therapist or physiotherapist	<b>Intervention group</b>	<b>Control group</b>	<b>Between groups comparison</b>
		<b>Pain intensity (0-10)</b>		
	<b>Control group</b> No intervention	<b>2 Weeks</b> Δ=-1.27	<b>2 Weeks</b> Δ=-0.69	<b>2 Weeks</b> -0.58 p=0.05* <i>in favour of the intervention group</i>
Sundstrup et al. 2014	<b>Intervention group</b> High intensity strength training	<b>Intervention group</b>	<b>Control group</b>	<b>Between groups comparison</b>
		<b>WAI Item 5: Sick leave (1-5)</b>		
	<b>Control group</b> Ergonomic training and education	<b>10 weeks</b> Δ=-0.2	<b>10 weeks</b> Δ=-0.5	<b>10 weeks</b> -2.3 p = 0.2
		<b>WAI Index Total (7-49)</b>		
	<b>10 weeks</b> Δ=-0.3	<b>10 weeks</b> Δ=-2.2 WAI decreased (i.e. worsened) in the ergonomic group p<0.01**	<b>10 weeks</b> -2.3 p = 0.012** <i>in favour of the intervention group</i>	

Zebis et al. 2011	Intervention group	Intervention group	Control group	Between groups comparison
High-intensity specific strength training at the workplace  <b>Control group</b>  Advice to stay physically active, weekly consultation	<b>Neck pain (0-9)</b>			
	<b>20 weeks</b> $\Delta = -1.8 (\pm 1.9)$	<b>20 weeks</b> $\Delta = -2.9 (\pm 2.3)$	<b>20 weeks</b> -1.1 <i>P &lt; 0.001** in favour of the intervention group</i>	
	<b>Shoulder pain (0-9)</b>			
	<b>20 weeks</b>	<b>20 weeks</b>	<b>20 weeks</b>	
	<b>Right Shoulder pain</b> $\Delta = -1.4 (\pm 1.7)$  <b>Left Shoulder pain</b> $\Delta = -0.9 (\pm 1.3)$	<b>Right Shoulder pain</b> $\Delta = -2.5 (\pm 2.6)$  <b>Left Shoulder pain</b> $\Delta = -2.2 (\pm 2.6)$	<b>Right Shoulder pain</b> -1.1 <b>Left Shoulder pain</b> $\Delta = -1.3$	
Results are presented in mean values and/or standard error, $\Delta$ demonstrates the difference between the baseline values and the time of the relevant measurement, * highlights significant difference of $p < 0.05$ , ** highlights significant difference of $p < 0.01$				

688



689 Appendix 1

690

691 *Words (and synonyms) used for Search Strategy*

randomized controlled trial	system*
worker*	improve
employe*	decrease
staff	cope
personnel	manage
workforce OR “work force”	prevent
“labour force”	control
strateg*	avoid
tactic*	reduce
intervention*	stop
practice	“deal with”
Policy	musculoskeletal
treatment*	MSK
plan*	chronic
approach*	condition*
method*	disease*
protocol*	disorder*
musculoskeletal disorders	“ill health”
process*	illness*

system*	pathosis
improve	complaint

692

693 *MEDLINE Search Strategy example*

Search ID#	Search Terms	Results
S1 AND S2 AND S3 AND S4 AND S5 AND S6 AND S7		203
S1 AND S2 AND S3 AND S4 AND S5 AND S6 AND S7 AND S8 AND S9		5
S9	(MM "Therapeutics+")	1,956,742
S8	(MM "Health Personnel+") OR (MM "Health Manpower")	320,520
S7	workplace OR work	826,220
S6	chronic	1,135,779
S5	condition* OR disease* OR disorder* OR "ill health "OR pathosis OR illness* OR complaint*	8,207,606

S4	MSK OR musculoskeletal OR "chronic MSK" or "chronic musculoskeletal"	64,994
S3	manage* OR Prevent* OR cope* OR decrease* OR improve* OR control* OR handle* OR avoid* OR reduce* OR stop* OR "deal with "	9,043,205
S2	(strateg* OR tactic*) OR intervention* OR practice* OR polic* OR treatment* OR plan* OR approach* OR method* OR protocol* OR process* OR system*	13,445,401
S1	employer OR employee* OR worker* OR (workforce OR workforce) OR staff OR personnel OR ("labour force" OR labor force )	908,429