

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/126059>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2019 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

# SYMPLECTIC AND ORTHOGONAL $K$ -GROUPS OF THE INTEGERS

MARCO SCHLICHTING

ABSTRACT. Nous calculons explicitement les groupes d'homotopie des espaces topologiques  $B\mathrm{Sp}(\mathbb{Z})^+$ ,  $BO_{\infty,\infty}(\mathbb{Z})^+$  et  $BO_{\infty}(\mathbb{Z})^+$ .

We explicitly compute the homotopy groups of the topological spaces  $B\mathrm{Sp}(\mathbb{Z})^+$ ,  $BO_{\infty,\infty}(\mathbb{Z})^+$  and  $BO_{\infty}(\mathbb{Z})^+$ .

## 1. ÉNONCÉ DES RÉSULTATS

Soient  $\mathrm{Sp}(\mathbb{Z})$ ,  $O_{\infty,\infty}(\mathbb{Z})$  et  $O_{\infty}(\mathbb{Z})$  le groupe symplectique infini, le  $\langle 1, -1 \rangle$ -groupe orthogonal infini et le groupe orthogonal hyperbolique sur l'anneau des entiers  $\mathbb{Z}$ . Ils sont obtenus comme réunion des sous-groupes  $\mathrm{Sp}_{2n}(\mathbb{Z})$ ,  $O_{n,n}(\mathbb{Z})$  et  $O_{2n}(\mathbb{Z})$  de  $GL_{2n}(\mathbb{Z})$  laissant invariant les formes bilinéaires de matrices de Gram

$$\begin{pmatrix} 0 & 1 & & \\ -1 & 0 & & \\ & & \ddots & \\ & & & 0 & 1 \\ & & & -1 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & & \\ 0 & -1 & & \\ & & \ddots & \\ & & & 1 & 0 \\ & & & 0 & -1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1 & & \\ 1 & 0 & & \\ & & \ddots & \\ & & & 0 & 1 \\ & & & 1 & 0 \end{pmatrix}.$$

Les groupes  $\mathrm{Sp}(\mathbb{Z})$ ,  $O_{\infty,\infty}(\mathbb{Z})$  et  $O_{\infty}(\mathbb{Z})$  ont des sous-groupes de commutateurs parfaits. Rappelons que pour un tel groupe  $G$  la construction plus de Quillen  $BG^+$  appliquée à l'espace classifiant  $BG$  de  $G$  est munie d'une application continue  $BG \rightarrow BG^+$  qui induit un isomorphisme sur les groupes d'homologie intégrale et vaut  $G \rightarrow G/[G, G]$  sur  $\pi_1$ .

Le but de cet article est de calculer explicitement les groupes d'homotopie des espaces topologiques  $B\mathrm{Sp}(\mathbb{Z})^+$ ,  $BO_{\infty,\infty}(\mathbb{Z})^+$  et  $BO_{\infty}(\mathbb{Z})^+$ . Ces espaces sont des espaces de lacets infinis puisqu'ils sont les composants connexes des espaces de la  $K$ -théorie  $K\mathrm{Sp}(\mathbb{Z})$ ,  $GW(\mathbb{Z})$  et  $KQ(\mathbb{Z})$  des formes non dégénérées symplectiques, bilinéaires symétriques et quadratiques sur  $\mathbb{Z}$ . On sait que les groupes d'homotopie de ces espaces sont des groupes abéliens de génération finie.

Pour un groupe abélien  $A$ , on note  $A_{\text{odd}}$  le sous-groupe des éléments d'ordre impaire fini.

**Theorem 1.1.** *Les groupes d'homotopie des espaces  $B\mathrm{Sp}(\mathbb{Z})^+$  et  $BO_{\infty,\infty}(\mathbb{Z})^+$  pour  $n \geq 1$  sont donnés dans le tableau du Theorem 2.1*

**Theorem 1.2.** *L'application qui envoie une forme quadratique sur sa forme bilinéaire symétrique associée induit un morphisme d'espaces de  $K$ -théorie  $KQ(\mathbb{Z}) \rightarrow GW(\mathbb{Z})$  qui est un isomorphisme*

$$\pi_n BO_{\infty}(\mathbb{Z})^+ \xrightarrow{\cong} \pi_n BO_{\infty,\infty}(\mathbb{Z})^+ \quad \text{en degré } n \geq 2$$

et le monomorphisme  $(\mathbb{Z}/2)^2 \subset (\mathbb{Z}/2)^3$  en degré  $n = 1$ .

*Date:* September 9, 2019.

This work was partially funded by the Leverhulme Trust.



conjectured to be zero [Wei05, Introduction]. For example,  $K_4(\mathbb{Z}) = 0$  [Rog00]. In particular for  $n \geq 1$  we have the following table of homotopy groups

$n \pmod 8$	0	1	2	3	4	5	6	7
$\pi_n BSp(\mathbb{Z})^+$	(0?)	0	$\mathbb{Z}$	$\mathbb{Z}/2d_n$	$\mathbb{Z}/2 \oplus (0?)$	$\mathbb{Z}/2$	$\mathbb{Z}$	$\mathbb{Z}/d_n$
$\pi_n BO_{\infty, \infty}(\mathbb{Z})^+$	$\mathbb{Z} \oplus \mathbb{Z}/2 \oplus (0?)$	$(\mathbb{Z}/2)^3$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/d_n$	$\mathbb{Z} \oplus (0?)$	0	0	$\mathbb{Z}/d_n$

where (0?) denotes a finite group of odd order conjectured to be zero.

### 3. PROOF PART 1: ODD TORSION

**Lemma 3.1.** *Let  $R$  be the ring of integers in a number field  $F$ . Then for all  $n \geq 0$  there are isomorphisms*

$$KQ_n(R)_{\text{odd}} \cong GW_n(R)_{\text{odd}} \cong KSp_n(R)_{\text{odd}} \cong (K_n(R)_{\text{odd}})^{C_2}$$

where the action of  $C_2$  on  $K$ -theory is induced by  $GL(R) \rightarrow GL(R) : M \mapsto {}^t M^{-1}$ .

*Proof.* The natural map  $KQ_n(R)_{\text{odd}} \rightarrow GW_n(R)_{\text{odd}}$  is an isomorphism with inverse the cup product with the quadratic space associated with the Leech lattice  $\Gamma_8$  [MH73, Ch. 2, §6]. Write  $GW^{[0]}(R)$  and  $GW^{[2]}(R)$  for  $GW(R)$  and  $KSp(R)$ ; see Section 4 below for general  $GW^{[n]}$ . The hyperbolic and forgetful maps factor as  $K^{[r]}(R)_{hC_2} \rightarrow GW^{[r]}(R) \rightarrow K^{[r]}(R)^{hC_2}$ ; see [Sch17, (7.3) and Lemma 7.4] which doesn't use  $1/2 \in R$ . Here  $K^{[n]}$  denotes the  $K$ -theory spectrum  $K$  with  $C_2$ -action induced by the  $n$ -th shifted duality  $\text{Hom}(\_, R[n])$ . On the spectrum level, this action depends on  $n = 0, 2$ . However, on homotopy groups the actions agree for  $n = 0, 2$ . Denote by  $L^{[r]}$  the homotopy cofibre of the map of spectra<sup>1</sup>  $K^{[r]}(R)_{hC_2} \rightarrow GW^{[r]}(R)$ , then  $L_i^{[r]} = L_{i-1}^{[r-1]}$  only depends on the difference  $n - i$ ,  $i \geq 1$  [Schc] and

$$GW_n^{[r]}(R)[1/2] \cong K_n^{[r]}(R)[1/2]^{C_2} \oplus L_n^{[r]}(R)[1/2]$$

since the composition  $K^{[r]}(R)[1/2]_{hC_2} \rightarrow GW^{[r]}(R)[1/2] \rightarrow K^{[r]}(R)[1/2]^{hC_2}$  is an equivalence [Sch17, Lemma B.14]. Strictly speaking we define a non-connective version of  $L^{[r]}$  as the homotopy colimit of the sequence

$$(3.1) \quad GW^{[r]} \rightarrow S^1 \wedge GW^{[r-1]} \rightarrow S^2 \wedge GW^{[r-2]} \rightarrow \dots$$

with appropriate delooping of  $GW^{[n]}$  as in [Sch17] using the definition of  $\mathcal{E}^{[n]}$  as below. The maps in (3.1) are the connecting maps of the homotopy fibration (4.1). Then we have *by definition*  $L_i^{[n]} = L_0^{[n-i]}$  and as in [Sch17] we formally obtain the homotopy fibration whose connected cover we used above:

$$(K^{[n]})_{hC_2} \rightarrow GW^{[n]} \rightarrow L^{[n]}.$$

By Lemma 4.4 below, the canonical map  $L_i^{[r]}(R)[1/2] \rightarrow L_i^{[r]}(F)[1/2]$  is an isomorphism for  $i \geq r$ . By [Sch17, Proposition 7.2] and [Bal01, Theorem 5.6], we

<sup>1</sup>All spectra in this paper are  $-1$ -connected, and all homotopy fibrations are in the category of  $-1$ -connected spectra unless otherwise stated. In particular, the second map of a homotopy fibration need not be surjective on  $\pi_0$

have

$$L_i^{[r]}(F)[1/2] = \begin{cases} W(F)[1/2] & r \equiv i \pmod{4} \\ 0 & \text{else} \end{cases}$$

where  $W(F)$  is the usual Witt group of  $F$ . But it is well-known that  $W(F)[1/2]$  is a free  $\mathbb{Z}[1/2]$ -module of rank the number of orderings of  $F$ . This proves the lemma for  $K_n Q$ ,  $GW_n$  for  $n \geq 0$  and  $K_n \text{Sp}$  for  $n \geq 2$ . From the Zariski local to global spectral sequence, we see  $L_1^{[2]}(R)[1/2] = L_0^{[1]}(R)[1/2] = H^1(R, L_0^0[1/2]) = 0$  since  $L_0^0[1/2]$  is constant (flasque) on a ring of integers  $R$  and  $L_0^{[1]}$  is Zariski-locally trivial. So,  $K_1 \text{Sp}(R)_{\text{odd}} = (K_1(R)_{\text{odd}})^{C_2}$ . Finally,  $L_0^{[2]}(R) = 0$  for a ring of integers since  $K_0 \text{Sp}(R) = H^0(R, \mathbb{Z})$ , by the Zariski spectral sequence, hence  $H : K_0(R) \rightarrow K_0 \text{Sp}(R)$  is surjective and  $L_0^{[2]} = 0$ .  $\square$

Continue to assume that  $R$  is a ring of integers in a number field. Let  $\ell \in \mathbb{Z}$  be an odd prime and set  $R' = R[1/\ell]$ . Then the inclusion  $R \subset R'$  induces an isomorphism:  $K_n(R)\{\ell\} \cong K_n(R')\{\ell\}$  on  $\ell$ -primary torsion subgroups for  $n \geq 1$ . For  $i \geq 1$  the abelian group  $K_{2i}(R')$  is finite and the group  $K_{2i-1}(R')$  is finitely generated. For all  $i \geq 1$  and large  $\nu$  we therefore have an exact sequence

$$(3.2) \quad 0 \rightarrow K_{2i}(R')\{\ell\} \rightarrow K_{2i}(R', \mathbb{Z}/\ell^\nu) \rightarrow K_{2i-1}(R')\{\ell\} \rightarrow 0$$

[Wei05, Lemma 68]. Since  $\ell$  is invertible in  $R'$  which has  $\text{cd}_\ell(R') \leq 2$ , the proved Quillen-Lichtenbaum conjecture says that the following change of topology map is an isomorphism  $K_{2i}(R', \mathbb{Z}/\ell^\nu) \cong K_{2i}^{\text{ét}}(R', \mathbb{Z}/\ell^\nu)$  for  $i \geq 1$ . The change of topology map is  $C_2$ -equivariant. From the étale local to global spectral sequence for  $K^{\text{ét}}$  we obtain the  $C_2$ -equivariant isomorphism

$$(3.3) \quad K_{2i}(R', \mathbb{Z}/\ell^\nu) \cong K_{2i}^{\text{ét}}(R', \mathbb{Z}/\ell^\nu) \cong H_{\text{ét}}^0(R', K_{2i}/\ell^\nu)$$

[Wei05, Proof of Theorem 70] on which the action on the left is  $GL(R) \rightarrow GL(R) : M \mapsto {}^t M^{-1}$  and on the right hand side it is multiplication with  $(-1)^i$ . Combining (3.2) and (3.3), Lemma 3.1 yields the following.

**Theorem 3.2.** *Let  $R$  be a ring of integers in a number field, and  $\ell \in \mathbb{Z}$  an odd prime. Then for all  $n \geq 1$  we have isomorphisms*

$$GW_n(R)\{\ell\} \cong K \text{Sp}_n(R)\{\ell\} \cong KQ_n(R)\{\ell\} \cong \begin{cases} K_n(R)\{\ell\} & n \equiv 0, 3 \pmod{4} \\ 0 & n \equiv 1, 2 \pmod{4}. \end{cases}$$

#### 4. PROOF PART 2: 2-ADIC COMPUTATIONS

For an exact category with weak equivalences and duality  $(\mathcal{E}, w, \#, \text{can})$ , denote by  $GW(\mathcal{E}, w, \#, \text{can})$  the associated Grothendieck-Witt space of symmetric bilinear forms [Sch10, Definition 3]. If  $\mathcal{E}$  has a strong symmetric cone [Sch10, Definition 4], [Sch] I denote by  $\mathcal{E}^{[1]} = (\text{Mor } \mathcal{E}, w_{\text{cone}}, \#, \text{can})$  the exact category with weak equivalences and duality of morphisms in  $\mathcal{E}$  with duality and double dual identification induced by functoriality of  $\#$  and  $\text{can}$  and weak equivalences those maps  $f \rightarrow g$  of arrows in  $\mathcal{E}$  such that  $\text{cone}(f) \rightarrow \text{cone}(g)$  is a weak equivalence in  $\mathcal{E}$ . By functoriality,  $\mathcal{E}^{[1]}$  also has a strong symmetric cone. Set  $GW^{[0]}(\mathcal{E}) = GW(\mathcal{E})$  and define inductively for  $r \geq 1$

$$GW^{[r+1]}(\mathcal{E}) = GW^{[r]}(\mathcal{E}^{[1]}).$$

By [Sch10, Theorem 6], the sequence

$$\mathcal{E} \xrightarrow{E \mapsto 1_E} \text{Mor } \mathcal{E} \xrightarrow{1} \mathcal{E}^{[1]}$$

induces a homotopy fibration  $GW(\mathcal{E}) \rightarrow K(\mathcal{E}) \rightarrow GW^{[1]}(\mathcal{E})$  of  $-1$ -connected spectra and by iteration the homotopy fibration

$$(4.1) \quad GW^{[r]}(\mathcal{E}) \rightarrow K(\mathcal{E}) \rightarrow GW^{[r+1]}(\mathcal{E});$$

compare [Sch17, Proof of Proposition 4.9]. For details and a generalisation; see [Schc]. For  $r < 0$ , we define  $GW^{[r]}(\mathcal{E})$  such that (4.1) holds for all  $r \in \mathbb{Z}$ . For a commutative ring  $R$ , we denote by  $GW^{[r]}(R)$  the space  $GW^{[r]}(\text{Ch}^b \mathcal{P}(R), \text{quis}, \text{Hom}(\_, R), \text{can})$  where  $\mathcal{P}(R)$  is the category of finitely generated projective  $R$ -modules and  $\text{quis}$  is the set of quasi-isomorphisms.

**Theorem 4.1** ([Schd]). *Let  $R$  be a commutative ring, then*

- (1)  $GW^{[0]}(R)$  is the  $K$ -theory space  $GW(R)$  of the category of non-degenerate symmetric bilinear forms over  $R$ ,
- (2)  $GW^{[2]}(R)$  is the  $K$ -theory space  $KSp(R)$  of the category of non-degenerate symplectic forms over  $R$ , and
- (3)  $GW^{[4]}(R)$  is the  $K$ -theory space  $KQ(R)$  of the category of non-degenerate quadratic forms over  $R$ .

In particular, by [Schb, Theorem 6.6, Example 3.11 and Remark 2.19] we have

$$\begin{aligned} GW^{[0]}(\mathbb{Z}) &= GW(\mathbb{Z}) \simeq \mathbb{Z} \times \mathbb{Z} \times BO_{\infty, \infty}(\mathbb{Z})^+, \\ GW^{[2]}(\mathbb{Z}) &= KSp(\mathbb{Z}) \simeq \mathbb{Z} \times BSp(\mathbb{Z})^+, \\ GW^{[4]}(\mathbb{Z}) &= KQ(\mathbb{Z}) \simeq \mathbb{Z} \times \mathbb{Z} \times BO_{\infty}(\mathbb{Z})^+. \end{aligned}$$

**Theorem 4.2** ([Scha]). *Let  $R$  be a Dedekind domain and  $S \subset R$  a multiplicative set of non-zero divisors. Then there is a natural homotopy fibration*

$$\bigoplus_{\mathfrak{p} \cap S \neq \emptyset} GW^{[-1]}(R/\mathfrak{p}) \rightarrow GW^{[0]}(R) \rightarrow GW^{[0]}(S^{-1}R).$$

Recall that Friedlander [Fri76] shows that  $K_n Sp(\mathbb{F}_2)$  is a finite group of odd order for  $n \geq 1$ . In particular its 2-adic completion  $K_n Sp(\mathbb{F}_2)_2^\wedge = 0$  for  $n \geq 1$ . Since the same is true for  $K(\mathbb{F}_2)$ , we obtain  $GW_n(\mathbb{F}_2)_2^\wedge = 0$  for  $n \geq 1$ ,  $GW_n^{[\pm 1]}(\mathbb{F}_2)_2^\wedge = 0$  for  $n \geq 0$  and the following from Theorems 4.1, 4.2 and the homotopy fibration (4.1).

**Theorem 4.3.** *Let  $\mathbb{Z}^! = \mathbb{Z}[1/2]$  then the ring homomorphism  $\mathbb{Z} \rightarrow \mathbb{Z}^!$  induces isomorphisms after 2-adic completion*

$$\begin{aligned} K_n Sp(\mathbb{Z})_2^\wedge &\cong K_n Sp(\mathbb{Z}^!)_2^\wedge, & n \geq 0, \\ GW_n(\mathbb{Z})_2^\wedge &\cong GW_n(\mathbb{Z}^!)_2^\wedge, & n \geq 1, \\ KQ_n(\mathbb{Z})_2^\wedge &\cong KQ_n(\mathbb{Z}^!)_2^\wedge, & n \geq 2. \end{aligned}$$

Finally, the 2-adic homotopy groups of  $K\mathrm{Sp}(\mathbb{Z}')$  and  $GW(\mathbb{Z}') = KQ(\mathbb{Z}')$  can be found in [Kar05, 4.7.2]. This proves the theorems in Section 2 apart from the following which was needed in the proof of Lemma 3.1.

**Lemma 4.4.** *Let  $R$  be the ring of integers in a number field  $F$ . Then the inclusion  $R \subset F$  induces an isomorphism*

$$L_i^{[r]}(R)[1/2] \simeq L_i^{[r]}(F)[1/2], \quad i \geq r.$$

*Proof.* It suffices to prove the case  $r = 0$  since  $L_i^{[r]} = L_{i-r}^{[0]}$ . From Theorem 4.2 we deduce the homotopy fibration of  $-1$ -connected spectra

$$\bigoplus_{\mathfrak{p} \neq (0)} GW^{[-1]}(R/\mathfrak{p})[1/2] \rightarrow GW^{[0]}(R)[1/2] \rightarrow GW^{[0]}(F)[1/2]$$

in which the right horizontal map is also surjective on  $\pi_0$ , by the computations in [MH73]. Using the analogous statement for  $K$ -theory, we obtain the homotopy fibration of spectra

$$\bigoplus_{\mathfrak{p} \neq (0)} L^{[-1]}(R/\mathfrak{p})[1/2] \rightarrow L^{[0]}(R)[1/2] \rightarrow L^{[0]}(F)[1/2].$$

The left term in that fibration is trivial since for a finite field  $\mathbb{F}_q$ , we have

$$L^{[-1]}(\mathbb{F}_q)[1/2] \simeq 0.$$

This is well-known for  $q$  odd, and for  $q$  even,  $L^{[-1]}(\mathbb{F}_q)$  is a module spectrum over  $L^{[0]}(\mathbb{F}_2)$  whose homotopy groups are 2-primary torsion since on  $\pi_0$  it is

$$L_0^{[0]}(\mathbb{F}_2) = W(\mathbb{F}_2) = \mathbb{Z}/2.$$

□

## REFERENCES

- [Bal01] Paul Balmer. Triangular Witt groups. II. From usual to derived. *Math. Z.*, 236(2):351–382, 2001.
- [Fri76] Eric M. Friedlander. Computations of  $K$ -theories of finite fields. *Topology*, 15(1):87–109, 1976.
- [Kar05] Max Karoubi. Bott periodicity in topological, algebraic and Hermitian  $K$ -theory. In *Handbook of K-theory. Vol. 1, 2*, pages 111–137. Springer, Berlin, 2005.
- [MH73] John Milnor and Dale Husemoller. *Symmetric bilinear forms*. Springer-Verlag, New York-Heidelberg, 1973. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 73*.
- [Rog00] John Rognes.  $K_4(\mathbf{Z})$  is the trivial group. *Topology*, 39(2):267–281, 2000.
- [Sch10] Marco Schlichting. The Mayer-Vietoris principle for Grothendieck-Witt groups of schemes. *Invent. Math.*, 179(2):349–433, 2010.
- [Sch17] Marco Schlichting. Hermitian  $K$ -theory, derived equivalences and Karoubi’s fundamental theorem. *J. Pure Appl. Algebra*, 221(7):1729–1844, 2017.
- [Schb] Marco Schlichting. Higher  $K$ -theory of forms I. From rings to exact categories. Accepted for publication in *J. Inst. Math. Jussieu*. DOI: 10.1017/S1474748019000410.
- [Schc] Marco Schlichting. Higher  $K$ -theory of forms II. From exact categories to chain complexes. *In preparation*.
- [Schd] Marco Schlichting. Higher  $K$ -theory of forms III. From chain complexes to derived categories. *In preparation*.
- [Scha] Marco Schlichting. Higher  $K$ -theory of forms for Dedekind domains. *In preparation*.
- [Wei05] Charles Weibel. Algebraic  $K$ -theory of rings of integers in local and global fields. In *Handbook of K-theory. Vol. 1, 2*, pages 139–190. Springer, Berlin, 2005.

MARCO SCHLICHTING, MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK,  
COVENTRY CV4 7AL, UK  
*E-mail address:* `m.schlichting@warwick.ac.uk`