

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/127112>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

[Click here to view linked References](#)

# Is robotic right colectomy economically sustainable? A multicentre retrospective comparative study and cost analysis.

**Running Head:** Robotic right colectomy economic sustainability

*Authors: Giovanni Merola<sup>1</sup> MD, Antonio Sciuto<sup>2,3</sup>MD, Felice Pirozzi<sup>2</sup>MD, Jacopo Andreuccetti<sup>4</sup>MD, Giusto Pignata<sup>4,5</sup>MD, Francesco Corcione<sup>1</sup>MD, Marco Milone<sup>1</sup>MD PhD, Giovanni Domenico De Palma<sup>1</sup> Prof, Rossana Castaldo<sup>8</sup>MD PhD, Leandro Pecchia<sup>8</sup> Prof, Graziano Ceccarelli<sup>6,7</sup>MD and Umberto Bracale<sup>1</sup> Prof.*

1. Department of Gastroenterology, Endocrinology and Surgical Endoscopy, University Hospital Federico II of Naples, Italy
2. Department of General Surgery, Santa Maria delle Grazie Hospital, Pozzuoli, Italy
3. Department of Abdominal Surgery, Casa Sollievo della Sofferenza Research Hospital, San Giovanni Rotondo, Italy
4. Department of General and Minimally Invasive Surgery, San Camillo Hospital of Trento, Trento Italy
5. Department of General Surgery II, Spedali Civili, Brescia, Italy
6. Department of Surgery, Division of General Surgery, San Donato Hospital, Arezzo, Italy
7. Department of Surgery, Division of General Surgery San Matteo Hospital, Spoleto, Italy
8. School of Engineering, University of Warwick, Coventry, CV4 7AL, UK.

## Corresponding Author:

Name: Umberto  
Surname: Bracale  
Address: Via Pansini 5, Naples, 80131, Italy  
Email: [umbertobracale@gmail.com](mailto:umbertobracale@gmail.com)  
Phone: +39 339 5720072  
Fax: +39 081 7462640

## ORCID:

Giovanni Merola:	0000-0002-3237-677X
Antonio Sciuto:	0000-0003-3234-4580
Felice Pirozzi:	0000-0002-6385-5394
Jacopo Andreuccetti:	0000-0003-0474-9775
Giusto Pignata:	0000-0002-2381-1055
Francesco Corcione:	0000-0002-7157-5157
Marco Milone:	0000-0001-5612-1926
Giovanni Domenico De Palma:	0000-0001-7474-7180
Rossana Castaldo:	0000-0003-4143-1678
Leandro Pecchia:	0000-0002-7900-5415
Graziano Ceccarelli:	0000-0002-7859-6815
Umberto Bracale:	0000-0002-9868-0889

The study was conceived, funded, and executed entirely by authors' institutions. There was no industry support of any kind. The authors have no conflicts of interest.

## Abstract

### Background

Following the Food and Drug Administration approval, robot-assisted colorectal surgery has gained more acceptance among surgeons. One of the open issues about robotic surgery is the economic sustainability. The aim of our study is to evaluate the economic sustainability of robotic as compared to laparoscopic right colectomy for the Italian National Health System.

### Methods

We performed a retrospective multicentre case-matched study including 94 patients for each group from four different Italian surgical departments. An economic evaluation gathered from a real world data was performed to assess the sustainability of the robotic approach for right colectomy in the Italian National Health System.

In particular, a differential cost analysis between the two procedures was performed.

### Results

No statistical differences were found between the two groups for postoperative outcomes. After a careful review of the literature on the cost assessment for the operative room, medical devices and hospital stay according with our data, we estimated the followings: (a) the mean operative room cost for robotic group was  $2179 \pm 476$  € vs.  $1376 \pm 322$  € for laparoscopic group; (b) the mean hospital stay cost for robotic group was  $3143 \pm 1435$  € vs.  $3292 \pm 1123$  € for laparoscopic group; (c) the mean cost for instruments was 6280 € for robotic group vs. 1504 € for laparoscopic group. The total mean cost of robotic right colectomy was  $11,576 \pm 1915$  € vs.  $6196 \pm 1444$  € for laparoscopic right colectomy.

### Conclusion

In conclusion, to date, robotic right colectomy with intracorporeal anastomosis does not provide any significant clinical advantages, which may justify the additional costs, as compared to its laparoscopic counterpart. Further evolution of robotic technology and experience may lead to a reduction of costs, especially if the robotic platform is used in an appropriate healthcare setting.

**Keywords:** Robotic surgery, Right Colectomy, Cost Analysis, Laparoscopic surgery, Real World Data.

## Introduction

In the last decades, surgical technology has gone through a great evolution, especially in the minimally invasive field. Robot-assisted surgery represents a new trend in minimally invasive surgery and surgical literature.

Nowadays, the minimally invasive approach represents the gold standard for many surgical procedures, for both benign and malignant diseases [1]. Although technically demanding, minimally invasive right colectomy with intracorporeal anastomosis may be the preferred approach for right colon cancer treatment. Several advantages including reduced short-term morbidity and faster recovery have been associated to intracorporeal as compared to extracorporeal anastomosis [2, 3].

Following the Food and Drug Administration approval, robot-assisted colorectal surgery has gained more acceptance among surgeons [4, 5]. In the field of colon surgery, the robotic approach has shown no significant benefits for patients as compared to the laparoscopic approach, but some advantages have been demonstrated for surgeons, including a better view of the operative field and a better instrument for precision and easiness of movement [6, 7]. Thus, robotic assistance could enhance the ability of the surgeon to perform a totally minimally invasive right colectomy with intracorporeal anastomosis.

One of the critical issues about robotic surgery is the economic sustainability [6]. In fact, despite the advantages for the surgeon and the possibility to easily tutor more surgeons due to the dual console, further investigation is warranted when considering routine use of the robotic platform for colectomies [8].

In our multicentre study, we performed a retrospective case-matched analysis of 94 patients who underwent robot-assisted right colectomy (RRC) compared with a group of patients who underwent laparoscopic right colectomy (LRC). The main aim of our study is to evaluate the economic sustainability of RRC as compared to LRC for the Italian National Health System.

## Materials and Methods

After the ethical committee approval, all consecutive patients who underwent elective LRC or RRC for cancer from January 2012 to August 2017 at four Italian surgical departments (*Department of General and Minimally Invasive Surgery, San Camillo Hospital, Trento; Department of Surgical Specialities and Nephrology, University of Naples Federico II; Department of Surgery, Division of General Surgery, Hospital of Arezzo; Department of Abdominal Surgery, Casa Sollievo della Sofferenza Research Hospital, San Giovanni Rotondo*), were included in the study. Using prospective colorectal surgery unit databases, 275 patients were identified.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Among these, 159 patients underwent LRC and 116 had RRC. In order to reduce the risk of bias due to the patient selection in the robotic group, we performed a case-matched analysis using the Mantel-Haenszel method. The two groups were matched according to their biometric features. After matching, a total of 188 patients were analysed, namely 94 patients in each group.

Preoperative (age, sex, BMI, previous abdominal surgery, ASA score and tumour location), intraoperative (operative time, complications, conversion rate) and postoperative (pathologic stage according to Wittekind et al. [9], length of specimen, number of harvested lymph nodes, postoperative return of bowel function, hospital stay, complications according to the Clavien-Dindo classification [10], reoperation and mortality rates) data were reviewed.

Conversion was defined as the unplanned change from laparoscopy to the open procedure or from the robotic approach to either the laparoscopic or the open approach. Anastomotic leakage was considered along with all conditions with clinical or radiologic features of anastomotic dehiscence in accordance with the UK Surgical Infection Study Group [11]. We considered operative time as the time from the first skin incision until the last scar was sutured.

All procedures in both groups were carried out by four highly experienced laparoscopic colorectal surgeons [12]. All surgeons had completed the learning curve of robotic colorectal surgery before the inclusion of cases in this series [13].

Follow-up was conducted by a clinical examination at 7, 30, 180 days and then each year after surgery.

#### *Cost Analysis*

The differential cost analysis was conducted from an Italian hospital perspective, including medical costs (use of operating room, length of stay and medical devices) only, as performed in previous studies [14, 15], and from a National Health System (NHS)/Health care payer perspective.

According to the existing literature [16], the differential cost analysis is widely used for healthcare decisions. In particular, differential cost analysis was used to assess the difference between the cost of two alternative decisions (RRC vs. LRC). The concept is usually used when there are two or multiple possible options to pursue, and a choice must be made to select one option and drop the others.

1 The costing search was based on the economic assessment methodology adopted in Health Technology  
2 Assessment and was performed by two clinicians (UB and GM) and two biomedical engineers (RC and LP).

3  
4 **Postoperative costs were also included in the cost analysis.**

5  
6 To assess the cost of the two procedures from the hospital perspective, we considered the mean operative time,  
7  
8 mean length of stay and mean cost of medical devices in both groups. According to our previous study, we  
9  
10 estimated that the cost of theatre per minute and the cost of length of stay per day were 10.7 € and 780 €,  
11  
12 respectively [14]. **The cost of hospital stay also included the cost of both daily drugs and health staff.** We  
13  
14 calculated the mean costs for medical devices (e.g., trocar) from the National Health Price List (NHPL) and as  
15  
16 reported by Ho et al. [17].  
17

18  
19 To evaluate the differential costs from the NHS perspective, we estimated that the mean reimbursement for  
20  
21 right colectomy for cancer was  $6838 \pm 900$  €, which was calculated performing the average reimbursement of  
22  
23 Diagnosis Related Group (DRG) codes for each Italian region. These results did not consider the fixed  
24  
25 maintenance costs of equipment, which are extremely variable in structure and must be distributed on the total  
26  
27 amount of procedures performed.  
28

### 29 *Statistical Analysis*

30  
31 The statistical analysis was carried out using IBM SPSS Statistics 25. Continuous data were expressed as mean  
32  
33  $\pm$  standard deviation (SD). To compare continuous variables, an independent sample T-test was performed.  
34  
35 The Chi-square test was employed to analyse categorical data. Logistic regression was performed in order to  
36  
37 understand if some factors could influence operative time or length of stay. Regarding non-parametric variables  
38  
39 (such as time to flatus and length of stay), we used a Mann-Whitney U test.  
40  
41

42  
43 All the results are presented herein as 2-tailed values with statistical significance if the *p*-values were below  
44  
45 0.05.  
46

### 47 *Perioperative Management*

48  
49 **Mechanical bowel preparation was not administered, and no diet restriction was applied before surgery.**  
50  
51 **Intravenous antibiotic prophylaxis was given to all patients.** Antiemetics were administered regularly for 72  
52  
53 hours postoperatively. On the first postoperative day the patient could drink, if tolerated, and a normal diet was  
54  
55 offered from day 2 onwards. Early mobilization and low molecular weight heparin were used for deep venous  
56  
57 thrombosis prophylaxis. Pain management was achieved by a peridural catheter and administration of  
58  
59  
60  
61  
62  
63  
64  
65

1 nonsteroidal anti-inflammatory drugs, if required. Discharge criteria included tolerance of oral intake, absence  
2 of nausea or vomiting, return of bowel function, absence of abdominal distention, no evidence of  
3 complications, adequate mobility, patient acceptance [14]. When a postoperative complication required  
4 reoperation, a minimally invasive approach was preferred [18].  
5  
6

### 7 *Surgical Technique*

#### 8 *LRC*

9  
10 The patient was placed supine in the Trendelenburg position (30°) and right flank rotation. Pneumoperitoneum  
11 was established with the Veress technique [19]. We used the same technique described in our previous study  
12 [12]. After abdominal exploration, in order to exclude the presence of metastases, the ileo-colic vessels were  
13 ligated and divided. A medial-to-lateral dissection was conducted between the Toldt's fascia and the Gerota's  
14 fascia. If present, the right colic vessels were isolated and divided. Once the transverse colon was pulled up,  
15 its mesentery was dissected from the root and the right branches of the middle colic vessels were identified  
16 and divided. Then, we mobilized the right colon by dissecting the gastro-colic and parieto-colic ligaments.  
17 Complete division of the ileal mesentery allowed for full mobilization of the right colon. The transverse colon  
18 and terminal ileum were transected intracorporeally by a blue- and white-cartridge linear stapler, respectively.  
19 A side-to-side isoperistaltic ileo-colic anastomosis was performed by a 45-mm linear stapler with a blue  
20 cartridge. The enterotomy was closed with a double layer running suture [12, 20]. The mesentery was closed  
21 by absorbable sutures or fibrin glue. The specimen was retrieved through a Pfannenstiel incision or umbilical  
22 scar enlargement [21].  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

#### 41 *RRC*

42 For the RRC, we performed the same technique described for the laparoscopic approach. Trocars' position is  
43 reported by Spinoglio et al. [22]. In the first 35 robotic cases, a Da Vinci Si surgical system was used, while a  
44 DaVinci Xi platform was used in the subsequent cases.  
45  
46  
47  
48  
49  
50

### 51 **Results**

52 We performed a case-matched analysis comparing two homogeneous group of 94 patients each. Patient  
53 features are reported in Table 1.  
54  
55  
56

57 Operative time was significantly longer for RRC than LRC ( $135.5 \pm 33.9$  min in the LRC group vs.  $207.5 \pm$   
58  $44.9$  min in the RRC group;  $p < 0.05$ ). As reported in Table 2, no differences were recorded in terms of intra-  
59  
60  
61  
62  
63  
64  
65

1  
2 or postoperative complications and length of stay ( $4.2 \pm 1.4$  days in the LRC group vs.  $4.0 \pm 1.8$  days in the  
3 RRC group,  $p = 0.475$ ) between the two groups. Three cases in the RRC group required conversion to open  
4 surgery compared to none in the laparoscopic group ( $p = 0.080$ ). They were all due to advanced colon cancer.

5  
6 As reported in Table 2, only 3 Clavien-Dindo III complications were recorded in each group. In particular,  
7  
8 there were 2 anastomotic bleedings requiring endoscopic management and one anastomotic leakage in both  
9  
10 groups.

11  
12 At the 30-day follow-up, there were neither reinterventions nor deaths, and only one readmission was recorded  
13  
14 in the LRC group due to heart failure. At a mean follow-up of  $51.7 \pm 12.6$  months, 5 deaths were recorded,  
15  
16 two in the RRC group and three in the LRC group. Logistic regression did not show any association between  
17  
18 operative time or length of stay and the other variables analysed, including age, BMI, time to flatus and  
19  
20 pathologic stage.  
21

22  
23 After a careful review of the literature on the cost assessment for operative room, medical devices and hospital  
24  
25 stay according with our data, we estimated the followings: (a) the mean operative room cost for the RRC group  
26  
27 was  $2179 \pm 476$  € vs.  $1376 \pm 322$  € for the LRC group; (b) the mean hospital stay cost for the RRC group was  
28  
29  $3143 \pm 1435$  € vs.  $3292 \pm 1123$  € for the LRC group; (c) the mean cost for instruments was 6280 € for the RRC  
30  
31 group vs. 1504 € for the LRC group. The total mean cost of RRC was  $11,576 \pm 1915$  € vs.  $6196 \pm 1444$  € for  
32  
33 LRC. Data from cost analysis are reported in Table 3.  
34  
35  
36

## 37 38 Discussion

39  
40 Over the last years, there has been a rapid increase in the use of the robotic approach to colorectal surgery,  
41  
42 although the real benefits over laparoscopy in terms of outcomes still remain unclear [23, 24]. Thus, the  
43  
44 adoption of robotic surgery must be examined against a backdrop of burdening an already expensive healthcare  
45  
46 system [25].

47  
48 In our evaluation, we found that postoperative short-term outcomes of RRC and LRC performed with  
49  
50 intracorporeal anastomosis are similar. To date, a few studies included intracorporeal anastomosis when  
51  
52 comparing RRC and LRC. A recent retrospective multicentre analysis of 389 patients did not find any  
53  
54 difference in terms of time-to-first flatus, postoperative complications and hospital stay, although RRC was  
55  
56 associated with a lower 90-day readmission rate. In this series, LRC group comprised only 84 of 389 cases,  
57  
58 and data on costs of both procedures are not presented [23]. The unbalanced number between the two groups  
59  
60  
61  
62  
63  
64  
65



1 is consistent with the results of a recent meta-analysis, which found a higher rate of intracorporeal anastomosis  
2 in the robotic group [26]. Indeed, the robotic platform decreases the difficulty of intracorporeal suturing [27].  
3  
4 Since intracorporeal anastomosis is associated with faster recovery and lower morbidity as compared to  
5  
6 extracorporeal anastomosis, the use of the robotic approach aiming to increase the number of intracorporeal  
7  
8 anastomosis may be advisable [23]. Moreover, for surgeons early in their career, robotic assistance may shorten  
9  
10 the learning curve of minimally invasive right colectomy as compared to the laparoscopic approach [24].

11  
12 In agreement with published studies, operative time was significantly longer in the RRC group also in this  
13  
14 cohort. This could be due to the docking time, which may prolong the duration of the robotic procedure [23].

15  
16 The learning curve effect has also been reported to play a role in increasing operative time. As in most  
17  
18 published series, all surgeons involved in our study were more highly experienced in laparoscopic rather than  
19  
20 robotic colorectal surgery, and this may have partially affected length of surgical time. However, they had  
21  
22 completed the learning curve of robotic colorectal surgery at the beginning of the study. Although not analysed  
23  
24 in our series, operative time may decrease over time, as the number of performed procedures increases. In a  
25  
26 series of 101 procedures, a statistically significant difference was found between the earlier and the later robotic  
27  
28 series (329 vs. 266 min). The authors also reported a significant reduction in the conversion rate, claiming a  
29  
30 benefit in performing both dissection and intracorporeal anastomosis by robotic assistance [22]. This was not  
31  
32 confirmed in the present study, where a significantly higher conversion rate was observed in the RRC group  
33  
34 than in the LRC group. A more meticulous dissection of tissues has also been suggested as a potential reason  
35  
36 for the increased duration of the robotic procedure [23].  
37  
38  
39  
40  
41

42 Pathologic outcomes were comparable between RRC and LRC. Some authors suggested that the number of  
43  
44 harvested lymph nodes may be higher with robotic assistance [23]. Moreover, RRC appears to achieve similar  
45  
46 long-term survival rates as compared with LRC [28].

47  
48 Robot-assisted procedures are increasing since laparoscopic surgeons are incorporating robotics into their  
49  
50 normal practice [25]. Experienced laparoscopic surgeons may be able to overcome – at least partly – the  
51  
52 technical drawbacks of conventional laparoscopy and be equally proficient at both procedures [24]. In the  
53  
54 future, it might be expected that a better robotic experience will show considerable clinical advantages of RRC  
55  
56 compared to LRC.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Although the costs of RRC exceed those of LRC, data are still based on a limited number of procedures  
2 available from different health systems [26, 27]. In a meta-analysis of five studies, Solaini et al. found that the  
3 mean total cost was US \$10,335 for LRC vs. US \$12,299 for RRC (SMD – 0.52, 95% CI – 1 to – 0.04,  $p =$   
4 0.035). Moreover, based on the results of three studies, surgery-only related costs were higher in the robotic  
5 group (\$5953 vs. \$3930, SMD – 2.8, 95% CI – 5.53 to 0.02,  $p = 0.051$ ) [26]. Indeed, the higher costs of RRC  
6 are primarily attributable to surgery, including consumables and longer operative times [28].  
7  
8  
9

10 A retrospective analysis of the 2012-2014 Healthcare Cost and Utilization Project-National Inpatient Sample  
11 (NIS) was recently conducted in the United States. In a cohort of 7685 colectomies, RRC was 20.1% more  
12 costly than LRC ( $\$15,027 \pm 6049$  vs.  $\$12,516 \pm 5281$ ), even if the length of stay was  $4.3 \pm 2$  in the laparoscopic  
13 group versus  $3.8 \pm 1.6$  in the robotic group. This may suggest that, to date, the additional costs related to robotic  
14 procedures are not compensated by the savings from shorter hospital stay. However, these costs may be lower  
15 in centres with higher volume cases, where more experienced surgeons and staffs can optimize the operating  
16 room time. Studies using costs from single hospital systems may be more informative for a detailed cost  
17 analysis as compared to national database analysis. In fact, the latter provides total charges for each admission  
18 but not the actual cost, and details of costs related to supplies and instruments are not reported [29]. Similarly,  
19 a matched analysis of 1066 patients from the Premier Hospital Database found significantly higher hospital  
20 costs and longer operative times in the robotic group, while length of stay was similar for both surgical  
21 approaches [8].  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 A review of a hospital cost-accounting database including 111 patients who underwent elective minimally  
41 invasive colectomy (18 robotic surgeons) over a 1-year period showed that the cost of robotic colectomy was  
42 53% greater than laparoscopic colectomy ( $\$7806$  vs.  $\$5096$ ,  $p < 0.001$ ). No significant difference in overall  
43 costs between different surgeons was demonstrated despite varied training, experience levels and operative  
44 techniques. This demonstrates that total costs are relatively institutionally fixed and minimally influenced by  
45 variations in individual surgeon practice preferences [30]. There are no data about cost-analysis of RRC from  
46 the Italian healthcare system. The present evaluation from Italian centers found that RRC was nearly 70%  
47 more costly than LRC, and this increased cost was primary attributable to consumables, while the costs for  
48 hospital stay were similar. Furthermore, the initial acquisition cost, depreciation, and service contract for the  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 robotic and laparoscopic systems should be also considered [31], and they were not included in the present  
2 analysis.

3  
4 A previous cost-analysis comparing laparoscopic versus open surgery for colorectal cancer in three Italian  
5 high-volume centres, found that the national DRG tariff is insufficient to remunerate the providers' activity,  
6  
7 irrespective of the type of disease and surgical technique adopted [32]. This issue is even more compelling  
8  
9 given the growing interest for robotic colorectal surgery. In our series, the mean reimbursement for right  
10 colectomy was 50% of total estimated costs for RRC.  
11  
12

13  
14 Our study includes comparison of equivalent surgical procedures in a quite large sample size population from  
15 different centres, with adequate follow-up. Although it is a retrospective study, a case-matched analysis has  
16  
17 been performed in order to reduce the risk of bias due to the selection of patients in the robotic group.  
18  
19

20  
21  
22 Some limitations should be considered. There is no full accounting for the costs of postoperative complications.  
23  
24 However, the length of the hospital stay itself may be an indirect indicator of the increased cost of complicated  
25  
26 patients. Furthermore, the incidence of complications – especially those requiring additional procedures and/or  
27  
28 ICU management – was low and comparable between the two groups (3 vs. 3). Therefore, we considered that  
29  
30 they did not significantly affect the differential cost analysis between the two procedures. Enhanced recovery  
31  
32 after surgery (ERAS) protocols have been increasingly applied to colorectal surgery over the last years. As  
33  
34 reported by many studies, such protocols seem to reduce drastically the economic impact of colorectal surgery,  
35  
36 although the breakdowns of the costs and statistical methods are ambiguous and inconsistent among studies  
37  
38 and institutions [33]. A structured enhanced recovery pathway has not been used in our cohort. Thus, the costs  
39  
40 associated with implementing an ERAS program have not been evaluated. The perioperative care included  
41  
42 only some items of the ERAS protocol and it has not significantly changed over the study period. Therefore,  
43  
44 it did not impact on both length of stay and costs. According to the ERAS principles, the patients in this cohort  
45  
46 did not receive preoperative MBP, although more recent evidence suggests that combination of MBP and oral  
47  
48 antibiotics can reduce the risk of surgical site infections [33].  
49  
50

51  
52  
53 In conclusion, to date, RRC with intracorporeal anastomosis does not provide tangible benefits to the patients,  
54  
55 which may justify the additional costs as compared to its laparoscopic counterpart. However, at a time when  
56  
57 taking care of optimal resource utilization is essential, every effort should be made to support the development  
58  
59 of robotic surgery, in the hope of developing more clinically effective tools and skills. Indeed, further  
60  
61  
62  
63  
64  
65

1 development of robotics may lead to reduction of costs and improvement of outcomes, as occurred in the past  
2 for laparoscopic surgery. Moreover, although it is beyond the evaluations of our paper, the robotic approach  
3  
4 may be particularly effective to reduce the learning curve for young surgeons [35]. The advantages of robotic  
5  
6 surgery can be better explored and developed in any institutions with additional resources and research  
7  
8  
9 mandate. Further prospective studies comparing both surgical approaches with cost analysis are needed to  
10  
11 draw robust conclusions.

## 14 Disclosure

16 Giovanni Merola: no conflict of interest

18 Antonio Sciuto: no conflict of interest

20 Felice Pirozzi: no conflict of interest

22 Jacopo Andreuccetti: no conflict of interest,

24 Giusto Pignata: no conflict of interest

26 Francesco Corcione: no conflict of interest

28 Marco Milone: no conflict of interest

30 Giovanni Domenico De Palma: no conflict of interest

32 Rossana Castaldo: no conflict of interest

34 Leandro Pecchia: no conflict of interest

36 Graziano Ceccarelli: no conflict of interest

38 Umberto Bracale: no conflict of interest

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Bracale, U., et al., *Single access laparoscopic left hemicolectomy with or without inferior mesenteric artery preservation: our preliminary experience*. *Minerva Chir*, 2013. **68**(3): p. 315-20.
2. Bracale, U., et al., *Reply to Early colon cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up*. *ESMO Open*, 2016. **1**(6).
3. Milone, M., et al., *Intracorporeal versus extracorporeal anastomosis. Results from a multicentre comparative study on 512 right-sided colorectal cancers*. *Surg Endosc*, 2015. **29**(8): p. 2314-20.
4. Casciola, L., et al., *[Laparoscopic right hemicolectomy with intracorporeal anastomosis. Technical aspects and personal experience]*. *Minerva Chir*, 2003. **58**(4): p. 621-7.
5. Damle, A., et al., *Diffusion of technology: Trends in robotic-assisted colorectal surgery*. *Am J Surg*, 2017. **214**(5): p. 820-824.
6. Fung, A.K. and E.H. Aly, *Robotic colonic surgery: is it advisable to commence a new learning curve?* *Dis Colon Rectum*, 2013. **56**(6): p. 786-96.
7. Ballantyne, G.H. and F. Moll, *The da Vinci telerobotic surgical system: the virtual operative field and telepresence surgery*. *Surg Clin North Am*, 2003. **83**(6): p. 1293-304, vii.
8. Davis, B.R., et al., *Robotic-assisted versus laparoscopic colectomy: cost and clinical outcomes*. *JSLS*, 2014. **18**(2): p. 211-24.
9. Wittekind, C. and B. Oberschmid, *[TNM classification of malignant tumors 2010: General aspects and amendments in the general section]*. *Pathologe*, 2010. **31**(5): p. 333-4, 336-8.
10. Clavien, P.A., et al., *The Clavien-Dindo classification of surgical complications: five-year experience*. *Ann Surg*, 2009. **250**(2): p. 187-96.
11. Sciuto, A., et al., *Predictive factors for anastomotic leakage after laparoscopic colorectal surgery*. *World J Gastroenterol*, 2018. **24**(21): p. 2247-2260.
12. Bracale, U., et al., *The Use of Barbed Suture for Intracorporeal Mechanical Anastomosis During a Totally Laparoscopic Right Colectomy: Is It Safe? A Retrospective Nonrandomized Comparative Multicenter Study*. *Surg Innov*, 2018. **25**(3): p. 267-273.
13. Shaw, D.D., et al., *Robotic Colorectal Surgery Learning Curve and Case Complexity*. *J Laparoendosc Adv Surg Tech A*, 2018. **28**(10): p. 1163-1168.
14. Bracale, U., et al., *Single-access laparoscopic rectal resection versus the multiport technique: a retrospective study with cost analysis*. *Surg Innov*, 2015. **22**(1): p. 46-53.
15. Bracale, U., et al., *Beneficial effects of fibrin glue (Quixil) versus Lichtenstein conventional technique in inguinal hernia repair: a randomized clinical trial*. *Hernia*, 2014. **18**(2): p. 185-192.
16. Rayburn, L. and J. Rayburn, *Using the differential costing concept for healthcare decisions*. *Healthcare computing & communications*, 1984. **1**(11): p. 26.
17. Edelman, D.S. and J.P. Hodde, *Update on Bioactive Prosthetic Material for the Treatment of Hernias*. *Surg Technol Int*, 2011. **21**: p. 135-9.
18. Cuccurullo, D., et al., *Relaparoscopy for management of postoperative complications following colorectal surgery: ten years experience in a single center*. *Surg Endosc*, 2015. **29**(7): p. 1795-803.
19. Milone, M., et al., *Colorectal resection in deep pelvic endometriosis: Surgical technique and post-operative complications*. *World J Gastroenterol*, 2015. **21**(47): p. 13345-51.
20. Reggio, S., et al., *Single-layer versus double-layer closure of the enterotomy in laparoscopic right hemicolectomy with intracorporeal anastomosis: a single-center study*. *Tech Coloproctol*, 2015. **19**(12): p. 745-50.
21. Roscio, F., et al., *Totally laparoscopic versus laparoscopic assisted right colectomy for cancer*. *Int J Surg*, 2012. **10**(6): p. 290-5.
22. Spinoglio, G., et al., *Robotic Versus Laparoscopic Right Colectomy with Complete Mesocolic Excision for the Treatment of Colon Cancer: Perioperative Outcomes and 5-Year Survival in a Consecutive Series of 202 Patients*. *Ann Surg Oncol*, 2018. **25**(12): p. 3580-3586.
23. Solaini, L., et al., *Robotic versus laparoscopic right colectomy with intracorporeal anastomosis: a multicenter comparative analysis on short-term outcomes*. *Surg Endosc*, 2018.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

24. de'Angelis, N., et al., *Laparoscopic extended right colectomy versus laparoscopic left colectomy for carcinoma of the splenic flexure: a matched case-control study*. *Int J Colorectal Dis*, 2016. **31**(3): p. 623-30.

25. Armijo, P.R., et al., *Growth in robotic-assisted procedures is from conversion of laparoscopic procedures and not from open surgeons' conversion: a study of trends and costs*. *Surg Endosc*, 2018. **32**(4): p. 2106-2113.

26. Solaini, L., et al., *Robotic versus laparoscopic right colectomy: an updated systematic review and meta-analysis*. *Surg Endosc*, 2018. **32**(3): p. 1104-1110.

27. Petrucciani, N., et al., *Robotic right colectomy: A worthwhile procedure? Results of a meta-analysis of trials comparing robotic versus laparoscopic right colectomy*. *J Minim Access Surg*, 2015. **11**(1): p. 22-8.

28. Park, J.S., et al., *Randomized clinical trial of robot-assisted versus standard laparoscopic right colectomy*. *Br J Surg*, 2012. **99**(9): p. 1219-26.

29. Khorgami, Z., et al., *The cost of robotics: an analysis of the added costs of robotic-assisted versus laparoscopic surgery using the National Inpatient Sample*. *Surg Endosc*, 2018.

30. Tillou, J., et al., *The impact of surgeon choices on costs associated with uncomplicated minimally invasive colectomy: you are not as important as you think*. *Gastroenterol Rep (Oxf)*, 2018. **6**(2): p. 108-113.

31. Higgins, R.M., et al., *Cost analysis of robotic versus laparoscopic general surgery procedures*. *Surg Endosc*, 2017. **31**(1): p. 185-192.

32. Berto, P., et al., *Cost of laparoscopy and laparotomy in the surgical treatment of colorectal cancer*. *Surg Endosc*, 2012. **26**(5): p. 1444-53.

33. Lemanu, D.P., et al., *A systematic review to assess cost effectiveness of enhanced recovery after surgery programmes in colorectal surgery*. *Colorectal Dis*, 2014. **16**(5): p. 338-46.

34. Toh, J.W.T., et al., *Association of Mechanical Bowel Preparation and Oral Antibiotics Before Elective Colorectal Surgery With Surgical Site Infection: A Network Meta-analysis*. *JAMA Netw Open*, 2018. **1**(6): p. e183226.

35. de'Angelis, N., et al., *Robotic Versus Laparoscopic Right Colectomy for Colon Cancer: Analysis of the Initial Simultaneous Learning Curve of a Surgical Fellow*. *J Laparoendosc Adv Surg Tech A*, 2016. **26**(11): p. 882-892.

**Table 1.** Patient characteristics

	<b>All patients (n=188)</b>	<b>Laparoscopic group (n=94)</b>	<b>Robotic group (n=94)</b>	<b>p-value</b>
<b>Age</b> years, mean±SD <sup>a</sup>	70.75±10 (range 30-88)	72.09±9.54	69.41±10.31	0.066
<b>Sex (M/F)</b> n	121/67	61/33	60/34	0.879
<b>BMI<sup>b</sup></b> Kg/m <sup>2</sup> , mean±SD	27.45±5.21 (range 16.5-42)	27.97±5.73	26.94±4.61	0.178
<b>Previous abdominal surgery, n</b>	32	18	14	0.438
<b>ASA<sup>c</sup> score, n</b>				0.474
<b>I</b>	18	11	7	
<b>II</b>	101	52	49	
<b>III</b>	68	31	37	
<b>IV</b>	1	0	1	
<b>Cancer location, n</b>				0.813
<b>Caecum</b>	56	27	29	
<b>Ascending colon</b>	112	58	54	
<b>Hepatic flexure</b>	20	9	11	
<b>Stage<sup>d</sup>, n</b>				0.560
<b>I</b>	23	13	10	
<b>II</b>	108	56	52	
<b>III</b>	54	23	31	
<b>IV</b>	3	2	1	

<sup>a</sup>Standard deviation; <sup>b</sup>Body mass index; <sup>c</sup>American Society of Anesthesiologists; <sup>d</sup>according to Wittekind et al. [9]

**Table 2.** Intra- and postoperative data

	<b>All patients (n=188)</b>	<b>Laparoscopic group (n=94)</b>	<b>Robotic group (n=94)</b>	<b>p-value</b>
<b>Operative time</b> min, mean±SD <sup>a</sup>	171.47±53.64 [range 85-330]	135.46±33.86	207.47±44.93	<0.05
<b>Length of stay</b> days, median±IQR <sup>b</sup>	4±2 [range 4-25]	4±2	4±2	0.475
<b>Intraoperative complications</b> n, mean±SD	0	0	0	1
<b>Passage of flatus</b> days, mean±SD	2.21±1.01 [range 1-7]	2.20±1.24	2.23±0.7	0.178
<b>Harvested lymph nodes</b> n, mean±SD	22.12±4.95 [range 12-58]	22.32±3.81	21.91±5.88	0.567
<b>Specimen length</b> cm, mean±SD	38.43±4.69 [range 27-64]	38.92±4.75	37.94±4.61	0.154
<b>Postoperative complications<sup>c</sup>, n</b>	32	15	17	0.784
<b>Grade I</b>	14	7	7	
<b>Grade II</b>	11	4	7	
<b>Grade III</b>	6	3	3	
<b>Grade IV</b>	0	0	0	
<b>Grade V</b>	1	1	0	
<b>Conversion (n)</b>	3	0	3	0.08

<sup>a</sup>Standard deviation; <sup>b</sup>Interquartile range; <sup>c</sup>According to the Clavien-Dindo classification [10]



**Table 3.** Cost-analysis<sup>a</sup>

	<b>Robotic group (n=94)</b>	<b>Laparoscopic group (n=94)</b>	<b>Difference</b>	<b><i>p</i>-value</b>
<b>Cost of instruments</b>	6280 €	1504 €	4776 €	
<b>Cost of operative room</b>	2179 ± 476 €	1376 ± 322 €	803 ± 575 €	<0.005
<b>Cost of length of stay</b>	3143 ± 1435 €	3292 ± 1123 €	-146 ± 315 €	0.475
<b>Final cost</b>	11,576 ± 1915€	6196 ± 1444 €	5380 ± 471 €	<0.005

<sup>a</sup>Numbers are represented as mean ± standard deviation (SD)

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.



# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

# Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting [http://www.adobe.com/go/reader\\_download](http://www.adobe.com/go/reader_download).

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.