

**Title:** Gastrointestinal peptides and small bowel hypomotility are possible causes for fasting and postprandial symptoms in active Crohn's disease.

**Authors:** Asseel Khalaf, Caroline L. Hoad, Alex Menys, Adam Nowak, Shellie Radford, Stuart A. Taylor, Khalid Latief, Melanie Lingaya, Yirga Falcone, Gulzar Singh, Robin C. Spiller, Penny A. Gowland, Luca Marciani and Gordon W. Moran

**Names for Pubmed Indexing:** Khalaf, Hoad, Menys, Nowak, Radford, Taylor, Latief, Lingaya, Falcone, Singh, Spiller, Gowland, Marciani, Moran.

**AFFILIATIONS:**

Nottingham Digestive Diseases Centre, School of Medicine, University of Nottingham UK (AK, AN, SR, ML, YF, GZ, RCS, LM, GWM).

National Institute of Health Research Nottingham Biomedical Research Centre at Nottingham University Hospitals NHS Trust and the University of Nottingham (AK, CLH, AN, SR, ML, YF, GZ, RCS, PAG, LM, GWM).

Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham UK (CLH, PAG)

Motilent Ltd, London (AM)

Centre for Medical Imaging, Division of Medicine, UCL UK (SAT)

Department of Radiology, Nottingham University Hospitals (KL)

New affiliation: Radiologic Sciences, Allied Health Sciences, Kuwait University, Kuwait (AK)

**DISCLAIMERS:**

Dr Moran has received educational support from Abbvie, Janssen, NAPP, Takeda Pharmaceuticals, Merck Sharp & Dohme Ltd, Ferring and Dr Falk. He has received speaker honoraria from Merck Sharp & Dohme Ltd, Abbvie, Janssen, Ferring and Takeda Pharmaceuticals. He attended advisory boards for Abbvie, Takeda Pharmaceuticals, Janssen, Medtronic, Phebra Pharmaceuticals, Servertus Associates Ltd and Dr Falk.

Professor Stuart Taylor is a research consultant for Robarts Clinical Trials.

Professor Robin Spiller has received research funding from Norgine and Zespri. He has also acted as a consultant on advisory boards for Allergan, Commonwealth Diagnostics International, Napo Pharmaceuticals, Ipsen and Yuhan, and received speakers' fees from Menarini and Alfawasserman.

Dr Alex Menys is the CEO of Motilent, a medical imaging analysis company.

**CORRESPONDING AUTHOR:**

Gordon W. Moran

Clinical Associate Professor in Gastroenterology

NIHR Nottingham Biomedical Research Centre,

Nottingham University Hospitals NHS Trust & The University of Nottingham, Nottingham,  
United Kingdom

E-mail: Gordon.Moran@nottingham.ac.uk

Telephone no: +44 (0)115 9249924 ext 60608

**SOURCES OF SUPPORT:** This study was funded by a scholarship from the Kuwait University and from a research grant from the Nottingham University Hospitals' Charity

(PP.Gordon Moran.Nov14).

**SHORT RUNNING HEAD:** Motility in Crohn's disease

**Abbreviations**

Analysis of Variance (ANOVA); Area under the Curve (AUC); au (arbitrary units); Biomedical Research Centre (BRC); Body Mass Index (BMI); Cholecystokinin (CCK); Crohn's disease (CD); C-Reactive protein (CRP); Enteroendocrine cells (EC); Enzyme-linked immunosorbent assay (ELISA); gastrointestinal (GI); Glucagon-like peptide-1 (GLP-1); Healthy volunteer (HV); Interactive data language (IDL); Inflammatory Bowel Disease (IBD); Magnetic Resonance Imaging (MRI); Magnetic Resonance Index of Activity (MaRIA); National Institute of Health Research (NIHR); Polypeptide YY (PYY); Radioimmunoassay (RIA); Region of Interest (ROI); Small Bowel Water Content (SBWC); Sir Peter Mansfield Imaging Centre (SPMIC); Standard error of the Mean (SEM); Visual Analogue Scores (VAS).

**Clinical trial registry:** Clinical trials.gov NCT03052465.

1  
2  
3  
4  
5  
6  
7

## 8 ABSTRACT

9 **Background:** Crohn's disease (CD) patients suffer postprandial aversive symptoms which  
10 may lead to anorexia and malnutrition. Changes in the regulation of gut hormones and gut  
11 dysmotility are believed to play a role.

12 **Objective:** This study aims to investigate small bowel motility and gut peptides responses to a  
13 standard test meal in CD by using Magnetic Resonance Imaging (MRI).

14 **Design:** Fifteen CD patients with active disease (age  $36\pm 3$  years, BMI  $26\pm 1$  kg/m<sup>2</sup>) and 20  
15 healthy volunteers (HV) (age  $31\pm 3$  years; body mass index, BMI  $24\pm 1$  kg/m<sup>2</sup>) were studied.  
16 They underwent baseline and postprandial MRI scans, symptom questionnaires, and blood  
17 sampling following a 400g soup meal (204 kcal). Small bowel motility and other MRI  
18 parameters and glucagon-like peptide-1 (GLP-1), polypeptide YY (PYY), and cholecystokinin  
19 (CCK) peptides were measured. Data is presented as mean $\pm$ standard deviation of the mean  
20 (SEM).

21 **Results:** HV had significantly higher fasting motility index ( $106\pm 13$  a.u.) compared to CD  
22 participants ( $70\pm 8$  a.u.,  $p\leq 0.05$ ). Postprandial small bowel water content showed a significant  
23 time by group interaction ( $p<0.05$ ) with CD showing higher levels from  $t=210$  min  
24 postprandially. Fasting levels of GLP-1 and PYY were significantly greater in CD compared  
25 to healthy volunteers (GLP-1, CD  $50\pm 8$   $\mu$ g/mL versus HV  $13\pm 3$   $\mu$ g/mL,  $p\leq 0.0001$  and PYY,  
26 CD  $236\pm 16$  pg/mL versus HV  $118\pm 12$  pg/mL,  $p\leq 0.0001$ ). The meal challenge induced a  
27 significant postprandial increase in aversive symptoms scores (fullness, distention, bloating,  
28 abdominal pain and sickness) in CD participants compared to healthy volunteers ( $p\leq 0.05$ ).

29 **Conclusions:** The decrease in fasting small bowel motility noted in CD may be ascribed to the  
30 increased fasting gut peptides. A better understanding of the etiology of aversive symptoms in  
31 CD will facilitate identification of better therapeutic targets to improve nutritional status.

32 **KEYWORDS:** Gastrointestinal motility; gut peptides; Crohn's disease; anorexia

### 33 INTRODUCTION

34 Malnutrition is common in patients with Crohn's disease (CD). A recent systematic review  
35 reported a reduction in body mass index (BMI) in 37% of CD patients with a reduction in fat-  
36 free mass and fat mass in 28% and 31% respectively (1).

37 Multiple reasons could account for the malnutrition but an abnormal eating behaviour is  
38 common in CD, with up to 37% of patients showing abnormal eating patterns (2), with an  
39 associated significant reduction in protein intake. Ingestion of food plays a predominant role in  
40 symptom generation in Inflammatory Bowel Disease (IBD) patients. In a large prospective  
41 multi-centre study, 77% of patients avoided some foods to prevent disease relapse, with 86%  
42 of patients avoiding foods when the disease is activity to prevent worsening symptoms (3).

43 The aetiology for this is still unclear but postprandial dysmotility of the inflamed gut could be  
44 an aetiological factor. A delayed gastric emptying or attenuated small bowel motility in the  
45 fasting or postprandial state could be an important aetiological factor in the patient symptom  
46 generation and eventual adoption of a diverse eating behaviour to improve symptom control.  
47 Recent technological advances in image analysis have allowed us to use magnetic resonance  
48 imaging (MRI) to measure CD activity through intestinal motility (4). A significant negative  
49 correlation is observed between terminal ileal motility and histological (4), biochemical (5)  
50 and radiological measures of disease activity (4). The exact aetiology of this  
51 pathophysiological finding is still unclear.

52 Enteroendocrine cells (EC) are intraluminal nutrient sensors. They play a pivotal role in  
53 orchestrating physiological functions in the gastrointestinal tract. Exposure of the ileum to  
54 fatty acids decreases jejunal (6) and duodenal motility (7) through polypeptide YY (PYY)-  
55 mediated mechanisms. Glucagon-like peptide-1 (GLP-1) has been shown to decrease small

56 bowel motility (8, 9). Moreover, cholecystokinin (CCK) has been shown to delay gastric  
57 emptying by increasing pyloric motility (10, 11), with CCK modulation (12) affecting food  
58 intake underlying its anorectic role in eating behaviour. CD patients with active small bowel  
59 inflammation have significant up-regulation of EC cells with an increase in ileal expression of  
60 GLP-1 (13), PYY (2-fold) (14) and (CCK) (3-fold) (15) levels. This increase in plasma  
61 peptide levels is associated with the symptoms of nausea and anorexia, with both symptoms,  
62 and tissue and plasma EC-peptide expression decreasing to normality in remission (14). We  
63 hypothesise that an increase in CCK, GLP-1 or PYY-EC activity could potentially lead to  
64 symptoms of nausea through delayed gastric emptying and attenuated post-prandial small  
65 bowel motility. To our knowledge, this mechanistic link has never been described.  
66 We have shown that a soup test meal challenge is effective at inducing a change in multiple  
67 physiological quantified end-points and in monitoring markers of gastrointestinal motility in a  
68 single MRI study acceptable to participants (16, 17). Our developed technique for the  
69 assessment of small bowel motility showed excellent inter and intra observer agreement (17).  
70 We now aim to investigate fasting and postprandial changes in gut motility, intestinal  
71 physiology and related symptoms in a cohort of CD patients with active disease and healthy  
72 volunteers (HV) to investigate the aetiology of patient symptoms.

## 73 **SUBJECTS AND METHODS**

### 74 **Study population**

75 This is a single-centre open-label clinical trial conducted between November 2015 and  
76 February 2017 at the National Institute of Health Research Nottingham Biomedical Research  
77 Centre (NIHR Nottingham BRC). CD patients (18-75 years) with active disease were  
78 recruited from the Inflammatory Bowel Disease clinic at Nottingham University Hospitals

79 Trust. Active disease was defined as ulceration seen at ileocolonoscopy, intestinal  
80 inflammation or deep ulceration seen on computer tomography or MR enterography, with the  
81 disease activity quantified via the Magnetic Resonance Index of Activity (MaRIA) score of  
82  $>7$  (18) or faecal calprotectin of  $>250\mu\text{g/g}$  (19, 20) or C-Reactive protein (CRP)  $>5\text{mg/dl}$ .  
83 These measures of disease activity were to be quantified within four weeks of recruitment.  
84 Stable doses of immunosuppressive and biological agents were permitted. No changes in  
85 medication were allowed at inclusion until data collection of the outcome measures has been  
86 completed. All participants had a good command of the English language and had the  
87 capacity to give informed written consent.

88 Age-, body mass index- and gender-matched HV participants were recruited from an existing  
89 participant database in the NIHR Nottingham BRC and from the local healthy populations of  
90 Nottingham University Hospitals and the University of Nottingham. Potential participants  
91 with a history of inflammatory bowel disease, smokers (21), a history of bowel resections or  
92 any gastrointestinal surgery, history of pancreatic insufficiency, thyroid disease, diabetes,  
93 protein-pump inhibitor usage or any medication that affects gastric emptying or small bowel  
94 transit and any potential participants scoring very highly on the depression scale questionnaire  
95 were excluded. Standard MRI exclusion criteria were applied.

96 This study was approved by the Research Ethics Committee (NRES approval 15/EM/0003) of  
97 the Health Research Authority. This study is registered on clinical trials.gov with identifier  
98 NCT03052465. All participants gave informed written consent.

### 99 **Outcome measures**

100 The primary outcome of this study was to compare fasting and postprandial small bowel  
101 motility between CD and HV participants as measured through MRI. Secondary outcomes



102 were gall bladder contraction, gastric volumes, small bowel water content, plasma GLP-1,  
103 PYY and CCK, symptoms scores and MRI disease activity scores as quantified by the MaRIA  
104 score (18).

### 105 **Experimental protocol**

106 This study was open label. Participants were asked to fast from 2000 h the previous evening  
107 and to avoid alcohol, caffeine, strenuous exercise and any medication that could affect gut  
108 function for 18 h before the experiment. On the day of the scan, participants attended the 1.5T  
109 Philips Achieva MRI scanner (Philips Healthcare, Best, the Netherlands) at the Sir Peter  
110 Mansfield Imaging Centre (SPMIC) at the University Park Campus, University of  
111 Nottingham at 0800.

112 Participants underwent a baseline fasting scan (defined at  $t = -20$  min time point), together  
113 with a fasting baseline blood sample. They were then asked to consume all their test meal  
114 within a maximum time of 20 min. The participants then underwent a first immediate  
115 postprandial scan (defined as  $t = 0$  min).

116 This was followed with data collection (MRI, questionnaire data and 10 ml blood samples)  
117 time points every 15 min for the first 60 min and then every 30 min up to 270 min. At each  
118 time point, participants filled a 100mm Visual analogue scale (VAS) symptoms questionnaire  
119 scoring their feeling of fullness, bloating, distension, abdominal pain/discomfort and nausea  
120 (14). At the end of the 270 min participants were given a volume (750mls-1250mls) of oral  
121 contrast agent (2.5% Mannitol, 0.2 % locust bean gum) to drink (within 60 minutes) and a  
122 further MRI scan (within 30 minutes) was undertaken to quantify disease activity. See  
123 **supplementary figure 1.**

**124 Test meal**

125 The test meal consisted of: cream of chicken soup (400g) (or mushroom for vegetarians)  
126 (Heinz, Wigan, UK) (14, 22), The nutrient content of this meal/100g was: energy (kcal) 51,  
127 protein 1.5 g (1.5%), carbohydrate 4.7 g (4.5%), fat 2.9 g (2.9%). For a complete list of  
128 contents please see <https://www.heinz.co.uk/products/soup/ranges/classics/cream-of-chicken>.

**129 Magnetic Resonance Imaging**

130 Participants were scanned using a range of sequences (22). At each time point scans were  
131 acquired to assess gastric volume (23), gall bladder volume (24), small bowel water content  
132 (25) and small bowel motility (26).

133 Gastric emptying was assessed using a balanced gradient echo sequence to yield a good  
134 contrast between the stomach contents and other abdominal organs (23).

135 The content of apparent freely mobile water in the small bowel was assessed as previously  
136 described (25). This sequence yields high-intensity signals from areas with freely mobile fluid  
137 and dark signals from poorly mobile or bound water and all other body tissues.

138 Small bowel motility was assessed using a single slice cine-MRI acquisition set at six  
139 contiguous parallel coronal planes through the small bowel (22).

140 Gall bladder volume was measured pre- and post-prandial, at every acquisition time point  
141 using the same images as for the gastric volumes as previously shown (24). These functional  
142 MRI measures were acquired prior to the use of any anti-spasmodic agents administered  
143 during serial image acquisition.

**144 Plasma collection and peptides assays analysis**

145 Fasting 10 ml blood sample was drawn and collected. After the test meal, data were acquired

146 every 15 min for the first 60 min and every 30 min thereafter to 270 min. Plasma peptides  
147 (total GLP-1, total PYY) were analysed through enzyme-linked immunosorbent assay  
148 (ELISA) techniques (Millipore, UK) as previously shown (14). The concentrations of serum  
149 CCK were measured by radioimmunoassay (RIA) (Euro Diagnostic Products, Sweden) as  
150 previously shown (27).

### 151 **MRI measures of disease activity**

152 Small bowel was scanned before and 10 minutes after 40mg of hyoscine butyl bromide was  
153 injected intravenously to reduce small bowel motility. Participants were scanned within 30  
154 minutes. Initially a true fast imaging with a steady sequence was acquired in the coronal  
155 plane. Axial T1 sequences were acquired before and 70 s after intravenous administration of  
156 0.2 ml/kg body weight of gadolinium chelate (gadodiamide 0.5 mmol/l) at a rate of 2 ml/s.

### 157 **Data analysis**

#### 158 *Motility assessment*

159 All dynamic data was processed with Dual Registration of Abdominal Motion (Motilent,  
160 London, UK). Registration results were further analysed using a customised graphical user  
161 interface written in MATLAB (MathWorks, Natick, MA, USA) (17).

162 The technique used in this study quantified the motility of the bowel using the pixel signal  
163 changes through the time series, within a defined region of interest (ROI) placed over all  
164 visible small bowel loops (17) as previously described. This method utilizes changes in signal  
165 intensities that occur when the small bowel contents move between segments in regions  
166 showing bolus movement of contents as well as those showing more oscillatory motion, rather  
167 than looking for continuous motion throughout the time series. Motility measures are

168 presented as arbitrary units (a.u.) and are calculated as the mean across the total small bowel  
169 ROI.

170 *Small bowel water content (SBWC), Gall bladder and gastric volumes*

171 These parameters were quantified as previously validated (25), using in house software  
172 written in interactive data language (IDL) (Research Systems Inc. Boulder, Colorado, USA).

173 *Visual analogue scale (VAS)*

174 Symptoms regarding appetite, satiety and abdominal pain were scored at each time point  
175 using a previously validated questionnaire (14).

#### 176 **MARIA score**

177 MRE variables were evaluated by a Nottingham University Hospitals clinical gastrointestinal  
178 (GI) MRI radiologist with > 10 years' experience (Dr. Khalid Latief) in each segment  
179 including: bowel wall thickening, enhancement of the bowel wall after administration of  
180 intravenous contrast with gadolinium (relative contrast enhancement), presence of ulcers,  
181 mural oedema, regional enlarged lymph nodes (>10 mm), peri-enteric vascularization (comb  
182 sign), peri-enteric fluid, fat stranding, and fibro-fatty proliferation. The MaRIA score in each  
183 segment was calculated according to a formula, as previously defined (18).

#### 184 **Sample size and Statistical analyses**

185 Previous literature showed 2-fold higher plasma PYY in CD (area under the curve, AUC 22990  
186  $\pm$  5585) vs HV (10700  $\pm$  1886) and a higher plasma GLP-1 in CD (AUC 1027  $\pm$  220) vs HV  
187 (1347  $\pm$  350) (28). Assuming  $\alpha$  of 0.05, power of 80%, a maximum sample of 15 participants  
188 in each group was needed to show a difference. Similar comparisons for CCK showed  
189 significant differences between CD and HV in 10 participants/group with a strong correlation  
190 ( $r=0.6$ ) between mean CCK plasma concentration and gastric emptying half-life (15). Assuming

191  $\alpha$  of 0.05, power of 80% a total sample size of 19 participants was needed to show a difference.  
192 To allow for missing data (~10%), we aimed to recruit a maximum of 20 participants in each  
193 of the CD and HV cohorts.

194 The primary analysis was an across-group analysis with further sub-analyses undertaken if the  
195 primary comparison is significant. Normality of the data was assessed using Shapiro–Wilk’s  
196 test. Parametric data was presented as mean  $\pm$  standard error of the mean (SEM) or median  $\pm$   
197 interquartile range if non-parametric. All statistical analyses were performed using GraphPad  
198 Prism 7.01 (La Jolla, USA). A p-value less than 0.05 was considered statistically significant.  
199 Analyses of variance (ANOVA) was used to assess the significance of differences between  
200 and within each group with different time points. When the analysis of variance was  
201 significant, post hoc test assessments of the individual time points were performed using the  
202 Dunnett’s (for parametric data) or Dunn’s test (for nonparametric data) for within group  
203 comparison and Sidak’s test was used for between groups to account for multiple  
204 comparisons. The Pearson correlation coefficient was used to measure the strength of  
205 correlation between MaRIA scores and the different variables (measured as AUC).

## 206 **RESULTS**

### 207 **Participants**

208 Nineteen CD participants with active disease (age  $36\pm 3$  years, BMI  $26\pm 1$  kg/m<sup>2</sup>) as well as 20  
209 HV participants (age  $31\pm 3$  years, BMI  $24\pm 1$  kg/m<sup>2</sup>) were recruited. One CD participant was  
210 lost to follow-up and the data from two CD participants were excluded from the final  
211 analyses: one because of a low MaRIA score of 3.39 and another because of an unanalysable  
212 MRI data set. One other participant was excluded because of a high score on the hospital

213 anxiety and depression questionnaire. See supplementary figure 2, table 1 and supplementary  
214 table 1. CD participants had a mean disease duration of  $12.3 \pm 1.7$  years, mean HBI of  $6 \pm 1$ ,  
215 mean CRP of  $18.2 \pm 3.4$  mg/dl and mean fecal calprotectin of  $787.2 \pm 146.8$   $\mu$ g/g. All patients  
216 had ileal involvement. Four patients were being prescribed anti-tumor necrosis factor therapy,  
217 one patient vedolizumab, three patients immunomodulators, with the rest on no medication at  
218 the time of recruitment. Eight patients were surgically naïve.  
219 All HV participants and CD participants completed the study and tolerated the experimental  
220 procedures well without any adverse event.

#### 221 **Small bowel motility**

222 The HV participants started with a significantly higher ( $106 \pm 13$  a.u.,  $p < 0.05$ ) fasting motility  
223 index compared to CD participants ( $70 \pm 8$  a.u.) (**Figure 1**). No significant difference in the  
224 postprandial time by group interaction was observed in small bowel motility between the CD  
225 and HV groups.

#### 226 **Gallbladder volumes**

227 Two CD patients had a cholecystectomy prior to recruitment. No difference in the fasting gall  
228 bladder volumes were observed between the two groups. No significant difference in the  
229 postprandial time by group interaction was observed in gall bladder volumes between the CD  
230 and HV groups. The difference in gallbladder volumes between HV and CD participants from  
231 fasting to 150 min are shown in **Figure 2**.

#### 232 **Gastric volumes**

233 The baseline gastric volumes showed a small amount of fasting gastric secretions (**Figure 3**,

234 HV:  $29 \pm 5$  mL, CD:  $25 \pm 4$  mL) with no differences observed between the groups. No  
235 significant difference in the postprandial time by group interaction was observed in gastric  
236 volumes between the CD and HV groups. Gastric volume increased upon feeding (HV:  
237  $388 \pm 18$  mL, CD:  $324 \pm 26$  mL). The average time to empty half of the stomach contents ( $T_{1/2}$ )  
238 in HV and CD participants was  $43 \pm 4$  min,  $63 \pm 7.5$  min respectively, with no significant  
239 difference observed between the groups.

#### 240 **Small bowel water content**

241 The data in **Figure 4** shows a small amount of fasting small bowel water content in both  
242 groups (HV:  $44 \pm 6$  mL, CD:  $36 \pm 9$  mL). No difference in the fasting small bowel water content  
243 was visualised between groups. A significant increase ( $p < 0.05$ ) was seen in CD participants  
244 compared to HV (measured as area under the curve (AUC) CD:  $19778 \pm 2119$  mL/min, HV:  
245  $14197 \pm 1249$  mL/min). A significant difference in the time by group interaction was observed  
246 in small bowel water content between the CD and HV groups ( $p = 0.0352$ ). An increase in  
247 postprandial water volume in CD when compared to healthy volunteers was observed at  
248  $t = 210$  min ( $p = 0.0388$ ),  $t = 240$  min ( $p = 0.0168$ ) and  $270$  min ( $p = 0.0048$ ).

#### 249 **Total GLP-1**

250 **Figure 5** show higher ( $p < 0.0001$ ) fasting GLP-1 levels in CD participants compared to HV  
251 (CD:  $50 \pm 8$   $\mu\text{g/mL}$ , HV:  $13 \pm 3$   $\mu\text{g/mL}$ ). The test meal did not induce a significant postprandial  
252 response in either group. No significant difference in the time by group interaction was  
253 observed between HV and CD groups. Significantly higher ( $p < 0.0001$ ) postprandial GLP-1  
254 levels were reported in CD participants compared to HV (AUC CD:  $12293 \pm 1586$   $\mu\text{g/mL}$ , HV:  
255  $3317 \pm 762$   $\mu\text{g/mL}$ ).

256 **Total PYY**

257 The CD participants showed higher ( $236\pm 16$  pg/mL,  $p<0.0001$ ) fasting PYY plasma levels  
258 when compared to HV ( $118\pm 12$  pg/mL) (**Figure 6**). The test meal did not induce a significant  
259 postprandial response in either group. No significant difference in the time by group  
260 interaction was observed between HV and CD groups. CD participants exhibited a  
261 significantly higher postprandial PYY response compared to the HV (AUC CD:  $62782\pm 4313$   
262 pg /mL, HV:  $34744\pm 3169$  pg/mL,  $p<0.0001$ ).

263 **CCK**

264 No significant difference was seen in fasting and postprandial levels of CCK between the two  
265 groups (**Figure 7**). No significant difference in the time by group interaction was observed in  
266 postprandial CCK between the CD and HV groups. ~~In CD group, a significant difference~~  
267 ~~( $p<0.0001$ ) was seen across the different time points with a significant increase ( $p<0.05$ ) from~~  
268 ~~time 0 to 90 min compared to fasting plasma levels. Within HV group, plasma CCK levels~~  
269 ~~showed a significant increase ( $p<0.05$ ) immediately after feeding to 60 min in comparison to~~  
270 ~~fasting concentrations.~~

271 **Symptom VAS data**

272 Fasting and postprandial VAS scores recorded from CD participants and HV are shown in  
273 **Figure 8**. CD participants showed a significantly higher ( $p<0.01$ ) fasting fullness and  
274 abdominal pain scores compared to HV (CD:  $21\pm 6$  mm,  $18\pm 5$  mm HV:  $5\pm 3$  mm,  $0.5\pm 0.3$   
275 mm). CD participants also showed a significantly higher ( $p<0.05$ ) fasting distention scores  
276 compared to HV (CD:  $14\pm 5$  mm HV:  $2\pm 1$  mm).

277 No significant difference in the time by group interaction was observed in VAS between the



278 CD and HV groups. The meal induced a significantly ( $p<0.05$ ) higher postprandial fullness  
279 scores in CD participants compared to HV (AUC CD:  $6795\pm1440$  mm/min HV:  $2907\pm703$   
280 mm/min). A significantly higher postprandial ( $p<0.0001$ ) VAS scores of bloating, distention  
281 and abdominal pain were noted in CD participants compared to HV (AUC CD:  $5558\pm1293$   
282 mm/min,  $5071\pm1253$  mm/min,  $3187\pm873$  mm/min HV:  $565\pm257$  mm/min,  $303\pm191$  mm/min,  
283  $7\pm5$  mm/min). The CD participants showed a significantly ( $p<0.01$ ) higher sickness scores  
284 compared to HV healthy volunteers (AUC, CD:  $2024\pm927$  mm/min HV:  $75\pm75$  mm/min).

### 285 **MaRIA scores**

286 The mean value of the MaRIA score was  $20.3\pm1.9$  (see the supplementary table for individual  
287 scores). There was moderate non-significant correlation between fasting small bowel motility  
288 and disease activity as measured through MaRIA scores ( $r=0.52$  (95% confidence interval -  
289  $0.050, 0.8303$ ,  $p=0.07$ ). There was no significant correlation between MaRIA scores and  
290 postprandial small bowel motility, small water content and GLP-1, PYY or CCK levels.

### 291 **DISCUSSION**

292 The primary aim of this study was to understand the altered intestinal physiology and aversive  
293 patient symptoms observed in active CD in the fasted and the postprandial state. Any  
294 alterations might help better understand symptom generation that may be leading to altered  
295 eating behaviour and malnutrition.

296 Participants with CD had a lower fasting small bowel intestinal motility, with otherwise no  
297 difference observed in the postprandial phase when compared to HV. We have shown  
298 significantly higher fasting and postprandial levels for both GLP-1 and PYY with associated  
299 aversive symptoms being reported to a significantly greater extent in CD than in HV.

300 Although small bowel hypomotility, elevated plasma GI peptide levels and aversive patient  
301 symptoms could be interlinked in the fasting state; our data suggests that this hypothesis may  
302 not hold true in the postprandial phase. Possibly, symptom generation in the postprandial  
303 phase could be related to GI-peptide mediated alterations in the gut-brain axis rather than  
304 altered intestinal physiology. A decrease in MR motility in CD, in the prepared bowel, was  
305 previously described, with motility correlated to histological and biochemical measures of  
306 disease activity and patient symptoms (29). Global small bowel hypomotility involving  
307 normal-looking bowel has been described in CD, with motility variance negatively correlating  
308 to key patient symptoms like diarrhoea, pain and clinical symptom scores (29, 30).

309 The fasting and postprandial plasma levels of GLP-1 were significantly higher in CD. This  
310 was confirmed in previous CD studies (31). Similarly, plasma fasting and postprandial PYY  
311 levels were significantly elevated in CD group as we previously described (14, 31). The  
312 higher fasting GLP-1 and PYY observed in CD could be the cause of the fasting small bowel  
313 hypomotility observed in CD. We observed no difference in the fasting and postprandial  
314 plasma CCK levels, although the observed postprandial CCK response was only significant in  
315 the CD rather than the healthy group despite the absolute levels not being significant in  
316 between groups. The CD location in this recruited cohort was predominantly ileal or  
317 ileocolonic rather than proximal in the duodenum where the CCK-secreting I cells are located.

318 Although earlier data in IBD murine models suggested that EC upregulation occurs  
319 irrespective of the anatomical location of intestinal inflammation (32), this observation was  
320 later refuted in CD (13). This might explain the lack of difference in CCK concentration  
321 between both groups.

322 No significant difference was seen in the fasting and postprandial gall bladder volumes in CD

323 compared to healthy volunteers. The lack of difference in CCK levels between the two groups  
324 explains the lack of difference in the gall bladder volumes as CCK is integral for the gall  
325 bladder contraction in response to a fat stimulus. Additionally, previous data showed a similar  
326 non-significant fasting gall bladder volumes in both healthy volunteers and patients with  
327 small bowel CD (33, 34).

328 Similarly to postprandial small bowel motility, a delay in gastric emptying was expected in  
329 the CD cohort with no difference in gastric emptying observed between CD and HV in this  
330 study. Both PYY and GLP-1 may delay gastric emptying (35). In a previous study, delayed  
331 gastric emptying was observed in an inflammatory bowel disease cohort with similar plasma  
332 CCK and elevated plasma GLP-1 to healthy participants (31). In that study, gastric emptying  
333 was measured through a breath test so the findings are not comparable to our study.

334 Postprandial small bowel water content typically showed two peaks. The first peak is the  
335 gastric phase and represents emptying of the gastric contents in the proximal small bowel. The  
336 second peak is called the intestinal phase (36) which represents the increase in small bowel  
337 water content due pancreatico-biliary and enterocyte secretion. A bimodal postprandial peak  
338 was also seen in the small bowel water content in CD group, with a significant difference in  
339 time by group interaction in postprandial small bowel water content between CD and HV. The  
340 noted increase could be due to the significant postprandial CCK response in the CD group  
341 which leads to an increase in bile acid production and pancreatic secretion. Additionally, the  
342 meal may have acted as an osmotic stimulus causing the increase in the small bowel water  
343 content seen in the intestinal phase in CD cohort (37). Moreover, the observed increase in  
344 small bowel water content in CD group might have induced an increase in small bowel  
345 motility by stimulating smooth muscle contraction due to the increase in small bowel

346 distention thus nullifying any postprandial differences between groups. Additionally, the  
347 increase in small bowel distention might have made small bowel motility more easily  
348 quantifiable in this unprepared MRI visit hence decreasing any possible difference to healthy  
349 participants. None of the CD participants had predominant stricture which may have caused a  
350 delay in intestinal transit and might explain the increase in small bowel water content. Finally,  
351 the increase in measureable small bowel water content in CD might merely represent a delay  
352 in the distal small bowel emptying its contents in the ascending colon.

353 The CD group demonstrated significantly higher fasting and postprandial symptom VAS  
354 scores, with more predominant symptoms of fullness, distention and abdominal pain scores  
355 compared to HV. The additional small bowel distention from the increase in water content in  
356 CD might explain the significant difference in the measured symptoms. However, the actual  
357 difference in small bowel water content is small and accounts to only 40 mL which might not  
358 been enough to account for such a drastic difference in symptoms. Another possible mechanism  
359 for these exaggerated symptoms might be from the upregulated gut-brain axis which we have  
360 not investigated in this study.

361 There were possible limitations to this work. The test meal used was small in volume and  
362 nutrients and did not induce a significant change in postprandial GLP-1, PYY and CCK.  
363 However, as we had previously shown (14), it was well tolerated by all the patients and acted  
364 as a good stimulus to gastrointestinal physiological responses. Another limitation is the  
365 relatively small participant population, but such detailed phenotyping made recruiting a larger  
366 cohort for this exploratory study unfeasible. We as well assayed total PYY and GLP-1 rather  
367 than the active peptides. In vivo, active GLP-1 is rapidly degraded to GLP-1 (9-36). Similarly,

368 PYY (1-36) is metabolised to the active PYY (3-36). Both of these activities are undertaken  
369 through DPP-4 (38, 39). We have previously shown that DPP-4 expression is decreased in CD  
370 (40). In this present study, we used a protease inhibitor to further minimise DPP4 activity and  
371 peptide degradation. In previous work we also have shown no significant difference between  
372 active and total GLP-1 in small bowel CD and HV (28). Similarly, total PYY, has been shown  
373 to have a temporal pattern similar to that of PYY (3–36) after meals (41). For these reasons, we  
374 assayed total rather than the active peptides.

375 In this work we successfully quantified fasting and postprandial small bowel motility and  
376 possibly ascribed a putative role for EC peptides in the aetiology of disordered intestinal  
377 motility and anorectic symptoms in CD. A better understanding of the role EC peptides in the  
378 altered eating behaviour and malnutrition has pharmacological relevance. EC peptide  
379 modulators (Exendin 9-39 (42) and dexloxiglumide (43)) are now available. Further work is  
380 now needed to deconstruct the gut-brain axis and possibly open a new therapeutic pathway in  
381 CD therapy, thus improving nutritional status, disease outcomes and quality of life.

## **ACKNOWLEDGMENTS**

We acknowledge the support of the National Institute of Health Research Nottingham Biomedical Research Centre in the conduct of this study.

## **CONFLICT OF INTEREST**

None

## **AUTHORS' CONTRIBUTION**

AK, CLH, LM, RCS, PAG, SAT and GWM designed research; AK, CLH, LM, AN, SR, ML,

YF, GS, GWM conducted research; AK, AM, CLH, ML, YF, GS and KL analyzed data or performed statistical analysis; AK, CLH, PAG, RCS, SAT, AM, LM, and GWM wrote the paper; GWM had primary responsibility for final content.

## REFERENCES

1. Bryant RV, Trott MJ, Bartholomeusz FD, Andrews JM. Systematic review: body composition in adults with inflammatory bowel disease. *Aliment Pharmacol Ther* 2013;38(3):213-25. doi: 10.1111/apt.12372.
2. Wardle RA, Thapaliya G, Nowak A, Radford S, Dalton M, Finlayson G, Moran GW. An Examination of Appetite and Disordered Eating in Active Crohn's Disease. *Journal of Crohn's & colitis* 2018;12(7):819-25. doi: 10.1093/ecco-jcc/jjy041.
3. Casanova MJ, Chaparro M, Molina B, Merino O, Batanero R, Duenas-Sadornil C, Robledo P, Garcia-Albert AM, Gomez-Sanchez MB, Calvet X, et al. Prevalence of Malnutrition and Nutritional Characteristics of Patients With Inflammatory Bowel Disease. *Journal of Crohns & Colitis* 2017;11(12):1430-9. doi: 10.1093/ecco-jcc/jjx102.
4. Menys A, Puylaert C, Tutein Nolthenius CE, Plumb AA, Makanyanga J, Tielbeek J, Pendse D, Brosens LA, Rodriguez-Justo M, Atkinson D, et al. Quantified Terminal Ileal Motility during MR Enterography as a Biomarker of Crohn Disease Activity: Prospective Multi-Institution Study. *Radiology* 2018;180100. doi: 10.1148/radiol.2018180100.
5. Bickelhaupt S, Pazahr S, Chuck N, Blume I, Froehlich JM, Cattin R, Raible S, Bouquet H, Bill U, Rogler G, et al. Crohn's disease: small bowel motility impairment correlates with inflammatory-related markers C-reactive protein and calprotectin. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society* 2013;25(6):467-73. doi: 10.1111/nmo.12088.
6. Spiller RC, Trotman IF, Adrian TE, Bloom SR, Misiewicz JJ, Silk DB. Further characterisation of the 'ileal brake' reflex in man--effect of ileal infusion of partial digests of fat, protein, and starch on jejunal motility and release of neurotensin, enteroglucagon, and peptide YY. *Gut* 1988;29(8):1042-51.
7. Wen J, Phillips SF, Sarr MG, Kost LJ, Holst JJ. PYY and GLP-1 contribute to feedback inhibition from the canine ileum and colon. *The American journal of physiology* 1995;269(6 Pt 1):G945-52.
8. Schirra J, Nicolaus M, Roggel R, Katschinski M, Storr M, Woerle HJ, Goke B. Endogenous glucagon-like peptide 1 controls endocrine pancreatic secretion and antro-pyloro-duodenal motility in humans. *Gut* 2006;55(2):243-51. doi: 10.1136/gut.2004.059741.

9. Hellstrom PM, Naslund E, Edholm T, Schmidt PT, Kristensen J, Theodorsson E, Holst JJ, Efendic S. GLP-1 suppresses gastrointestinal motility and inhibits the migrating motor complex in healthy subjects and patients with irritable bowel syndrome. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society* 2008;20(6):649-59. doi: 10.1111/j.1365-2982.2007.01079.x.
10. MacIntosh CG, Andrews JM, Jones KL, Wishart JM, Morris HA, Jansen JB, Morley JE, Horowitz M, Chapman IM. Effects of age on concentrations of plasma cholecystokinin, glucagon-like peptide 1, and peptide YY and their relation to appetite and pyloric motility. *The American journal of clinical nutrition* 1999;69(5):999-1006. doi: 10.1093/ajcn/69.5.999.
11. MacIntosh CG, Morley JE, Wishart J, Morris H, Jansen JB, Horowitz M, Chapman IM. Effect of exogenous cholecystokinin (CCK)-8 on food intake and plasma CCK, leptin, and insulin concentrations in older and young adults: evidence for increased CCK activity as a cause of the anorexia of aging. *The Journal of clinical endocrinology and metabolism* 2001;86(12):5830-7. doi: 10.1210/jcem.86.12.8107.
12. Beglinger C, Degen L, Matzinger D, D'Amato M, Drewe J. Loxiglumide, a CCK-A receptor antagonist, stimulates calorie intake and hunger feelings in humans. *American journal of physiology Regulatory, integrative and comparative physiology* 2001;280(4):R1149-54. doi: 10.1152/ajpregu.2001.280.4.R1149.
13. Moran GW, Pennock J, McLaughlin JT. Enteroendocrine cells in terminal ileal Crohn's disease. *J Crohns Colitis* 2012;6(9):871-80. doi: 10.1016/j.crohns.2012.01.013.
14. Moran GW, Leslie FC, McLaughlin JT. Crohn's disease affecting the small bowel is associated with reduced appetite and elevated levels of circulating gut peptides. *Clin Nutr* 2013;32(3):404-11. doi: 10.1016/j.clnu.2012.08.024.
15. Keller J, Beglinger C, Holst JJ, Andresen V, Layer P. Mechanisms of gastric emptying disturbances in chronic and acute inflammation of the distal gastrointestinal tract. *American journal of physiology Gastrointestinal and liver physiology* 2009;297(5):G861-8. doi: 10.1152/ajpgi.00145.2009.
16. Khalaf A, Hoad CL, Menys A, Nowak A, Taylor SA, Paparo S, Lingaya M, Falcone Y, Singh G, Spiller RC, et al. MRI assessment of the postprandial gastrointestinal motility and peptide response in healthy humans. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society* 2018;30(1). doi: 10.1111/nmo.13182.
17. Khalaf A, Nowak A, Menys A, Marciani L, Taylor SA, Spiller RC, Gowland PA, Moran GW, Hoad CL. Cine MRI assessment of motility in the unprepared small bowel in the fasting and fed state: Beyond the breath-hold. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society* 2019;31(1):e13466. doi: 10.1111/nmo.13466.
18. Rimola J, Rodriguez S, Garcia-Bosch O, Ordas I, Ayala E, Aceituno M, Pellise

- M, Ayuso C, Ricart E, Donoso L, et al. Magnetic resonance for assessment of disease activity and severity in ileocolonic Crohn's disease. *Gut* 2009;58(8):1113-20. doi: 10.1136/gut.2008.167957.
19. Schoepfer AM, Beglinger C, Straumann A, Trummeler M, Vavricka SR, Bruegger LE, Seibold F. Fecal Calprotectin Correlates More Closely With the Simple Endoscopic Score for Crohn's Disease (SES-CD) than CRP, Blood Leukocytes, and the CDAI. *Am J Gastroenterol* 2010;105(1):162-9. doi: 10.1038/ajg.2009.545.
  20. Turvill J. Mapping of Crohn's disease outcomes to faecal calprotectin levels in patients maintained on biologic therapy. *Frontline gastroenterology* 2014;5(3):167-75. doi: 10.1136/flgastro-2014-100441.
  21. Kadota K, Takeshima F, Inoue K, Takamori K, Yoshioka S, Nakayama S, Abe K, Mizuta Y, Kohno S, Ozono Y. Effects of smoking cessation on gastric emptying in smokers. *Journal of clinical gastroenterology* 2010;44(4):e71-e5. doi: 10.1097/MCG.0b013e3181be9a0f.
  22. Khalaf A, Hoad CL, Menys A, Nowak A, Taylor SA, Paparo S, Lingaya M, Falcone Y, Singh G, Spiller RC, et al. MRI assessment of the postprandial gastrointestinal motility and peptide response in healthy humans. *Neurogastroenterol Motil* 2018;30(1). doi: 10.1111/nmo.13182.
  23. Hoad CL, Parker H, Hudders N, Costigan C, Cox EF, Perkins AC, Blackshaw PE, Marciani L, Spiller RC, Fox MR, et al. Measurement of gastric meal and secretion volumes using magnetic resonance imaging. *Phys Med Biol* 2015;60(3):1367-83. doi: 10.1088/0031-9155/60/3/1367.
  24. Marciani L, Cox EF, Hoad CL, Totman JJ, Costigan C, Singh G, Shepherd V, Chalkley L, Robinson M, Ison R, et al. Effects of various food ingredients on gall bladder emptying. *Eur J Clin Nutr* 2013;67(11):1182-7. doi: 10.1038/ejcn.2013.168.
  25. Hoad CL, Marciani L, Foley S, Totman JJ, Wright J, Bush D, Cox EF, Campbell E, Spiller RC, Gowland PA. Non-invasive quantification of small bowel water content by MRI: a validation study. *Phys Med Biol* 2007;52(23):6909-22. doi: 10.1088/0031-9155/52/23/009|issn 0031-9155.
  26. Menys A, Taylor SA, Emmanuel A, Ahmed A, Plumb AA, Odille F, Alam A, Halligan S, Atkinson D. Global Small Bowel Motility: Assessment with Dynamic MR Imaging. *Radiology* 2013;269(2):442-9. doi: 10.1148/radiol.13130151.
  27. Astbury NM, Taylor MA, Macdonald IA. Breakfast consumption affects appetite, energy intake, and the metabolic and endocrine responses to foods consumed later in the day in male habitual breakfast eaters. *The Journal of nutrition* 2011;141(7):1381-9. doi: 10.3945/jn.110.128645.
  28. Moran GW, Leslie FC, McLaughlin JT. Crohn's disease affecting the small bowel is associated with reduced appetite and elevated levels of circulating gut peptides. *Clinical nutrition (Edinburgh, Scotland)* 2013;32(3):404-11. doi:



- 10.1016/j.clnu.2012.08.024.
29. Gollifer RM, Menys A, Makanyanga J, Puylaert CAJ, Vos FM, Stoker J, Atkinson D, Taylor SA. Relationship between MRI quantified small bowel motility and abdominal symptoms in Crohn's disease patients-a validation study. *Br J Radiol* 2018;91(1089). doi: 10.1259/bjr.20170914.
  30. Menys A, Makanyanga J, Plumb A, Bhatnagar G, Atkinson D, Emmanuel A, Taylor SA. Aberrant Motility in Unaffected Small Bowel is Linked to Inflammatory Burden and Patient Symptoms in Crohn's Disease. *Inflamm Bowel Dis* 2016;22(2):424-32. doi: 10.1097/mib.0000000000000601.
  31. Keller J, Binnewies U, Rosch M, Juul Holst J, Beglinger C, Andresen V, Layer P. Gastric emptying and disease activity in inflammatory bowel disease. *European journal of clinical investigation* 2015;45(12):1234-42. doi: 10.1111/eci.12542.
  32. O'Hara JR, Lomax AE, Mawe GM, Sharkey KA. Ileitis alters neuronal and enteroendocrine signalling in guinea pig distal colon. *Gut* 2007;56(2):186-94. doi: 10.1136/gut.2006.102780.
  33. Murray FE, McNicholas M, Stack W, O'Donoghue DP. Impaired fatty-meal-stimulated gallbladder contractility in patients with Crohn's disease. *Clinical science* 1992;83(6):689-93.
  34. Vu MK, Gielkens HA, van Hogezaand RA, van Oostayen JA, Lamers CB, Masclee AA. Gallbladder motility in Crohn disease: influence of disease localization and bowel resection. *Scand J Gastroentero* 2000;35(11):1157-62.
  35. Marathe CS, Rayner CK, Jones KL, Horowitz M. Relationships Between Gastric Emptying, Postprandial Glycemia, and Incretin Hormones. *Diabetes Care* 2013;36(5):1396-405. doi: 10.2337/dc12-1609.
  36. Marciani L, Pritchard SE, Hellier-Woods C, Costigan C, Hoad CL, Gowland PA, Spiller RC. Delayed gastric emptying and reduced postprandial small bowel water content of equicaloric whole meal bread versus rice meals in healthy subjects: novel MRI insights. *Eur J Clin Nutr* 2013;67(7):754-8. doi: 10.1038/ejcn.2013.78.
  37. Kellum JM, Albuquerque FC, Stoner MC, Harris RP. Stroking human jejunal mucosa induces 5-HT release and Cl<sup>-</sup> secretion via afferent neurons and 5-HT<sub>4</sub> receptors. *The American journal of physiology* 1999;277(3 Pt 1):G515-20.
  38. Deacon CF, Nauck MA, Toft-Nielsen M, Pridal L, Willms B, Holst JJ. Both subcutaneously and intravenously administered glucagon-like peptide I are rapidly degraded from the NH<sub>2</sub>-terminus in type II diabetic patients and in healthy subjects. *Diabetes* 1995;44(9):1126-31. doi: 10.2337/diab.44.9.1126.
  39. Grandt D, Schimiczek M, Beglinger C, Layer P, Goebell H, Eysselein VE, Reeve JR, Jr. Two molecular forms of peptide YY (PYY) are abundant in human blood: characterization of a radioimmunoassay recognizing PYY 1-36 and PYY 3-36. *Regulatory peptides* 1994;51(2):151-9.
  40. Moran GW, O'Neill C, Padfield P, McLaughlin JT. Dipeptidyl peptidase-4

- expression is reduced in Crohn's disease. *Regulatory peptides* 2012;177(1-3):40-5. doi: 10.1016/j.regpep.2012.04.006.
41. Batterham RL, Heffron H, Kapoor S, Chivers JE, Chandarana K, Herzog H, Le Roux CW, Thomas EL, Bell JD, Withers DJ. Critical role for peptide YY in protein-mediated satiation and body-weight regulation. *Cell metabolism* 2006;4(3):223-33. doi: 10.1016/j.cmet.2006.08.001.
  42. Serre V, Dolci W, Schaerer E, Scrocchi L, Drucker D, Efrat S, Thorens B. Exendin-(9-39) is an inverse agonist of the murine glucagon-like peptide-1 receptor: implications for basal intracellular cyclic adenosine 3',5'-monophosphate levels and beta-cell glucose competence. *Endocrinology* 1998;139(11):4448-54. doi: 10.1210/endo.139.11.6295.
  43. Lassman DJ, McKie S, Gregory LJ, Lal S, D'Amato M, Steele I, Varro A, Dockray GJ, Williams SC, Thompson DG. Defining the role of cholecystokinin in the lipid-induced human brain activation matrix. *Gastroenterology* 2010;138(4):1514-24. doi: 10.1053/j.gastro.2009.12.060.

**Tables 1: Patient demographics**

	<b>HV participants</b>	<b>CD participants</b>
<b>Number</b>	20	15
<b>Age</b>	31±3 years	36±3 years
<b>Gender</b>	10F:10M	8F:8M
<b>BMI</b>	24±1 kg/m <sup>2</sup>	26±1 kg/m <sup>2</sup>
<b>Smoking Status</b>	Non-smokers	Non-smokers
<b>Mean CRP (SEM)</b>	N/A	18.2±3.4mg/dl
<b>Mean Faecal Calprotectin (SEM)</b>	N/A	787.2±146.8µg/g

**Figure legends:**

**Figure 1:** Time courses of the small bowel motility in Crohn's disease (CD) and in healthy volunteers. A. Significantly lower fasting small bowel motility in CD when compared to healthy volunteers. B. Time courses of the small bowel motility for CD and healthy volunteer. No significant difference in the time by group interaction was observed between CD and healthy volunteer groups. Data presented as mean  $\pm$  SEM (healthy volunteer: n=20, CD: n=15). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 2:** Time courses of the gallbladder volumes in Crohn's disease (CD) and in healthy volunteer groups. A. No significant difference in the fasting gall bladder volumes between CD and healthy volunteer groups. B. Time courses of the gall bladder volumes in CD and healthy volunteer groups. No significant difference in the time by group interaction was observed between CD and healthy volunteer groups. Data presented as mean  $\pm$  SEM (healthy volunteer: n=20, CD: n=13). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 3:** Time courses of the stomach content volumes in Crohn's disease (CD) and in healthy volunteers. A. No significant difference in the fasting stomach volumes between CD and healthy volunteers. B. Time courses of the stomach volumes for CD and healthy volunteers. No significant difference in the time by group interaction was observed between CD and healthy volunteer groups. Data presented as mean  $\pm$  SEM (healthy volunteers: n=20, CD: n=15). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 4:** Time courses of the small bowel water content in Crohn's disease (CD) and in healthy

volunteers. A. No significant difference in the fasting small bowel water content for CD and healthy volunteers. B. Time courses of the small bowel water content for CD and healthy volunteers showing a significant difference in time by group interaction ( $p=0.0352$ ) with an increase in postprandial water volume in CD when compared to healthy volunteers at  $t=210$  min ( $p=0.0388$ ),  $t=240$  min ( $p=0.0168$ ) and  $270$  min ( $p=0.0048$ ). Data presented as mean  $\pm$  SEM (healthy volunteers:  $n=20$ , CD:  $n=15$ ). ANOVA was used to assess the significance of differences between and within each group with different time points. Sidak's test was used for the assessments of the individual time points between groups.

**Figure 5:** Time courses of the GLP-1 concentrations in Crohn's disease (CD) and in healthy volunteers. A. Significantly higher fasting GLP-1 concentrations for CD when compared to healthy volunteers. B. Time courses of the GLP-1 concentrations for CD and healthy volunteers. No significant difference in the time by group interaction was observed between CD and healthy volunteer groups. CD participants exhibited a significantly higher postprandial GLP-1 response compared to healthy volunteers (AUC CD:  $12293 \pm 1586$   $\mu\text{g/mL}$ , healthy volunteers:  $3317 \pm 762$   $\mu\text{g/mL}$ ,  $p < 0.0001$ ). Data presented as mean  $\pm$  SEM (healthy volunteers:  $n=20$ , CD  $n=15$ ). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 6:** Time courses of the PYY concentrations in Crohn's disease (CD) and in healthy volunteers. A. Significantly higher fasting PYY concentrations in CD when compared to healthy volunteers. B. Time courses of the PYY concentrations for CD and healthy volunteers. No significant difference in the time by group interaction was observed between CD and healthy volunteer groups. CD participants exhibited a significantly higher postprandial PYY response compared to healthy volunteers (AUC CD:  $62782 \pm 4313$   $\text{pg/mL}$ ,

healthy volunteers:  $34744 \pm 3169$  pg/mL,  $p < 0.0001$ ). Data presented as mean  $\pm$  SEM (healthy volunteers:  $n=20$ , CD  $n=15$ ). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 7:** Time courses of the CCK concentrations in Crohn's disease (CD) and in healthy volunteers. A. No significant difference in fasting CCK concentrations between CD and healthy volunteers. B. Time courses of the CCK concentrations for CD and healthy volunteers. No significant time by group interaction observed. Data are mean  $\pm$  SEM (healthy volunteers:  $n=20$ , CD  $n=15$ ). ANOVA was used to assess the significance of differences between and within each group for different time points.

**Figure 8:** Time courses of the fasting and postprandial VAS scores in in Crohn's disease (CD) and in healthy volunteers. CD participants showed a significantly higher fasting fullness ( $p < 0.01$ ), abdominal pain ( $p < 0.01$ ) and distention scores ( $p < 0.05$ ) compared to healthy volunteers. Data presented as mean  $\pm$  SEM (HV:  $n=20$ , CD  $n=15$ ). ANOVA was used to assess the significance of differences between and within each group for different time points.


## **Supplementary Material**

**Gastrointestinal peptides and small bowel hypomotility are possible causes for fasting and postprandial symptoms in active Crohn's disease.**

**Asseel Khalaf, Caroline L. Hoad, Alex Menys, Adam Nowak, Shellie Radford, Stuart A. Taylor, Khalid Latief, Melanie Lingaya, Yirga Falcone, Gulzar Singh, Robin C. Spiller, Penny A. Gowland, Luca Marciani and Gordon W. Moran**

## Supplementary Material

Supplementary figure 1: Diagram of MIC2's study protocol.





**Gastrointestinal peptides and small bowel hypomotility are possible causes for fasting and postprandial symptoms in active Crohn's disease.**

Asseel Khalaf, Caroline L. Hoad, Alex Menys, Adam Nowak, Shellie Radford, Stuart A. Taylor, Khalid Latief, Melanie Lingaya, Yirga Falcone, Gulzar Singh, Robin C. Spiller, Penny A. Gowland, Luca Marciani and Gordon W. Moran

Supplementary table 1

	Age years	Sex	Disease Duration years	Montreal Classification	Con Meds	Surgical History	MaRIA	HBI	CRP mg/dl	fC µg/g
1	24	F	12	A1L1B2	Adalimumab	Right hemicolectomy	16.10	2	41	933
2	29	F	4	A2L1B2	Adalimumab	No	11.37	2	7	177
3	39	F	14	A2L1B2	Adalimumab	No	16.72	13	<5	1435
4	57	M	8	A3L3B1	Methotrexate	No	27.48	1	15	393
5	20	F	3	A2L1B1	Nil	No	7.00	6	<5	275
6	52	F	25	A2L1B1p	Nil	Right hemicolectomy	27.07	7	<5	107
7	30	M	8	A2L1B3p	Nil	Right hemicolectomy	24.54	9	32	u/a
8	33	M	3	A2L1B1	Nil	Right hemicolectomy	13.65	2	<5	577
9	46	M	15	A2L1B3	Nil	No	17.50	2	10	u/a
10	33	M	4	A2L1B2	Adalimumab	No	33.75	2	<5	1023
11	51	F	37	A1L1B2	Methotrexate Vedolizumab	3 small bowel resections	17.00	5	9	535
12	30	F	15	A1L3B1	nil	Subtotal colectomy	17.06	6	<5	604
13	53	M	27	A2L1B2	Methotrexate	Right hemicolectomy, small bowel resection, ileocolic resection	23.68	2	<5	1800
14	23	F	3	A2L1B1	Azathioprine	No	24.49	6	13	1587
15	19	M	6	A1L1B1	Nil	No	27.48	18	<5	u/a

Table summarising the demographics and key clinical variables of the CD subjects; Con Meds=concomitant medication at the time of recruitment, MaRIA=Magnetic Resonance Index of Activity, HBI-Harvey-Bradshaw Index, CRP=C-Reactive Protein, fC=Faecal Calprotectin, u/a=unavailable.