

Analysis of the X-Factor and X-Factor stretch during the completion of a golf practice session in low-handicap golfers

Graeme G. Sorbie
Yaodong Gu
Julien S. Baker
Ukadike C. Ugbolue

Sorbie, G.G., Gu, Y., Baker, J.S. & Ugbolue, U.C.
Analysis of the X-Factor and X-Factor stretch during the completion of a golf practice session in low-handicap golfers. *International Journal of Sports Science and Coaching*.

© 2018 the authors

Reprinted by permission of SAGE Publications.

DOI: <https://doi.org/10.1177/1747954118791330>

1 Original Research

2 Corresponding Author:

3 Graeme G Sorbie, Biomechanics Laboratory, Division of Sport and Exercise Sciences,
4 Abertay University, Dundee, DD1 1HG, United Kingdom, Phone: +44 (0)1382308015,
5 Email: g.sorbie@uad.ac.uk

6 **Analysis of the X-Factor and X-Factor stretch during the**
7 **completion of a golf practice session in low-handicap golfers**

Graeme G. Sorbie,^{1,2} Yaodong Gu,³ Julien S. Baker,^{1,3} and Ukadike C. Ugbolue,^{1,4}

8 ¹ School of Science and Sport, Institute for Clinical Exercise & Health Science, University of
9 the West of Scotland, Hamilton, ML3 0JB, United Kingdom

10 ² Division of Sport and Exercise Sciences, Abertay University, Dundee, DD1 1HG, United
11 Kingdom

12 ³ Faculty of Sports Science, Ningbo University, China

13 ⁴ Department of Biomedical Engineering, University of Strathclyde, Glasgow, United
14 Kingdom

15

16

17

18

19

20

21

22

23

24

25

1 **Abstract**

2 The X-Factor and X-Factor stretch have been positively correlated with golf long game
3 performance. The aim of this study was to compare the X-Factor, X-Factor stretch and long
4 game performance variables pre and following a golf practice session. A group of male golfers
5 ($n = 15$, handicap = 3.3 ± 1.7) participated in the laboratory-based-study. Movement and
6 performance variables were collected from five golf swings performed pre and following a golf
7 practice session using a motion capture system and launch monitor respectively. Following the
8 practice session, significant increases were observed in the X-Factor ($p = 0.00$, $d = 0.22$) and
9 the X-Factor stretch ($p = 0.02$, $d = 0.25$). Specifically, the X-Factor increased from $52.82 \pm$
10 5.64° to $54.06 \pm 5.61^\circ$ following the practice session. The X-Factor stretch increased from
11 $1.54 \pm 1.05^\circ$ to $1.90 \pm 1.41^\circ$ following the practice session. Significant differences were
12 displayed in club head velocity ($p = 0.00$, $d = 0.35$), ball velocity ($p = 0.01$, $d = 0.21$) and actual
13 carry distance ($p = 0.00$, $d = 0.29$) following the practice session. These findings suggest that
14 performing multiple golf shots is not detrimental in terms of muscular fatigue in the long game
15 performance. In actual fact, the findings demonstrate that performing 100 golf shots increases
16 the X-Factor, X-Factor stretch patterns and performance variables which, in turn, increases
17 long game performance. These findings can help PGA golf Professionals improve teaching
18 practices and formulation of golf programmes and warm-up sessions.

19

Key words: Flexibility, ROM, Performance, Movement, Biomechanics

20

21

22

23

1 **Introduction**

2 Similar to many other sports, there are several different ways to improve performance in golf;
3 including technique changes and physiological improvements.¹ The importance of long game
4 performance within the game of golf is widely recognised within the golfing literature. Long
5 game performance takes into consideration golf shots that are performed with a full swing. This
6 has recently intensified due to the rise in specific physical conditioning and warm-up routines
7 at the elite level.²⁻⁵ These physical conditioning and warm-up studies have mainly focused on
8 improving range of motion (ROM), stability and balance in order to improve movement
9 patterns, including the X-Factor, and in turn aim to increase long game performance.^{3,4}

10 Many of the studies that have focused on the physical conditioning of golfers have investigated
11 key components of long game performance including: club head velocity,^{2,3,5,6} ball velocity,^{7,8}
12 carry distance “distance travelled by the ball in the air”⁹ and kinematic changes during the golf
13 swing.¹⁰⁻¹³ These studies have reported increases in long game performance variables as a
14 result of performing golf specific conditioning and warm-up programmes.^{1,4-6} Specifically,
15 Thompson et al.³ reported a significant increase in club head velocity following a functional
16 training intervention. Similarly, Lephart et al.⁴ investigated the effect of an 8-week golf
17 stretching exercise programme. These authors reported a significant increase in: club head
18 velocity, ball velocity, carry distance, pelvis rotation and X-Factor velocity following the
19 intervention.

20 The X-Factor and X-Factor stretch have been extensively used within the golf scientific
21 research.^{4,9,11,14,15} The X-Factor describes the rotation of the torso relative to the pelvis at the
22 top of the backswing.¹⁶ Golfers that are able to exhibit a larger X-Factor are shown to increase
23 club head velocity and ball velocity at impact.^{4,11} Within the studies that reported X-Factor,
24 angles typically range between 30 - 55°, however, this is dependent on the skill level of the

1 golfer as previously discussed.^{4,15} The X-Factor stretch has been used to describe the additional
2 rotation that occurs at the start of the downswing.¹⁵ The X-Factor stretch is caused by the
3 initiation of the pelvic rotation prior to the rotation of the torso.¹⁵ It has been noted previously
4 that golfers who exhibit this stretch at the beginning of the downswing through dynamic tension
5 of the torso muscles have a greater club head velocity.⁴ Within highly skilled golfers, X-Factor
6 stretch angles typically range between 1 - 5°.⁹

7 In order to exhibit an increased X-Factor, golfers must demonstrate a good level of ROM¹⁷,
8 which is promoted through flexible muscles and joints.¹⁸ Flexibility has been found to decrease
9 resistance to swing plane and decrease stretch reflex.¹⁸ These factors enable an increased ROM
10 in the backswing¹⁹ and a reduction in injury.²⁰ It has been previously reported that performing
11 warm-up techniques before a competition can increase ROM and, therefore, increase the
12 performance of the athlete.^{21,22} In addition to these findings, Joyce⁸ reported that axial rotation
13 flexibility variables of the trunk were associated with club head velocity, which may help
14 facilitate a greater ROM and may create a greater X-Factor. Although previous research has
15 shown that performing warm-up techniques prior to a competition can increase ROM and in
16 turn increase performance, the golf literature is somewhat contrasting.^{23,24} Henry et al.¹⁵
17 reported that performing a golf specific dynamic rotational warm-up prior to golf competition
18 had no effect on the X-Factor or X-Factor stretch. In contrast, Gergley²³ reported a dynamic
19 golf club warm-up significantly increased club head velocity and distance in comparison to a
20 passive static stretching warm-up. These improvements are thought to be a result of increased
21 ROM and a more compliant muscle tendon unit. The contrasting findings within the literature
22 may be a result of the different warm-up routines examined. Although Henry et al.¹⁵ examined
23 a dynamic rotational warm-up, which has similarities to the golf swing, Gergley²³ examined a
24 warm-up routine that incorporated golf swings within the process. This may have a greater
25 effect on long game performance due to the greater specificity of the task being performed.²³

1 With the contrasting findings within the golf literature and the limited research on performing
2 multiple golf shots, it is unknown if the X-Factor at the top of the backswing or the X-Factor
3 stretch during the initiation of the downswing changes as a result of performing these golf
4 shots. During golf practice or competition, golfers may perform the golf swing 100 times or
5 more,²⁵ which includes golf shots when hitting the ball and practice swings (based on golf
6 handicap). It is possible that multiple golf swings during golf practice or warm-up routine may
7 increase ROM of the torso and pelvis regions at the top of the backswing and, therefore,
8 increase the X-Factor and X-Factor stretch. These swings, however, may also have a negative
9 effect on the separation between these segments by inducing fatigue in the muscles which are
10 highly active during the golf swing, resulting in a reduced ROM, which in turn reduces the X-
11 Factor, X-Factor stretch and long game performance variables.²⁶

12 Due to the limited knowledge regarding the effect that multiple golf swings can have on the X-
13 factor and X-Factor stretch, the purpose of the current study was to assess what effect the golf
14 practice session has on the X-factor, X-Factor stretch and performance variables. The aim of
15 the current study was to describe the X-Factor and X-Factor stretch patterns during the golf
16 swings performed pre and post golf practice session. Additionally, the current study aimed to
17 describe the changes, if any, in club head velocity, ball velocity and actual carry distance from
18 the golf swings performed pre and post golf practice session. A null hypothesis for all variables
19 was adopted due to the conflicting views on the topic area.

20 **Methods**

21 **Participants**

22 Fifteen right-handed male golfers participated in this study (height: 186.0 ± 5.3 cm, weight:
23 80.9 ± 6.9 kg, age: 23.8 ± 2.9 years, British Golf Association handicap: 3.3 ± 1.7). Additional
24 anthropometric measurements (leg length, knee width, ankle width, shoulder offset, elbow

1 width, wrist width, and hand thickness) were recorded during the initial stages of the testing
2 process in order to run a customised Golf Model. Measurements were recorded to the nearest
3 mm. Participants were required to have no history of lower back pain and/or persistent
4 musculoskeletal disorders. These factors may limit flexibility and golf swing attributes.²⁰
5 Participants were also required to undertake no conditioning or resistance training 48 hours
6 before the testing session. In addition to these specific requirements, participants were required
7 to exhibit a 'modern' golf swing rather than a 'classic swing'. This was assessed by a
8 Professional Golfers' Association (PGA) member. The main differences between the two
9 swing types is that the 'classic' swing exhibits lead heel raise during the backswing, resulting
10 in different pelvic movements, hence the reason why this swing was excluded from the current
11 study.⁸ All participants completed a consent form and physical readiness questionnaire before
12 participating in the current study. The University of the West of Scotland, School of Science
13 and Sport Ethics Committee granted ethical approval for the study to take place (Approval
14 number: 5-3-14-002).

15 **Experimental Procedure**

16 Prior to data collection, participants performed a standardised golf specific warm-up. Dynamic
17 stretches were performed which targeted specific musculature highly active during the golf
18 swing and was followed by five air swings. Following the warm-up, participants performed
19 five golf shots using the TaylorMade driver^{8,27,28} (TaylorMade, Basingstoke, UK) and a
20 premium golf ball. After these initial five golf shots were completed, participants then
21 completed a typical practice session, hitting 50 golf shots with the driver and 50 golf shots with
22 the TaylorMade 7-iron (TaylorMade, Basingstoke, UK). Prior to the testing all participants
23 were given five minutes to familiarise themselves with the TaylorMade golf clubs.¹ Utilizing
24 the identical clubs throughout the study removed the effect of the golf club. After the practice
25 session, participants hit five golf shots with the driver again.²⁶ Before hitting shots, participants

1 were informed of what the practice session would entail and advised to take into consideration
2 their average distance when using the 7-iron and driver.²⁹ During the practice session, all golf
3 shots in the session were hit at a rate of one shot every 30 s. During an initial pilot study, golfers
4 stated that this was a comfortable pace to perform the golf shots. In order for all golf shots to
5 be hit safely during the testing session, shots were hit from an artificial golf mat (Longridge,
6 United Kingdom), which was placed in the centre of the laboratory, towards an enclosed golf
7 net (Sports Net Company, United Kingdom). When performing golf shots, participants were
8 advised to aim towards a red target pole, which was situated behind the enclosed golf net.

9 **Data Collection**

10 During the five golf shots that were performed pre and post the practice session, three
11 dimensional (3-D) data were collected using an 8-camera Vicon Bonita (Oxford Metrics Ltd,
12 United Kingdom) Motion Analysis System operating at 250 Hz positioned around the golfer.
13 The system was calibrated at the start of the testing session and this calibration was used
14 throughout. Spherical retro-reflective markers (1.4 cm) were adhered to the skin on anatomical
15 regions throughout the testing session according to the adapted version of the Vicon Plug-in-
16 Gait Model (Vicon Motion Systems Ltd, Oxford, UK) using double-sided tape. This model
17 was adapted by adding an additional marker to the 4th spinous process of the lumbar region due
18 to the occlusion of jugular notch marker. This marker is often occluded from the camera view
19 during the golf swing due to the head position and the left and right upper section of the arm
20 during the backswing and downswing. Spherical markers were adhered to the golf club in four
21 locations including the base of the grip, halfway down the club, the hosel of the club, and the
22 club head. The four markers on the club were used to determine the top of the backswing and
23 the initial phase of the downswing. To ensure accurate 3-D data collection, participants were
24 asked to wear tight fitting shorts and their own golf shoes when performing all golf shots. In
25 addition to kinematic data collection, the Voice Caddie Swing Launch Monitor SC 100 GPS

1 (La Mirada, CA, USA) was used to calculate club head velocity, ball velocity and actual carry
2 distance of each golf shot. The Launch Monitor was previously validated in-house against the
3 Vicon Bonita Motion Analysis System; Trackman™ III Golf Swing and Ball Flight Analysis
4 System (Brighton, MI, USA). (CHS: ICC = 0.92, BS: ICC = 0.99). The Launch Monitor was
5 required to be positioned 1 m directly behind the golf ball and positioned towards the target
6 line of the golfer. After each golf shot, the performance variables were logged using Microsoft
7 Excel (Excel 2016 (v16.0)).

8 **Data Analysis**

9 For the five swings performed pre and post practice session with the driver, the trials were
10 individually analysed and compared between the sessions.⁷ All Raw 3D co-ordinate data was
11 filtered using a fourth-order low-pass Butterworth filter with a cut-off frequency of 6 Hz. A
12 multi-segment model used to analyse the X-Factor in the current study was developed using
13 BodyBuilder (Oxford, UK) and used in Vicon version 2.2.2. The X-Factor was calculated by
14 measuring the orientation of the torso in relation to the pelvis.¹⁷ The X-Factor stretch was
15 calculated by subtracting the X-Factor at the top backswing from the maximum X-Factor value
16 during the downswing. The backswing was defined as the point in which the clubhead reached
17 its most lateral position, towards the intended target, before changing direction. The
18 downswing is defined as the period from top of the backswing to ball contact.

19 **Statistical Analysis**

20 Normal distribution for all variables was assessed using the Shapiro-Wilk test. If normal
21 distribution was not granted, a log transformation was conducted on the specific data sets.
22 Following this, an ANCOVA was used to determine significant differences, if any, between
23 the X-Factor and the X-Factor stretch from the five shots performed pre and post practice
24 session. Additionally, an ANCOVA was used to determine significant differences, if any,
25 between the club head velocity, ball velocity and actual carry distance from the five shots

1 performed pre and post practice session. Following a p value correction, the level of
2 significance was set at $p \leq 0.02$ and effect sizes were calculated using the Cohen's d method.³⁰
3 Effect sizes of less than 0.2 were considered negligible effects and between 0.2 and 0.5 were
4 considered small effects. Values between 0.5 and 0.8 were considered medium effects and
5 greater than 0.8 represented large effects.³⁰ All calculations were performed in SPSS (version
6 23) and Microsoft Excel.

7 **Results**

8 **Biomechanical Variables**

9 Significant differences were observed for the X-Factor after the completion of the golf practice
10 session in comparison to the golf swings performed before the practice session ($p = 0.00$, $d =$
11 0.22). On average participants X-Factor was $52.82 \pm 5.64^\circ$ at the start of the golf practice
12 compared to $54.06 \pm 5.61^\circ$ at the end of the golf practice session.

13 Significant differences were observed for the X-Factor stretch after the completion of the golf
14 practice session in comparison to the golf swings performed before the practice session ($p =$
15 0.02 , $d = 0.25$). On average participants X-Factor stretch was $1.54 \pm 1.05^\circ$ at the start of the
16 golf practice compared to $1.90 \pm 1.41^\circ$ at the end of the golf practice session.

17 **Performance Variables**

18 Significant changes were displayed in club head velocity after the golf practice session in
19 comparison to the swings performed before the practice session when using the driver ($p =$
20 0.00 , $d = 0.35$). On average participants club head velocity was 166.79 ± 7.81 km/h at the start
21 of the golf practice compared to 169.49 ± 7.43 km/h at the end of the golf practice session.

22 Significant changes were displayed in ball velocity after the golf practice session in comparison
23 to the swings performed before the practice session when using the driver ($p = 0.01$, $d = 0.21$).

1 On average participants ball velocity was 237.81 ± 14.95 km/h at the start of the golf practice
2 compared to 240.83 ± 13.36 km/h at the end of the golf practice session.

3 Significant changes were displayed in actual carry distance after the golf practice session in
4 comparison to the swings performed before the practice session when using the driver ($p =$
5 0.00 , $d = 0.29$). On average participants actual carry distance was 235.44 ± 17.77 m at the start
6 of the golf practice compared to 240.14 ± 13.88 m at the end of the golf practice session.

7 **Discussion**

8 The aim of the current study was to describe the changes, if any, in the X-Factor at the top of
9 the backswing and X-Factor stretch during the initiation of the downswing during the golf
10 swings performed pre and post golf practice session. Additionally, the current study aimed to
11 describe the changes, if any, in club head velocity, ball velocity and actual carry distance from
12 the golf swings performed pre and post golf practice session. A null hypothesis for all variables
13 was adopted due to the conflicting views on the topic area.

14 The findings from the current study show that the X-Factor at the top of the backswing
15 increased significantly as a result of performing multiple golf swings, therefore rejecting the
16 initial hypothesis. It must be highlighted, however, that the effect sizes for these X-Factor
17 comparisons were small. Although the magnitude of change was small when comparing pre
18 and post conditions, the results demonstrate that the incorporation of multiple golf swings
19 during a practice session could be important in producing increased coordination patterns,
20 which can result in greater movement patterns and in turn can increase the performance of the
21 golfer. These findings suggest that when PGA Professionals are designing golf practice and
22 warm-up sessions for elite golfers, they should advocate that golfers hit 100 balls. As
23 previously discussed, limited research has been conducted regarding the effect that performing
24 multiple golf swings has on movement patterns during the golf swing. Researchers have,

1 however, investigated the effect that various warm-up routines have on the X-Factor at the top
2 of the backswing.

3 In relation to warm-up routines, it has been previously reported that performing a golf specific
4 dynamic rotational warm-up prior to golf play does not significantly increase the X-Factor.¹⁵
5 The study conducted by Henry and colleagues has similarities to the current study. Both studies
6 have examined movements directly related to the golf swing and what effect repeating these
7 over time has on the X-Factor. The contrasting findings between the studies may be a result of
8 the multiple golf swings performed in the current study increasing ROM of the upper limbs to
9 a greater extent than performing a golf specific dynamic warm-up routine. Previous research
10 has shown that performing multiple golf swings as part of a warm-up routine significantly
11 increases long game performance in comparison to passive static stretching.²³ Gergley²³
12 reported club head velocity significantly increased by 6.19 km/h when performing a warm-up
13 protocol comprising of only golf swings, compared to performing passive static stretching.
14 Taking these previous findings into consideration, the findings of the current study suggests
15 that performing multiple golf shots as a warm-up method may be more beneficial than dynamic
16 and passive static stretching. Furthermore, although the protocols have similarities in relation
17 to increasing ROM, there are distinct differences in the participants handicap range between
18 the two studies, which could be a reason for the conflicting findings. The mean handicap within
19 the study conducted by Henry and colleagues was 13.9 ± 6.4 , whereas the handicap in the
20 current study was 3.3 ± 1.7 . The higher skilled golfers participating in the current study may
21 respond differently to the intervention in comparison to the higher handicap golfers due to their
22 increased practice hours. With regards to the X-Factor results at the top of the backswing
23 between the two studies, Henry and colleagues reported a mean angle of $31.0 \pm 9.2^\circ$ prior to
24 the dynamic stretch routine and $34.1 \pm 9.0^\circ$ after the warm-up. The values within the current
25 study are considerably higher which are to be expected due to the higher skill level of golfer.

1 It has been previously reported that there is a correlation between handicap and the X-Factor
2 at the top of the swing. As the handicap of the golfers increases, the X-Factor decreases.^{11,31}

3 In relation to the X-Factor stretch, the current findings show significant differences between
4 the pre and post golf practice session, however, it must be highlighted that the effect sizes for
5 these comparisons were small. Although small in magnitude, the results suggest that the X-
6 Factor stretch increases following multiple golf swing. This finding is conflicting with
7 previously published literature where warm-up techniques have been analysed. Henry et al.¹⁵
8 found no significant difference in the X-Factor stretch between the pre and post golf swing
9 during a golf specific dynamic warm-up routine. These conflicting findings may be a result of
10 the previously discussed participant differences between the current study and the study
11 conducted by Henry and colleagues. It has been noted previously that an increase in the X-
12 Factor stretch at the beginning of the downswing can increase the performance of the golfer.⁴
13 The findings from the current study support these views by displaying a significant increase in
14 club head velocity, ball velocity and actual carry distance when the X-Factor stretch
15 significantly increased after the completion of the practice session.

16 In addition to the changes in the X-Factor and X-Factor stretch, the current study displayed
17 significant changes in club head velocity, ball velocity and actual carry distance following
18 completion of the practice session. Similar to the X-Factor and X-Factor stretch, the effect sizes
19 for all performance variables were small. Although the magnitude of change between pre and
20 post practice session swings are small, these results are in agreement with the previously
21 reported correlation between club head velocity and the X-Factor at the top of the
22 backswing.^{4,11} Previous studies have reported the effect a golf practice session has on long
23 game performance variables. Horton et al.⁷ reported no significant changes in ball velocity after
24 the completion of the golf practice session. Although the authors did report an increase in ball
25 velocity after the golf practice session. Sorbie et al.²⁶ reported no significant changes to club

1 head velocity, ball velocity and actual carry distance. Contrasting to the study by Horton and
2 colleagues, Sorbie et al.²⁶ reported a decrease in club head velocity, ball velocity and actual
3 carry distance after the golf practice session was complete. The current study and the study
4 conducted by Horton et al.⁷ tested low handicap golfers and professional respectively. In
5 contrast, Sorbie et al.²⁶ examined higher handicap golfers (15.2 ± 5.7). These contrasting
6 findings may be a result of the higher handicap golfers conducting less practice time compared
7 to skilled golfers and as a result display signs of muscular fatigue during the repetitive golf
8 shots.²⁶

9 To the researchers knowledge, only Horton et al.⁷ and Sorbie et al.²⁶ have investigated the effect
10 that the golf practice session has on injury prevention and specific performance variables
11 associated with the long game. In terms of injury, Horton et al.⁷ found that performing practice
12 sessions is unlikely to increase injury in the trunk area during the golf swing. Similarly, Sorbie
13 et al.²⁶ reported no change in erector spinae muscle activation during the golf swing when
14 performing a golf practice session. The current study adds to this topic by displaying increases
15 in the X-Factor, X-Factor stretch and long game performance variables during the completion
16 of a golf practice session within low-handicap golfers. These findings can help PGA
17 Professionals, who teach highly skilled golfers, to improve movement patterns in relation to
18 the X-Factor and X-Factor stretch and, in turn, improve the long game performance of the
19 golfer. The findings of the current study can also help PGA Professionals to design golf practice
20 sessions and warm-up sessions that can improve the long game of the golfer. Specifically, as a
21 result of the findings from the current study, PGA Professionals can promote that golfers should
22 hit 100 balls during a practice session or warm-up session in order to optimize movement
23 patters and long game performance. Implementing this prior to golf play or competition may
24 enable highly skilled golfers to be at an optimal level at the beginning of the golf round.

1 However, future studies should examine the performance of the golf round following the
2 practice session in the current study.

3 Following the reported increase in the X-Factor at the top of the backswing, and marginal
4 increases in club head velocity, ball velocity and actual carry distance, further investigation is
5 required to ascertain the point where the X-Factor and performance variables are at their
6 optimal throughout the golf practice session. This could be an important finding for
7 practitioners when designing golf practice sessions and golf specific warm-up sessions for
8 professional and amateur golfers. It is important for future research to be conducted using
9 participants with a higher handicap and also female golfers in order to increase the
10 understanding of the topic area. In addition to this, it is also important for future search to
11 investigate additional movement patterns throughout the golf swing that are associated with
12 increased performance.

13 It is important to acknowledge the limitations within the current study. All golf shots performed
14 by the participants were hit with the identical clubs and shaft lengths. Although participants
15 were given time to familiarise themselves with the two clubs and shaft lengths used within the
16 current study, it has been previously suggested swing kinematics are influenced by a golfer
17 hitting a non-familiar club and shafts.³² However, standardising the golf clubs throughout the
18 study removes the effect of the golf club from the results.

19 **Conclusion**

20 The results of the current study indicate that performing multiple golf swings can increase the
21 X-Factor at the top of the backswing and can increase the X-Factor stretch during the initiation
22 of the downswing. Furthermore, the results from the current study indicate that performing
23 multiple golf swings increases long game performance variables. These results demonstrate
24 that performing multiple golf swings could be important in order to produce increased

1 coordination patterns, which can result in greater movement patterns of the golfer and therefore
2 increase long game performance. These findings can be important for PGA Professionals in
3 terms of better understanding of the efficacy of golf practice and warm-up sessions in relation
4 to golf performance.

5 **Funding**

6 The author(s) received no financial support for the research, authorship, and/or publication of
7 this article.

8 **Disclosure statement**

9 No potential conflict of interest was reported by the authors.

10 **References**

- 11 1. Doan BK, Newton RU, Kwon Y-H, et al. Effects of physical conditioning on
12 intercollegiate golfer performance. *J Strength Cond Res* 2006; 20: 62–72.
- 13 2. Fradkin A, Sherman C, Finch C. How well does club head speed correlate with golf
14 handicaps? *J Sci Med Sport* 2004; 7: 465–472.
- 15 3. Thompson C, Blackwell J. Functional training improves club head speed and
16 functional fitness in older golfers. *J Strength Cond Res* 2007; 21: 131–137.
- 17 4. Lephart S, Smoliga J, Myers J, et al. An eight-week golf-stretching exercise program
18 improves physical characteristics, swing mechanics, and golf performance in
19 recreational golfers. *J Strength Cond Res* 2007; 21: 860–869.
- 20 5. Fletcher IM, Hartwell M. Effect of an 8-week combined weights and plyometrics
21 training program on golf drive performance. *J strength Cond Res* 2004; 18: 59–62.
- 22 6. Hetu F, Christie C, Faigenbaum A. Effects of conditioning on physical fitness and club
23 head speed in mature golfers. *Percept Mot Skills* 1998; 86: 811–815.
- 24 7. Horton J, Lindsay D, Macintosh B. Abdominal muscle activation of elite male golfers
25 with chronic low back pain. *Med Sci Sport Exerc* 2001; 1647–1654.
- 26 8. Joyce C. An examination of the correlation amongst trunk flexibility, x-factor and
27 clubhead speed in skilled golfers. *J Sports Sci* 2016; 1–7.
- 28 9. An J, Wulf G, Kim S. Increased carry distance and X-Factor stretch in golf through an
29 external focus of attention. *J Mot Learn Dev* 2013; 1: 2–11.
- 30 10. Tinmark F, Hellström J, Halvorsen K, et al. Elite golfers' kinematic sequence in full-
31 swing and partial-swing shots. *Sport Biomech* 2010; 9: 236–44.
- 32 11. Myers J, Lephart S, Tsai Y-S, et al. The role of upper torso and pelvis rotation in

- 1 driving performance during the golf swing. *J Sports Sci* 2008; 26: 181–188.
- 2 12. Nesbit SM. A three dimensional kinematic and kinetic study of the golf swing. *J*
3 *Sports Sci Med* 2005; 4: 499–519.
- 4 13. Healy A, Moran KA, Dickson J, et al. Analysis of the 5 iron golf swing when hitting
5 for maximum distance. *J Sports Sci* 2011; 29: 1079–88.
- 6 14. Hume PA, Keogh J, Reid D. The role of biomechanics in maximising distance and
7 accuracy of golf shots. *Sport Med* 2005; 35: 429–449.
- 8 15. Henry E, Berglund K, Millar L, et al. Immediate effects of a dynamic rotation-specific
9 warm-up on x-factor and x-factor stretch in the amateur golfer. *Int J Sports Phys Ther*
10 2015; 10: 998–1007.
- 11 16. Mclean J, Andrisani J. *The X-factor swing: And other secrets to power and distance*.
12 New York: Harper Collins, 1996.
- 13 17. Brown SJ, Selbie WS, Wallace ES. The X-Factor: An evaluation of common methods
14 used to analyse major inter-segment kinematics during the golf swing. *J Sports Sci*
15 2013; 31: 1156–63.
- 16 18. Chettle DK, Neal RJ. Optimising performance in golf. In: Ltd. AAP (ed) *Strength and*
17 *conditioning for golf*. Brisbane, 2001, pp. 207–223.
- 18 19. Keogh JW, Marnewick MC, Maulder PS, et al. Are anthropometric, flexibility,
19 muscular strength, and endurance variables related to clubhead velocity in low- and
20 high-handicap golfers? *J Strength Cond Res* 2009; 23: 1841–1850.
- 21 20. Lindsay DM, Horton JF. Trunk rotation strength and endurance in healthy normals and
22 elite golfers with and without lower back pain. *North Am J Sport Phys Ther* 2006; 1:
23 80–89.
- 24 21. Radford JA, Burns J, Buchbinder R, et al. Does stretching increase ankle dorsiflexion
25 range of motion? A systematic review. *Br J Sports Med* 2006; 40: 870–875.
- 26 22. Stewart IB, Sleivert GG. The effect of warm-up intensity on range of motion and
27 anaerobic performance. *J Orthop Sport Phys Ther* 1998; 27: 154–161.
- 28 23. Gergley J. Acute effects of passive stratic stretching during warm-up on driver
29 clubhead speed, distance, accuracy, and consistent ball contact in young male
30 competitive golfers. *J strength Cond Res* 2009; 23: 863–867.
- 31 24. Tilley NR, Macfarlane A, Tilley N. Effects of different warm-up programs on golf
32 performance in elite male golfers. *Int J fo Sport Phys Ther* 2012; 7: 388–395.
- 33 25. Higdon NR, Finch WH, Leib D, et al. Effects of fatigue on golf performance. *Sport*
34 *Biomech* 2012; 11: 190–196.
- 35 26. Sorbie GG, Baker JS, Gu Y, et al. Comparison of thoracic and lumbar erector spinae
36 muscle activation before and after a golf practice session. *J Appl Biomech* 2017; 33:
37 288–293.
- 38 27. Okuda I, Gribble P, Armstrong C. Trunk rotation and weight transfer patterns between
39 skilled and low skilled golfers. *J Sport Sci Med* 2010; 9: 127–133.
- 40 28. Sorbie GG, Grace FM, Gu Y, et al. Electromyographic analyses of the erector spinae

- 1 muscles during golf swings using four different clubs. *J Sports Sci* 2018; 36: 717–723.
- 2 29. Silva L, Vaz JR, Castro MA, et al. Recurrence quantification analysis and support
3 vector machines for golf handicap and low back pain EMG classification. *J*
4 *Electromyogr Kinesiol* 2015; 25: 637–647.
- 5 30. Cohen J. A Power Primer. *Psychol Bullitin* 1992; 112: 155–159.
- 6 31. Zheng N, Barrentine S, Fleisig G, et al. Kinematic analysis of swing in pro and
7 amateur golfers. *Int J Sports Med* 2008; 29: 487–493.
- 8 32. Joyce C, Burnett A, Cochrane J, et al. A preliminary investigation of trunk and wrist
9 kinematics when using drivers with different shaft properties. *Sport Biomech* 2016; 15:
10 61–75.
- 11