

## Article

## Investigating Permeation Behaviour of Flufenamic Acid Cocrystals using A Dissolution and Permeation System

Minshan Guo, Ke Wang, Ning Qiao, Vanessa Yardley, and Mingzhong Li

*Mol. Pharmaceutics*, **Just Accepted Manuscript** • DOI: 10.1021/acs.molpharmaceut.8b00670 • Publication Date (Web): 06 Aug 2018Downloaded from <http://pubs.acs.org> on August 15, 2018

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Investigating Permeation Behaviour of Flufenamic Acid Cocrystals Using A Dissolution and Permeation System

*Minshan Guo,<sup>†</sup> Ke Wang,<sup>†</sup> Ning Qiao,<sup>‡</sup> Vanessa Yardley,<sup>†§</sup> and Mingzhong Li<sup>\*†</sup>*

<sup>†</sup> School of Pharmacy, De Montfort University, Leicester LE1 9BH, UK


<sup>‡</sup> College of Materials Science and Engineering, North China University of Science  
and Technology, Tangshan, 063210, Hebei, China

<sup>†§</sup> Department of Infection & Immunity, Faculty of Infectious & Tropical Diseases,  
London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E  
7HT, UK

1  
2  
3 \*Address: School of Pharmacy, De Montfort University, Leicester, LE1 9BH, U.K.  
4

5 Tel: +44-1162577132; Email: mli@dmu.ac.uk  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Table of Contents (TOC)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **ABSTRACT:** The dissolution and permeation of the cocrystals, Flufenamic acid –  
4 nicotinamide (FFA-NIC) and Flufenamic acid – theophylline (FFA-TP), have been  
5 investigated in the presence of two polymers, polyvinylpyrrolidone (PVP) and  
6 copolymer of vinyl pyrrolidone/vinyl acetate (PVP-VA), using a  
7 dissolution/permeation (D/P) system. It showed that the types and concentrations of  
8 the polymers and their interactions with the coformers had significant effects on the  
9 dissolution and permeation of the FFA cocrystals. The role of PVP as a stabilizing  
10 agent was not altered in spite of its interaction with the coformer of NIC or TP, which  
11 was supported by the proportional flux rate of FFA to the dissolution performance  
12 parameter (DPP). With an appropriate PVP concentration, the maximal flux rate of  
13 FFA could be obtained for a given FFA cocrystal. The situation was complicated in  
14 the presence of PVP-VA. The role of PVP-VA could change due to its association  
15 with the coformers, i.e., from a stabilizing agent to a solubilisation agent. In addition,  
16 PVP-VA reduced the flux rate of FFA contrasting to its DPP for FFA cocrystals.  
17 Finally,  $^1\text{H}$  NMR provided evidence regarding the molecular interactions between  
18 FFA, coformers and polymers at atomic level and gave insight into the mechanism  
19 underlying the supersaturated solution and subsequent permeation behaviour of the  
20 cocrystals.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41  
42  
43 **KEYWORDS:** *cocrystal, flufenamic acid,  $^1\text{H}$  NMR, polymer, dissolution,*  
44 *permeation*  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1. INTRODUCTION

Over the last decade, pharmaceutical cocrystals have been studied extensively as potential tools to optimise the drug product performance, in particular for compounds with limited aqueous solubility/dissolution rates.<sup>1-3</sup> However, the supersaturated solution generated by dissolution of the highly soluble cocrystals was thermodynamically unstable and the parent drug had a tendency of rapid precipitation.<sup>4-6</sup> It was shown that the parent drug tended to precipitate directly onto the surface of the dissolving cocrystals during dissolution, acting as a coating layer to diminish the dissolution and solubility of the cocrystals.<sup>4,5</sup> In sight of this, inclusion of effective crystallisation inhibitors in the formulations have been explored to maintain the supersaturated conditions of API (active pharmaceutical ingredient).<sup>4,6-11</sup> On one occasion, an excess use of coformer or micellar solubilisation decreased the solubility of cocrystals, leading to a lower degree of supersaturation and subsequent prevention of drug precipitation.<sup>7,8</sup> Maintaining supersaturated drug concentration was also achieved by introducing stabilizing agents in the formulation, that the equilibrium solubility of the drug was not enhanced but the nucleation and/or crystal growth of the drug were inhibited by the intermolecular interactions between the drug molecules and the stabilizing agents. Although polymeric crystallisation inhibitors have been widely used in amorphous solid dispersion formulations to stabilize the amorphous state of the drug substance and to maintain its supersaturated status in solution,<sup>12,13</sup> such studies have rarely been carried out in cocrystal based formulations.<sup>4,9,10,14-16</sup> The selection of a polymeric excipient in a cocrystal formulation seemed to be much more complicated than those used in the solid dispersion system because the inhibitory ability of the polymers could be undermined significantly by the competitions between the drug, the coformer and the polymer in

1  
2  
3 terms of hydrogen bondings.<sup>9</sup> Furthermore, a dissolved polymer could alter the  
4 dissolution of the cocrystals and subsequently affect its inhibitory ability on the  
5 precipitation of the parent drug.<sup>4</sup>  
6  
7

8  
9 For orally administered solid dosage forms of a drug compound, the elevation of  
10 drug concentration in the gastrointestinal (GI) tract did not always improve the  
11 absorption as the drug molecules have to permeate across the GI membrane. It was  
12 reported that, despite of an increased solubility due to the presence of co-solvent,  
13 micelle, cyclodextrin, hydrotrophy and emulsifier, a reduced fraction of free drug for  
14 membrane permeation by drug encapsulation could eventually lead to very poor drug  
15 absorption.<sup>17-25</sup> Therefore, the balance between the solubility and the permeability had  
16 to be optimised to achieve the maximal overall absorption of a poorly water soluble  
17 drug. The drug permeability of an amorphous solid dispersion formulation was highly  
18 polymer dependent.<sup>20, 26-29</sup> A few evidence showed that the cocrystals alone could  
19 alter both the solubility and the permeability of the APIs.<sup>30-32</sup> Though it was not clear  
20 that, in a cocrystal based formulation, how the permeability of the parent drug was  
21 affected by additional polymeric excipients, some speculated that the interplay of the  
22 solubility and permeability was the determining factor for a successful formulation.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Our previous investigations found that inclusion of the polymers of  
40 polyvinylpyrrolidone (PVP) and copolymer of vinyl pyrrolidone/vinyl acetate (PVP-  
41 VA) altered the dissolution of cocrystals of flufenamic acid-nicotinamide (FFA-NIC)  
42 and flufenamic acid-theophylline (FFA-TP) and maintained the supersaturation of  
43 FFA in solution.<sup>33</sup> In this work we pioneered a systematic examination of key  
44 processes occurred during cocrystal dissolution of FFA-NIC and FFA-TP in the  
45 absence and presence of PVP or PVP-VA at different concentrations to reveal the  
46 kinetics of drug supersaturation and permeation. The molecular structures of FFA  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 cocrystals and the monomer units of the polymers were shown in Table 1 and a  
4 detailed description could be found from previous work.<sup>4, 9, 34, 35</sup> Simultaneous  
5 evaluations of the dissolution of FFA cocrystals in the absence and presence of PVP  
6 or PVP-VA as well as the permeation of FFA through a dialysis membrane were  
7 achieved using a dissolution/permeation (D/P) system (Fig. 1). The dialysis  
8 membrane modelled the intestinal epithelium. The D/P system represented a  
9 simplified permeation model for quantitative analysis because of the negligible effect  
10 of polymer on the membrane.<sup>28, 36, 37</sup> And it had been proved to be a useful tool for  
11 examining the absorption of oral drugs after administration.<sup>36-41</sup> In the D/P system,  
12 the powdered drug substances were applied to the donor compartment, which was  
13 filled with dissolution media in the absence or presence of the polymer at various  
14 concentrations, under nonsink conditions to evaluate the ability of the polymer to  
15 maintain the supersaturated drug solution. Meanwhile, the amount of FFA molecules  
16 permeated into the acceptor compartment, which was filled with the dissolution  
17 media, was monitored. The temperature of the whole D/P system was maintained at  
18 37°C using a circulating water bath. The molecular interactions of the FFA, cofomers  
19 and polymers were analysed by one-dimensional proton nuclear magnetic resonance  
20 (<sup>1</sup>H NMR) spectroscopy. The <sup>1</sup>H NMR analysis was conducted in low polarity solvent  
21 deuterated chloroform (CDCl<sub>3</sub>) solutions containing singular, binary and ternary  
22 components of FFA, cofomers and polymers. The changes of the chemical  
23 environment surrounding the molecules would be reflected by the characteristic peak  
24 shifts in <sup>1</sup>H NMR spectra and the molecular mobility would be reflected by the peak  
25 widths.<sup>42-51</sup>



## 2. MATERIALS AND METHODS

**2.1. Materials.** Flufenamic acid form I (FFA), Nicotinamide (NIC) ( $\geq 99.5\%$  purity), Theophylline (TP) ( $\geq 99.5\%$  purity), Potassium dihydrogen phosphate ( $\text{KH}_2\text{PO}_4$ ) and sodium hydroxide (NaOH) were purchased from Sigma-Aldrich (Dorset, UK). Plasdone K-29/32 (PVP) and Plasdone S-630 (PVP-VA) were gifts from Ashland Inc. (Schaffhausen, Switzerland). Methanol (HPLC grade) and acetonitrile (HPLC grade) were purchased from Fisher Scientific (Loughborough, UK) and used as received. Double distilled water was generated from a Bi-Distiller (WSC044.MH3.7, Fistreem International Limited, Loughborough, UK) and used throughout the study. Deuterated chloroform ( $\text{CDCl}_3$ ) was purchased from Goss Scientific Instruments Ltd (Crewe, UK).

### 2.2. Methods.

*2.2.1. Preparation of 0.01M Phosphate Buffer Solution (PBS), pH 4.5.* PBS solution (0.01M, pH 4.5) was used as the dissolution medium. It was prepared according to British Pharmacopeia 2010: 1.361g of potassium phosphate monobase was dissolved in double distilled water and the pH of the solution was adjusted to 4.5 using 1 M of sodium hydroxide solution.

*2.2.2. Preparation of FFA Cocrystals.* As previously described,<sup>9</sup> the cocrystals of FFA-NIC and FFA-TP were prepared with the solvent evaporation and the cooling crystallization methods, respectively. The formation of cocrystals was confirmed by Differential Scanning Colorimetry (DSC), Fourier Transform Infrared Spectroscopy (FTIR) and Powder X-ray Diffraction (PXRD).

*2.2.3. The Apparent Equilibrium Solubility Measurement for FFA in the Absence and Presence of Polymers and/or Coformers.* To investigate the effect of coformers and/or polymers on the solubility of FFA, 20 mL of 0.01 M PBS (pH 4.5) was used as

1  
2  
3 the solvent to prepare the solutions of PVP or PVP-VA at concentrations of 200  
4  $\mu\text{g/mL}$ , 500  $\mu\text{g/mL}$ , 1000  $\mu\text{g/mL}$ , 2000  $\mu\text{g/mL}$ , 4000  $\mu\text{g/mL}$ , 8000  $\mu\text{g/mL}$ , 12000  
5  $\mu\text{g/mL}$  and 16000  $\mu\text{g/mL}$ . Appropriate amount of the powdered FFA crystals was  
6 then added to these solutions for the apparent equilibrium solubility test. The FFA  
7 crystals were slightly grounded by a mortar and pestle and passed through a 60 mesh  
8 sieve so that the particle size of the crystals was no more than 250  $\mu\text{m}$ . The test was  
9 carried out for 24 h at a temperature of  $37 \pm 0.5^\circ\text{C}$  with constant mixing at 150 r.p.m.  
10 The clear solution was then separated from the remaining solid by centrifugation  
11 (MSB 010.CX2.5 centrifuge, MSE Ltd, London, U.K.) at 13,000 r.p.m. for 1 min and  
12 the concentration of FFA was determined by High Performance Liquid  
13 Chromatography (HPLC).  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26  
27 To investigate the combined effects of coformer and polymer on the solubility of  
28 FFA, the above series of PVP or PVP-VA solutions were added with NIC or TP at a  
29 concentration of 47.8  $\mu\text{g/mL}$  or 71.1  $\mu\text{g/mL}$ , respectively. The coformer concentration  
30 selected in the experiments was based on the flux rate experiments, which 1.43 mg of  
31 FFA-NIC and 1.64 mg of FFA-TP were corresponding to 47.8  $\mu\text{g/mL}$  of NIC and  
32 71.1  $\mu\text{g/mL}$  of TP, under the assumption of complete dissolving of the FFA  
33 cocrystals. An appropriate amount of the FFA crystals was then added to these  
34 solutions for the apparent equilibrium solubility test.  
35  
36  
37  
38  
39  
40  
41  
42  
43

44  
45 *2.2.4. The Dissolution and Permeation Measurements.* The D/P system was used to  
46 evaluate the dissolution and permeation of cocrystals. It consisted of a donor and an  
47 acceptor compartment, which was separated by a regenerated cellulose membrane  
48 with a Molecular Weight Cut-Off (MWCO) of 6-8 KDa (Spectrum Labs Inc. Rancho  
49 Dominguez, CA 90220, USA) (Fig. 1). Either the donor or acceptor compartment has  
50 a capacity of 10 mL. The orifice diameter of the compartment was 0.9 cm,  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 corresponding to 0.671 cm<sup>2</sup> surface area of the membrane. Like the FFA crystals, the  
4  
5 crystalline substance used in this experiment were prepared to sizes that no bigger  
6  
7 than 250 μm. 1 mg of single component crystals of FFA or cocrystals with equivalent  
8  
9 amount of FFA were firstly added to the donor compartment and this was followed by  
10  
11 the addition of 9 mL of 0.01 M PBS (pH 4.5) to the receptor compartment. 9 mL of  
12  
13 0.01 M PBS (pH 4.5) in the absence or presence of PVP or PVP-VA (200 μg/mL,  
14  
15 4000 μg/mL, and 16000 μg/mL) was then added to the donor compartment which  
16  
17 contained the crystals. The temperature of the whole system was maintained at 37°C  
18  
19 by a circulating water bath. 0.5 mL of the sample was withdrawn from the donor  
20  
21 compartment at the time intervals of 5, 10, 15, 30, 60, 120 and 240 mins using a  
22  
23 syringe and any volume change due to the withdrawal was immediately compensated  
24  
25 with 0.01 M PBS (pH 4.5). The concentrations of FFA were then determined by  
26  
27 HPLC.  
28  
29  
30

31 The dissolution performance parameter (DPP)<sup>52</sup> was used to evaluate the dissolution  
32  
33 of the FFA cocrystals, as detailed previously,<sup>4</sup> and the dissolution of FFA crystals was  
34  
35 used as the reference. The difference of the DPP of the FFA cocrystals measured in  
36  
37 the absence and presence of PVP or PVP-VA would give indication of the effect of  
38  
39 the polymers on dissolution and on the maintenance of drug supersaturation.  
40

41 The flux of a drug through the membrane, which was defined as the amount of drug  
42  
43 crossing a unit area perpendicular to its flow direction per unit time *t*, was calculated  
44  
45 using the following equation:  
46  
47

$$48 \quad J(t) = \frac{(C_{t_2} - C_{t_1})}{A(t_2 - t_1)} V \quad (1)$$

49 where *J(t)* was the flux of a drug; *C*<sub>*t*1</sub> was the drug concentration (μg/mL) at *t*<sub>1</sub>; *C*<sub>*t*2</sub>  
50  
51 was the drug concentration (μg/mL) at *t*<sub>2</sub>; *V* was the solution volume and *A* is the  
52  
53 area of exposed membrane.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3       2.2.5. *NMR Measurements.*  $^1\text{H}$  NMR was used to identify the interactions between  
4 the drug, coformer and polymer in solution. The measurements were carried out with  
5 a Bruker AV400 NMR Spectrometer (Bruker UK Limited, Coventry, UK) and the  
6 same settings were used throughout the measurements: 64 scans with a relaxation  
7 delay of 1 s, a spectral of 8278Hz, a time domain of 32k data and analysing software  
8 Mestrenova V11.0 (Mestrelab Research, Escondido, CA 92027, USA).

9  
10  
11  
12  
13  
14  
15  
16       All the samples were prepared in deuterated chloroform ( $\text{CDCl}_3$ ) using the standard  
17 5 mm NMR tubes and the spectra of tetramethylsilane (TMS) was used as an internal  
18 standard. Detailed measurements were shown in Table S1 in the Supporting  
19 Information, which included spectra of the singular component (FFA, NIC, TP, PVP,  
20 PVP-VA), the binary components (FFA/NIC, FFA/TP, FFA/PVP, FFA/PVP-VA,  
21 NIC/PVP, NIC/PVP-VA, TP/PVP, TP/PVP-VA) and the ternary components  
22 (FFA/NIC/PVP, FFA/NIC/PVP-VA, FFA/TP/PVP, FFA/TP/PVP-VA).

23  
24  
25  
26  
27  
28  
29  
30  
31       To mimic a cocrystal system, the FFA and its coformers were included in the  
32 solution at 1:1 molar ratio. Three sets of concentrations were prepared for the  
33 simulated cocrystal system: 215  $\mu\text{g}/\text{mL}$  NIC/ 320  $\mu\text{g}/\text{mL}$  TP and 500  $\mu\text{g}/\text{mL}$  FFA; 430  
34  $\mu\text{g}/\text{mL}$  NIC/640  $\mu\text{g}/\text{mL}$  TP and 1000  $\mu\text{g}/\text{mL}$  FFA; 2150  $\mu\text{g}/\text{mL}$  NIC/3200  $\mu\text{g}/\text{mL}$  TP  
35 and 5000  $\mu\text{g}/\text{mL}$  FFA. For investigation of the effect of polymer on the simulated  
36 cocrystal system, PVP or PVP-VA was included in the solution at a concentration of  
37 200  $\mu\text{g}/\text{mL}$  or 5000  $\mu\text{g}/\text{mL}$ , respectively.

38  
39  
40  
41  
42       2.2.6. *High Performance Liquid Chromatography (HPLC) Analysis.* The  
43 concentration of FFA, NIC or TP was determined by HEWLETT PACKARD series  
44 1100 automatic HPLC with a Luna@ Omega PS C18 100A LC column (5  $\mu\text{m}$ , 150  $\times$   
45 4.6 mm) (Phenomenex, Inc., Macclesfield, UK). The sample temperature was  
46 maintained at 40°C. The conditions used for measurements were listed in Table 2.

### 3. RESULTS

**3.1 Effects of Polymer and/or Coformer on the Equilibrium Solubility of FFA.** The effects of polymer and/or coformer on the equilibrium solubility of FFA were shown in Fig. 2. The equilibrium solubility of FFA at 37°C measured in PBS was  $9.0 \pm 0.7$   $\mu\text{g/mL}$ . PVP had a moderate effect, which the solubility of FFA was slightly increased with the PVP concentration of 1000  $\mu\text{g/mL}$  and it reached the maximum of  $14.9 \pm 1.0$   $\mu\text{g/mL}$  at the PVP concentration of 2000  $\mu\text{g/mL}$ . In contrast, the effect of PVP-VA could be seen at the concentration of 1000  $\mu\text{g/mL}$  and it became more prominent with the increase of the concentration. In fact, the solubility of FFA reached  $17.2 \pm 1.6$   $\mu\text{g/mL}$  at a PVP-VA concentration of 4000  $\mu\text{g/mL}$  and  $37.6 \pm 0.5$   $\mu\text{g/mL}$  at a PVP-VA concentration of 16000  $\mu\text{g/mL}$ .

Whereas the presence of NIC seemed to have no effect, the presence of TP at a concentration of 71.1  $\mu\text{g/mL}$  increased the equilibrium solubility of FFA to  $10.24 \pm 0.71$   $\mu\text{g/mL}$ . Neither NIC nor TP altered the equilibrium solubility of FFA in the solution containing PVP. In contrast, the presence of NIC or TP further enhanced the solubility FFA observed in solutions containing PVP-VA.

**3.2. Effect of Polymers on the Dissolution and Permeation of FFA Cocrystals.** The dialysis membrane acted as a selective barrier for the substances to pass through based on their sizes. The MWCO of the dialysis membrane used in the study was 6-8 KDa and it would thus allow the transfer of FFA, which has a molecular weight (MW) of 281 Da, from the donor compartment

*3.2.1. Dissolution and Permeation Performances of FFA, FFA-NIC and FFA-TP in PBS.* Fig. 3 (a) showed the dissolution of FFA, FFA-NIC and FFA-TP obtained from the D/P system. It could be seen that FFA-NIC had the fastest dissolution and FFA reached its maximum concentration,  $8.8 \pm 0.2$   $\mu\text{g/mL}$ , within the first 5 min. After

1  
2  
3 that the FFA concentration measured for FFA-NIC gradually dropped and the  
4  
5 minimum concentration,  $7.5 \pm 0.3 \mu\text{g/mL}$ , was obtained at 120 min. In contrast, the  
6  
7 concentrations measured for FFA and FFA-TP slowly increased to the equilibrium  
8  
9 solubility of FFA,  $9.0 \pm 0.3 \mu\text{g/mL}$ , at 60 min and remained that concentration  
10  
11 afterwards. The quantitative comparison of DPPs<sup>4</sup> were shown in Fig. 4 (a), which  
12  
13 FFA-NIC had a lower DDP than that of FFA and FFA-TP.  
14  
15

16 Fig. 3 (b) showed the permeation of FFA, FFA-NIC and FFA-TP obtained from the  
17  
18 D/P system. In the first 60 min, the gradually increased FFA concentrations measured  
19  
20 for all three solids overlapped. After that, the concentrations of FFA measured at all  
21  
22 three occasions increased steadily and FFA and FFA-TP had a slightly greater rate  
23  
24 than that of FFA-NIC. This was in consistent with the mass flux rates shown in Fig. 4  
25  
26 (b).  
27  
28

29 *3.2.2. Effect of PVP or PVP-VA on the Dissolution and Permeation of FFA, FFA-*  
30  
31 *NIC and FFA-TP.* Fig. 5 demonstrated the effect of PVP on the dissolution of FFA,  
32  
33 FFA-NIC and FFA-TP in the donor compartment and the subsequent permeation of  
34  
35 FFA to the acceptor compartment. Both FFA-NIC and FFA-TP showed an enhanced  
36  
37 dissolution in comparison to their parent drug FFA when PVP was added to the  
38  
39 dissolution media. A supersaturated state of FFA was observed for FFA-TP and FFA-  
40  
41 NIC in the dissolution media with PVP at  $200 \mu\text{g/mL}$ , where the concentration of  
42  
43 FFA reached  $16.7 \mu\text{g/mL}$  for FFA-TP at 5 min and  $12.2 \mu\text{g/mL}$  for FFA-NIC at 15  
44  
45 min [Fig. 5(a1)]. The concentrations of FFA observed for FFA-NIC and FFA-TP were  
46  
47 then gradually reduced to the parent drug's solubility at the 60 min point [Fig. 5(a1)].  
48  
49 Correspondingly, the presence of PVP increased the amount of FFA detected in the  
50  
51 acceptor compartment to 1.23-fold for FFA-TP and 1.17-fold for FFA-NIC [Fig. 4  
52  
53 (b)], but decreased the amount of FFA detected at the acceptor compartment by 15.4%  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 for the parent FFA [Fig. 5 (b1) and Fig. 4 (b)]. All these phenomena suggested that  
4 the inclusion of PVP in the dissolution media increased the dissolution and  
5 permeation of FFA cocrystals, whereas it slowed down the dissolution as well as the  
6 permeation of the parent FFA.  
7  
8  
9

10  
11 When PVP was added to a concentration of 4000  $\mu\text{g/mL}$ , the concentration of FFA  
12 detected in the donor compartment for FFA-TP reached 40  $\mu\text{g/mL}$  at 15 min, which  
13 was approximately 4.5-fold of the solubility of FFA [Fig. 5 (a2)]. And the  
14 concentration of FFA detected for FFA-NIC at the same time was 23.6  $\mu\text{g/mL}$ . These  
15 indicated that FFA-TP had a better dissolution than FFA-NIC. On the other hand, the  
16 concentration of FFA detected for FFA-NIC in the donor compartment quickly  
17 decreased to 19.5  $\mu\text{g/mL}$  at 15 min and remained constant afterwards. In contrast, the  
18 concentration of FFA detected in the donor compartment for FFA-TP decreased  
19 slowly from 40  $\mu\text{g/mL}$  and stabilized to 28.8  $\mu\text{g/mL}$  after 60 min. Consequently, the  
20 DPP of FFA-TP was 228% higher than that of FFA and was 128% higher than that of  
21 FFA-NIC [Fig. 4 (a)]. In accordance with these, the amount of FFA detected at the  
22 acceptor compartment increased by 1.69-fold in the case of FFA-TP and by 1.27-fold  
23 in the case of FFA-NIC [Fig. 5 (b2) and Fig. 4 (b)].  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 Interestingly, a further increased concentration of PVP to 16000  $\mu\text{g/mL}$  diminished  
41 the dissolution of FFA-TP and FFA-NIC, as shown in Figs. 5 (a3) and Fig. 4 (a). The  
42 maximal concentration of FFA detected in the donor compartment was 27.5  $\mu\text{g/mL}$   
43 for FFA-TP and 19.7  $\mu\text{g/mL}$  for FFA-NIC. Meanwhile, the amount of FFA permeated  
44 cross the membrane reduced significantly for FFA-TP and FFA-NIC [Fig. 5 (b3)].  
45 And the flux rate of the FFA-TP was 0.62-fold of the FFA observed in 0.01 M PBS  
46 and 0.45-fold for the FFA-NIC [Fig. 5 (b3) and Fig. 4 (b)]. In addition, in comparison  
47 to those measured in PBS, 16000  $\mu\text{g/mL}$  of PVP significantly slowed down the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 dissolution of FFA [Fig. 5 (a3)], as demonstrated by a 9% reduction of DPP [Fig. 4  
4  
5 (a)] and a just 0.34-fold of flux rate [Fig. 5 (b3) and Fig. 4 (b)].  
6

7 The dissolution and permeation measured for FFA and FFA cocrystals in the  
8 presence of PVP-VA were shown in Fig. 6. The dissolution of FFA cocrystals was  
9 significantly enhanced by PVP-VA. At a PVP-VA concentration of 200  $\mu\text{g/mL}$ , a  
10 33.8% increase of DPP was observed for FFA-NIC and a 42.6% increase of DPP was  
11 observed for FFA-TP [Fig. 6 (a1) and Fig. 4 (a)]. At a PVP-VA concentration of  
12 16000  $\mu\text{g/mL}$ , the DPP observed was increased to 460.5% for FFA-NIC and 409.4%  
13 for FFA-TP [Fig. 6 (a3) and Fig. 4 (a)]. However, the amount of FFA detected at the  
14 acceptor compartment seemed not to be proportional to the corresponding DPPs [Fig.  
15 6 (b3) and Fig. 4 (b)]. Actually, the FFA flux rate of FFA and FFA cocrystals were  
16 reduced with the increased concentration of PVP-VA. In spite of that, the FFA flux  
17 rates measured for FFA-NIC and FFA-TP at the PVP-VA concentration of 200  $\mu\text{g/mL}$   
18 or 4000  $\mu\text{g/mL}$  were still higher than that measured for FFA in the absence of PVP-  
19 VA [Fig. 4 (b)]. Whereas, when the added PVP-VA reached a concentration of 16000  
20  $\mu\text{g/mL}$  the flux rate measured for FFA-NIC or FFA-TP was lower than that measured  
21 for FFA in the absence of PVP-VA [Fig. 4 (b)].  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

### 40 **3.3. NMR Analysis of Interactions Between FFA, Coformers and Polymers.**

41 The  $^1\text{H}$  NMR experiments were conducted to investigate the interactions between  
42 FFA, coformers (NIC and TP) and polymers (PVP and PVP-VA) in solution, and the  
43 tests were summarized in Table S1 of the Supplementary Information. Assignments of  
44 the  $^1\text{H}$  chemical shifts of FFA, NIC, TP, PVP and PVP-VA were carried out  
45 previously.<sup>49, 53-57</sup> The focus of this work was to examine the characteristic chemical  
46 shifts of the protons in each of the molecules of FFA, TP, and NIC. These shifts could  
47 be due to the hydrogen bonding or self-aggregation in solution, including i) the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 chemical shifts of singlet peak of H<sub>j</sub> and two doublet peaks of H<sub>h</sub> in FFA; ii) the  
4  
5 chemical shifts of H<sub>a</sub>, H<sub>b</sub> and H<sub>c</sub> in NIC; and iii) the chemical shifts of H<sub>a</sub>, H<sub>b</sub>, H<sub>c</sub> and  
6  
7 H<sub>d</sub> in TP. The detailed information regarding the protons in each molecule could be  
8  
9 found in Table 1 and the full <sup>1</sup>H NMR spectra of the experiments could be found in  
10  
11 Fig. S1 of the Supplementary Information.

12  
13 *3.3.1. <sup>1</sup>H NMR Analysis of Single Component of FFA, NIC or TP.* Fig. 7 (a) showed  
14  
15 the <sup>1</sup>H NMR chemical shifts of H<sub>j</sub> and H<sub>h</sub> in FFA and they seemed to be related to the  
16  
17 concentrations of FFA. The existence of the hydrogen bond between the NH<sub>j</sub> group  
18  
19 and the carbonyl O of the carboxylic acid within FFA had been well illustrated.<sup>54</sup>  
20  
21 With an increased concentration, the self-association of FFA could also occur through  
22  
23 the inter carboxylic acid hydrogen bondings, which was demonstrated by a downfield  
24  
25 chemical shift of H<sub>h</sub> from 8.049 ppm at 500 μg/mL to 8.061 ppm at 1000 μg/mL, and  
26  
27 to 8.082 ppm at 5000 μg/mL. This could be due to proton deshielding by the  
28  
29 hydrogen bonds, which the lengthening of the O-H bond reduced the valence electro  
30  
31 density around the proton of H<sub>h</sub>. The intermolecular association between the dimer of  
32  
33 FFAs did not affect the molecular conformation in solution and the intramolecular N-  
34  
35 H<sub>j</sub>⋯O=C was not disrupted.<sup>54</sup> Therefore, at an increased concentration of FFA, the  
36  
37 peaks of H<sub>j</sub> displayed an upfield shift from 9.463 ppm at 500 μg/mL to 9.443 ppm at  
38  
39 1000 μg/mL, and to 9.410 ppm at 5000 μg/mL. The self-association of NIC in  
40  
41 solution occurred through inter-amide hydrogen bonding but not the stacking of the  
42  
43 pyridine rings.<sup>58</sup> Therefore, with an increased NIC concentration, the downfield shifts  
44  
45 of H<sub>a</sub> and H<sub>b</sub> were observed. This was in contrast to H<sub>c</sub>, whose spectra was almost  
46  
47 constant ( $\Delta\delta = 0.001$  at 430 μg/mL and  $\Delta\delta = 0.005$  at 2150 μg/mL) [Fig. 7(b)]. The  
48  
49 self-association of TP occurred through the asymmetric dimer involving hydrogen  
50  
51 bonding of H<sub>b</sub>⋯O of C-CO-N-CH<sub>d</sub> and C=N⋯H<sub>a</sub>N-C.<sup>59</sup> Subsequently, all of the <sup>1</sup>H  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 spectra of  $H_a$  and  $H_b$  of TP moved to a lower field with an increased concentration,  
4  
5 whereas the  $^1H$  chemical shifts of  $H_c$  and  $H_d$  remained constants ( $\Delta\delta = 0.003$  of  $H_c$   
6  
7 and  $\Delta\delta = 0.005$  of  $H_d$  at  $640 \mu\text{g/mL}$  and  $\Delta\delta = 0.013$  of  $H_c$  and  $\Delta\delta = 0.020$  of  $H_d$  at  
8  
9  $3200 \mu\text{g/mL}$ ) [Fig. 7(c)].

10  
11 The  $^1H$  NMR chemical shifts of the polymer PVP or PVP-VA are shown in  
12  
13 supporting information in Fig. S1, which were in the range from 4.000 ppm to 1.000  
14  
15 ppm and did not overlap with the peaks of FFA, NIC/TP. Therefore the change of  
16  
17 chemical shifts of FFA, NIC/TP in the presence of a polymer was caused by the  
18  
19 interaction among molecules.  
20  
21

22  
23 *3.3.2.  $^1H$  NMR Analysis of Binary Components.* Samples of FFA, NIC and TP were  
24  
25 analysed for  $^1H$  NMR in the presence of PVP or PVP-VA to identify their self-  
26  
27 association properties and their interactions with PVP or PVP-VA (Fig. 8). The  
28  
29 concentrations used were  $1000 \mu\text{g/mL}$  for FFA,  $430 \mu\text{g/mL}$  for NIC and  $640 \mu\text{g/mL}$   
30  
31 for TP.  
32

33  
34 In the presence of PVP or PVP-VA at a concentration of  $200 \mu\text{g/mL}$ , the peaks of  
35  
36  $H_h$  in FFA had a downfield change, indicating the existence of hydrogen bonding  
37  
38 between O-H of the carboxylic acid function group in FFA and the carbonyl O group  
39  
40 in PVP or PVP-VA. As O=C of the carboxylic acid function group in FFA did not  
41  
42 interact with PVP or PVP-VA, the intramolecular attraction of  $N-H_j \cdots O=C$  in FFA  
43  
44 was not disrupted. This was indicated by a constant spectra of  $H_j$  in FFA in the  
45  
46 presence or absence of PVP or PVP-VA [Fig. 8 (a)]. When the concentration of PVP  
47  
48 or PVP-VA was increased to  $5000 \mu\text{g/mL}$ , the peaks of  $H_j$  and  $H_h$  in FFA moved in  
49  
50 opposite directions: an upfield (shielding) shift for  $H_h$  and a down field shift for  $H_j$ .  
51  
52 This suggested a different mechanism for the interactions between the FFA and the  
53  
54 PVP or PVP-VA polymer. The upfield shift of  $H_h$  might associate with the transition  
55  
56  
57  
58  
59  
60

1  
2  
3 of a solution environment from hydrophilic to hydrophobic as a result of the increased  
4 concentration of PVP/PVP-VA, leading to hydrophobic interactions between FFA and  
5 PVP/PVP-VA.<sup>57, 60</sup> Obviously, the hydrophobic interaction would affect the  
6 conformation of FFA in solution, resulting in change of the intramolecular attraction  
7 of N-H<sub>j</sub>...O=C and a consequent observation of the downfield shift of H<sub>j</sub>. Furthermore,  
8 the broader <sup>1</sup>H NMR peaks observed for FFA in solution containing PVP was in  
9 contrast to those observed in solution containing PVP-VA, indicating that PVP could  
10 significantly suppress the mobility of FFA molecules in comparison to PVP-VA.  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 The peaks of NIC were not affected by PVP or PVP-VA at a concentration of 200  
21 μg/mL [Fig. 8 (b)], implying that there was no or weak interaction between NIC and  
22 PVP or PVP-VA. When the concentration of PVP or PVP-VA was increased to 5000  
23 μg/mL, the hydrogen bonding formed between NIC and PVP or PVP-VA was  
24 demonstrated as the downfield shifts of <sup>1</sup>H NMR peaks of NIC.  
25  
26  
27  
28  
29  
30

31 <sup>1</sup>H NMR spectra of TP were complex. It could be affected by the interactions  
32 between solute and solvent, solute and solute and solute and polymer. Moreover,  
33 dimerization of TP molecules was found to be the dominating factor that affected the  
34 <sup>1</sup>H chemical shifts of TP in CDCl<sub>3</sub>.<sup>59</sup> TP was recognised as good proton donor and the  
35 dimers of TPs could be disrupted by PVP or PVP-VA in solution. This was proven by  
36 the downfield shifts of <sup>1</sup>H NMR peaks of TP at a polymer concentration of 200  
37 μg/mL [Fig. 8 (c)]. Interestingly, further increased concentration of the polymers  
38 caused an upfield (shielding) shifts of H<sub>b</sub>, H<sub>d</sub> and H<sub>c</sub> in TP.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Within the structure of a solid crystalline FFA-NIC, the hydrogen bonds formed  
50 between two FFA and two NIC molecules. In detail, the H<sub>i</sub> of amide acid in FFA  
51 formed hydrogen bond with the aromatic N of one of the NIC, and the carbonyl O  
52 atom of the acid of the same FFA formed another hydrogen bond with the amide H<sub>a</sub> or  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  $H_b$  of the other NIC.<sup>34</sup> It was speculated that, as those formed in solid crystalline, the  
4 same interactions might form between FFA and NIC in  $CDCl_3$  solution. This was  
5 based on the facts of the deshielded chemical shifts of the  $H_h$  in FFA and the  
6 downfield  $H_c$ ,  $H_a$  and  $H_b$  in NIC [Fig. 9 (a)]. As a result of the formation of hydrogen  
7 bond between the carbonyl O atom of the acid in FFA and NIC, the intramolecular  
8 attraction of  $N-H_j \cdots O=C$  was disrupted and the chemical shift of  $H_j$  in FFA had a  
9 downfield movement.  
10  
11  
12  
13  
14  
15  
16  
17

18 It was suggested that FFA and TP could form a dimer through the  $H_i$  of amide acid  
19 in FFA and the aromatic N in TP.<sup>35</sup> This was in consistent with the finding of the  
20 significant downfield chemical shift of the  $H_h$  in FFA [Fig. 9(b)]. Because the  
21 carbonyl O atom of the acid of FFA was not involved in the intermolecular interaction  
22 with TP, the intramolecular attraction of  $N-H_j \cdots O=C$  of FFA was not disrupted, as  
23 shown by the constant spectra of  $H_j$ .  
24  
25  
26  
27  
28  
29  
30

31 *3.3.3.  $^1H$  NMR Analysis of Ternary Components.*  $^1H$  NMR spectra of the mixed  
32 samples of FFA, NIC/TP and PVP/PVP-VA were shown in Fig. 10. When the  
33 concentration of PVP was set to 200  $\mu g/mL$ , the characteristic peaks of the  $H_h$  and  $H_j$   
34 in FFA moved upfield, suggesting that, as a result of the interactions between the FFA  
35 and the PVP, the hydrogen bonds formed between the FFA and NIC were disrupted  
36 [Fig. 10 (a)]. In contrast, PVP-VA at 200  $\mu g/mL$  had no effect on the spectra of the  $H_h$   
37 and  $H_j$  in FFA, indicating that the interaction between the FFA and NIC played a  
38 dominant role over the interaction between the FFA and PVP-VA [Fig. 10 (b)].  
39 Increasing PVP or PVP-VA to a concentration of 5000  $\mu g/mL$  rendered the peaks of  
40  $H_j$  and  $H_h$  moved to opposite directions [Fig. 10 (a) and Fig. 10-(b)]. This might  
41 reflect the dominant effect caused by the hydrophobic interaction between the FFA  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 and PVP or PVP-VA in comparison to the hydrogen bonds formed between the FFA,  
4  
5 NIC and PVP or PVP-VA.  
6

7  
8 When 200  $\mu\text{g/ml}$  of PVP/PVP-VA was mixed with FFA and TP, the spectra of the  
9  
10  $H_h$  in FFA slightly moved upfield while that of the  $H_j$  almost had no change. It  
11  
12 suggested that the hydrogen bonds formed between the FFA and TP were disturbed by  
13  
14 the PVP or PVP-VA [Fig. 10 (c) and Fig. 10 (d)]. At a concentration of 5000  $\mu\text{g/mL}$ ,  
15  
16 the PVP or PVP-VA rendered the peaks of the  $H_j$  and the  $H_h$  to opposite directions  
17  
18 [Fig. 10 (c) and Fig. 10 (d)]. This was similar to those observed for the mixture of  
19  
20 FFA and NIC and, once more, demonstrated the dominating hydrophobic interactions  
21  
22 between the FFA and PVP or PVP-VA.  
23

#### 24 **4. DISCUSSION**

25  
26 This research demonstrated that the supersaturated drug concentration generated by  
27  
28 the cocrystal dissolution could be maintained through inclusion of a polymeric  
29  
30 excipient in the formulation, for example, the supersaturated concentration of FFA  
31  
32 generated by dissolution of FFA-NIC or FFA-TP was maintained by PVP or PVP-  
33  
34 VA [Fig. 4 (a)].<sup>4,9,10</sup> On the other hand, in the presence of PVP or PVP-VA, the DPP  
35  
36 detected for FFA-NIC or FFA-TP was proportional to the concentration of the  
37  
38 polymers and it was much higher than that detected without PVP or PVP-VA [Fig. 4  
39  
40 (a)]. However, as shown in the D/P system, an elevated concentration of FFA in the  
41  
42 donor compartment was not accompanied by an increased flux rate of FFA to the  
43  
44 acceptor compartment [Fig. 4 (b)]. This erratic drug permeation could lead to poor  
45  
46 drug bioavailability, which was crucially related to the amount of drug molecules that  
47  
48 could pass through the GI membrane. Therefore, it was important to explore the  
49  
50 interplay of the solubility and permeability in the cocrystal based formulation, that  
51  
52 enhanced solubility/dissolution might have positive, negative or no effect on the  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 subsequent permeability.<sup>18-20, 29</sup> The mechanism of how a polymer affected the  
4  
5 dissolution of the cocrystals had been investigated, and it seemed that the polymer  
6  
7 could interact with the surface of the cocrystal to alter both the dissolution and  
8  
9 precipitation.<sup>4</sup> As a further investigation, this research revealed the interactions  
10  
11 between the drug substance (FFA), the coformer (NIC or TP) and the polymer (PVP  
12  
13 or PVP-VA) in solution and how these interactions maintain the supersaturated drug  
14  
15 concentration and the subsequent permeation of the drug substance.  
16  
17

18 Both PVP and PVP-VA are water soluble polymers. However, due to their  
19  
20 structural difference (Table 1), the intermolecular interactions between PVP/PVP-VA  
21  
22 and the drug substance FFA or coformer (NIC or TP) were different. This could affect  
23  
24 not only the nucleation and growth kinetics<sup>9</sup> but also the permeation of FFA in  
25  
26 solution [Fig. 4 (b)]. Interestingly, at the same concentration, PVP-VA always  
27  
28 rendered a higher DPP for FFA-NIC and FFA-TP than PVP [Fig. 4(a)]. In contrary,  
29  
30 the effect of PVP-VA on the flux rate of FFA-NIC or FFA-TP could be greater or  
31  
32 smaller than PVP, which was seemingly concentration dependent (Fig. 4). In fact, for  
33  
34 FFA-NIC and FFA-TP, while the FFA flux rates and DPPs measured were in  
35  
36 proportion to the concentration of PVP, the FFA flux rates measured were in contrast  
37  
38 to the DPPs in the presence of PVP-VA (Fig. 4). Therefore, it was essential to take  
39  
40 into account both the flux rate and DPP when evaluating a cocrystal based  
41  
42 formulation.  
43  
44  
45

46 Based on the results in Fig.2, PVP could act as a stabilizing agent in the formulation  
47  
48 as the equilibrium solubility of FFA was not increased significantly across the whole  
49  
50 range of the concentrations of PVP. The hydrogen bonds between FFA and PVP  
51  
52 should be the dominant force when the concentration of PVP was below 4000 µg/mL,  
53  
54 which was confirmed by the downfield chemical shifts of  $H_h$  in FFA [Fig. 8 (a)]. The  
55  
56  
57  
58  
59  
60

1  
2  
3 flux rate was determined by the degree of supersaturation (ratio of dissolved amount  
4 of unchanged API to its thermodynamic solubility), which was the driving force for  
5 transporting the drug substance across the membrane.<sup>29</sup> PVP might inhibit the  
6 nucleation of FFA to maintain its supersaturated concentration [Fig. 5 (a)]<sup>9</sup> and to  
7 increase the flux rate of FFA across the regenerated cellulose membrane [Fig.5 (b)].<sup>29</sup>  
8  
9  
10  
11  
12  
13 Due to the bulk precipitation dissolution mechanism, the higher supersaturation  
14 concentration of FFA generated by FFA-TP led to a higher flux rate in comparison to  
15 FFA-NIC.<sup>4</sup>  
16  
17  
18  
19

20  
21 When the concentration of PVP was above 4000  $\mu\text{g/mL}$ , the attraction between  
22 FFA and PVP changed into hydrophobic interaction due to the confined spaces  
23 between the FFA and PVP molecules. This was confirmed by the upfield chemical  
24 shifts of  $H_h$  in FFA [Fig. 8 (a)]. It must be stressed that, at such a concentration,  
25 micellization of PVP did not occur due to its rigorous backbone structure and,  
26 therefore, PVP had no significant effect on the equilibrium solubility of FFA (Fig. 2).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

20  
21 The reduced flux rate of FFA at a PVP concentration of 16000  $\mu\text{g/mL}$  [Fig. 4 (b)]  
22 was the result of a combined effects of the reduced dissolution of the cocrystals [Fig.  
23 5 (a)] and the reduced solute activity coefficient.<sup>37</sup> Although the interaction between  
24 FFA and PVP was disturbed by the interaction between PVP and the cofomers (Fig.  
25 9 and Fig. 10), the role of PVP as a stabilizing agent did not change, as demonstrated  
26 by the minor change of FFA solubility with or without the presence of the cofomers  
27 (Fig. 2). As a consequence, the flux rate of FFA solely depended on the DPP of the  
28 cocrystals, which the maximal flux rate of FFA was observed for FFA-TP in the  
29 presence of 4000  $\mu\text{g/mL}$  of PVP.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 As seen in Fig.2, PVP-VA had a transitional role in the cocrystal formulation,  
4 depending on its concentrations. At a concentration of up to 1000  $\mu\text{g/mL}$  and the  
5 absence of NIC or TP, PVP-VA acted as a stabilizing agent as it had no effect on the  
6 equilibrium solubility of FFA (Fig. 2). The downfield shifts of the  $H_h$  in FFA  
7 indicated the formation of hydrogen bond between the FFA and the PVP-VA [Fig. 8  
8 (a)]. Compared to PVP, PVP-VA was more hydrophobic and flexible, leading to  
9 micellization at a higher concentration.<sup>61</sup> This research set the critical micelle  
10 concentration (CMC) of PVP-VA at 4000  $\mu\text{g/mL}$ . Therefore, at a concentration over  
11 the CMC, PVP-VA significantly increased the solubility of FFA. The  $^1\text{H}$  NMR  
12 spectra of FFA showed the upfield shifts of the  $H_h$  at PVP-VA concentration of 5000  
13  $\mu\text{g/mL}$  [Fig. 8 (a)], from which the conclusion could be drawn that the encapsulation  
14 of FFA took place in PVP-VA micelles. In other words, the role of PVP-VA changed  
15 to a solubilisation agent in the formulation.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 In the presence of NIC or TP, PVP-VA demonstrated a significant solubilisation  
32 capacity (Fig. 2). It was likely that the association of PVP-VA with a cofomer , as  
33 shown in Fig. 8 (b) and Fig. 8(c), provided an expanded region of the inner core for  
34 drug solubilization.<sup>62</sup> The CMC of PVP-VA reduced significantly around 200  $\mu\text{g/mL}$ ,  
35 at which point the equilibrium solubility of FFA started to increase (Fig. 2). Taken  
36 together, it seemed that PVP-VA behaved as a solubilisation agent in the cocrystal  
37 formulations regardless of its concentrations.  
38  
39  
40  
41  
42  
43  
44  
45

46 Corresponding to the increased equilibrium solubility of FFA, the DPPs of the FFA  
47 cocrystals were proportional to the concentration of PVP-VA [Fig. 4 (a)]. From the  
48 perspective of thermodynamics, an elevated drug concentration due to the presence of  
49 PVP-VA was fundamentally different from the drug supersaturation generated by the  
50 stabilizing agent of PVP. Solubilizing agent, such as PVP-VA, affected the  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 concentrations by increasing the equilibrium solubility rather than increasing the  
4  
5 chemical potential (degree of supersaturation), as shown by the reduced flux rate of  
6  
7 FFA across the membrane [Fig. 4 (b)]. This was in consistent with the findings from  
8  
9 previous studies.<sup>18-20, 29</sup>  
10

11 Overall, the permeation of a FFA cocrystal could be affected by a series of factors,  
12  
13 such as the types of polymers, the concentrations of the polymers and cofomers. In  
14  
15 particular, the interactions between the polymer and the cofomer might play roles in  
16  
17 improving the absorption of the drug substance.  
18  
19

## 20 5. CONCLUSION

21  
22 The dissolution and permeation of the cocrystals, FFA-NIC and FFA-TP, in the  
23  
24 presence of polymers, PVP and PVP-VA, were examined using a  
25  
26 dissolution/permeation (D/P) system. It showed that the types of polymers, their  
27  
28 concentrations and interactions with cofomers were the determining factors. The role  
29  
30 of PVP as a stabilizing agent was not altered in spite of its interactions with the  
31  
32 cofomers of NIC and TP, which was supported by the fact that the flux rate of FFA  
33  
34 solely depended on the DPP of the cocrystals. With an appropriate PVP concentration,  
35  
36 the maximal flux rate of FFA could be obtained for a given FFA cocrystal. The role of  
37  
38 PVP-VA could change due to its association with the cofomers, such as from a  
39  
40 stabilizing agent to a solubilisation agent. It was also found that PVP-VA rendered the  
41  
42 measurement of flux rate contrasting to the measurement of the corresponding DPP.  
43  
44 The maximal flux rate of FFA could only be obtained at a very low concentration of  
45  
46 PVP-VA. At last, <sup>1</sup>H NMR provided evidence regarding the molecular interactions  
47  
48 between the FFA, the cofomers and the polymers at the atomic level. In conclusion,  
49  
50 understanding the relationships between the drug substance, cofomers and polymers  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 at molecular level was the key to optimise the solubility and permeability to maximize  
4 the performance of cocrystal based oral drug products.  
5

## 6 7 ASSOCIATED CONTENT

### 8 9 10 Supporting Information

- 11  
12  
13 1) **Table S1.** List of  $^1\text{H}$  NMR samples.
- 14  
15  
16 2) **Figure S1.** All  $^1\text{H}$  NMR results.  
17

## 18 19 AUTHOR INFORMATION

### 20 21 22 Corresponding Author

23  
24 \*E-mail: mli@dmu.ac.uk; Tel: +44-1162577132.  
25  
26

### 27 28 Notes

29  
30 The authors declare no competing financial interest.  
31  
32

## 33 34 ACKNOWLEDGEMENTS

35  
36 We would like to thank the financial support of the work by UK Engineering and  
37 Physical Sciences Research Council (EPSRC, EP/R021198/1) and Science and  
38 Technology Project of Hebei Province of China (NO. 16211505). De Montfort  
39 University and the Great Britain-China Educational Trust are gratefully acknowledged  
40 for providing scholarships for Miss Minshan Guo to conduct her PhD study. We also  
41 thank Dr. Ketan Ruparelia at De Montfort University for his help in  $^1\text{H}$  NMR  
42 experiments.  
43  
44  
45  
46  
47  
48  
49  
50

## 51 52 REFERENCES

1. Duggirala, N. K.; Perry, M. L.; Almarsson, O.; Zaworotko, M. J., Pharmaceutical cocrystals: along the path to improved medicines. *Chem. Commun. (Camb.)* **2016**, *52*, 640-55.
2. Kuminek, G.; Cao, F.; da Rocha, A. B. d. O.; Cardoso, S. G.; Rodríguez-Hornedo, N., Cocrystals to facilitate delivery of poorly soluble compounds beyond-rule-of-5. *Adv. Drug. Deliv. Rev.* **2016**, *101*, 143-166.
3. Qiao, N.; Li, M.; Schlindwein, W.; Malek, N.; Davies, A.; Trappitt, G., Pharmaceutical cocrystals: an overview. *Int. J. Pharm.* **2011**, *419*, 1-11.
4. Guo, M.; Wang, K.; Qiao, N.; Fábíán, L.; Sadiq, G.; Li, M., Insight into Flufenamic Acid Cocrystal Dissolution in the Presence of a Polymer in Solution: from Single Crystal to Powder Dissolution. *Molecular Pharmaceutics* **2017**, *14*, 4583-4596.
5. Qiao, N.; Wang, K.; Schlindwein, W.; Davies, A.; Li, M., In situ monitoring of carbamazepine–nicotinamide cocrystal intrinsic dissolution behaviour. *European Journal of Pharmaceutics and Biopharmaceutics* **2013**, *83*, 415-426.
6. Remenar, J. F.; Peterson, M. L.; Stephens, P. W.; Zhang, Z.; Zimenkov, Y.; Hickey, M. B., Celecoxib:Nicotinamide Dissociation: Using Excipients To Capture the Cocrystal's Potential. *Molecular Pharmaceutics* **2007**, *4*, 386-400.
7. Yamashita, H.; Sun, C. C., Harvesting Potential Dissolution Advantages of Soluble Cocrystals by Depressing Precipitation Using the Common Cofomer Effect. *Crystal Growth & Design* **2016**, *16*, 6719-6721.
8. Cao, F.; Amidon, G. L.; Rodriguez-Hornedo, N.; Amidon, G. E., Mechanistic Analysis of Cocrystal Dissolution as a Function of pH and Micellar Solubilization. *Molecular Pharmaceutics* **2016**, *13*, 1030-1046.
9. Guo, M.; Wang, K.; Hamill, N.; Lorimer, K.; Li, M., Investigating the Influence of Polymers on Supersaturated Flufenamic Acid Cocrystal Solutions. *Molecular Pharmaceutics* **2016**, *13*, 3292-3307.
10. Qiu, S.; Lai, J.; Guo, M.; Wang, K.; Lai, X.; Desai, U.; Juma, N.; Li, M., Role of polymers in solution and tablet-based carbamazepine cocrystal formulations. *CrystEngComm* **2016**, *18*, 2664-2678.
11. Li, M.; Qiu, S.; Lu, Y.; Wang, K.; Lai, X.; Rehan, M., Investigation of the Effect of Hydroxypropyl Methylcellulose on the Phase Transformation and Release Profiles of Carbamazepine-Nicotinamide Cocrystal. *Pharm Res* **2014**, *31*, 2312-2325.
12. Crowley, K. J.; Zografí, G., The Effect of Low Concentrations of Molecularly Dispersed Poly(Vinylpyrrolidone) on Indomethacin Crystallization from the Amorphous State. *Pharm Res* **2003**, *20*, 1417-1422.
13. DiNunzio, J. C.; Miller, D. A.; Yang, W.; McGinity, J. W.; Williams, R. O., Amorphous Compositions Using Concentration Enhancing Polymers for Improved Bioavailability of Itraconazole. *Mol. Pharm.* **2008**, *5*, 968-980.
14. Ullah, M.; Raza Shah, M.; Asad, H. B.; Hassham, M.; Farid Hasan, S.; Hussain, I., Improved in vitro and in vivo performance of carbamazepine enabled by using a succinic acid cocrystal in a stable suspension formulation. *Pak. J. Pharm. Sci.* **2017**, *30*.
15. Ullah, M.; Ullah, H.; Murtaza, G.; Mahmood, Q.; Hussain, I., Evaluation of influence of various polymers on dissolution and phase behavior of carbamazepine-succinic acid cocrystal in matrix tablets. *BioMed research international* **2015**, *2015*.
16. Ullah, M.; Hussain, I.; Sun, C. C., The development of carbamazepine-succinic acid cocrystal tablet formulations with improved in vitro and in vivo performance. *Drug Dev. Ind. Pharm.* **2016**, *42*, 969-976.
17. Beig, A.; Lindley, D.; Miller, J. M.; Agbaria, R.; Dahan, A., Hydrotropic Solubilization of Lipophilic Drugs for Oral Delivery: The Effects of Urea and Nicotinamide on Carbamazepine Solubility–Permeability Interplay. *Frontiers in Pharmacology* **2016**, *7*, 379.

18. Dahan, A.; Beig, A.; Lindley, D.; Miller, J. M., The solubility–permeability interplay and oral drug formulation design: Two heads are better than one. *Advanced Drug Delivery Reviews* **2016**, *101*, 99-107.
19. Dahan, A.; Miller, J. M., The Solubility–Permeability Interplay and Its Implications in Formulation Design and Development for Poorly Soluble Drugs. *The AAPS Journal* **2012**, *14*, 244-251.
20. Beig, A.; Miller, J. M.; Lindley, D.; Carr, R. A.; Zocharski, P.; Agbaria, R.; Dahan, A., Head-To-Head Comparison of Different Solubility-Enabling Formulations of Etoposide and Their Consequent Solubility–Permeability Interplay. *Journal of Pharmaceutical Sciences* **2015**, *104*, 2941-2947.
21. Miller, J. M.; Beig, A.; Carr, R. A.; Webster, G. K.; Dahan, A., The solubility–permeability interplay when using cosolvents for solubilization: revising the way we use solubility-enabling formulations. *Mol. Pharm.* **2012**, *9*, 581-90.
22. Beig, A.; Miller, J. M.; Dahan, A., Accounting for the solubility–permeability interplay in oral formulation development for poor water solubility drugs: The effect of PEG-400 on carbamazepine absorption. *European Journal of Pharmaceutics and Biopharmaceutics* **2012**, *81*, 386-391.
23. Beig, A.; Miller, J. M.; Lindley, D.; Dahan, A., Striking the Optimal Solubility–Permeability Balance in Oral Formulation Development for Lipophilic Drugs: Maximizing Carbamazepine Blood Levels. *Molecular Pharmaceutics* **2017**, *14*, 319-327.
24. Fine-Shamir, N.; Beig, A.; Zur, M.; Lindley, D.; Miller, J. M.; Dahan, A., Toward Successful Cyclodextrin Based Solubility-Enabling Formulations for Oral Delivery of Lipophilic Drugs: Solubility–Permeability Trade-Off, Biorelevant Dissolution, and the Unstirred Water Layer. *Molecular Pharmaceutics* **2017**, *14*, 2138-2146.
25. Miller, J. M.; Beig, A.; Carr, R. A.; Spence, J. K.; Dahan, A., A Win–Win Solution in Oral Delivery of Lipophilic Drugs: Supersaturation via Amorphous Solid Dispersions Increases Apparent Solubility without Sacrifice of Intestinal Membrane Permeability. *Molecular Pharmaceutics* **2012**, *9*, 2009-2016.
26. Frank, K. J.; Rosenblatt, K. M.; Westedt, U.; Hölig, P.; Rosenberg, J.; Mägerlein, M.; Fricker, G.; Brandl, M., Amorphous solid dispersion enhances permeation of poorly soluble ABT-102: True supersaturation vs. apparent solubility enhancement. *International Journal of Pharmaceutics* **2012**, *437*, 288-293.
27. Frank, K. J.; Westedt, U.; Rosenblatt, K. M.; Hölig, P.; Rosenberg, J.; Mägerlein, M.; Fricker, G.; Brandl, M., What Is the Mechanism Behind Increased Permeation Rate of a Poorly Soluble Drug from Aqueous Dispersions of an Amorphous Solid Dispersion? *Journal of Pharmaceutical Sciences* **2014**, *103*, 1779-1786.
28. Beig, A.; Fine-Shamir, N.; Lindley, D.; Miller, J. M.; Dahan, A., Advantageous Solubility-Permeability Interplay When Using Amorphous Solid Dispersion (ASD) Formulation for the BCS Class IV P-gp Substrate Rifaximin: Simultaneous Increase of Both the Solubility and the Permeability. *The AAPS Journal* **2017**, *19*, 806-813.
29. Borbás, E.; Sinkó, B. I.; Tsinman, O.; Tsinman, K.; Kiserdei, E. v.; Démuth, B. z.; Balogh, A.; Bodák, B.; Domokos, A. s.; Dargó, G., Investigation and mathematical description of the real driving force of passive transport of drug molecules from supersaturated solutions. *Mol. Pharm.* **2016**, *13*, 3816-3826.
30. Sanphui, P.; Devi, V. K.; Clara, D.; Malviya, N.; Ganguly, S.; Desiraju, G. R., Cocrystals of Hydrochlorothiazide: Solubility and Diffusion/Permeability Enhancements through Drug–Cofomer Interactions. *Molecular Pharmaceutics* **2015**, *12*, 1615-1622.
31. Yan, Y.; Chen, J.-M.; Lu, T.-B., Simultaneously enhancing the solubility and permeability of acyclovir by crystal engineering approach. *CrystEngComm* **2013**, *15*, 6457-6460.

- 1  
2  
3 32. Ferretti, V.; Dalpiaz, A.; Bertolasi, V.; Ferraro, L.; Beggiato, S.; Spizzo, F.; Spisni, E.;  
4 Pavan, B., Indomethacin Co-Crystals and Their Parent Mixtures: Does the Intestinal Barrier  
5 Recognize Them Differently? *Molecular Pharmaceutics* **2015**, *12*, 1501-1511.
- 6 33. Guo, M.; Wang, K.; Qiao, N.; Fábíán, L.; Sadiq, G.; Li, M., Insight into Flufenamic Acid  
7 Cocrystal Dissolution in the Presence of a Polymer in Solution: from Single Crystal to Powder  
8 Dissolution. *Molecular Pharmaceutics* **2017**.
- 9 34. Fábíán, L.; Hamill, N.; Eccles, K. S.; Moynihan, H. A.; Maguire, A. R.; McCausland, L.;  
10 Lawrence, S. E., Cocrystals of Fenamic Acids with Nicotinamide. *Crystal Growth & Design*  
11 **2011**, *11*, 3522-3528.
- 12 35. Aitipamula, S.; Wong, A. B. H.; Chow, P. S.; Tan, R. B. H., Cocrystallization with  
13 flufenamic acid: comparison of physicochemical properties of two pharmaceutical cocrystals.  
14 *CrystEngComm* **2014**, *16*, 5793-5801.
- 15 36. Raina, S. A.; Zhang, G. G. Z.; Alonzo, D. E.; Wu, J.; Zhu, D.; Catron, N. D.; Gao, Y.;  
16 Taylor, L. S., Enhancements and Limits in Drug Membrane Transport Using Supersaturated  
17 Solutions of Poorly Water Soluble Drugs. *Journal of Pharmaceutical Sciences* **2014**, *103*,  
18 2736-2748.
- 19 37. Raina, S. A.; Zhang, G. G. Z.; Alonzo, D. E.; Wu, J.; Zhu, D.; Catron, N. D.; Gao, Y.;  
20 Taylor, L. S., Impact of Solubilizing Additives on Supersaturation and Membrane Transport of  
21 Drugs. *Pharm Res* **2015**, *32*, 3350-3364.
- 22 38. Kataoka, M.; Masaoka, Y.; Yamazaki, Y.; Sakane, T.; Sezaki, H.; Yamashita, S., In Vitro  
23 System to Evaluate Oral Absorption of Poorly Water-Soluble Drugs: Simultaneous Analysis on  
24 Dissolution and Permeation of Drugs. *Pharm Res* **2003**, *20*, 1674-1680.
- 25 39. Kataoka, M.; Masaoka, Y.; Sakuma, S.; Yamashita, S., Effect of Food Intake on the  
26 Oral Absorption of Poorly Water-Soluble Drugs: In Vitro Assessment of Drug Dissolution and  
27 Permeation Assay System. *Journal of Pharmaceutical Sciences* **2006**, *95*, 2051-2061.
- 28 40. Buch, P.; Langguth, P.; Kataoka, M.; Yamashita, S., IVIVC in oral absorption for  
29 fenofibrate immediate release tablets using a dissolution/permeation system. *Journal of*  
30 *Pharmaceutical Sciences* **2009**, *98*, 2001-2009.
- 31 41. Kataoka, M.; Sugano, K.; da Costa Mathews, C.; Wong, J. W.; Jones, K. L.; Masaoka, Y.;  
32 Sakuma, S.; Yamashita, S., Application of Dissolution/Permeation System for Evaluation of  
33 Formulation Effect on Oral Absorption of Poorly Water-Soluble Drugs in Drug Development.  
34 *Pharm. Res.* **2012**, *29*, 1485-1494.
- 35 42. Abu-Diak, O. A.; Jones, D. S.; Andrews, G. P., An Investigation into the Dissolution  
36 Properties of Celecoxib Melt Extrudates: Understanding the Role of Polymer Type and  
37 Concentration in Stabilizing Supersaturated Drug Concentrations. *Molecular Pharmaceutics*  
38 **2011**, *8*, 1362-1371.
- 39 43. Maniruzzaman, M.; Morgan, D. J.; Mendham, A. P.; Pang, J.; Snowden, M. J.;  
40 Douroumis, D., Drug-polymer intermolecular interactions in hot-melt extruded solid  
41 dispersions. *International Journal of Pharmaceutics* **2013**, *443*, 199-208.
- 42 44. Ueda, K.; Higashi, K.; Limwikrant, W.; Sekine, S.; Horie, T.; Yamamoto, K.; Moribe, K.,  
43 Mechanistic Differences in Permeation Behavior of Supersaturated and Solubilized Solutions  
44 of Carbamazepine Revealed by Nuclear Magnetic Resonance Measurements. *Molecular*  
45 *Pharmaceutics* **2012**, *9*, 3023-3033.
- 46 45. Ueda, K.; Higashi, K.; Yamamoto, K.; Moribe, K., Inhibitory Effect of Hydroxypropyl  
47 Methylcellulose Acetate Succinate on Drug Recrystallization from a Supersaturated Solution  
48 Assessed Using Nuclear Magnetic Resonance Measurements. *Molecular Pharmaceutics* **2013**,  
49 *10*, 3801-3811.
- 50 46. Ueda, K.; Higashi, K.; Yamamoto, K.; Moribe, K., Equilibrium State at Supersaturated  
51 Drug Concentration Achieved by Hydroxypropyl Methylcellulose Acetate Succinate:  
52 Molecular Characterization Using 1H NMR Technique. *Molecular Pharmaceutics* **2015**, *12*,  
53 1096-1104.
- 54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 47. Ueda, K.; Higashi, K.; Moribe, K., Direct NMR Monitoring of Phase Separation  
4 Behavior of Highly Supersaturated Nifedipine Solution Stabilized with Hypromellose  
5 Derivatives. *Molecular Pharmaceutics* **2017**, *14*, 2314-2322.
- 6 48. Ueda, K.; Higashi, K.; Yamamoto, K.; Moribe, K., In situ molecular elucidation of drug  
7 supersaturation achieved by nano-sizing and amorphization of poorly water-soluble drug.  
8 *European Journal of Pharmaceutical Sciences* **2015**, *77*, 79-89.
- 9 49. Roscigno, P.; Asaro, F.; Pellizer, G.; Ortona, O.; Paduano, L., Complex Formation  
10 between Poly(vinylpyrrolidone) and Sodium Decyl Sulfate Studied through NMR. *Langmuir*  
11 **2003**, *19*, 9638-9644.
- 12 50. Prasad, D.; Chauhan, H.; Atef, E., Role of Molecular Interactions for Synergistic  
13 Precipitation Inhibition of Poorly Soluble Drug in Supersaturated Drug–Polymer–Polymer  
14 Ternary Solution. *Molecular Pharmaceutics* **2016**, *13*, 756-765.
- 15 51. Chen, Y.; Wang, S.; Wang, S.; Liu, C.; Su, C.; Hageman, M.; Hussain, M.; Haskell, R.;  
16 Stefanski, K.; Qian, F., Sodium Lauryl Sulfate Competitively Interacts with HPMC-AS and  
17 Consequently Reduces Oral Bioavailability of Posaconazole/HPMC-AS Amorphous Solid  
18 Dispersion. *Molecular Pharmaceutics* **2016**, *13*, 2787-2795.
- 19 52. Chen, Y.; Liu, C.; Chen, Z.; Su, C.; Hageman, M.; Hussain, M.; Haskell, R.; Stefanski, K.;  
20 Qian, F., Drug–Polymer–Water Interaction and Its Implication for the Dissolution  
21 Performance of Amorphous Solid Dispersions. *Molecular Pharmaceutics* **2015**, *12*, 576-589.
- 22 53. Singha, N. C.; Sathyanarayana, D. N., <sup>1</sup>H and <sup>13</sup>C NMR spectral studies of  
23 conformation of some N-(2-pyridinyl)-3-pyridinecarboxamides. *Journal of Molecular*  
24 *Structure* **1998**, *449*, 91-98.
- 25 54. Munro, S. L. A.; Craik, D. J., NMR conformational studies of fenamate non-steroidal  
26 anti-inflammatory drugs. *Magnetic Resonance in Chemistry* **1994**, *32*, 335-342.
- 27 55. You, Z. Y.; Chen, Y. J.; Wang, Y. Y.; Chen, C., Synthesis of Deuterium Labeled  
28 Standards of 1-Benzylpiperazine, Fenetylline, Nicocodeine and Nicomorphine. *Journal of the*  
29 *Chinese Chemical Society* **2008**, *55*, 663-667.
- 30 56. Ma, J.-h.; Guo, C.; Tang, Y.-l.; Zhang, H.; Liu, H.-z., Probing Paeonol–Pluronic Polymer  
31 Interactions by <sup>1</sup>H NMR Spectroscopy. *The Journal of Physical Chemistry B* **2007**, *111*, 13371-  
32 13378.
- 33 57. Ma, J.-h.; Guo, C.; Tang, Y.-l.; Chen, L.; Bahadur, P.; Liu, H.-z., Interaction of Urea with  
34 Pluronic Block Copolymers by <sup>1</sup>H NMR Spectroscopy. *The Journal of Physical Chemistry B*  
35 **2007**, *111*, 5155-5161.
- 36 58. Charman, W.; Lai, C.; Craik, D.; Finnin, B.; Reed, B., Self-Association of Nicotinamide  
37 in Aqueous-Solution: N.M.R. Studies of Nicotinamide and the Mono- and Di-methyl-  
38 Substituted Amide Analogs. *Australian Journal of Chemistry* **1993**, *46*, 377-385.
- 39 59. Bobrovs, R.; Seton, L.; Dempster, N., The reluctant polymorph: investigation into the  
40 effect of self-association on the solvent mediated phase transformation and nucleation of  
41 theophylline. *CrystEngComm* **2015**, *17*, 5237-5251.
- 42 60. Otsuka, N.; Ueda, K.; Ohyagi, N.; Shimizu, K.; Katakawa, K.; Kumamoto, T.; Higashi, K.;  
43 Yamamoto, K.; Moribe, K., An Insight into Different Stabilization Mechanisms of Phenytoin  
44 Derivatives Supersaturation by HPMC and PVP. *Journal of Pharmaceutical Sciences* **2015**, *104*,  
45 2574-2582.
- 46 61. Humpolíčková, J.; Štěpánek, M.; Procházka, K.; Hof, M., Solvent Relaxation Study of  
47 pH-Dependent Hydration of Poly(oxyethylene) Shells in Polystyrene-block-poly(2-  
48 vinylpyridine)-block-poly(oxyethylene) Micelles in Aqueous Solutions. *The Journal of Physical*  
49 *Chemistry A* **2005**, *109*, 10803-10812.
- 50 62. Oliveira, C. P.; Ribeiro, M. E.; Ricardo, N. M.; Souza, T. V.; Moura, C. L.; Chaibundit, C.;  
51 Yeates, S. G.; Nixon, K.; Attwood, D., The effect of water-soluble polymers, PEG and PVP, on  
52 the solubilisation of griseofulvin in aqueous micellar solutions of Pluronic F127. *Int. J. Pharm.*  
53 **2011**, *421*, 252-7.
- 54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Molecular structures of FFA, coformers (NIC and TP), FFA cocrystals (FFA-NIC and FFA-TP) and polymers (PVP and PVP-VA).


|         |  |
|---------|--|
| FFA     |  |
| NIC     |  |
| TP      |  |
| FFA-NIC |  |
| FFA-TP  |  |
| PVP     |  |
| PVP-VA  |  |




**Table 2.** HPLC methods

|     | Mobile phase                                      | Flow rate<br>(mL/min) | Detection wavelength<br>(nm) |
|-----|---|-----------------------|------------------------------|
| FFA | 14.5% water (with 0.5% formic acid), 85% methanol | 1.5                   | 286                          |
| NIC | 10% methanol, 90% water                           | 1                     | 265                          |
| TP  | 40% methanol, 60% water                           | 1                     | 265                          |


**Figure 1.** Diagram of the dissolution/permeation (D/P) system.




**Figure 2.** Effect of polymer and/or conformer on the equilibrium solubility of FFA. The apparent equilibrium solubility of FFA were measured in 0.01 M PBS (pH 4.5) with or without PVP or PVP-VA. The concentrations of NIC or TP used were 47.8  $\mu\text{g/mL}$  and 71.1  $\mu\text{g/mL}$ .



**Figure 3.** The dissolution (a) and permeation (b) of the FFA, FFA-NIC and FFA-TP were measured using the dissolution/permeation (D/P) system. The solvent used was 0.01 M PBS.




(a)




(b)

**Figure 4.** The D/P system was used for the DPP and flux rate measurement for FFA, FFA-NIC and FFA-TP. The donor compartment was filled with 0.01 M PBS (pH 4.5) containing either PVP or PVP-VA at concentrations of 200  $\mu\text{g/mL}$ , 4000  $\mu\text{g/mL}$  or 16000  $\mu\text{g/mL}$ , and the receptor compartment was filled with 0.01 M PBS (pH 4.5).




(a)




(b)


**Figure 5.** The dissolution and permeation of FFA, FFA-NIC and FFA-TP were investigated using the D/P system. The donor compartment was filled with 0.01 M PBS (pH 4.5) containing PVP at concentrations of 200  $\mu\text{g/mL}$ , 4000  $\mu\text{g/mL}$  or 16000  $\mu\text{g/mL}$ , and the receptor compartment was filled with 0.01 M PBS (pH 4.5).



**Figure 6.** The dissolution and permeation of FFA, FFA-NIC and FFA-TP were investigated using the D/P system. The donor compartment was filled with 0.01 M PBS (pH 4.5) containing PVP-VA at concentrations of 200  $\mu\text{g/mL}$ , 4000  $\mu\text{g/mL}$  or 16000  $\mu\text{g/mL}$ , and the receptor compartment was filled with 0.01 M PBS (pH 4.5).




**Figure 4.**  $^1\text{H}$  NMR spectra of FFA, NIC and TP. Samples of FFA (500  $\mu\text{g/mL}$ , 1000  $\mu\text{g/mL}$  and 5000  $\mu\text{g/mL}$ ), NIC (215  $\mu\text{g/mL}$ , 430  $\mu\text{g/mL}$  and 2150  $\mu\text{g/mL}$ ) and TP (320  $\mu\text{g/mL}$ , 640  $\mu\text{g/mL}$  and 3200  $\mu\text{g/mL}$ ) were prepared in deuterated chloroform ( $\text{CDCl}_3$ ) using the standard 5 mm NMR tubes and the spectrum of tetramethylsilane (TMS) was used as an internal standard.







**Figure 5.** Effect of PVP or PVP-VA on the  $^1\text{H}$  NMR spectra of FFA, NIC and TP.

Samples of FFA (1000  $\mu\text{g/mL}$ ), NIC (430  $\mu\text{g/mL}$ ) and TP (640  $\mu\text{g/mL}$ ) were prepared in deuterated chloroform ( $\text{CDCl}_3$ ) using the standard 5 mm NMR tubes and the spectrum of tetramethylsilane (TMS) was used as an internal standard. PVP or PVP-VA was included in the solution of  $\text{CDCl}_3$  at a concentration of 200  $\mu\text{g/mL}$  or 5000  $\mu\text{g/mL}$ .



**Figure 6.**  $^1\text{H}$  NMR spectra of mixture FFA with NIC or TP. Samples were prepared in deuterated chloroform ( $\text{CDCl}_3$ ) using the standard 5 mm NMR tubes and the spectrum of tetramethylsilane (TMS) was used as an internal standard. The FFA and its conformers were included in the  $\text{CDCl}_3$  solution at 1:1 molar ratio. The concentration of FFA, NIC and TP were  $1000\ \mu\text{g/mL}$ ,  $430\ \mu\text{g/mL}$ , and  $640\ \mu\text{g/mL}$ .





**Figure 7.** Effect of PVP or PVP-VA on the  $^1\text{H}$  NMR spectra of mixed FFA with NIC or TP.  $^1\text{H}$  NMR spectra of FFA in an equal molar mixture of FFA and NIC or TP at difference concentrations of a polymer in  $\text{CDCl}_3$ . Samples were prepared in deuterated chloroform ( $\text{CDCl}_3$ ) using the standard 5 mm NMR tubes and the spectrum of tetramethylsilane (TMS) was used as an internal standard. The FFA and its cofomers were included in the  $\text{CDCl}_3$  solution at 1:1 molar ratio to obtain the concentrations of 430  $\mu\text{g/mL}$  of NIC, 640  $\mu\text{g/mL}$  of TP and 1000  $\mu\text{g/mL}$  of FFA. PVP or PVP-VA was included in the solution of  $\text{CDCl}_3$  at a concentration of 200  $\mu\text{g/mL}$  or 5000  $\mu\text{g/mL}$ .

