


## Article

# Human Cytomegalovirus Upregulates Expression of HCLS1 Resulting in Increased Cell Motility and Transendothelial Migration during Latency



Yusuf Aslam,  
James Williamson,  
Veronika  
Romashova, ...,  
Paul Lehner, John  
Sinclair, Emma  
Poole

js152@cam.ac.uk (J.S.)  
elp27@cam.ac.uk (E.P.)

**HIGHLIGHTS**

HCLS1 is upregulated in cells latently infected with HCMV

HCLS1 modulates monocyte motility

Latent infection with HCMV increases monocyte motility

Aslam et al., iScience 20, 60–72  
October 25, 2019 © 2019 The Author(s).  
<https://doi.org/10.1016/j.isci.2019.09.016>

## Article

# Human Cytomegalovirus Upregulates Expression of HCLS1 Resulting in Increased Cell Motility and Transendothelial Migration during Latency

Yusuf Aslam,<sup>1</sup> James Williamson,<sup>1</sup> Veronika Romashova,<sup>1</sup> Elizabeth Elder,<sup>1</sup> Benjamin Krishna,<sup>2</sup> Mark Wills,<sup>1</sup> Paul Lehner,<sup>1</sup> John Sinclair,<sup>1,\*</sup> and Emma Poole<sup>1,3,\*</sup>

## SUMMARY

**Human cytomegalovirus establishes a lifelong, latent infection in the human host and can cause significant morbidity and mortality, particularly, in immunocompromised individuals. One established site of HCMV latency and reactivation is in cells of the myeloid lineage. In undifferentiated myeloid cells, such as CD14+ monocytes, virus is maintained latently. We have recently reported an analysis of the total proteome of latently infected CD14+ monocytes, which identified an increase in hematopoietic lineage cell-specific protein (HCLS1). Here we show that this latency-associated upregulation of HCLS1 occurs in a US28-dependent manner and stabilizes actin structure in latently infected cells. This results in their increased motility and ability to transit endothelial cell layers. Thus, latency-associated increases in monocyte motility could aid dissemination of the latently infected reservoir, and targeting this increased motility could have an impact on the ability of latently infected monocytes to distribute to tissue sites of reactivation.**

## INTRODUCTION

Following primary infection with human cytomegalovirus (HCMV), the virus is never cleared but is carried for life. This lifelong viral persistence is underpinned by true latent carriage in certain cell types *in vivo*, which sporadically reactivate subclinically (Poole and Sinclair, 2015; Sinclair and Poole, 2014). In normal healthy carriers, HCMV primary infection or reactivation is rarely symptomatic, but it does cause significant morbidity and mortality in the immunocompromised, immunosuppressed, or the immunonaive. One established site of HCMV latency *in vivo* is known to be CD34+ progenitor cells of the myeloid lineage (Poole and Sinclair, 2015; Sinclair and Poole, 2014). For instance, CD34+ progenitor cells from the bone marrow or from granulocyte colony-stimulating factor-mobilized donors have been shown to carry viral genome in the absence of detectable virus production (Poole and Sinclair, 2015; Reeves et al., 2005a, 2005b; Sinclair and Poole, 2014), an accepted hallmark of latent infection. It is now also clear that CD14+ monocytes, which are derived from CD34+ progenitors, also carry latent viral genomes. However, as these myeloid cells differentiate to tissue macrophages and dendritic cells (DCs), virus reactivates resulting in lytic infection and the *de novo* production of infectious virions. The effect of latent infection on myeloid cells has now become a topic of considerable interest, and, far from the view that latency is a passive carriage of quiescent viral genomes, more recent studies suggest that latent infection imparts important changes on the cell, which support maintenance of latency and enable efficient virus reactivation (Krishna et al., 2016; Lau et al., 2016; Mason et al., 2012; Poole et al., 2014; Poole et al., 2015; Poole et al., 2011; Poole and Sinclair, 2015; Rossetto et al., 2013; Slobedman and Cheung, 2008). For instance, studies using experimental models of latency have shown that latent infection of myeloid cells with HCMV profoundly modulates the cell secretome, apoptome, and microRNAome (Mason et al., 2012; Poole et al., 2011, 2014, 2015; Poole and Sinclair, 2015; Rossetto et al., 2013; Slobedman and Cheung, 2008). Recently, we reported an analysis of total latency-associated changes in the cell proteome of latently infected CD14+ monocytes using Tandem Mass Tag technology and identified robust changes in cellular proteins resulting from latent infection (Elder et al., 2019). Besides the secreted cellular proteins S100A8 and A9, which we have already reported on (Elder et al., 2019), one of the other most highly upregulated proteins was hematopoietic cell lineage-specific protein 1 (HCLS1). HCLS1 has been implicated in a number of cellular processes, but its role in cell motility, centered on actin rearrangement, is well established. For instance, HCLS1 is a cortactin homolog and can increase the stability of actin filaments (Cavnar et al., 2012; Dehring et al., 2011; Gomez et al., 2006; Hao et al., 2005; Mukherjee et al., 2015; Uruno et al.,

<sup>1</sup>Cambridge University, Department of Medicine, Level 5, Addenbrooke's Hospital, Hills Road, Cambridge CB2 0QQ, UK

<sup>2</sup>Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44106, USA

<sup>3</sup>Lead Contact

\*Correspondence: js152@cam.ac.uk (J.S.), elp27@cam.ac.uk (E.P.)

<https://doi.org/10.1016/j.isci.2019.09.016>



2003). Interestingly, HCMV is known to regulate actin at a number of points in lytic infection, and this helps to mediate viral egress (Wilkie et al., 2016), restructure lipid rafts (Low et al., 2016), impair immune recognition (Fielding et al., 2014; Gabaev et al., 2014), and promote cell migration (Dehring et al., 2011; Reinhardt et al., 2014; Streblow et al., 1999; Tseliou et al., 2016). However, little is known about the effect of latent infection on actin, although it is known that virus binding to monocytes can cause immediate effects on paxillin protein, which regulates actin filament networks and enhances motility (Chan et al., 2008; Nogalski et al., 2011). Here, we now show that, subsequent to virus binding and in response to the latency-associated upregulation of HCLS1, latent infection of monocytes results in increased stability of filamentous actin, which, in turn, enhances monocyte migration.

A number of studies have linked the actin filament association of HCLS1 with cell motility of natural killer (NK) cells, DCs, and neutrophils (Dehring et al., 2011; Hao et al., 2005; Latasiewicz et al., 2017; Mukherjee et al., 2015; Uruno et al., 2003). Depletion of HCLS1 from NK cells renders them less motile (Mukherjee et al., 2015). Furthermore, knockout of HCLS1 in the mouse model system decreases neutrophil rolling, adhesion, and migration across the endothelial cell layer. Although it is established that the rolling, adhesion, and migration properties of monocytes, like other leukocytes, help them extravasate across the endothelial cell layer (Martin et al., 2007), it is not known whether HCLS1 plays a role in such monocyte migration and endothelial cell layer transit.

Our analyses now show that latently infected monocytes, in which HCLS1 is profoundly upregulated, have increased motility as well as increased ability to adhere to endothelial cells in a vascular flow system and that they are also more able to cross endothelial cell layers. We confirmed that these effects directly involved latency-associated HCLS1 upregulation by showing that latently infected cells, in which HCLS1 was knocked down by RNAi, no longer showed latency-associated increases in adherence to vascular layers under flow conditions.


Finally, we also show that mycophenolate mofetil (MMF), which modulates actin polymerization (Chaigne-Delalande et al., 2008; Dubus et al., 2003) and prevents monocyte adhesion and transmigration (Glomsda et al., 2003; Park et al., 2016), also prevented latently infected CD14<sup>+</sup> monocytes from adhering to the endothelial layer during physiological flow conditions. This is consistent with the view that it is the increased stability of filamentous actin in latently infected monocytes, mediated by HCLS1 upregulation, which enhances their motility and crossing of endothelial cell layers.

We suggest that the increased endothelial adherence and increased motility of latently infected monocytes, mediated by latency-associated upregulation of HCLS1, could help monocytes latently infected with HCMV to extravasate from blood vessels into surrounding tissues to aid dissemination of the latent monocyte reservoir to tissue sites of reactivation.

## RESULTS

### Analysis of HCMV Latently Infected CD14<sup>+</sup> Monocytes Shows a Marked Increase in Cellular Protein HCLS1

Recently, we carried out a total proteomic screen of HCMV latently infected CD14<sup>+</sup> monocytes, which we had confirmed were truly latently infected by RNA analysis and virion release assays, and identified proteins that were changed as a result of latent infection (Elder et al., 2019). Further analysis of these data identified that one of the most highly upregulated proteins associated with latent infection was HCLS1 (Figure 1A), which was not upregulated following infection with a UV-inactivated virus (Figure 1B). Therefore, we validated this upregulation of HCLS-1 during latent infection. To do this, we infected CD14<sup>+</sup> monocytes with an SV40GFP-tagged TB40E HCMV, which allowed us to enrich for GFP<sup>+</sup> latently infected cells, and used western blot analysis to analyze levels of HCLS1. This confirmed that HCLS1 is substantially increased during latent infection at both the protein (western blot, Figure 1C) and mRNA (RT-qPCR, Figure 1C) levels, but not in response to UV-inactivated virus (Figures 1B and 1C). We further analyzed this latency-associated increase in HCLS1 in latently infected cells by indirect immunofluorescence (Figure 1D). To do this, monocytes were latently infected with a second recombinant TB40E isolate expressing mCherry, which allows latently infected cells to be identified for much longer times after infection and overcomes the waning of GFP expression, which is known to occur when using a GFP-tagged virus (Krishna et al., 2016), and then stained these cells for HCLS1. This confirmed that latently infected cells (mCherry positive) had increased expression of HCLS1 (Figure 1D). Taken together these data show



**Figure 1. HCLS1 Levels Are Increased during HCMV Latency which Is Regulated by Src Phosphorylation**

(A–H) CD14<sup>+</sup> monocytes were either left uninfected or infected with TB40E-SV40GFP to allow sorting of the latently infected cells on the basis of GFP expression. Cells latently infected for 4 days, or after 4-day infection with UV-inactivated virus, were harvested for total cell proteomics analysis (A and B); western blot analysis or RT-qPCR analysis for HCLS1 and actin (C). Alternatively, CD14<sup>+</sup> primary monocytes were infected with TB40E-GATA2mCherry and then co-stained for HCLS1 using primary HCLS1 antibody followed by fluorescein isothiocyanate-conjugated secondary antibody. Red,

**Figure 1. Continued**

green, and blue channels as well as bright-field images are shown (scale bar, 30  $\mu$ M), virally infected cells are highlighted with arrows (D). CD14<sup>+</sup> monocytes were also either mock infected or infected with TB40E-SV40GFP and either treated with Src inhibitor PP2 or left untreated. Then they were sorted for GFP expression, and the levels of HCLS1 mRNA were analyzed by RT-qPCR (E, top panel); ATF1 occupancy on the HCLS1 promoter in these cells was determined by chromatin immunoprecipitation assay using either an ATF1-specific antibody or an isotype-matched control antibody (E, bottom panel). In addition, wild-type (WT) and US28-expressing THP-1 cells were also treated or untreated with PP2 and then analyzed for levels of HCLS1 RNA by RT-qPCR relative to GAPDH housekeeping gene (F, top panel), and ATF1 occupancy on the HCLS1 promoter was determined using ATF1-specific antibody or isotype-matched control antibody (F, bottom panel). RT-qPCR graphs represent three replicates with standard deviations shown (E and F). Finally, WT THP-1 (empty vector-transduced "control") cells or THP-1 cells overexpressing US28 ("US28") were either treated with PP2 or left untreated and analyzed by western blot for phosphorylated Src and actin (as a loading control). Bands were analyzed by densitometry as shown graphically (G). Finally, undifferentiated monocytes, monocytes that had been terminally differentiated into DCs (using granulocyte-macrophage colony-stimulating factor/interleukin-4 and lipopolysaccharide), and fibroblasts were either left uninfected (–) or infected with TB40E-SV40GFP (+) (left panel) or with the Titan strain of HCMV, which expresses a UL32-GFP fusion protein (WT) or the US28 deletion version of the same virus (US28-KO) (right panel). Myeloid cells were then sorted at 4 d post-infection (dpi), based on SV40-GFP expression, and fibroblasts were also sorted at 4 dpi based on GFP-UL32 expression; then all cells were harvested for HCLS1 and actin western blot analysis (H).

that cells latently infected with a number of different virus isolates increase expression of HCLS1 at both the protein and RNA levels.

**Upregulation of HCLS1 during Latent Infection Is Mediated by ATF1 in an Src-Dependent Manner**

To try to address the mechanism by which HCLS1 is upregulated during latency we examined the HCLS1 enhancer regions for known activatory transcription factor-binding sites using Genecard analysis and confirmed this using Physbinder, as described in the [Transparent Methods](#). One transcription factor that was predicted to bind to the HCLS1 promoter was ATF1, and this particularly came to our attention because it is known to be upregulated by Src signaling, a signaling pathway known to be involved in infection of monocytes by HCMV (Nogalski et al., 2011) (Shi et al., 2011) (Hokari et al., 2005).

Therefore, we interrogated whether increased levels of HCLS1 in latently infected cells occurred via the phosphorylation of Src and subsequent signaling to allow recruitment of ATF1 to the HCLS1 promoter. To do this, we treated latently infected cells with a known inhibitor of Src signaling, PP2. [Figure 1E](#) (top panel) shows that consistent with Src signaling playing a role in the induction of HCLS1 in infected latently infected monocytes, the latency-associated increase in HCLS1 is abrogated by the PP2 Src signaling inhibitor.

We pursued this further by testing whether the latency-associated increase in HCLS1 expression resulted from increased ATF1 binding to the HCLS1 promoter. [Figure 1E](#) (bottom panel) shows that significant ATF1 occupancy on the HCLS1 promoter only occurred in latently infected cells and that this was prevented by PP2 Src inhibitor. These data, therefore, suggest that, during latency, HCLS1 is upregulated via Src signaling, which mediates ATF1 binding to the HCLS1 promoter.

Previous work has shown that during HCMV latency, the viral protein US28 is able to induce a number of cellular signaling changes in undifferentiated myeloid cells, quite different to those induced in cells undergoing a lytic infection (Krishna et al., 2017). Interestingly, these changes during latent infection include alterations to Src signaling. Therefore, we tested whether the viral US28 gene product could be involved in HCLS1 upregulation in an Src phosphorylation-dependent manner to increase ATF1 occupancy at the HCLS1 promoter. [Figure 1F](#) (top panel) shows that, in myelomonocytic THP-1 cells, overexpression of US28 in isolation results in upregulation of HCLS1 at the RNA level. Furthermore, this US28-mediated upregulation of HCLS1 is inhibited if these cells are treated with the Src inhibitor PP2. [Figure 1F](#) (bottom panel) also shows that US28-expressing THP-1 cells have increased ATF1 occupancy on the HCLS1 promoter and that, as with CD14<sup>+</sup> cells during HCMV latency, this is also inhibited by PP2.

Finally, Src phosphorylation, itself, was analyzed in the US28-expressing cells by western blot. [Figure 1G](#) shows that Src phosphorylation increased after overexpression of US28 in THP-1 cells, and this was inhibited by treatment with Src inhibitor (PP2).

As US28 is also known to target Src phosphorylation during lytic infection (Streblov et al., 1999), we tested whether US28 could also mediate HCLS1 upregulation during a lytic infection. To do this, two cell types that support lytic infection, fibroblasts and DCs, were infected alongside monocytes for 4 days and analyzed for HCLS1 and actin content by western blot.

First, we were unable to detect HCLS1 expression in untreated fibroblasts consistent with the known low levels of HCLS1 in this cell type (Shin et al., 2012), and their lytic infection with HCMV failed to show any induction of HCLS1 (Figure 1H, right panels).

As before (Figure 1), latent infection of monocytes showed an approximate 10-fold increase in HCLS1 (Figure 1H, left panels). In contrast, lytic infection of monocyte-derived DCs resulted in less than a 2-fold increase in HCLS1, and this was above an already high background level of HCLS1 (Figure 1H, left panels). To test whether this minor increase in HCLS1 during lytic infection was US28 dependent, we repeated this analysis using a US28 deletion virus but still observed an increase in HCLS1 expression in infected DCs (Figure 1H, right panels).

Consequently, lytic infection of DCs results in only a small increase in HCLS1 over an already high basal level of HCLS1 expression in these cells, and this small increase is only partially US28 dependent.

Taken together, these data suggest that latent infection of monocytes results in a substantial increase in HCLS1, which is, at least in part, mediated by US28. In contrast, lytic infection of monocyte-derived DCs showed only a minor increase in HCLS1 over already high background levels of HCLS1 and argues that the most profound functional effects of HCLS1 induction are likely to be during latent infection.

### HCMV Latent Infection Results in Monocytes with Stabilized F-actin


We next addressed the potential role of HCLS1 induction during latent infection. As a cortactin homolog, HCLS1 is known to stabilize F-actin and increase cell motility (Dehring et al., 2011; Martin et al., 2007; Uruno et al., 2003). Consequently, we reasoned that the increased expression of HCLS1 during latency could prevent destabilization of F-actin, which can result from cytochalasin D (CD) treatment (Aunis and Bader, 1988; Sontag et al., 1988; Thieler et al., 1995).

Figure 2A shows that, treatment of control untreated monocytes with CD resulted in a substantial decrease in detectable F-actin (CD+). In contrast, overexpression of HCLS1 in isolation in monocytes was able to prevent this F-actin disruption by CD treatment (Figure 2B), which has only been shown for fully differentiated DCs previously (Dehring et al., 2011). We next tested whether this protective effect of increased HCLS1 on CD-mediated destabilization of F-actin also occurred in latently infected monocytes. Figure 2C shows that uninfected monocytes (mCherry negative) treated with CD lost F-actin staining. In contrast, this CD-mediated decrease in F-actin staining was not observed in latently infected (mCherry positive) monocytes (Figure 2C).

Taken together, these results show that the upregulation of HCLS1 during HCMV latency results in increased stabilization of F-actin structure.

### Overexpression of HCLS1 as well as Latent Infection of CD14+ Monocytes Leads to Increased Cell Motility

One consequence of increased HCLS1, and subsequent stabilized F-actin can be increased motility of cells in certain cell types (Dehring et al., 2011; Mukherjee et al., 2015). Therefore, we overexpressed HCLS1 by electroporation with HCLS1-GFP or GFP expression plasmids into monocytes, a cell type in which HCMV latency resides, and analyzed the frequency of actin protrusions, as a surrogate for motility, as determined by cells that phenotypically contained protrusions that stained for F-actin (Melak et al., 2017). Initially, we analyzed latently infected cells for the presence of actin protrusions. Figure 3A clearly shows that latently infected cells had substantial increase in actin protrusions (an example of an actin protrusion is highlighted in phase contrast, top panel, Figure 3A), suggesting increased motility, yet the monocyte protrusions were inhibited if the cells were treated with PP2, a drug known to affect actin-based motility (Figure 3A right panel). To test this directly, these HCLS1-GFP-positive cells were also analyzed by Cellomics time-lapse imaging at 24 h post electroporation. Figure 3B shows that cells that had been electroporated with HCLS1-GFP were substantially more motile and moved further than those transfected with GFP alone. Still



### Figure 2. HCLS1 Upregulation during Latency Leads to Increased and Stable Actin Filament Formation

(A–C) (A) CD14<sup>+</sup> monocytes in which actin filament formation had been disrupted using CD were analyzed by fluorescence for F-actin (green) and Hoechst (blue). Alternatively, CD14<sup>+</sup> monocytes overexpressing GFP or GFP-HCLS1 were analyzed for actin filament formation in the presence and absence of CD and analyzed by fluorescence. GFP (green), actin (red), and Hoechst (blue) stains are shown with cells enumerated in three experiments of triplicate wells. Percentage of cells that are both GFP and actin positive are represented; \*\*\* $p < 0.0001$ ; NS, not significant (B). Finally, actin filament formation in the presence of HCMV latency was assessed in the presence and absence of CD. CD14<sup>+</sup> monocytes were infected with TB40E-GATA2-mCherry virus, which expresses mCherry under the control of the GATA2 promoter (red), and stained for actin filaments (green) and Hoechst (blue) and analyzed by fluorescence, the virally infected cell is highlighted with arrows (C) (scale bars, 30  $\mu\text{m}$ ).


images and videos of these analyses are also shown in [Figure S1B](#). Therefore, monocytes, like other cell types, also increase their motility in response to overexpression of HCLS1.

Next, to test whether latently infected monocytes also have increased motility, we again used Cellomics time-lapse live cell imaging to assess cell movement in latently infected populations of CD14<sup>+</sup> monocytes. We employed TB40E-GATA2-mCherry virus for these analyses to allow long-term tracking of latently infected cells. [Figure 3C](#) shows the quantification of movement and distance traveled by latently infected monocytes compared with uninfected bystander cells in the same population. Latently infected CD14<sup>+</sup> monocytes appeared more pleomorphic ([Figure 3A](#)) and were substantially more motile over the time course studied than uninfected bystander cells. Still images and videos of these analyses are shown in [Figure S1](#) and [Videos S1, S2, and S3](#).

These data argue that latent infection with HCMV enhances the motility of CD14<sup>+</sup> monocytes.


### Latently Infected CD14<sup>+</sup> Monocytes Adhere to Endothelial Cells More Efficiently and Are More Able to Cross Endothelial Cell Layers

HCLS1 expression is also known to mediate neutrophil rolling, adhesion, and migration across the endothelium. Similarly, the rolling, adhesion, and migration properties of monocytes also help them extravasate across endothelial cell layers ([Martin et al., 2007](#)). Consequently, we reasoned that latency-associated HCLS1 increases in monocytes might affect monocyte motility, adhesion, and transmigration across endothelial cell monolayers.



**Figure 3. Monocytes Overexpressing HCLS1 Are More Motile, and This Is Recapitulated after Latent Infection**  
 (A–C) CD14<sup>+</sup> cells were latently infected for 4 days with HCMV and then analyzed for membranous protrusions by phase contrast microscopy and cell counting (scale bar, 30  $\mu$ m) (A). The frequency of membranous protrusions during HCMV latency compared with mock infected cells or latently infected cells that have been treated with PP2 1 h post infection are shown graphically where data represent three experiments with standard deviation; \*\*\*  $p < 0.0001$ , \*\*  $p < 0.001$  (A). CD14<sup>+</sup> monocytes were transfected with either GFP or HCLS1-GFP, and then Cellomics analysis was used to assess distance moved and speed of movement of the cells (B). Alternatively, CD14<sup>+</sup> monocytes were either left uninfected or infected with the TB40E-GATA2mCherry HCMV virus (to allow visualization of latent cells by red fluorescence) and latency established for 4 days before they were analyzed using Cellomics for cell motility by assessing distance moved and their speed of movement (C). Data represent triplicate experiments, and motility was analyzed for 100 cells per sample. Standard deviation error bars and significance determined using the Student's t test; \*\*  $p < 0.001$ , \*\*\*  $p < 0.0001$  (B and C).





**Figure 4. Monocytes Latently Infected with HCMV Show Enhanced Adhesion to, and Transmigration through, Endothelial Cell Layers in an HCLS1-Dependent and Src Phosphorylation-Dependent Manner**

(A–G) CD14<sup>+</sup> cells that had been transfected with either HCLS1-GFP or GFP (A) or that had been left uninfected (mock) or infected with TB40E-GATA2mCherry (B) were assessed for their ability to adhere to endothelial cell layers. CD14<sup>+</sup> cells that had been either mock infected or infected with TB40-GATA2mCherry virus were also assessed for their ability to migrate across endothelial cell layers in transwell assays (C). CD14<sup>+</sup> cells that had been infected with TB40E-GATA2mCherry were treated with either scramble siRNA or siRNA specific to HCLS1 and similarly assessed for their ability to adhere to the endothelial cell layer (D). CD14<sup>+</sup> cells that had been latently infected with TB40E-GATA2mCherry were also assessed for the ability to adhere to the endothelial cell layer in the presence of either Src inhibitor PP2 (E) or the drug MMF (F). Finally, “WT-THP1” (empty vector transduced) cells or THP1 cells expressing US28 “US28-THP1” were assessed for the ability to adhere to the endothelial cell layer (G). Experiments were carried out in replicate, and data represent triplicate experiments and show standard deviation error bars and significance determined using the Student’s t test; \* $p < 0.01$ , \*\* $p < 0.001$ , \*\*\* $p < 0.0001$ .

To test this, we first analyzed whether monocytes overexpressing HCLS1 were more able to adhere to endothelial cells under *in vitro* flow simulation. In this system (Butler et al., 2011; Zhao et al., 2016), endothelial cells were allowed to adhere to an artificial vessel and monocytes overexpressing GFP or HCLS1-GFP were perfused through the vessel at a physiological flow rate of 0.4 mm/mL and then fixed and analyzed by fluorescence. Figure 4A clearly shows that monocytes overexpressing HCLS1 showed increased adherence to the endothelial cell layer under physiological flow conditions (the electroporation efficiency for these plasmids was between 40% and 80% based on GFP expression as shown in Figure S2A). Therefore, as with neutrophils, increased expression of HCLS1 in monocytes can also mediate adherence to endothelial cell layers.

We then repeated these analyses with monocytes that had been not only latently infected with GATA2-mCherry virus but also co-stained with calcein (to allow differential detection of bystander monocytes and latently infected monocytes). Figure 4B shows that latently infected monocytes also adhered to the endothelial layer under flow conditions more efficiently than uninfected monocytes.

Next, we addressed whether latently infected monocytes were more adept at crossing endothelial cell layers. To do this, we sorted mCherry-positive latently infected monocytes and assessed their ability to cross a transwell membrane, which had been overlaid with endothelial cells. Figure 4C shows that latently infected monocytes were more efficient at crossing these endothelial-coated transwell membranes than uninfected monocytes.

Our data, so far, clearly showed that overexpression of HCLS1 in monocytes increased their motility and adhesion to endothelial cells as well as their ability to cross endothelial cell layers under flow conditions and that latently infected cells showed similar enhanced capabilities. However, this did not provide direct evidence that it was the latency-associated increase in HCLS1 that was mediating these effects in latently infected monocytes. To address this directly, we repeated the analysis of the ability of latently infected monocytes to adhere to endothelial monolayers, but in monocytes in which we had ablated the latency-associated upregulation of HCLS1 by RNA interference.

To do this, CD14<sup>+</sup> monocytes were infected with HCMV and, subsequently, HCLS1 expression was inhibited by electroporation of these latently infected cells with HCLS1-specific small interfering RNA (siRNA). This resulted in good levels of knockdown of HCLS1 protein (Figure S2B). Figure 4D shows that knockdown of HCLS1 in latently infected CD14<sup>+</sup> monocytes resulted in a significant reduction in their ability to adhere to endothelial cell layers under physiological flow conditions compared with those electroporated with a scramble control siRNA. To further validate the role of HCLS1 in these effects we also tested whether inhibition of Src, which we showed in Figure 1 to be important for HCLS1 induction in latently infected cells, had any effect on the ability of monocytes to adhere to endothelial cell layers under physiological flow conditions. Adherence of latently infected monocytes was decreased when Src phosphorylation was inhibited (Figure 4E), demonstrating that Src signaling was indeed playing a role in the HCLS1-mediated motility observed during latent infection.

MMF, besides a general immune suppressant, is also known to modulate actin polymerization (Chaigne-Delalande et al., 2008), thereby blocking monocytes from transmigrating across endothelial cell layers (Juttner et al., 2009). Consequently, we reasoned that if increased adherence and transendothelial

migration of latently infected monocytes were due to latency-associated stabilization of F-actin through HCLS1 upregulation, MMF should inhibit adhesion of latently infected monocytes to endothelial cells. To test this, latently infected CD14<sup>+</sup> monocytes were pre-incubated with MMF before perfusion over endothelial cell layers. [Figure 4F](#) shows that, following treatment with MMF, latently infected CD14<sup>+</sup> monocytes were significantly less able to adhere to the endothelial cell layer.

Finally, on the basis of the fact that our data showed that US28-mediated activation of Src drives overexpression HCLS1 and subsequent HCLS1-mediated increase in endothelial cell adherence, we tested whether monocytic cells overexpressing US28 had increased endothelial cell adherence. [Figure 4G](#) shows that THP1 cells that overexpress US28 are, indeed, more able to adhere to endothelial cell layers.

Taken together, these data argue that latent infection of CD14<sup>+</sup> monocytes results in increased levels of cellular HCLS1, which stabilizes actin filaments and concomitantly results in enhanced monocyte motility. This, in turn, aids monocyte adhesion to, and transmigration through, the endothelial cell layer. We believe that such effects could play an important role in virus dissemination from the peripheral blood compartment to tissue sites of reactivation.

## DISCUSSION

The work presented here shows that the latency-associated upregulation of HCLS1, which we previously identified in an unbiased analysis of changes in the cell proteome upon HCMV latent infection ([Elder et al., 2019](#)), modulates cell motility of latently infected cells and increases their ability to extravasate across endothelial cell layers.

HCLS1 is a substrate of antigen receptor-coupled tyrosine kinases, and it is only expressed in tissues and cells of hematopoietic origin ([Yamanashi et al., 1993](#)). In myeloid cells, HCLS1 has a number of established functions including the regulation lymphocyte trafficking ([Scielzo et al., 2010](#)), neutrophil chemotaxis ([Cavnar et al., 2012](#)), and growth arrest and apoptosis ([Fukuda et al., 1995](#)). Importantly, it also binds F-actin and can increase the rate of actin assembly ([Hao et al., 2005](#); [Uruno et al., 2003](#)). Interestingly, such changes in actin assembly are known to be important for cell motility of NK cells, DCs, and neutrophils as well as neutrophil rolling, adhesion, and migration across the endothelial cell layer ([Dehring et al., 2011](#); [Hao et al., 2005](#); [Latasiewicz et al., 2017](#); [Mukherjee et al., 2015](#); [Uruno et al., 2003](#)).

Our analyses also show that the mechanism by which HCLS1 is upregulated during latency is by increasing ATF1 association with the HCLS1 promoter by Src signaling and is consistent with previous observations in other conditions ([Hokari et al., 2005](#); [Shi et al., 2011](#)). This is particularly pertinent because HCMV lytic infection is known to regulate Src signaling ([Streblow et al., 1999](#)) and this upregulation during lytic infection is mediated by the viral US28 gene product. Importantly, US28 is also a latency-associated viral gene product ([Bego et al., 2005](#); [Poole et al., 2013](#)), which has been suggested to increase Src phosphorylation in latently infected cells ([Krishna et al., 2017](#)). In addition, Src signaling is also known to increase upon monocyte binding and entry of HCMV ([Nogalski et al., 2011](#)), a known site of HCMV latency. Therefore, not only does initial binding of HCMV to monocytes alter Src phosphorylation, which then modulates actin filament formation via paxillin regulation ([Nogalski et al., 2011](#)), but also this is continued, once latency has been fully established, via regulation of HCLS1. As Src phosphorylation has also been published as an effect of US28 during lytic infection, we also addressed whether US28 can mediate HCLS1 upregulation during lytic infection. We found that, in contrast to latent infection, lytic infection of DCs resulted in only minor increases in HCLS1, over an already high basal level of HCLS1 expression and that US28 likely played only a minor role in this upregulation. Consequently, we believe that the most profound functional effects of HCLS1 induction are likely to be during latent infection.

Consistent with the known effects of HCLS1 in regulating F-actin filaments, which can lead to enhanced motility, latently infected monocytes have more stable actin and are more mobile. One consequence of increased actin filaments and enhanced motility in monocytes is an increased ability to adhere to, and cross, the endothelial cell layer. Again, consistent with this, latently infected cells showed enhanced adherence to endothelial cell layers and increased ability to migrate across them. In all cases, our analyses showed that it was increased HCLS1 that mediated these latency-associated changes in monocyte motility, adherence, and trafficking as inhibition of the latency-associated increase in HCLS1 by RNAi prevented them.

One obvious advantage of these effects on monocytes during latent infection could be in viral dissemination in the myeloid lineage. *In vivo*, latent HCMV can be carried in peripheral blood monocytes (Reeves et al., 2005b; Taylor-Wiedeman et al., 1991). Consequently, any increased ability of a latently infected cell in the peripheral blood to adhere to and cross the endothelial cell layer would very likely enhance dissemination of the latent virus to tissue sites in the host conducive to monocyte differentiation and subsequent reactivation. HCMV is not the only herpes virus to increase endothelial cell motility during latency. The gamma herpesviruses Epstein-Barr virus (Tugizov et al., 2007) and Kaposi's sarcoma virus also promote endothelial transmigration following latent infection of endothelial cells (DiMaio et al., 2011; Giffin et al., 2015), although, in contrast to HCMV, lytic infection of the endothelial layer prevents neutrophil migration (Butler et al., 2011). Similarly, HIV-1, which can establish latency in resting CD4+ cells, promotes endothelial transmigration of infected monocytes (He et al., 2016; Maslin et al., 2005; Noguchi et al., 2007; Westhorpe et al., 2009), suggesting that this may be a common feature of blood-borne viruses, which establish latent life cycles.

In conclusion, we believe that changes in myeloid-specific HCLS1 imparted by latent infection of monocytes with HCMV can alter their adhesion to endothelial cells and promote transendothelial migration and that this could aid efficient dissemination of HCMV latently infected cells from the peripheral blood to potential tissue sites of reactivation.

### Limitations of the Study

As with all studies on species-specific pathogens, the inherent lack of an animal model is a limitation to any study. However, with increasingly more effective experimental latency models for the study of HCMV these types of studies are becoming more informative and directly relevant to the *in vivo* system.

### METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

### SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2019.09.016>.

### ACKNOWLEDGMENTS

We would like to thank Linda Teague and Roy Whiston for technical assistance. We also thank Jing Garland and Neda Farahi for help with the endothelial flow setup. This work was funded by the British Medical Research Council, Grant from the British Medical Research Council (Grant G0701279), and the Cambridge National Institute for Health Research Biomedical Research Council (NIHR BRC) Cell Phenotyping Hub.

### AUTHOR CONTRIBUTIONS

Y.A., J.W., V.R., B.K., E.E. and E.P. performed experiments. M.W., P.L., J.S. and E.P. designed experiments. J.S and E.P wrote the manuscript.

### DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: February 14, 2019

Revised: August 6, 2019

Accepted: September 11, 2019

Published: October 25, 2019

### REFERENCES

- Aunis, D., and Bader, M.F. (1988). The cytoskeleton as a barrier to exocytosis in secretory cells. *J. Exp. Biol.* 139, 253–266.
- Bego, M., Maciejewski, J., Khaiboullina, S., Pari, G., and St Jeor, S. (2005). Characterization of an antisense transcript spanning the UL81-82 locus of human cytomegalovirus. *J. Virol.* 79, 11022–11034.
- Butler, L.M., Jeffery, H.C., Wheat, R.L., Rae, P.C., Townsend, K., Alkharsah, K.R., Schulz, T.F., Nash, G.B., and Blackbourn, D.J. (2011). Kaposi's sarcoma-associated herpesvirus infection of endothelial cells inhibits neutrophil recruitment through an interleukin-6-dependent mechanism: a new paradigm for viral immune evasion. *J. Virol.* 85, 7321–7332.
- Cavnar, P.J., Mogen, K., Berthier, E., Beebe, D.J., and Huttenlocher, A. (2012). The actin regulatory

- protein HS1 interacts with Arp2/3 and mediates efficient neutrophil chemotaxis. *J. Biol. Chem.* **287**, 25466–25477.
- Chaigne-Delalande, B., Guidicelli, G., Couzi, L., Merville, P., Mahfouf, W., Bouchet, S., Molimard, M., Pinson, B., Moreau, J.F., and Legembre, P. (2008). The immunosuppressor mycophenolic acid kills activated lymphocytes by inducing a nonclassical actin-dependent necrotic signal. *J. Immunol.* **181**, 7630–7638.
- Chan, G., Bivins-Smith, E.R., Smith, M.S., Smith, P.M., and Yurochko, A.D. (2008). Transcriptome analysis reveals human cytomegalovirus reprograms monocyte differentiation toward an M1 macrophage. *J. Immunol.* **181**, 698–711.
- Dehring, D.A., Clarke, F., Ricart, B.G., Huang, Y., Gomez, T.S., Williamson, E.K., Hammer, D.A., Billadeau, D.D., Argon, Y., and Burkhardt, J.K. (2011). Hematopoietic lineage cell-specific protein 1 functions in concert with the Wiskott-Aldrich syndrome protein to promote podosome array organization and chemotaxis in dendritic cells. *J. Immunol.* **186**, 4805–4818.
- DiMaio, T.A., Gutierrez, K.D., and Lagunoff, M. (2011). Latent KSHV infection of endothelial cells induces integrin beta3 to activate angiogenic phenotypes. *PLoS Pathog.* **7**, e1002424.
- Dubus, I., L'Azou, B., Gordien, M., Delmas, Y., Labouyrie, J.P., Bonnet, J., and Combe, C. (2003). Cytoskeletal reorganization by mycophenolic acid alters mesangial cell migration and contractility. *Hypertension* **42**, 956–961.
- Elder, E., Krishna, B., Williamson, J., Aslam, Y., Farahi, N., Wood, A., Romashova, V., Roche, K., Murphy, E., Chilvers, E., et al. (2019). Monocytes latently infected with human cytomegalovirus evade neutrophil killing. *iScience* **12**, 13–26.
- Fielding, C.A., Aicheler, R., Stanton, R.J., Wang, E.C., Han, S., Seirafian, S., Davies, J., McSharry, B.P., Weekes, M.P., Antrobus, P.R., et al. (2014). Two novel human cytomegalovirus NK cell evasion functions target MICA for lysosomal degradation. *PLoS Pathog.* **10**, e1004058.
- Fukuda, T., Kitamura, D., Taniuchi, I., Maekawa, Y., Benhamou, L.E., Sarthou, P., and Watanabe, T. (1995). Restoration of surface IgM-mediated apoptosis in an anti-IgM-resistant variant of WEHI-231 lymphoma cells by HS1, a protein-tyrosine kinase substrate. *Proc. Natl. Acad. Sci. U S A* **92**, 7302–7306.
- Gabaev, I., Elbasani, E., Ameres, S., Steinbruck, L., Stanton, R., Doring, M., Lenac Rovis, T., Kalinke, U., Jonjic, S., Moosmann, A., et al. (2014). Expression of the human cytomegalovirus UL11 glycoprotein in viral infection and evaluation of its effect on virus-specific CD8 T cells. *J. Virol.* **88**, 14326–14339.
- Giffin, L., West, J.A., and Damania, B. (2015). Kaposi's sarcoma-associated herpesvirus interleukin-6 modulates endothelial cell movement by upregulating cellular genes involved in migration. *MBio* **6**, e01499–01415.
- Glomsda, B.A., Blaheta, R.A., and Hailer, N.P. (2003). Inhibition of monocyte/endothelial cell interactions and monocyte adhesion molecule expression by the immunosuppressant mycophenolate mofetil. *Spinal Cord* **41**, 610–619.
- Gomez, T.S., McCarney, S.D., Carrizosa, E., Labno, C.M., Comiskey, E.O., Nolz, J.C., Zhu, P., Freedman, B.D., Clark, M.R., Rawlings, D.J., et al. (2006). HS1 functions as an essential actin-regulatory adaptor protein at the immune synapse. *Immunity* **24**, 741–752.
- Hao, J.J., Zhu, J., Zhou, K., Smith, N., and Zhan, X. (2005). The coiled-coil domain is required for HS1 to bind to F-actin and activate Arp2/3 complex. *J. Biol. Chem.* **280**, 37988–37994.
- He, X., Shi, X., Puthiyakunnon, S., Zhang, L., Zeng, Q., Li, Y., Boddu, S., Qiu, J., Lai, Z., Ma, C., et al. (2016). CD44-mediated monocyte transmigration across *Cryptococcus neoformans*-infected brain microvascular endothelial cells is enhanced by HIV-1 gp41-190 ectodomain. *J. Biomed. Sci.* **23**, 28.
- Hokari, R., Lee, H., Crawley, S.C., Yang, S.C., Gum, J.R., Jr., Miura, S., and Kim, Y.S. (2005). Vasoactive intestinal peptide upregulates MUC2 intestinal mucin via CREB/ATF1. *Am. J. Physiol. Gastrointest. Liver Physiol.* **289**, G949–G959.
- Juttner, B., Bencel, M., Weissig, A., Studzinski, A., Stenger, K., and Scheinichen, D. (2009). Mycophenolic acid inhibits PMA-induced activation of the neutrophil respiratory burst. *Transpl. Infect. Dis.* **11**, 235–240.
- Krishna, B.A., Lau, B., Jackson, S.E., Wills, M.R., Sinclair, J.H., and Poole, E. (2016). Transient activation of human cytomegalovirus lytic gene expression during latency allows cytotoxic T cell killing of latently infected cells. *Sci. Rep.* **6**, 24674.
- Krishna, B.A., Poole, E.L., Jackson, S.E., Smit, M.J., Wills, M.R., and Sinclair, J.H. (2017). Latency-associated expression of human cytomegalovirus US28 attenuates cell signaling pathways to maintain latent infection. *MBio* **8**, e01754-17.
- Latasiewicz, J., Artz, A., Jing, D., Blanco, M.P., Currie, S.M., Avila, M.V., Schnoor, M., and Vestweber, D. (2017). HS1 deficiency impairs neutrophil recruitment in vivo and activation of the small GTPases Rac1 and Rap1. *J. Leukoc. Biol.* **101**, 1133–1142.
- Lau, B., Poole, E., Krishna, B., Sellart, I., Wills, M.R., Murphy, E., and Sinclair, J. (2016). The expression of human cytomegalovirus MicroRNA MiR-UL148D during latent infection in primary myeloid cells inhibits activin A-triggered secretion of IL-6. *Sci. Rep.* **6**, 31205.
- Low, H., Mukhamedova, N., Cui, H.L., McSharry, B.P., Avdic, S., Hoang, A., Ditiatkovski, M., Liu, Y., Fu, Y., Meikle, P.J., et al. (2016). Cytomegalovirus restructures lipid rafts via a US28/CDC42-mediated pathway, enhancing cholesterol efflux from host cells. *Cell Rep.* **16**, 186–200.
- Martin, J., Collot-Teixeira, S., McGregor, L., and McGregor, J.L. (2007). The dialogue between endothelial cells and monocytes/macrophages in vascular syndromes. *Curr. Pharm. Des.* **13**, 1751–1759.
- Maslin, C.L., Kedzierska, K., Webster, N.L., Muller, W.A., and Crowe, S.M. (2005). Transendothelial migration of monocytes: the underlying molecular mechanisms and consequences of HIV-1 infection. *Curr. HIV Res.* **3**, 303–317.
- Mason, G.M., Poole, E., Sissons, J.G., Wills, M.R., and Sinclair, J.H. (2012). Human cytomegalovirus latency alters the cellular secretome, inducing cluster of differentiation (CD)4+ T-cell migration and suppression of effector function. *Proc. Natl. Acad. Sci. U S A* **109**, 14538–14543.
- Melak, M., Plessner, M., and Grosse, R. (2017). Actin visualization at a glance. *J. Cell Sci.* **130**, 525–530.
- Mukherjee, S., Kim, J., Mooren, O.L., Shahan, S.T., Cohan, M., and Cooper, J.A. (2015). Role of cortactin homolog HS1 in transendothelial migration of natural killer cells. *PLoS One* **10**, e0118153.
- Nogalski, M.T., Chan, G., Stevenson, E.V., Gray, S., and Yurochko, A.D. (2011). Human cytomegalovirus-regulated paxillin in monocytes links cellular pathogenic motility to the process of viral entry. *J. Virol.* **85**, 1360–1369.
- Noguchi, K., Albarracin, D., Durantini, M.R., and Glasman, L.R. (2007). Who participates in which health promotion programs? A meta-analysis of motivations underlying enrollment and retention in HIV-prevention interventions. *Psychol. Bull.* **133**, 955–975.
- Park, S.W., Um, S.H., Lee, H.A., Kim, S.H., Sim, Y., Yim, S.Y., Seo, Y.S., and Ryu, H.S. (2016). Mycophenolate mofetil as an alternative treatment for autoimmune hepatitis. *Clin. Mol. Hepatol.* **22**, 281–285.
- Poole, E., Avdic, S., Hodkinson, J., Jackson, S., Wills, M., Slobedman, B., and Sinclair, J. (2014). Latency-associated viral interleukin-10 (IL-10) encoded by human cytomegalovirus modulates cellular IL-10 and CCL8 Secretion during latent infection through changes in the cellular microRNA hsa-miR-92a. *J. Virol.* **88**, 13947–13955.
- Poole, E., Lau, J.C., and Sinclair, J. (2015). Latent infection of myeloid progenitors by human cytomegalovirus protects cells from FAS-mediated apoptosis through the cellular IL-10/PEA-15 pathway. *J. Gen. Virol.* **96**, 2355–2359.
- Poole, E., McGregor Dallas, S.R., Colston, J., Joseph, R.S., and Sinclair, J. (2011). Virally induced changes in cellular microRNAs maintain latency of human cytomegalovirus in CD34 progenitors. *J. Gen. Virol.* **92**, 1539–1549.
- Poole, E., and Sinclair, J. (2015). Sleepless latency of human cytomegalovirus. *Med. Microbiol. Immunol.* **204**, 421–429.
- Poole, E., Walther, A., Raven, K., Benedict, C.A., Mason, G.M., and Sinclair, J. (2013). The myeloid transcription factor GATA-2 regulates the viral UL144 gene during human cytomegalovirus latency in an isolate-specific manner. *J. Virol.* **87**, 4261–4271.
- Reeves, M.B., Lehner, P.J., Sissons, J.G., and Sinclair, J.H. (2005a). An in vitro model for the regulation of human cytomegalovirus latency and reactivation in dendritic cells by chromatin remodelling. *J. Gen. Virol.* **86**, 2949–2954.
- Reeves, M.B., MacAry, P.A., Lehner, P.J., Sissons, J.G., and Sinclair, J.H. (2005b). Latency, chromatin remodeling, and reactivation of human cytomegalovirus in the dendritic cells of healthy carriers. *Proc. Natl. Acad. Sci. U S A* **102**, 4140–4145.

- Reinhardt, B., Godfrey, R., Fellbrich, G., Frank, H., Luske, A., Olieslagers, S., Mertens, T., and Waltenberger, J. (2014). Human cytomegalovirus infection impairs endothelial cell chemotaxis by disturbing VEGF signalling and actin polymerization. *Cardiovasc. Res.* *104*, 315–325.
- Rossetto, C.C., Tarrant-Elorza, M., and Pari, G.S. (2013). Cis and trans acting factors involved in human cytomegalovirus experimental and natural latent infection of CD14 (+) monocytes and CD34 (+) cells. *PLoS Pathog.* *9*, e1003366.
- Scielzo, C., Bertilaccio, M.T., Simonetti, G., Dagklis, A., ten Hacken, E., Fazi, C., Muzio, M., Caiolfa, V., Kitamura, D., Restuccia, U., et al. (2010). HS1 has a central role in the trafficking and homing of leukemic B cells. *Blood* *116*, 3537–3546.
- Shi, G., Fu, Y., Jiang, W., Yin, A., Feng, M., Wu, Y., Kawai, Y., Miyamori, I., and Fan, C. (2011). Activation of Src-ATF1 pathway is involved in upregulation of Nox1, a catalytic subunit of NADPH oxidase, by aldosterone. *Endocr. J.* *58*, 491–499.
- Shin, J.W., Suzuki, T., Ninomiya, N., Kishima, M., Hasegawa, Y., Kubosaki, A., Yabukami, H., Hayashizaki, Y., and Suzuki, H. (2012). Establishment of single-cell screening system for the rapid identification of transcriptional modulators involved in direct cell reprogramming. *Nucleic Acids Res.* *40*, e165.
- Sinclair, J., and Poole, E. (2014). Human cytomegalovirus latency and reactivation in and beyond the myeloid lineage. *Future Virol.* *6*, 7.
- Slobedman, B., and Cheung, A.K. (2008). Microarrays for the study of viral gene expression during human cytomegalovirus latent infection. *Methods Mol. Med.* *141*, 153–175.
- Sontag, J.M., Aunis, D., and Bader, M.F. (1988). Peripheral actin filaments control calcium-mediated catecholamine release from streptolysin-O-permeabilized chromaffin cells. *Eur. J. Cell Biol.* *46*, 316–326.
- Streblov, D.N., Soderberg-Naucler, C., Vieira, J., Smith, P., Wakabayashi, E., Ruchti, F., Mattison, K., Altschuler, Y., and Nelson, J.A. (1999). The human cytomegalovirus chemokine receptor US28 mediates vascular smooth muscle cell migration. *Cell* *99*, 511–520.
- Taylor-Wiedeman, J., Sissons, J.G., Borysiewicz, L.K., and Sinclair, J.H. (1991). Monocytes are a major site of persistence of human cytomegalovirus in peripheral blood mononuclear cells. *J. Gen. Virol.* *72* (Pt 9), 2059–2064.
- Thieler, H., Kohler, I., Achenbach, H., Goetz, K.H., Kraatz, G., Osten, B., and Smit, H. (1995). Further advances of chronic renal replacement therapy in eastern Germany, 1994 versus 1989. *Clin. Nephrol.* *44*, 108–112.
- Tseliou, M., Al-Qahtani, A., Alarifi, S., Alkahtani, S.H., Stourmaras, C., and Sourvinos, G. (2016). The role of RhoA, RhoB and RhoC GTPases in cell morphology, proliferation and migration in human cytomegalovirus (HCMV) infected glioblastoma cells. *Cell Physiol. Biochem.* *38*, 94–109.
- Tugizov, S., Herrera, R., Veluppillai, P., Greenspan, J., Greenspan, D., and Palefsky, J.M. (2007). Epstein-Barr virus (EBV)-infected monocytes facilitate dissemination of EBV within the oral mucosal epithelium. *J. Virol.* *81*, 5484–5496.
- Uruno, T., Zhang, P., Liu, J., Hao, J.J., and Zhan, X. (2003). Haematopoietic lineage cell-specific protein 1 (HS1) promotes actin-related protein (Arp) 2/3 complex-mediated actin polymerization. *Biochem. J.* *371*, 485–493.
- Westhorpe, C.L., Zhou, J., Webster, N.L., Kalionis, B., Lewin, S.R., Jaworowski, A., Muller, W.A., and Crowe, S.M. (2009). Effects of HIV-1 infection in vitro on transendothelial migration by monocytes and monocyte-derived macrophages. *J. Leukoc. Biol.* *85*, 1027–1035.
- Wilkie, A.R., Lawler, J.L., and Coen, D.M. (2016). A role for nuclear F-actin induction in human cytomegalovirus nuclear egress. *MBio* *7*, e01254-16.
- Yamanashi, Y., Okada, M., Semba, T., Yamori, T., Umemori, H., Tsunasawa, S., Toyoshima, K., Kitamura, D., Watanabe, T., and Yamamoto, T. (1993). Identification of HS1 protein as a major substrate of protein-tyrosine kinase(s) upon B-cell antigen receptor-mediated signaling. *Proc. Natl. Acad. Sci. U S A* *90*, 3631–3635.
- Zhao, J., Mitrofan, C.G., Appleby, S.L., Morrell, N.W., and Lever, A.M. (2016). Disrupted endothelial cell layer and exposed extracellular matrix proteins promote capture of late outgrowth endothelial progenitor cells. *Stem Cells Int.* *2016*, 1406304.

ISCI, Volume 20

## **Supplemental Information**

### **Human Cytomegalovirus Upregulates Expression of HCLS1 Resulting in Increased Cell Motility and Transendothelial Migration during Latency**

**Yusuf Aslam, James Williamson, Veronika Romashova, Elizabeth Elder, Benjamin Krishna, Mark Wills, Paul Lehner, John Sinclair, and Emma Poole**

## **Supplementary figures and methods**

### **Supplementary Figure Legends**

Supplementary figure 1 related to figure 3

A) CD14<sup>+</sup> monocytes were infected with TB40E-GATA2mCherry and latency established for 4 days before analysing by cellomics over a period of 18h, as shown. Video images were also acquired of the cells over this period (A). Alternatively, monocytes were electroporated with a plasmid expressing GFP or HCLS1-GFP for 24h before analysing by cellomics over a period of 18h as shown. Videos were also taken of the cells over this time period (B).

Supplementary figure 2 related to figure 4

Monocytes were electroporated with plasmids expressing GFP, HCLS1-GFP or an siRNA conjugated to FAM (for visualisation to assess transfection efficiency) and then analysed for green fluorescence 24h later (A). Cells which had been electroporated with either a control siRNA or an siRNA to HCLS1 were harvested for the western blot of actin and HCLS1 proteins (B).

Supplementary figure 3 relating to figures 1, 2, 3 and 4



HCMV latency validation. Primary CD14<sup>+</sup> monocytes were either uninfected or infected with HCMV for 4 days before analysing by RT-qPCR (A) and then either leaving undifferentiated or differentiating into DCs using GM-CSF/IL-4. After this time supernatants were taken and transferred onto indicator fibroblasts to assess for virus release by infectious foci detection (B).

### **Supplementary figures**


Supplementary movies are in a separate file




Supplementary figure 1



Supplementary figure 2



## Supplementary figure 3



### Transparent Methods

#### *Cells and viruses*

Primary monocytes were isolated from apheresis cones or venous blood as described previously (Mason et al., 2012). HAEC (Promcell) and HUVEC (ATCC) were cultured as described by the suppliers. Human Foreskin Foetal Fibroblasts (HFFFs) were obtained from ATCC and cultured as described previously (Poole et al., 2006).

TB40E-SV40-GFP and TB40E-GATA2mCherry viruses have been described previously (Elder, 2019; Krishna et al., 2016).

Titan WT and Titan US28 deletion virus have been described previously (Krishna et al., 2017a)

## Ethical Statement

All research describing studies on primary human material with HCMV were assessed and approved by the Cambridge Local Research Ethics committee. Informed consent was received from blood donors with the Cambridge Local Research Ethics committee and the Cambridge Internal Review Board. Cells were harvested from healthy adult donors, and the decision to use tissue was not affected by gender and age, as this was not important to the studies performed.

### *Generation of CTCF shRNA and US28-expressing cell lines by lentiviral transduction*

pHRsinGKpuro and pHRsinUbEm lentiviral expression vectors were a kind gift from Dr D van den Boomen, University of Cambridge. The sequence encoding US28 was cloned into pHRsinUbEm using the BamHI and NotI sites. The lentiviral expression vector encoding CTCFshRNA and puromycin resistance (Santa Cruz Biotechnology).

To generate lentiviral particles, 293T cells were seeded into 6-well plates at  $5 \times 10^5$  cells/well. Approximately 6 hours later, 1250 ng of lentiviral expression vector, 625 ng of lentiviral packaging vector psPAX and 625 ng of lentiviral envelope vector pMD.2G (both gifts from S. Karniely, Weizmann Institute, Israel) were transfected into 293T cells using transfection reagent FuGene6 (Promega) according to the manufacturer's instructions. 24h post transduction, media was replaced with 2.5mL RPMI supplemented with 30% fetal calf serum. 24h post media change,  $2.5 \times 10^5$  THP-1 cells were pelleted, and then resuspended in the lentiviral or control supernatant in a 6-well plate. Polybrene was added to the cells at 2  $\mu\text{g}/\text{mL}$  and the cells were then centrifuged in the plate at 600xg for 45 minutes, and incubated overnight at 37°C/5% CO<sub>2</sub>. 5 days post transduction the transduced THP-1 cells were pelleted and resuspended in fresh RPMI supplemented with 10% fetal calf serum. 7 days post transduction, puromycin (Sigma) was added at 2  $\mu\text{g}/\text{mL}$  and the selective media was refreshed every 2 days GFP positive cells were sorted by FACS.

### *In silico transcription factor binding analysis*

The HCLS1 enhancer binding sequence was obtained from Genecards <https://www.genecards.org/cgi-bin/carddisp.pl?gene=HCLS1> and the identified transcription factor, ATF1, binding site was confirmed using Physbinder <http://bioit.dnbr.ugent.be/physbinder/>

### *HCMV latent infection*

To establish latency, primary monocytes or a monocytic cell line were infected for 3h with virus at a predicted MOI of 1 and then left for a minimum of 4 days before analysing for latent infection. This was determined by RTqPCR for the presence of the latency associated gene UL138 in the relative absence of the lytic gene IE. Additionally, supernatants were assessed for virus release pre- and post-differentiation into dendritic cells using GM-CSF and IL-4 as previously described (Poole et al., 2014a). An example analysis is shown in supplementary figure 3. We define latency as the presence of latency associated transcripts such as UL138 in the relative absence of the lytic gene IE (Supplementary figure 3A). Crucially, during latency there is no virion production as measured by co-culture of latently infected cell/cell supernatants on fully permissive indicator fibroblasts (Supplementary figure 3B). Latently infected cells used in all our experiments were consistent with this definition of latency.

#### *HCLS1 RT-qPCR*

HCLS1 and GAPDH commercial primers were obtained from Qiagen and then RNA was isolated from the indicated samples using the Qiagen RNA isolation kit followed by SYBR green RT-qPCR analysis using standard parameters (Qiagen).

#### *Plasmids, siRNAs and transfection*

Plasmids and siRNAs were electroporated into primary monocytes using the Amaxa monocyte transfection kit (Lonza). GFP plasmid was obtained from Lonza whereas HCLS1-GFP plasmid and siRNA scramble and siRNA-HCLS1 were obtained from OriGene and FAM labelled control siRNA from ThermoFisher. 1ug of each plasmid or 50nM of each siRNA was electroporated into  $3 \times 10^6$  monocytes by Amaxa.

*Western blotting, and immunofluorescence* Western blotting and immunofluorescence were carried out as described previously (Poole et al., 2012; Poole et al., 2014b). HCLS1 primary antibody was used for both western blotting and immunofluorescence (rabbit polyclonal, abcam). Actin antibody for western blotting was obtained from Abcam (rabbit polyclonal) both as directed by the manufacturer. Western blots were analysed using autoradiographic film and immunofluorescence computerized image analysis software (Image ProPlus and Image J). The immunofluorescence method has been described previously (Poole et al., 2006).

#### *Actin filament analysis and drug inductions*

To assess actin filament formation, cells were first fixed and then stained with phalloidin which is specific for F-actin using a previously described method (Melak et al., 2017). CD14+ monocytes in

which latency had been established for 6 days with either GATA2mCherryTB40E were either fixed and stained directly with the F-actin specific marker phalloidin 488 or phalloidin 594 (CytoPainter, Abcam) respectively, in the presence of Hoechst. Alternatively, cells were treated with 200nM cytochalasin D (a concentration that is reversible for actin filament formation; (Domnina et al., 1982)) for 1h before washing, staining and visualising by fluorescence microscopy using computerized image analysis software (Image ProPlus and Image J). MMF and PP2 were obtained from Sigma and used at the recommended concentrations from the supplier and as described previously (Melak et al., 2017) (Chan et al., 2012). Specifically, cells were treated with 1uM PP2 1h post and during the establishment of HCMV latency. The media was changed every 24h with fresh PP2 added to the media.

*Chromatin Immunoprecipitation (ChIP)* The HCLS1 promoter was analysed by ChIP using primers to the enhancer: Forward -TAAAAGAACTCACCCCTA and Reverse - AGATTAGGAAAGGCTAGGTC using the Imprint ChIP kit (Sigma) following the manufacturer instructions.

#### *Endothelial migration*

Monocyte to transmigration across the endothelial layer was analysed as previously described (Bentz et al., 2006). In brief, endothelial cells were seeded onto the filters of transwell plates before carrying out standard monocyte transwell analysis (Mason et al., 2012). The CD14+ monocytes in which HCMV latency had been established were isolated by live cell sorting (Krishna et al., 2017b; Lau et al., 2016).  $1 \times 10^4$  infected or uninfected cells were added to each transwell and after 2 hours the monocytes which had migrated through the endothelial cell layer were enumerated by bright field microscopy using computerized image analysis software (Image ProPlus and Image J).

#### *Endothelial Flow analysis*

Adhesion of monocytes to the endothelial cell layer was determined by flow analysis using previously published methods (Butler et al., 2011; Zhao et al., 2016). In brief, endothelial cells were seeded into Ibidi slides VI<sup>0.4</sup>. When the cells reached confluency,  $1 \times 10^6$  monocytes were perfused over the endothelial cell layer, within a Perspex chamber maintained at 37degrees, at the physiological flow rate of 0.4ml/min. The flow rate of 0.4 mL/min was pump-controlled. The equivalent shear stress ( $\tau$ ) exerted on the Ibidi  $\mu$ -Slide VI<sup>0.4</sup> slide surface at a flow rate of 0.4 mL/min ( $\Phi$ ) was  $0.7 \text{ dyn/cm}^2$ , calculated from the equation " $\tau = \eta 176.1\Phi$ ".  $\eta$  (dynamical viscosity) was

0.01 dyn·s/cm<sup>2</sup>. CD14+ monocytes were then washed in PBS before fixing in 4% PFA in PBS and visualising by fluorescence using computerized image analysis software (Image ProPlus and Image J).

### *Proteomic screen*

#### Lysis, digestion and clean-up of protein preps

From 6 well plates, cells were washed 2x with cold PBS before scraping cells into 200uL 8Murea/50mM TEAB pH 8.5. Samples were quantified by BCA assay and 28ug of each sample was taken and adjusted to the same volume with lysis buffer. Reduction and alkylation was achieved by addition of 10mM TCEP and 20mM iodoacetamide for 20mins at room temperature in the dark followed by quenching with 10mM DTT. 3ug of rLysC (Promega) was added to each sample and incubated at 30 degrees for 3h before diluting the 1:5 with 50mM TEAB and addition of 3ug trypsin (Proteomics Grade, Thermo Fisher Scientific) and incubation at 37 degrees overnight. Samples were acidified by adding 1 volume of 0.2% TFA and formic acid until pH was ~2. Samples were desalted using in-house prepared microcolumns consisting of Oligo R3 resin (SCIEX) packed behind a C18 plug (Empore, 3M) in p200 pipette tips. Columns were washed with 100uL ACN and equilibrated with 2x100uL 0.1% TFA before loading of samples (3 passes over the column). Columns were washed with 3x 50uL 0.1% TFA before sequential elution with 20uL each of 40% ACN, 70% ACN and 70% ACN with 1% FA. Samples were dried in a vacuum centrifuge.

#### TMT Labelling

Samples were resuspended in 20uL 100mM TEAB and to each tube 0.2ug of a unique TMT label for each sample was added in 8.5uL acetonitrile and incubated for 1h at room temperature. Labels were as follows: TMT reactions were quenched by addition of 3uL of 200mM ammonium formate, pooled and dried in a vacuum centrifuge. The sample was then Resuspended in 800uL 0.1% TFA and acidified to ~pH2 with formic acid before performing a C18-SPE cleanup using a Sep-Pak cartridge (Waters) attached to a vacuum manifold. C18 Eluate was dried in a vacuum centrifuge and resuspended in 40uL 200mM ammonium formate, pH10.

#### High pH Reversed Phase Fractionation

Sample was injected onto an Ultimate 3000 RSLC UHPLC system (Thermo Fisher Scientific) equipped with a 2.1 i.d x25cm, 1.7uM particle Kinetix Evo C18 column (Phenomenex). Mobile phase consisted of A: 3% ACN, B:ACN and C: 200mM ammonium formate pH 10. Isocratic conditions were 90% A/10%C and C was maintained at 10% throughout the gradient elution. Separations were carried out at 45 degrees. After loading at 200uL/min for 5 mins and ramping the flow rate to 400uL/min over

5mins the gradient elution proceed as follows: 0-19% B over 10 minutes (curve 3), 19-34%B over 14.25mins (curve 5), 34-50%B over 8.75mins (curve 5), followed by a 10 min wash at 90% B. UV absorbance was monitored at 280nm and 15s fractions were collected into 96 well microplates using the integrated fraction collector. Peptide containing fractions were then orthogonally recombined into 12 fractions and dried in a vacuum centrifuge and resuspended in 10 $\mu$ L 5% DMSO 0.5% TFA for analysis.

### LC-MS analysis

All samples were injected onto an Ultimate 3000 RSLC nano UHPLC equipped with a 300 $\mu$ m i.d. x 5mm Acclaim PepMap  $\mu$ -Precolumn (Thermo Fisher Scientific) and a 75 $\mu$ m i.d. x50cm 2.1 $\mu$ m particle Acclaim PepMap RSLC analytical column. Loading solvent was 0.1% TFA, analytical solvent A: 0.1% FA and B: ACN+0.1% FA. All separations are carried out at 55 degrees. Samples were loaded at 10 $\mu$ L/min for 5 mins in loading solvent before beginning the analytical gradient. For High pH RP fractions a gradient of 3-5.6% B over 4 mins, 5.6 – 32%B over 162mins, followed by a 5 minute wash at 80%B and a 5 minute wash at 90%B and equilibration at 3%B for 5mins. During the gradient the Orbitrap Fusion mass spectrometer (Thermo Fisher Scientific) was set to acquire spectra.

### Data Processing

All Raw files were searched by Mascot within Proteome Discoverer 2.1 (Thermo Fisher Scientific) against the Swissprot Human database and a database of common contaminants.

For TMT labelled samples the search parameters were as follows. Enzyme: Trypsin. MS1 tol: 10ppm. MS2 tol: 0.6Da. Fixed modifications: Carbamidomethyl Cysteine, TMT peptide N-termini and Lysine. Variable modification oxidised methionine. MS3 reporter ion tol: 20ppm, most confident centroid. Mascot Percolator was used to calculate PSM FDR.

Search results were further processed and filtered as follows: Peptides below a percolator FDR of 0.01% and proteins below the 0.01% protein FDR (calculated from a built in decoy database search) were rejected. Protein groups were then generated using the strict parsimony principle. Peptides both unique and razor with a co-isolation threshold of 50 and an average s/n threshold of 10 were used for quantification and a normalisation of these values to the total peptide amount in each channel was applied. Instances where a protein was identified but not quantified in all channels were rejected from further analysis. "Scaled" abundances of proteins provided by Proteome Discoverer were used to derive ratios of abundance.

Bentz, G.L., Jarquin-Pardo, M., Chan, G., Smith, M.S., Sinzger, C., and Yurochko, A.D. (2006). Human cytomegalovirus (HCMV) infection of endothelial cells promotes naive monocyte extravasation and transfer of productive virus to enhance hematogenous dissemination of HCMV. *J Virol* *80*, 11539-11555.

Butler, L.M., Jeffery, H.C., Wheat, R.L., Rae, P.C., Townsend, K., Alkharsah, K.R., Schulz, T.F., Nash, G.B., and Blackbourn, D.J. (2011). Kaposi's sarcoma-associated herpesvirus infection of endothelial cells inhibits neutrophil recruitment through an interleukin-6-dependent mechanism: a new paradigm for viral immune evasion. *J Virol* *85*, 7321-7332.

Chan, G., Nogalski, M.T., Stevenson, E.V., and Yurochko, A.D. (2012). Human cytomegalovirus induction of a unique signalsome during viral entry into monocytes mediates distinct functional changes: a strategy for viral dissemination. *J Leukoc Biol* *92*, 743-752.

Domnina, L.V., Gelfand, V.I., Ivanova, O.Y., Leonova, E.V., Pletjushkina, O.Y., Vasiliev, J.M., and Gelfand, I.M. (1982). Effects of small doses of cytochalasins on fibroblasts: preferential changes of active edges and focal contacts. *Proc Natl Acad Sci U S A* *79*, 7754-7757.

Elder, E., Krishna, B., Williamson, J., Aslam, Y., Farahi, N., Wood, A., Romashova, V., Roche, K., Murphy, E., Chilvers, E., Lehner, P., Sinclair, J., Poole, E. (2019). Monocytes latently infected with human cytomegalovirus evade neutrophil killing. *iScience*.

Krishna, B.A., Lau, B., Jackson, S.E., Wills, M.R., Sinclair, J.H., and Poole, E. (2016). Transient activation of human cytomegalovirus lytic gene expression during latency allows cytotoxic T cell killing of latently infected cells. *Sci Rep* *6*, 24674.

Krishna, B.A., Poole, E.L., Jackson, S.E., Smit, M.J., Wills, M.R., and Sinclair, J.H. (2017a). Latency-Associated Expression of Human Cytomegalovirus US28 Attenuates Cell Signaling Pathways To Maintain Latent Infection. *MBio* *8*.

Krishna, B.A., Spiess, K., Poole, E.L., Lau, B., Voigt, S., Kledal, T.N., Rosenkilde, M.M., and Sinclair, J.H. (2017b). Targeting the latent cytomegalovirus reservoir with an antiviral fusion toxin protein. *Nat Commun* *8*, 14321.

Lau, B., Poole, E., Krishna, B., Sellart, I., Wills, M.R., Murphy, E., and Sinclair, J. (2016). The Expression of Human Cytomegalovirus MicroRNA MiR-UL148D during Latent Infection in Primary Myeloid Cells Inhibits Activin A-triggered Secretion of IL-6. *Sci Rep* *6*, 31205.

Mason, G.M., Poole, E., Sissons, J.G., Wills, M.R., and Sinclair, J.H. (2012). Human cytomegalovirus latency alters the cellular secretome, inducing cluster of differentiation (CD)4<sup>+</sup> T-cell migration and suppression of effector function. *Proc Natl Acad Sci U S A*.

Melak, M., Plessner, M., and Grosse, R. (2017). Actin visualization at a glance. *J Cell Sci* *130*, 525-530.

Poole, E., Avdic, S., Hodkinson, J., Jackson, S., Wills, M., Slobedman, B., and Sinclair, J. (2014a). Latency-associated viral interleukin-10 (IL-10) encoded by human cytomegalovirus modulates cellular IL-10 and CCL8 Secretion during latent infection through changes in the cellular microRNA hsa-miR-92a. *J Virol* *88*, 13947-13955.

Poole, E., Bain, M., Teague, L., Takei, Y., Laskey, R., and Sinclair, J. (2012). The cellular protein MCM3AP is required for inhibition of cellular DNA synthesis by the IE86 protein of human cytomegalovirus. *PLoS One* *7*, e45686.

Poole, E., Juss, J.K., Krishna, B., Herre, J., Chilvers, E.R., and Sinclair, J. (2014b). Alveolar macrophages isolated directly from HCMV seropositive individuals are sites of HCMV reactivation in vivo. *J Infect Dis*.

Poole, E., King, C.A., Sinclair, J.H., and Alcami, A. (2006). The UL144 gene product of human cytomegalovirus activates NFkappaB via a TRAF6-dependent mechanism. *Embo J* *25*, 4390-4399.

Zhao, J., Mitrofan, C.G., Appleby, S.L., Morrell, N.W., and Lever, A.M. (2016). Disrupted Endothelial Cell Layer and Exposed Extracellular Matrix Proteins Promote Capture of Late Outgrowth Endothelial Progenitor Cells. *Stem Cells Int* *2016*, 1406304.