

## Prevalence of Hypertension in Renal Diseases in Iran: Systematic Review and Meta-Analysis

### Abstract

**Background:** Hypertension is a risk factor for renal disease. Therefore, this study was aimed at estimating the prevalence of hypertension in renal patients in Iran through meta-analysis. **Methods:** The search was carried out using authentic Persian and English keywords in national and international databases including IranMedex, SID, Magiran, IranDoc, Medlib, ScienceDirect, Pubmed, Scopus, Cochrane, Embase, Web of Science, Medline, and Google Scholar search engine without any time limitation until 2017. Heterogeneity of studies was assessed using the  $I^2$  index. Data were analyzed using STATA ver 11. **Results:** In 35 reviewed studies with a sample of 39,621 subjects, the prevalence of hypertension in renal patients was 35% (95% CI: 29%–41%) (25% in women and 18% in men). The prevalence of systolic hypertension in renal patients was 5%, diastolic hypertension 26%, and diabetes 23%. The prevalence of hypertension in hemodialysis patients was 34%, 27% in peritoneal dialysis, 43% in kidney transplantation, and 26% in chronic renal failure. In addition, meta-regression showed that the prevalence of hypertension in renal patients did not significantly decrease during the years 1988–2017. **Conclusions:** More than a third of kidney patients in Iran suffer from high blood pressure. The diastolic blood pressure of these patients is about five times higher than their systolic blood pressure. Moreover, the age group under 30 is a high-risk group. The prevalence of hypertension in women with kidney disease is higher than in men. In addition, patients who have kidney transplants are more likely to have high blood pressure than other kidney patients.

**Keywords:** Hypertension, Iran, kidney disease, meta-analysis, renal disease

### Introduction

Among the selected articles from 1999 to 2012, the prevalence of hypertension was 17%. The prevalence of hypertension among the people above 20 years was 24% and that among the people below 20 years was 5%.<sup>[1]</sup> In 32 studies with a sample of 34,714 subjects, the prevalence of hypertension in Iranian diabetic patients was 51% (95% CI: 43%–60%). The prevalence of hypertension was 55% in type I diabetics and 53% in type II diabetic patients.<sup>[2]</sup> In Haghdoost study, in 2006, the prevalence of hypertension in 29 studies was 21.9%.<sup>[3]</sup>

Today, chronic disease is one of the most important health problems affecting different dimensions of physical, psychological, economic, social, and quality of life of patients.<sup>[4–9]</sup> Hypertension is a common chronic disease that underlies several diseases such as heart attack, brain attack,

and advanced kidney disease,<sup>[10–17]</sup> which is defined as systolic blood pressure above 140 mmHg and diastolic blood pressure above 90 mmHg.<sup>[12]</sup> As a global problem, it is a common asymptomatic disease, often called a silent killer.<sup>[18]</sup> According to the World Health Organization, high blood pressure after smoking has been introduced as the second leading cause of death in developing countries and is the third most common cause of noncommunicable diseases. This disease has different prevalence in different societies and the prevalence ranges from 10% to more than 60% in different countries.<sup>[19,20]</sup> It has also been estimated that blood pressure has been responsible for 4.5% of the global burden of diseases.<sup>[21–24]</sup>

End-stage renal disease (ESRD) is the severe type of chronic kidney disease, which is the final stage of an irreversible progressive renal disorder. In this disease, the body's ability to maintain the balance between fluid and electrolyte disappears

**Morteza Motedayen,  
Diana Sarokhani<sup>1</sup>,  
Bahareh Ghiasi<sup>2</sup>,  
Alireza Khatony<sup>3</sup>,  
Ali Hasanpour  
Dehkordi<sup>4</sup>**

*Department of Cardiology,  
Faculty of Medicine, Zanjan  
University of Medical Sciences,  
Zanjan, Iran, <sup>1</sup>Psychosocial  
Injuries Research Center, Ilam  
University of Medical Sciences,  
Ilam, Iran, <sup>2</sup>Department  
of Nephrology, Faculty of  
Medicine, Ilam University of  
Medical Sciences, Ilam, Iran,  
<sup>3</sup>Social Development and Health  
Promotion Research Center,  
Kermanshah University of  
Medical Sciences, Kermanshah,  
Iran, <sup>4</sup>Social Determinants of  
Health Research Center, School  
of Allied Medical Sciences,  
Shahrekord University of  
Medical Sciences,  
Shahrekord, Iran*

### Address for correspondence:

*Dr. Alireza Khatony,  
Social Development and Health  
Promotion Research Center,  
Kermanshah University of  
Medical Sciences,  
Kermanshah, Iran.  
E-mail: Akhatony@gmail.com  
Dr. Bahareh Ghiasi,  
Department of Nephrology,  
Faculty of Medicine, Ilam  
University of Medical Sciences,  
Ilam, Iran.  
E-mail: bahar55gh@yahoo.co*

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

**How to cite this article:** Motedayen M, Sarokhani D, Ghiasi B, Khatony A, Hasanpour Dehkordi A. Prevalence of hypertension in renal diseases in Iran: Systematic review and meta-analysis. *Int J Prev Med* 2019;10:124.

### Access this article online

**Website:**  
[www.ijpvmjournal.net/www.ijpvm.ir](http://www.ijpvmjournal.net/www.ijpvm.ir)

**DOI:**  
10.4103/ijpvm.IJPVM\_522\_18

### Quick Response Code:



and leads to uremia or asthenia.<sup>[25–31]</sup> In fact, when more than 95% of the kidney tissue is destroyed for various reasons, the accumulation of toxins in the body increases considerably and life-threatening complications force the person to undergo Renal replacement therapies forever.<sup>[32,33]</sup> The population of patients with kidney failure in Iran is 320,000; 49% of them use renal transplantation therapy, 48% of them use the hemodialysis method, and 3% of them use the peritoneal dialysis method.<sup>[34]</sup> This disease can lead to death if the patient does not undergo dialysis or kidney transplantation.<sup>[25–27,35–37]</sup>

In 1990, renal failure was considered as the 27<sup>th</sup> mortality factor in the world, and reached the 18<sup>th</sup> rank in 2010.<sup>[38]</sup> According to the Centers for Disease Control and Prevention, in 2014, more than 20% of people with serious hypertension suffered from chronic kidney disease and were at risk for ESRD.<sup>[39]</sup> On the other hand, about 75%–80% of patients with ESRD have high blood pressure.<sup>[40]</sup> In all developed countries and in many developing countries, diabetes and hypertension are considered as the main cause of chronic kidney disease.<sup>[23,36,41–48]</sup>

Considering that published articles about the prevalence of hypertension in renal patients in Iran reported varied incidence of 9%–86% for the prevalence of high blood pressure, the need for a meta-analysis seemed necessary.

## Materials and Methods

### Study protocol

This study is a systematic review and meta-analysis study that the prevalence of hypertension in Iranian renal patients. This study was conducted on the basis of the PRISMA<sup>1</sup> statement that is concerned with systematic review and meta-analysis studies.<sup>[49]</sup> Based on this protocol, all stages of the research methodology such as search, selection of studies and qualitative assessment of studies, and data extraction from the studies were conducted by two researchers independently. If there was a difference in the report of the researchers, the third researcher investigated and resolved the dispute.

### Search strategy

This study is a meta-analysis that investigates the prevalence of hypertension in renal patients. The findings of this study were based on studies conducted in Iran. We searched national databases including IranMedex, SID, Magiran, IranDoc and Medlib, and international databases including ScienceDirect, PubMed, Scopus, Cochrane, Embase, Web of Science, and Medline. The mechanism for searching articles mainly based on the systematic search of related Persian keywords and their English equivalents (“Iran,” “Meta-analysis,” “Kidney disease,” “Renal disease,” “Hypertension”) was carried out independently by two researchers. For the final analysis, keywords were also searched in the Google Scholar search engine without time

limit until 2017. It should be noted that the keywords were also searched together using (OR/AND) operators.

### Study selection

First, all articles related to this subject in Iran were collected and a list of abstracts was prepared after the search was completed. This task was done independently by two researchers. Then, the articles with duplicate titles were excluded. Subsequently, the abstracts of articles were reviewed to find appropriate studies. Regarding international databases, the process was similar to that of national databases.

In the first stage of the search, 732 articles were found, and after reviewing the titles of articles, 285 overlapping articles were removed. Among the remaining 447 papers, 229 articles were excluded from the exit criteria. The following is a list of the exclusion reasons: (1) the study was performed on other patients, except for renal patients; (2) the study was conducted in a country other than Iran; (3) the articles looked at risk factors or blood pressure complications. The full text of 218 articles was reviewed. Among them, 182 articles were excluded, because they were incomplete or their full text was not available. In the remaining 36 articles, 1 article was removed because of low quality. Finally, 35 appropriate articles were selected to enter the meta-analysis stage [Figure 1].

### Quality evaluation


All studies that reported the prevalence of osteoporosis in Iranian elderly people entered the study. To assess the quality of the studies, STROBE (3) was used as the standard checklist. This checklist contains 22 items that cover different parts of a report (sampling, measurement of variables, study objectives, and statistical analysis). One point was given to each item, and some other items that were more important to us had more points. The STROBE checklist contains 22 sections that cover different parts of a report, and the maximum score of a report equals 44, so that a score of 1–15 indicates poor quality, 16–30 shows average quality, and 31–44 is considered to be excellent. Articles with a total score of less than 16 were excluded from the meta-analysis.

### Data extraction

To reduce reporting bias and error in data collection, two researchers independently extracted data from articles and entered the extracted data into a checklist containing the following items: first author’s name, study title, sample size, year and place of study, prevalence of hypertension in renal patients based on sex, prevalence of systolic and diastolic hypertension, prevalence of diabetes, Type of disease, age, etc.

### Statistical analysis

To analyze and combine the results of various studies, the prevalence of hypertension in renal patients in each study



**Figure 1:** Prevalence of hypertension in renal patients in Iran with a 95% confidence interval based on the author’s name and year of research according to the random effects model. The midpoint of each section shows the prevalence of hypertension in renal patients in each study. The lozenge shows the prevalence of hypertension in renal patients in Iran for all studies

was considered as a binomial probability distribution and its variance was calculated by binomial distribution. The heterogeneity of studies was measured using the Q test and I square ( $I^2$ ) index. Due to the heterogeneity of the studies, the random effects model was used to combine the results of the studies. The data were analyzed using STATA Ver. 11, and the significance level of the test was considered as 0.05.

Meta-regression was used to investigate the relationship between the prevalence of hypertension in renal patients and the sample size and year of research. Sensitivity analysis was used to determine the effect of removing each study on the final meta-analysis.

## Results

### Summary of how to enter the articles into the meta-analysis process

In 35 articles with a sample size of 39,621 people, the prevalence of hypertension in Iranian renal patients was 35% (95% CI: 29%–41%). The lowest and highest prevalence of hypertension were 9% in Brahimi’s study<sup>[50]</sup> and 86% in Fazelzadeh’s study,<sup>[51]</sup> respectively. Considering the heterogeneity of studies, the confidence interval (CI) for

each study based on the random effects model is presented in Diagram 1. The characteristics of the reviewed articles are presented in Table 1.

The prevalence of hypertension in women with renal disease was 25% (95% CI: 16%–35%) and in men was 18% (95% CI: 15%–22%). In addition, the prevalence of systolic hypertension in renal patients was 5% (95% CI: 0%–13%), the prevalence of diastolic hypertension was 26% (95% CI: 0%–77%), and the prevalence of diabetes was 23% (95% CI: 17%–29%). The prevalence of hypertension in women with renal disease is 7% higher than men [Table 2].

In an analysis based on the type of renal disease, we found that the prevalence of hypertension in hemodialysis patients was 34% (95% CI: 23%–45%) in 20 reviewed studies. In nine other studies performed on renal transplant patients, the prevalence of hypertension was 43% (95% CI: 19%–67%). Two studies were also performed on peritoneal dialysis patients with a hypertension prevalence of 27% (95% CI: 16%–38%). The three last studies were conducted on chronic renal failure patients and we found that the prevalence of hypertension was 26% (95% CI: 17%–34%). The

**Table 1: Characteristics of Studies Qualified for Systematic Review and Meta-Analysis**

Author	Age mean (SD)	Year of study	City of study	Type of disease	Sample size	Prevalence of hypertension (%)
[52] Bashardoost	53 (16.7)	2006	Ardebil	Hemodialysis Patients	80	64.9
[53] Hasan Zamani	48 (18.03)	2011	Mashhad	Hemodialysis Patients	66	22.7
[54] Pourmand	39.3	2015	Tehran	Kidney transplantation patients	400	26.3
[55] Basiratnia	17.4	2011	Shiraz	Kidney transplantation patients	66	25.7
[56] Shamsa	35.26	2003-2004	Mashhad	Kidney transplantation patients	15	40
[51] Fazel zadeh	45	1988-1992	Shiraz	Kidney transplantation patients	500	86
[57] Houli saz	40.26	2006	Tehran	Kidney transplantation patients	119	63
[58] Nikoubakht	40.45	2001-2002	Tehran	Kidney transplantation patients	119	50
[59] Noushad	48	2003-2005	Tabriz	Kidney transplantation patients	50	26
[60] Rostami	40.7	2006-2009	Tehran	Kidney transplantation patients	804	58.2
[61] Pak fetrat	38.5	2000-2001	Shiraz	Kidney transplantation patients	1354	11.6
[62] Rezaei ghalehchi	25-70	2013	Ardebil	Hemodialysis Patients	100	38.2
[63] Noubakht haghghi	30.43	2005-2008	Iran	Donor nephrectomies	1549	-
[64] Rouhani	61.3	2012	Tehran	Hemodialysis Patients	163	49.1
[65] Najafi	46.6	2001-2009	Iran	peritoneal dialysis	2302	24.4
[50] Barahimi	>30	2011	Shahreza	Chronic kidney disease	11720	9
[66] Gheisari	11.01	2001-2011	Esfahan	Chronic kidney disease	268	55
[67] Jalal zadeh	56.6	2009-2010	Zanjan	Hemodialysis Patients	80	68.8
[68] Neamati	62	2006	Tehran, Ahvaz, Kermanshah, Mashhad, shiraz	Hemodialysis Patients	337	23.7
[69] Safari nezhad	>14	2002-2005	Iran	Chronic kidney disease	16354	17.3
[70] Ayazi	51.2	2002-2003	Tehran	peritoneal dialysis	21	38
[71] Nakhjavani	32-89	2001-2002	Tehran	Hemodialysis Patients	620	82.5
[72] Barzin	49	1998	Sari	Hemodialysis Patients	41	17
[73] Monfarad		2001	Gilan	Hemodialysis Patients	414	17
[74] Shasti	62.2 (13.7)	2010	Tehran	Hemodialysis Patients	100	26.1
[75] Raeisi far	51.48	2005	Abadan	Hemodialysis Patients	59	40.4
[76] Ghafourifard	48.6 (19)	2008	Esfahan	Hemodialysis Patients	26	11.5
[77] Heidari	47	1998	Mazandaran	Hemodialysis Patients	353	24.6
[78] Mogharab	56-65	2007	Birjand	Hemodialysis Patients	40	12.5
[79] Eslami amirabadi	50.4 (15.2)	2014	Tehran	Hemodialysis Patients	189	23.3
[80] Rahimian	56.7	2005	Yazd	Hemodialysis Patients	60	12
[81] Ghorbani Moghaddam1	55.7 (15.63)	2014	Bushehr	Hemodialysis Patients	93	34.2
[82] Hadian	53.2 (16.4)	2011	Lorestan	Hemodialysis Patients	318	38.1
[83] Rezaein langroodi	56-65	2009	Hamedan	Hemodialysis Patients	455	22.1
[84] Khadem Vatan	55 (17.7)	2014	Orumie	Hemodialysis Patients	386	35.5

prevalence of hypertension in renal transplant patients is higher than that of hemodialysis, peritoneal, and chronic kidney dialysis patients [Table 2].

In an analysis performed based on the age group, the prevalence of hypertension was 41% (95% CI: 12%–69%) among 1–29-year-old renal patients, 22% (95% CI: 9%–36%) among 30–39-year-old patients, 39% (95% CI: 20%–57%) among 40–49-year-old renal patients, 35% (95% CI: 26%–44%) among 50–59-year-old renal patients, and 32% (95% CI: 16%–48%) among 60–70-year-old renal patients. Moreover, 1–29-year-old renal patients are more likely to have hypertension compared with other patients; however, given the varying number of studies in any age group, we cannot say with certainty [Table 2].


The prevalence of hypertension in renal patients in the north of Iran was 42% (95% CI: 28%–55%), in the south was 37% (95% CI: 29%–44%), in the center was 30% (95% CI: 10%–50%), and in the west was 38% (95% CI: 25%–50%). Renal patients in the north of Iran suffer more from high blood pressure compared with other regions of Iran [Table 2].

In sensitivity analysis, we find that the prevalence of hypertension in renal patients in Iran was reduced to 33.65% (95% CI: 28.55%–38.75%) by removing the study of Fazelzadeh<sup>[20]</sup> and increased to 36.16% (95% CI: 28.58%–43.77%) after removing Brahimi's study,<sup>[19]</sup> and these two studies are the most effective studies in the final meta-analysis [Figure 2].



**Table 2: Prevalence of hypertension in renal patients in the studied subgroups**


Subgroups		Number of study	Prevalence (95% CI)	P	I <sup>2</sup> (%)
Sex	Total	34	35 (29-41)	P<0.001	99.5
	Women	3	25 (16-35)	P<0.001	91.7
	Men	3	18 (15-22)	0.103	56.1
Type of hypertension	Systolic	3	5 (0-13)	P<0.001	99.2
	Diastolic	3	26 (0-77)	P<0.001	100
Type of disease	Hemodialysis Patients	20	34 (23-45)	P<0.001	98.4
	Kidney transplantation patients	9	43 (19-67)	P<0.001	99.6
	Chronic kidney disease	3	26 (17-34)	P<0.001	99.7
	Peritoneal dialysis	2	27 (16-38)	0.201	38.9
Age (Year)	<30	2	41 (12-69)	P<0.001	95.6
	30-39	3	22 (9-36)	P<0.001	95.3
	40-49	10	39 (20-57)	P<0.001	99.4
	50-59	11	35 (26-44)	P<0.001	94.0
	60-70	3	32 (16-48)	P<0.001	93.9
Region	North	14	42 (28-55)	P<0.001	98.7
	South	2	37 (29-44)	0.442	0
	Center	7	30 (10-50)	P<0.001	99.8
	West	5	38 (25-50)	P<0.001	95.3
	East	3	21 (9-33)	0.091	58.4



**Diagram 1: Flowchart of studies to the systematic review and meta-analysis**

**Discussion**

In 35 reviewed studies with a sample of 39,621 subjects, the prevalence of hypertension in renal patients was 35% (95% CI: 29%–41%) (25% in women and 18% in men). The prevalence of systolic hypertension in renal patients was 5%, diastolic hypertension 26%, and diabetes 23%. The prevalence of hypertension in hemodialysis patients was 34%, 27% in peritoneal dialysis, 43% in kidney transplantation, and 26% in chronic renal failure. In addition, meta-regression showed that the prevalence of



**Figure 2: Sensitivity analysis**

hypertension in renal patients did not significantly decrease during the years 1988–2017.

According to the initial care system surveys of noncommunicable disease risk factors in Iran in 2005, the prevalence of hypertension in individuals aged 15–64 was 17.1%.<sup>[19,20]</sup> In a general study, the prevalence of hypertension was 11% in Isfahan, 17% in Chaharmahal and Bakhtiari, 11.6% in Gilan, 17.5% in Zanjan, 6.25% in Kermanshah, and 18.9% in Arak.<sup>[85]</sup> In the study of Bashardost *et al.*, 44.9% of hemodialysis patients had high blood pressure.<sup>[86]</sup> In a study on renal transplant patients, Rostami *et al.* found that the prevalence of hypertension was 58.2%.<sup>[60]</sup> In another study by Rezaei Ghalechi to analyze depression in hemodialysis patients, the prevalence of hypertension

and diabetes was 38.2% and 34.2%, respectively.<sup>[62]</sup> The prevalence of hypertension was estimated to be 17.3% in a study conducted in Iran in 2002–2005 to investigate the risk factors associated with chronic kidney disease.<sup>[69]</sup> In the study of Malekmakan *et al.*, the most common causes of chronic renal failure in hemodialysis patients in Fars province were hypertension and diabetes.<sup>[87]</sup> In a study entitled “A 17-year history of peritoneal dialysis in Iran,” which was conducted on 2302 people, the prevalence of high blood pressure in these patients was 24.4%.<sup>[65]</sup> Due to the differences in the results of studies on the prevalence of high blood pressure in renal patients, this study was conducted through systematic review and meta-analysis.

In 35 articles with a sample size of 39,621 individuals, performed from 1988 to 2017, 34 articles outlined the prevalence of high blood pressure in renal patients, and the prevalence of hypertension in renal patients was estimated to be 35% (95% CI: 29%–41%) (25% in women and 18% in men). The prevalence of systolic hypertension in renal patients was 5%, diastolic hypertension was 26%, and diabetes was 23%.

Meta-regression showed that there was no significant relationship between the prevalence of hypertension in renal patients and the number of research samples ( $P = 0.128$ ), that is, with an increase in the number of samples, the prevalence of hypertension in renal patients decreased, but this was not statistically significant [Figure 3]. There was no significant relationship between the prevalence of hypertension in renal patients and the year of study ( $P = 0.102$ ). During the years 1988–2017, the prevalence of hypertension in renal patients in Iran decreased but was not statistically significant [Figure 4]. In Figures 5 and 6, we see that the effect of publication bias is not statistically significant.

In the study of Familoni *et al.*, 53.2% of the subjects referred to hypertension as one of the causes of brain attack, renal failure, and cardiac failure.<sup>[88]</sup> In a study on patients in developed countries, 11.1% of the cause of chronic kidney disease was diabetes and 10.6% was high blood pressure.<sup>[89]</sup> According to studies in the United States, the most common causes of kidney failure are diabetes (50%), and both diabetes and high blood pressure (50–80%).<sup>[90]</sup> In a study in Pakistan in 1998–2001, high blood pressure (19.45%) and diabetes (19.67%) were identified as the most important causes of kidney failure.<sup>[91]</sup> In a study in Australia in 1997, hypertension (12%) was the most common cause of chronic renal failure.<sup>[92]</sup> In another study in Japan, this was reported as 6.2%.<sup>[93]</sup> In a Saudi study on renal patients, the prevalence of hypertension was reported to be 22.1%.<sup>[94]</sup>

### Limitations of the study

Lack of uniform distribution of studies in different regions of Iran, different age groups, and different genders. Failure to report hypertension accurately in some studies.




Figure 3: Relationship between high prevalence of hypertension in renal patients in Iran and the sample size using meta-regression




Figure 4: Relationship between the prevalence of hypertension in renal patients in Iran and the year of research using meta-regression




Figure 5: Meta-bias

### Conclusions

The prevalence of hypertension in women with kidney disease is higher than men, and patients with kidney

Begg's Test						
adj. Kendall's Score (P-Q) = -61						
Std. Dev. of Score = 67.46						
Number of Studies = 34						
z = -0.90						
Pr >  z  = 0.366						
z = 0.89 (continuity corrected)						
Pr >  z  = 0.374 (continuity corrected)						
Egger's test						
Std_Eff	Coef.	Std.Err.	t	P> t	[95%Conf.	Interval]
slope	.1189642	.0262281	4.54	0.000	.0655394	.172389
bias	7.909413	2.545924	3.11	0.004	2.723536	13.09529

**Figure 6: Begg's test and Eggers test to examine the significance of bias**

transplantation are at higher risk of hypertension than other renal patients. The prevalence of diastolic hypertension in renal patients is about five times higher than systolic blood pressure.

### Ethical considerations

Ethical issues (including plagiarism, data fabrication, double publication) have been completely observed by the authors.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

**Received:** 13 Nov 18 **Accepted:** 12 Feb 19

**Published:** 05 Jul 19

### References

- Mohsenzadeh Y, Motedayen M, Hemmati F, Sayehmiri K, Sarokhani M, Sarokhani D. Investigating the prevalence rate of hypertension in Iranian men and women: A study of systematic review and meta-analysis *J Bas Res Med Sci* 2017;4:53-62.
- Motedayen M, Sarokhani D, Meysami A, Jouybari L, Sanagoo A, Hasanpour Dehkordi A. The prevalence of hypertension in diabetic patients in Iran; a systematic review and meta-analysis. *J Nephrothol* 2018;7:137-44.
- Von Elm E, Altman D, Egger M, Pocock S, Gotsche P, Vandenbroucke J. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Ann Intern Med* 2007;147:573-7.
- Ahmadi SM, Jalali A, Jalali R. Factors associated with the choice of peritoneal dialysis in Iran: Qualitative study. *Open Access Macedonian J Medical Sciences* 2018;6:1253.
- Rezaei Z, Jalali A, Jalali R, Khaledi-Paveh B. Psychological problems as the major cause of fatigue in clients undergoing hemodialysis: A qualitative study. *International J Nursing Sciences* 2018;5:262-7.
- Kalender B, Ozdemir A, Dervisoglu E, Ozdemir O. Quality of life in chronic kidney disease: Effects of treatment modality, depression, malnutrition and inflammation. *International J Clinical Practice* 2007;61:569-76.
- Noghabi AA, Zandi M, Mehran A, Alavian SM, Dehkordi AH. The effect of education on quality of life in patients under interferon therapy. *Hepatitis Monthly* 2010;10:218.
- Cruz MC, Andrade C, Urrutia M, Draibe S, Nogueira-Martins LA, Sesso RdCC. Quality of life in patients with chronic kidney disease. *Clinics* 2011;66:991-5.

- Hasanpour AD. Influence of yoga and aerobics exercise on fatigue, pain and psychosocial status in patients with multiple sclerosis: A randomized trial. *The J Sports Medicine and Physical Fitness* 2016;56:1417-22.
- Series WTR. Hypertension control. Report of a WHO Expert Committee, 1996.
- Burt V. Prevalence of hypertension in the US adult population: Results from the third National health and nutrition examination survey, 1988-1991. *Hypertension* 1995;25:305-13.
- Damir Chi A, Mehrbani J. Prevalence of obesity, overweight and hypertension and their associated risk factors in adult males. *Olympic J* 2009;17:87-103.
- Dalvand S, Salehi Z, Asl MT, Zahednezhad H, Ebrahimi N, Nazari M, *et al.* An examination of factors effecting systolic blood pressure, diastolic blood pressure, and total cholesterol simultaneously using mixed responses model. *Iranian Red Crescent Med J* 2017;19:e42309.
- Akbari R, Bahadoram M, Ghorbani A, Zarghami A. Campaigning for kidney health; an experience from kidney day in Iran. *Ann Res Dial* 2016;1:e02.
- Gadhavi BP, Makwana AH, Mehta HB, Shah CJ, Gokhale PA. Early screening of hypertension and cardiac dysautonomia in each hypertensive is needed-inference from a study of QTc interval in Gujarat, India. *Int J Prev Med* 2018;9:62.
- Tamadon M. On the occasion of world kidney day 2018 with the theme of kidney disease in women; vitamin D deficiency and kidney diseases in women. *J Renal Inj Prev* 2018;7:33-4.
- Sadeghi M, Roohafza H, Kelishadi R. High blood pressure and associated cardiovascular risk factors in Iran: Isfahan Healthy Heart Programme. *Med J Malaysia* 2004;59:460-7.
- Rafiei M, Sayfi A. Distribution of blood pressure values in menopausal women in Arak population in 2007. *Arak Univ Med Sci* 2006;12:47-56.
- Delavari A, Horri N, Alikhani S, Guya M, Mahdavi A, Hoseini M, *et al.* Prevalence of hypertension in Iranian urban and rural populations aged over 20 years in 2004. *J Mazandaran Univ Med Sci* 2007;17:79-86.
- Refahi S, Shamsi A, Ebadi A, Saeed Y, Moradi A. Comparison of military and civilian life style of people with hypertension. *J Health Promot Manag* 2012;1:43-50.
- Whitworth J. World Health Organization (WHO), International Society of Hypertension Writing Group (ISH). Statement on Management of Hypertension. *J Hypertens* 2003;21:1983-92.
- Flynn JT, Kaelber DC, Baker-Smith CM, Blowey D, Carroll AE, Daniels SR, *et al.* Clinical practice guideline for screening and management of high blood pressure in children and adolescents. *Pediatrics* 2017;140:e20171904.
- Mahmoodnia L, Beigrezaei S. Chronic kidney disease and obesity; a mini-review to the current knowledge. *J Nephropharmacol* 2017;6:30-2.
- Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L. Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm Hg, 1990-2015. *Jama* 2017;317:165-82.
- Bond M, Pitt M, Akoh J, Moxham T, Hoyle M, Anderson R. The effectiveness and cost-effectiveness of methods of storing donated kidneys from deceased donors: A systematic review and economic model. *Health Technol Assess* 2009;13:1-156.
- Raiiesifar A, Torabpour M, Mohsenizad P. Causes of chronic renal failure in hemodialysis patients of Abadan. *IJCCN* 2010;2:157-60.
- Tayebi A, Raiiesifar A, Ebadi A, Eynollahi B, Rafiyan Z, Keyvanloo F. Review the renal transplantation patients' quality

- of life by using kidney transplantation questionnaire (KTQ-25). *J Crit Care Nurs* 2010;3:13-4.
28. Hasanpour- Dehkordi A. Kidneys and women's health, the theme of world kidney day 2018. *J Negat Results Clin Exp Stud* 2018;1:e02.
  29. Freedman BI, Langefeld CD, Andringa KK, Croker JA, Williams AH, Garner NE, *et al.* End-stage renal disease in African Americans with lupus nephritis is associated with APOLI. *Arthritis and Rheumatology* 2014;66:390-6.
  30. Wheeler DC, Winkelmayer, WC. Improving, and G. Work, KDIGO 2017 Clinical Practice Guideline Update for the Diagnosis, Evaluation, Prevention, and Treatment of Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD). *Kidney Int Suppl* 2017;7:1.
  31. Wanner C, Lachin JM, Inzucchi SE, Fitchett D, Mattheus M, George J, *et al.* Empagliflozin and clinical outcomes in patients with type 2 diabetes mellitus, established cardiovascular disease, and chronic kidney disease. *Circulation* 2018;137:119-29.
  32. Sayin A, Mutluay R, Sindel S. Quality of life in hemodialysis, peritoneal dialysis, and transplantation patients. *Transplant Proc* 2007;39:3047-53.
  33. Eryilmaz M, Ozdemir C, Yurtman F, Cilli A, Karaman T. Quality of sleep and quality of life in renal transplantation patients. *Transplant Proc* 2005;37:2072-6.
  34. Ghaffari A. Seminar of kidney transplantation report. *Biomed Eng* 2008;6:73.
  35. Nasri H. Trends toward amelioration of renal inflammation and fibrosis in various kidney diseases. *J Inj Inflamm* 2016;1:e02.
  36. Jay CL, Dean PG, Helmick RA, Stegall MD. Reassessing preemptive kidney transplantation in the United States: Are we making progress? *Transplantation* 2016;100:1120.
  37. Smith JM, Martz K, Blydt-Hansen TD. Pediatric kidney transplant practice patterns and outcome benchmarks, 1987–2010: A report of the North American Pediatric Renal Trials and Collaborative Studies. *Pediatric Transplantation*. 2013;17:149-57.
  38. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, *et al.* Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128.
  39. Prevention CfDca. National Chronic Kidney Disease Fact Sheet. 2014.
  40. Vendemia F, Amico G. Antihypertensive therapy in dialysed patients. *Contrib Nephrol* 1988;61:161-70.
  41. Jha V, Garcia Garcia G, Iseki K, Li Z, Naicker S, Plattner B, *et al.* Chronic kidney disease: Global dimension and perspectives. *Lancet* 2013;382:260-72.
  42. Ezeonwu B, Nwafor I, Nnodim I, Ayodeji A, Ajaegbu O, Maduemem E, *et al.* Risk factors for chronic kidney disease in children attending pediatric outpatient clinic in federal medical center Asaba. *J Prev Epidemiol* 2016;1:e10.
  43. Lala M, Nazar C, Lala H, Singh J. Interrelation between blood pressure and diabetes. *J Ren Endocrinol* 2015;1:e05.
  44. Oluyombo R, Olamoyegun M, Ayodele O, Akinwusi P, Akinsola A. Clustering of chronic kidney disease and cardiovascular risk factors in South-West Nigeria. *J Nephropathol* 2017;6:196-203.
  45. Navaneethan SD, Schold JD, Jolly SE, Arrigain S, Winkelmayer WC, Nally Jr JV. Diabetes control and the risks of ESRD and mortality in patients with CKD. *American J Kidney Diseases* 2017;70:191-8.
  46. Liyanage T, Ninomiya T, Jha V, Neal B, Patrice HM, Okpechi I, *et al.* Worldwide access to treatment for end-stage kidney disease: A systematic review. *The Lancet* 2015;385:1975-82.
  47. Bah A, Lamine C, Balde M, Bah M, Rostaing L. Epidemiology of chronic kidney diseases in the Republic of Guinea; future dialysis needs. *J Nephropathol* 2015;4:127-33.
  48. Targher G, Bertolini L, Chonchol M, Rodella S, Zoppini G, Lippi G, *et al.* Non-alcoholic fatty liver disease is independently associated with an increased prevalence of chronic kidney disease and retinopathy in type 1 diabetic patients. *Diabetologia* 2010;53:1341-8.
  49. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev J* 2015;4:1-9.
  50. Barahimi H, Aghighi M, Aghayani K, Rahimi Foroushani A. Chronic kidney disease management program in Shahreza. *Iran J Kidney Dis* 2014 8:450-6.
  51. Fazelzadeh A, Mehdizadeh A, Ostovan M, Raiss Jalali G. Incidence of cardiovascular risk factors and complications before and after kidney transplantation. *Transplant Proc* 2006;38:506-8.
  52. Bashardoost B, Adib A, Faaalpoor Z, Gavami Nashr M. The study of hypertension relationship with weight gain in patients hemodialysis intervals. *J Ardabil Univ Med Sci* 2007;7:22-6.
  53. Hasanzamani B, Zeraati A, Sharifpour F, Paeizi R. Prevalence of hypertension in hemodialysis versus peritoneal patients. *Iran J Kidney Dis* 2011;5:53.
  54. Pourmand G, Dehghani S, Rahmati M, Mehrsai A, Gooran S, Alizadeh F, *et al.* Does hypertension remain after kidney transplantation?. *Acta Med Iran* 2015;53:297-300.
  55. Basiratnia M, Esteghamati M, Ajami G, Amoozgar H, Cheriki C, Soltani M, *et al.* Blood pressure profile in renal transplant recipients and its relation to diastolic function: Tissue Doppler echocardiographic study. *Pediatr Nephrol* 2011;26:449-57.
  56. Shamsa A, Motavalli M, Aghdam B. Erectile function in end-stage renal disease before and after renal transplantation. *Transplant Proc* 2005;37:3087-9.
  57. Hollisaaz M, Aghanassir M, Lorgard Dezfuli Nezaad M, Assari S, Hafezie R, Ebrahiminia M. Medical comorbidities after renal transplantation. *Transplant Proc* 2007;39:1048-50.
  58. Nikoobakht M, Mahboobi A, Saraji A, Mehrsai A, Emamzadeh A, Mahmoudi M. Pelvic nerve neuropathy after kidney transplantation. *Transplant Proc* 2007;39:1108-10.
  59. Noshad H, Ardalan M, Mortazavi M, Tayebi H, Safa J, Nezami N. Kidney transplantation candidates and cardiovascular risk factors. *Transplant Proc* 2007;39:871-4.
  60. Rostami Z, Shafiqhi N, Baghersad M, Einollahi B. Risk factors for immediate anemia in renal transplant recipients: A single-center experience. *Transplant Proc* 2011;43:581-3.
  61. Pakfetrata M, Roozbeha J, Nikooc M, Asemb Z, Malekmakanb L, Nikood M. Common echocardiography findings in pretransplant dialysis patients and their associations. *Hong Kong J Nephrol* 2013;15:68-74.
  62. Rezaei Ghalechi E, Kazemi R, Hagh Kerdar M. The evaluation of prevalence of depression in hemodialysis patients in north west of Iran. *Eur Psychiatry* 2013;28:1-4.
  63. Nobakht Haghighi A, Malakoutian T, Radfar M, Abdi E, Kamgar M, Broumand B, *et al.* A 4-year follow-up of living unrelated donors of kidney allograft in Iran. *Iran J Kidney Dis* 2015;9:273-8.
  64. Rohani M, Aghaei M, Jenabi A, Yazdanfar S, Mousavi D, Miri S. Restless legs syndrome in hemodialysis patients in Iran. *Neurol Sci* 2015;36:723-7.
  65. Najafi I, Alatab S, Atabak S, Nouri Majelan N, Sanadgol H, Makhdoomi K, *et al.* Seventeen years' experience of peritoneal dialysis in Iran: First official report of the Iranian peritoneal dialysis registry. *Perit Dial Int* 2014;34:636-42.



66. Gheissari A, Hemmatzadeh S, Merrikhi A, Fadaei Tehrani S, Madihi Y. Chronic kidney disease in children: A report from a tertiary care center over 11 years. *J Nephropathol* 2012;1:177-82.
67. Jalalzadeh M, Mohammadi R, Mirzamohammadi F, Ghadiani M. Prevalence of metabolic syndrome in a hemodialysis population. *Iran J Kidney Dis* 2011;5:248-54.
68. Nemati E, Ghanbrpour F, Taheri S, Einollahi B. Prevalence of hypertension among Iranian hemodialysis patients and risk factors: A nationwide multicenter study. *J Biol Sci* 2008;11:910-4.
69. Safarinejad M. The epidemiology of adult chronic kidney disease in a population-based study in Iran: Prevalence and associated risk factors. *J Nephrol* 2009;22:99-108.
70. Ayazi K, Atabak S, Saghebi R, Ayazi S, Aryasepehr S. Evaluation of efficacy, survival rate and complications of peritoneal catheter placement of patients with end-stage renal disease. *Saudi Med J* 2005;26:1391-3.
71. Nakhjavani M, Esfahanian F, Safavi M, Kalbasi Anaraki M, Zohrevand P. Prevalence of diabetic patients among under-Hemodialysis patients in ten hemodialysis centers of Tehran. *Iran J Diabetes Metab (ijdd)* 2004;4:39-46.
72. Barzin M, Tazikie O, Pezeshki M. Incidence of renal osteodystrophy in hemodialysis patients of fateme-Zahra hospital in Sari in 1998. *J Mazandaran Univ Med Sci* 2001;11:31-7.
73. Monfared A, Khosravi M, Orang poor R, Moosavian Roshan Zamir S, Aghajani Nargesi D. Reasons of chronic renal failure in hemodialysis patients in Guilan Province. *J Guilan Univ Med Sci* 2003;12:76-83.
74. Shasti S, Babahaji M. The assessment of dialysis adequacy among hemodialysis patients in Tehran City. *EBNESINA- Journal of Medical* 2011;14:23-7.
75. Raiisifar A, Tayyebi A, Ebadi A, Najafi S, Hashemi S, Asiyabi M. An investigation of quality of life in kidney transplant patients. *Iran J Crit Care Nurs* 2011;4:149-52.
76. Ghafourifard M, Rafieian M, Shahgholian N, Mortazavi M. Effect of sodium dialysate variation in combining with ultrafiltration on intradialytic hypotension and intradialytic weight gain for patients on hemodialysis. *J Mazandaran Univ Med Sci* 2009;19:19-26.
77. Heidari B. The etiology of kidney failure in hemodialysis patient in Mazandaran. *Mazandaran Univ Med Sci J* 2001;11:43-7.
78. Mogharab M, Rezaee N, Tahouri F, Taheri P, Jani H. Complications during hemodialysis in chronic hemodialysis patients using dialysis buffer solution with sodium acetate and sodium bicarbonate. *Mod Care J* 2007;4:21-8.
79. Eslami Amirabadi M, Hosein D, Nasrollahi A, Norouzian M, Bozorg B, Amjadi Kivi SM, *et al.* Cognitive dysfunction in hemodialysis patients and its related factors. *Res Med* 2014;38:53-9.
80. Rahimian M, Hasan Zadeh A, Sami R. The effects of erythropoietin in the management of anemia of chronic renal failure in patients on hemodialysis. *JSSU* 2005;13:12-5.
81. Ghorbani Moghadam Z, Sharifi S. The frequency of underlying conditions for chronic renal failure in patients undergoing dialysis in Shohada Hospital in Bushehr. *Nurs J Vulnerable* 2015;2:46-54.
82. Hadian B, Anbari K, Heidari R. Epidemiologic study of end stage renal disease and related risk factors in patients under hemodialysis in Lorestan province. *Yafte* 2014;16:44-53.
83. Rezaeian Langroodi R. Evaluation of risk factors for chronic renal failure in hemodialysis patients referring to hospitals affiliated to Hamedan University of Medical Sciences. *Aflak Q J* 2008;7:13-9.
84. KhademVatan K, Rahmani A, Ghasemi S, Ahangary H, Mehrpooya M. Evaluation of the effective risk factors for peripheral vascular disease in hemodialysis patients. *J Urmia Nurs Midwifery Fac* 2014;12:585-90.
85. Ebadi P, Shamsi A, Refahi A, Saied Y. Comparison of the Quality of Life in Men with and Without Hypertension. *Sci J Hamadan Nurs Midwifery* 2010;20:5-15.
86. Rocco M, Yan G, Heyka R, Benz R, Cheung A. Risk factors for hypertension in chronic hemodialysis patients: Baseline data from the HEMO study. *Am J Nephrol* 2001;21:280-8.
87. Malekmakan L, Haghpanah S, Pakfetrat M, Malekmakan A, Khajehdehi P. Causes of chronic renal failure among Iranian hemodialysis patients. *Saudi J Kidney Dis Transpl* 2009;20:501-4.
88. Familoni B, Ogun S, Aina A. Knowledge and awareness of hypertension among patients with hypertension. *J Natl Med Assoc* 2004;96:620-4.
89. Snouber S, Khader M, Alkhatib A, Nazzal Z, Dudin A. Prevalence of patients with end-stage renal disease on dialysis in the West Bank. *Saudi J Kidney Dis Transpl* 2013;24:832-7.
90. Goretti M, Janeza T, Croatia R, *al e.* Special problems in Hemodialysis patients 2011:123-6.
91. Adibul A, Rizvi K. Causes of chronic renal failure in Pakistan: A single large center experience. *Saudi Kidney Dis Transplant Proc* 2002;13:376-9.
92. Disney A, Russ G, Walker R, *al e.* Australia and New Zealand Dialysis and Transplant Registry. *Kidney Int* 1998;62:2281-7.
93. Shinzato T, Nakai S, Akiba T, Yamazaki C, Sasaki R, Kitaoka T, *et al.* Current status of renal replacement therapy in Japan: Results of the annual survey of the Japanese Society for Dialysis Therapy in 1996. *Kidney Int* 1999;55:700-12.
94. Faissal A, Shaheen N. Pre end stage renal failure: The Jeddah kidney center experience. *Saudi Kidney Dis Transplant* 2002;13:371-5.