

ID Design Press, Skopje, Republic of Macedonia  
Open Access Macedonian Journal of Medical Sciences. 2019 Jul 30; 7(14):2282-2286.  
<https://doi.org/10.3889/oamjms.2019.609>  
eISSN: 1857-9655  
*Clinical Science*



# Chest Pain Characteristics in Cardiac Syndrome X Compared to Coronary Artery Disease

Ramaze Farouke Elhakeem<sup>1</sup>, Mohamed Faisal Lutfi<sup>1,2\*</sup>, Ahmed Babiker Mohamed Ali<sup>3</sup>, Mohamed Yusif Sukkar<sup>2</sup>

<sup>1</sup>College of Medicine, Qassim University, KSA, Buraydah, Qassim, Saudi Arabia; <sup>2</sup>Nile College of Medicine, Khartoum, Sudan; <sup>3</sup>Faculty of Medicine, Al Neelain University, Khartoum, Sudan

## Abstract

**Citation:** Elhakeem RF, Lutfi MF, Ali ABM, Sukkar MY. Chest Pain Characteristics in Cardiac Syndrome X Compared to Coronary Artery Disease. Open Access Maced J Med Sci. 2019 Jul 30; 7(14):2282-2286. <https://doi.org/10.3889/oamjms.2019.609>

**Keywords:** Chest pain; Cardiac syndrome X; Coronary artery disease

**\*Correspondence:** Mohamed Faisal Lutfi, College of Medicine, Qassim University, KSA, Buraydah, Qassim, Saudi Arabia; Nile College of Medicine, Khartoum, Sudan. E-mail: mohamedfaisallutfi@gmail.com

**Received:** 30-Apr-2019; **Revised:** 23-Jun-2019; **Accepted:** 24-Jun-2019; **Online first:** 25-Jul-2019

**Copyright:** © 2019 Ramaze Farouke Elhakeem, Mohamed Faisal Lutfi, Ahmed Babiker Mohamed Ali, Mohamed Yusif Sukkar. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**AIM:** This study aimed to assess if clinical remarks gained by analysis of the present and past medical history of patients undergoing elective coronary angiography (ECA) due to typical chest pain can help to predict the outcome of ECA.

**MATERIAL AND METHODS:** One hundred and fifty-four ECA candidates with a history of typical chest were seen on the same day intended for ECA in the cardiac centre of AlShaab Teaching Hospital, Khartoum, Sudan. The details of the present complaints, characteristics of chest pain, past medical and socioeconomic history were recorded from each subject guided by a questionnaire. ECA confirmed CAD in 112 of the studied patients and were considered as the test group. The remaining patients (N = 42) were diagnosed as CSX after exclusion of significant narrowing of the coronary vessels and were considered as the control group.

**RESULTS:** Univariate analysis of pain characteristics among patients undergoing coronary angiography revealed that pain is less likely to radiate to the neck (OR = 0.44, 95% CI = 0.21 – 0.91, P = 0.027) and the back (OR = 0.48, 95% CI = 0.23 – 1.00, P = 0.049) in patients with CAD. Presence of shortness of breathing and/or dizziness significantly decrease the odds of having abnormal coronary angiography (OR = 0.30 and 0.48, 95% CI = 0.12 – 0.77 and 0.22 – 0.92, P = 0.013 and 0.030 respectively). Past history of diabetes mellitus significantly increases the odds of having abnormal coronary angiography (OR = 3.96, 95% CI = 1.68 – 9.30, P = 0.002). In contrast, past medical history of migraine decreases the odds of having positive finding in ECA (OR = 0.31, 95% CI = 0.13 – 0.72, P = 0.006).

**CONCLUSION:** Characteristics of chest pain are comparable in CAD and CSX. However, pain is less likely to radiate to the neck and/or the back in the first group. Presence of dyspnea and dizziness during angina attacks as well as the history of migraine significantly decreases the odds of having abnormal coronary angiography.

## Introduction

Patients with typical chest pain and positive stress electrocardiography (ECG) or other cardiac tests are not certainly suffering from coronary artery disease (CAD) [1], [2]. Cardiac syndrome X (CSX) is frequently used to diagnose patients with typical chest pain, positive cardiac stress test(s) and normal coronary macrocirculation [3]. Dysfunction of coronary microcirculation [4] and abnormal perception of pain [5], [6] are the most acceptable explanations for CSX in the literature so far. In clinical practice, the relatively

high percentage of patients with no significant angiographic findings following elective coronary angiography (ECA) raise a question whether ECA is overused in patients with suspected CAD [7]. Although there are a lot of studies investigating risk factors for CAD [8], [9], researches exploring the predictors of CSX are scarce [2], if any. This study aimed to assess if clinical remarks gained by analysis of the present and past medical history can help to predict outcome of ECA. Special care was given in comparing chest pain characteristics between patients with CSX and CAD.

## Material and Methods

The present study was approved from the ethics review committee (ERC), Faculty of Medicine, University of Khartoum, Sudan. All candidates who agreed to join this study signed a written informed consent before being evaluated.

One hundred and fifty-four ECA candidates with a history of typical chest pain were seen on the same day intended for ECA in the cardiac centre of AlShaab Teaching Hospital, Khartoum, Sudan. The details of the present complaints, characteristics of chest pain, past medical and socioeconomic history were recorded from each subject guided by a questionnaire. The body mass index (BMI) and mean arterial blood pressure (MABP) were calculated for each subject by the formulae:

$$\text{BMI (kg/m}^2\text{)} = \text{weight (kg)} / (\text{height (m)}^2)$$

and

$$\text{MABP} = \text{Diastolic blood pressure} + \frac{1}{3} (\text{Diastolic blood pressure} - \text{Systolic blood pressure})$$

respectively.

ECA confirmed CAD in 112 of the studied patients and were considered as the test group. The remaining patients (N = 42) were diagnosed as CSX after exclusion of significant narrowing of the coronary vessels and were considered as the control group.

Statistical analysis was performed using Statistical Package for the social sciences (SPSS) for Windows, version 16.0 (SPSS Inc., Chicago, IL, USA). Normal distribution of variables was examined using Shapiro-Wilk test. Unpaired T-test was used to assess statistical difference between mean (SD) of normally distributed scaled variables. Mann-Whitney U test was used to compare median (25<sup>th</sup> – 75<sup>th</sup> interquartile) of abnormally distributed scaled variables. Univariate analyses were carried out to evaluate characteristics of chest pain and past medical history as past predictors of CAD. Results of univariate analyses were expressed by odds ratios (OR) and their 95% CI. In all tables expressing results of univariate analysis, OR described the ratio of the odds of an event occurring in patients with CAD to the odds of the same event occurring in subject with CSX.  $P < 0.05$  was considered significant.

## Results

Coronary artery catheterization of the studied subjects (N = 154) revealed 112 patients with CAD ( $P$  (95% CI) = 72.7% (65.3 – 79.3%), mean (SD) of age = 60.58 (10.26) years) and 42 subjects with normal

coronary arteries ( $P$  (95% CI) = 27.3% (20.7–34.7%), mean (SD) of age = 50.95 (16.46) years).

Table 1 compares age, anthropometric measurements and blood pressures of subjects with CSX and CAD. Age was significantly higher in CAD patients compared to CSX ( $P < 0.001$ ). In contrast, BMI was significantly higher in CSX compared to CAD patients ( $P = 0.004$ ).

**Table 1: Comparison of age, anthropometric measurements and blood pressures among patients undergoing ECA**

	CSX	CAD	<i>P</i>
	N = 42	N = 112	
	Mean (SD)	Mean (SD)	
	Median (25 – 75 interquartile)	Median (25 – 75 interquartile)	
Age (years)	50.0 (42.5 – 53.3)	55.0 (60.0 – 68.8)	< 0.001*
Weight (Kg)	80.04 (13.77)	72.39 (13.67)	0.004*
Height (Cm)	164.65 (8.33)	165.38 (8.14)	0.640
BMI (kg/m <sup>2</sup> )	29.61 (5.06)	26.32 (4.52)	0.001*
SBP (mmHg)	132.83 (20.16)	129.41 (22.86)	0.379
DBP (mmHg)	79.29 (11.99)	78.21 (11.94)	0.627
MABP (mmHg)	97.14 (12.82)	95.28 (13.92)	0.446

Univariate analysis of chest pain characteristics among patients undergoing coronary angiography revealed that chest pain is less likely to radiate to the neck (OR = 0.44, 95% CI = 0.21 – 0.91,  $P = 0.027$ ) and the back (OR = 0.48, 95% CI = 0.23 – 1.00,  $P = 0.049$ ) in patients with CAD (Table 2). In addition, presence of shortness of breathing and / or dizziness significantly decrease the odds of having abnormal coronary angiography (OR = 0.30 and 0.48, 95% CI = 0.12 – 0.77 and 0.22 – 0.92,  $P = 0.013$  and 0.030 respectively).

**Table 2: Univariate analyses of pain in patients undergoing ECA**

	OR	95% CI	<i>P</i>
Duration since first chest pain attack > 2years	0.86	0.39 – 1.86	0.695
Duration of pain attack > 30 minutes	0.95	0.38 – 2.39	0.950
Severity of pain	0.92	0.56 – 1.52	0.750
Pain radiation			
Retrosternal	1.24	0.51 – 2.00	0.631
Left sided chest pain	0.70	0.32 – 1.55	0.378
Radiation of pain to the neck	0.44	0.21 – 0.91	0.027*
Radiation of pain to the right shoulder	0.74	0.35 – 1.53	0.410
Radiation of pain to the left shoulder	0.52	0.25 – 1.09	0.082
Radiation of pain to the right upper limb	1.19	0.52 – 2.70	0.686
Radiation of pain to the left upper limb	0.64	0.31 – 1.32	0.230
Radiation of pain to the Back	0.48	0.23 – 1.00	0.049*
Radiation of pain to other areas	1.16	0.55 – 2.42	0.702
Aggravating factors			
Exercise	1.34	0.57 – 3.15	0.503
Cold	0.56	0.26 – 1.23	0.148
Food intake	1.05	0.46 – 2.42	0.902
Others	0.75	0.13 – 2.24	0.743
Relieving factors			
Rest	1.07	0.35 – 3.25	0.905
Sublingual nitrates	1.05	0.52 – 2.12	0.883
Others	1.64	0.44 – 6.14	0.461
Associated factors			
Sweating	0.61	0.27 – 1.39	0.242
Nausea	0.80	0.38 – 1.67	0.552
Vomiting	0.92	0.42 – 2.02	0.841
Palpitation	0.58	0.28 – 1.20	0.143
Shortness of breathing	0.30	0.12 – 0.77	0.013*
Dizziness	0.45	0.22 – 0.92	0.030*
Loss of consciousness	0.84	0.34 – 2.11	0.714

Table 3 shows the results of univariate analyses of past medical and socioeconomic history in patients undergoing coronary angiography. Past history of diabetes mellitus significantly increases the odds of having abnormal coronary angiography (OR = 3.96, 95% CI = 1.68 – 9.30,  $P = 0.002$ ). In contrast, past medical history of migraine decreases the odds

of having positive finding in ECA (OR = 0.31, 95% CI = 0.13 – 0.72, P = 0.006).

**Table 3: Univariate analyses of past medical and socioeconomic history for patients undergoing ECA**

	OR	95% CI	P
Past medical history			
Hypertension	1.29	0.63 – 2.63	0.490
Diabetes mellitus	3.96	1.68 – 9.30	0.002*
Peptic ulcer	0.45	0.11 – 1.76	0.249
Esophageal disease	0.50	0.23 – 1.07	0.072
Respiratory disease	0.87	0.21 – 3.52	0.841
Migraine	0.31	0.13 – 0.72	0.006*
Raynaud's phenomenon	0.37	0.02 – 6.01	0.482
Chronic inflammatory disease	0.58	0.18 – 1.89	0.366
Socioeconomic history			
Smoking			
- Active smoking	1.64	0.80 – 3.39	0.178
- Passive smoking	1.20	0.55 – 2.64	0.650
Socioeconomic status	0.97	0.54 – 1.76	0.923

## Discussion

Univariate analysis of pain characteristics, namely onset and duration of pain attacks, severity, radiations, aggravating and relieving factors reveals only a few predictors for positive coronary angiography findings in patient undergoing ECA because of typical chest pain. According to the current results, distributions of chest pain are comparable in both studied groups. However, pain is less likely to radiate to the neck and/or the back in patients with CAD. Also, presence of shortness of breathing and/or dizziness significantly decreases the odds of having abnormal coronary angiography. Although previous reports on CSX patients suggest enhancement of their pain perception [10], the current study failed to demonstrate a significant difference in pain severity, duration or aggravating factors when CAD and CSX patients were compared. The special attention paid by cardiologists while evaluating the need of patients with chest pain for diagnostic coronary angiography may explain the limited difference in chest pain characteristics of studied groups. This is because typical features of angina are carefully evaluated by cardiologists while selecting patients who are in real need of diagnostic coronary angiography.

At least two previous studies explain the radiation of chest pain in patients with CSX [11], [12]. The first study was conducted by Lanza and his group in the late nineties of the last century [12]. Lanza *et al.* demonstrated cardiac adrenergic nerve dysfunction in 75% of patients with CSX patients suggesting cardiac origin of chest pain in these patients. Five years later, Rosen *et al.* used positron emission tomography (PET) and stress echocardiography studies to assess origin of pain in patients with CSX [11]. According to Rosen *et al.*, results, chest pain and ECG changes during attacks were not accompanied by demonstrable myocardial dysfunction. However, there was altered central neural handling of afferent signals suggesting that CSX might be a cortical pain

syndrome. The hypotheses suggested by either Lanza *et al.*, or Rosen *et al.* can explain the great similarities in the areas of chest pain radiation in patients with CSX and CAD targeted by this study.

According to the results of the present study, the presence of dyspnea and dizziness in patients with typical chest pain significantly increases the probability of having normal rather than abnormal coronary angiography. This finding is not necessarily contradictory to what was reported before that dyspnea and dizziness are common during angina attacks of CAD patients [13], [14]. In contrast, it may indicate the higher frequency of these symptoms in patients with a separate pathology that causes typical angina in the presence of patent coronary vessels. Patients with CSX are at higher risk of neuroticism like anxiety and depression [15], [16]. Neuroticism could perpetuate to somatoform disorder with physical symptoms like shortness of breathing and dizziness; however, however, the present literature lacks scientific proof for this hypothesis and further researches are desirable to explore this area.

It is worth mentioning that previous studies suggested that patients with angina and normal coronary angiography may have a diffuse disorder of smooth muscle tone [17]. Cannon *et al.*, studied forced expiratory volume in the first second (FEV1) in the basal state and after methacholine inhalation to determine whether the bronchial smooth muscle is affected in CSX patients [18]. Fourteen per cent of patients with CSX had a basal FEV1 of less than 70% of that predicted and did not receive methacholine. Also, the product of the methacholine dose inhaled and the magnitude of decline in FEV1 from baseline was significantly lower in patients with CSX than in normal volunteers suggesting airway smooth muscle hyperresponsiveness in CSX patients. Cannon *et al.* findings may explain the shortness of breathing in patients with angina and normal coronary angiography during chest pain attacks. This is because the initiator of microvascular smooth muscle dysfunction, and hence angina, may at the same time trigger shortness of breathing by inducing airway hyperresponsiveness.

Previous researches that compare features of chest pain in patients with obstructive, non-obstructive and normal coronary vessels are scarce [19], [20]. The current study findings on the major differences of chest pain in patients classified according to ECA outcome are probably naive and deserved to be explored further by additional researches.

Evaluation of past medical history in patients undergoing ECA shows that the odds of having CAD increases about four times in the presence of diabetes mellitus but decreases to about one-third if migraine exists. The results of the present study failed to link positive angiographic finding and common CAD risk factors like past medical history of hypertension [21], [22], Raynaud's phenomenon [23], chronic inflammatory diseases [24], smoking [25] and

socioeconomic status [26]. The current data also failed to demonstrate decreased odds of having positive angiographic finding in patients with past medical history of peptic ulcer, oesophageal or respiratory diseases, whose clinical presentations may mimic angina pain [27], [28].

The prevalence of migraine is significantly increased in either subject with spastic or normal coronary arteries compared with patients with CAD [29], [30]. Koh *et al.* conducted a prospective study on the prevalence of migraine in Korean patients with proven variant angina compared to two control groups: one group with CAD and another one with subjects without heart disease [30]. According to the results of Koh *et al.*, migraine was diagnosed in 40.0%, 20.0% and 38.7% of patients with variant angina, CAD and no heart diseases respectively. Five years later, a comparable study was conducted by Nakamura *et al.*, in Japanese [29]. The data of Nakamura *et al.*, showed that the prevalence of migraine in Japanese patients with vasospastic angina, angina with effort and subjects without known ischemic heart disease are 23%, 4% and 11% respectively. Although it is evident from the works of Koh *et al.*, and Nakamura *et al.*, that there is possible link between migraine and normal coronary arteries, gender and age-specific prevalence of migraine mismatch with that of CAD, which could explain low risk of CAD in migraineurs [23]. This possible explanation for low prevalence of migraine among CAD patients warrants additional investigations and studies.

In conclusion, characteristics of chest pain are comparable in CAD and CSX, however, pain is less likely to radiate to the neck and / or the back in the first group. Presence of shortness of breathing and / or dizziness during angina attacks as well as history of migraine significantly decreases the odds of having abnormal coronary angiography.

## References

- Makharova NV, Voevoda MI, Lyutova FF, Pinigina IA, Tarasova VE. Comparative analysis of clinical, electrocardiographic, angiographic and echocardiographic data of indigenous and non-indigenous residents of Yakutia with coronary artery atherosclerosis. *Int J Circumpolar Health*. 2013; 72(SUPPL.1). <https://doi.org/10.3402/ijch.v72i0.21219> PMID:23967413 PMCID:PMC3748439
- Levitt K, Guo H, Wijesundera HC, et al. Predictors of normal coronary arteries at coronary angiography. *Am Heart J*. 2013; 166(4):694-700. <https://doi.org/10.1016/j.ahj.2013.07.030> PMID:24093849
- Vermeltoort IAC, Rajmakers PGHM, Riphagen II, et al. Definitions and incidence of cardiac syndrome X: review and analysis of clinical data. *Clin Res Cardiol*. 2010; 99(8):475-481. <https://doi.org/10.1007/s00392-010-0159-1> PMID:20407906 PMCID:PMC2911526
- Panting JR, Gatehouse PD, Yang G-Z, et al. Abnormal Subendocardial Perfusion in Cardiac Syndrome X Detected by Cardiovascular Magnetic Resonance Imaging. *N Engl J Med*. 2002; 346(25):1948-1953. <https://doi.org/10.1056/NEJMoa012369> PMID:12075055
- Chauhan A, Mullins PA, Thuraingham SI, Taylor G, Petch MC, Schofield PM. Abnormal cardiac pain perception in syndrome X. *J Am Coll Cardiol*. 1994; 24(2):329-335. [https://doi.org/10.1016/0735-1097\(94\)90284-4](https://doi.org/10.1016/0735-1097(94)90284-4)
- Shintani S, Nishiyama Y, Yamamoto K, Koga Y. Different Long-Term Course Between Chest Pain and Exercise-Induced ST Depression in Syndrome X. *Jpn Heart J*. 2003; 44(4):471-479. <https://doi.org/10.1536/jhj.44.471> PMID:12906029
- Reid CJ, Tanner M, Murphy C. Is angiography overused for the investigation of suspected coronary disease? A single-centre study. *Br J Cardiol*. 2014; 21(2):77.
- Y. W, L. W, X. L, et al. Genetic variants associated with myocardial infarction and the risk factors in Chinese population. *PLoS One*. 2014; 9(1):e86332. <https://doi.org/10.1371/journal.pone.0086332> PMID:24475106 PMCID:PMC3903528
- Petretta M, Acampa W, Evangelista L, et al. Reclassification of cardiovascular risk by myocardial perfusion imaging in diabetic patients with abnormal resting electrocardiogram. *Nutr Metab Cardiovasc Dis*. 2014; 24(6):588-593. <https://doi.org/10.1016/j.numecd.2013.11.009> PMID:24472632
- Pasceri V, Lanza G a, Buffon a, Montenero a S, Crea F, Maseri a. Role of abnormal pain sensitivity and behavioral factors in determining chest pain in syndrome X. *J Am Coll Cardiol*. 1998; 31(1):62-66. [https://doi.org/10.1016/S0735-1097\(97\)00421-X](https://doi.org/10.1016/S0735-1097(97)00421-X)
- Rosen SD, Paulesu E, Wise RJS, Camici PG. Central neural contribution to the perception of chest pain in cardiac syndrome X. *Heart*. 2002; 87(6):513-519. <https://doi.org/10.1136/heart.87.6.513> PMID:12010930 PMCID:PMC1767119
- Lanza GA, Giordano A, Pristipino C, et al. Abnormal cardiac adrenergic nerve function in patients with syndrome X detected by [<sup>123</sup>I]Metaiodobenzylguanidine myocardial scintigraphy. *Circulation*. 1997; 96(3):821-826. <https://doi.org/10.1161/01.CIR.96.3.821> PMID:9264488
- Zimmerman L, Barnason S, Young L, Tu C, Schulz P, AA A. Symptom profiles of coronary artery bypass surgery patients at risk for poor functioning outcomes. *J Cardiovasc Nurs*. 2010; 25(4):292-300. <https://doi.org/10.1097/JCN.0b013e3181cfba00> PMID:20498614
- Miller KH, Grindel CG. Comparison of symptoms of younger and older patients undergoing coronary artery bypass surgery. *Clin Nurs Res*. 2004; 13(3):178-179. <https://doi.org/10.1177/1054773804265693> PMID:15245634
- Altintas E, Yigit F, Taskintuna N. The impact of psychiatric disorders with cardiac syndrome X on quality of life: 3 months prospective study. *International journal of clinical and experimental medicine*. 2014; 7(10):3520.
- Piegza M, Pudlo R, Badura-Brzoza K, Hese RT. Cardiac syndrome X from a psychosomatic point of view. *Arch Psychiatry Psychother*. 2009; 11(2):23-27.
- Lekakis JP, Papamichael CM, Vemmos CN, Voutsas AA, Stamatelopoulos SF, Mouloupoulos SD. Peripheral vascular endothelial dysfunction in patients with angina pectoris and normal coronary arteriograms. *J Am Coll Cardiol*. 1998; 31(3):541-546. [https://doi.org/10.1016/S0735-1097\(97\)00542-1](https://doi.org/10.1016/S0735-1097(97)00542-1)
- Cannon RO, Peden DB, Berkebile C, Schenke WH, Kaliner MA, Epstein SE. Airway hyperresponsiveness in patients with microvascular angina. Evidence for a diffuse disorder of smooth muscle responsiveness. *Circulation*. 1990; 82(6):2011-2017. <https://doi.org/10.1161/01.CIR.82.6.2011> PMID:2242525
- Patel MR, Peterson ED, Dai D, et al. Low diagnostic yield of elective coronary angiography. *NEJM*. 2010; 362(10):886-895. <https://doi.org/10.1056/NEJMoa0907272> PMID:20220183 PMCID:PMC3920593
- Kimble LP, McGuire DB, Dunbar SB, et al. Gender differences

- in pain characteristics of chronic stable angina and perceived physical limitation in patients with coronary artery disease. *Pain*. 2003; 101(1-2):45-53. [https://doi.org/10.1016/S0304-3959\(02\)00319-6](https://doi.org/10.1016/S0304-3959(02)00319-6)
21. Denardo SJ, Gong Y, Nichols WW, et al. Blood pressure and outcomes in very old hypertensive coronary artery disease patients: An INVEST substudy. *Am J Med*. 2010; 123(8):719-726. <https://doi.org/10.1016/j.amimed.2010.02.014> PMID:20670726 PMCID:PMC3008373
22. Gierach GL, Johnson BD, Bairey Merz CN, et al. Hypertension, menopause, and coronary artery disease risk in the Women's Ischemia Syndrome Evaluation (WISE) study. *J Am Coll Cardiol*. 2006; 47(3 SUPPL.). <https://doi.org/10.1016/j.jacc.2005.02.099> PMID:16458172
23. Rosamond W. Are migraine and coronary heart disease associated? An epidemiologic review. *Headache*. 2004; 44(SUPPL. 1). <https://doi.org/10.1111/j.1526-4610.2004.04103.x> PMID:15149488
24. Roifman I, Beck PL, Anderson TJ, Eisenberg MJ, Genest J. Chronic inflammatory diseases and cardiovascular risk: A systematic review. *Can J Cardiol*. 2011; 27(2):174-182. <https://doi.org/10.1016/j.cjca.2010.12.040> PMID:21459266
25. Huxley RR, Woodward M. Cigarette smoking as a risk factor for coronary heart disease in women compared with men: A systematic review and meta-analysis of prospective cohort studies. *Lancet*. 2011; 378(9799):1297-1305. [https://doi.org/10.1016/S0140-6736\(11\)60781-2](https://doi.org/10.1016/S0140-6736(11)60781-2)
26. Franks P, Tancredi DJ, Winters P, Fiscella K. Including socioeconomic status in coronary heart disease risk estimation. *Ann Fam Med*. 2010; 8(5):447-453. <https://doi.org/10.1370/afm.1167> PMID:20843887 PMCID:PMC2939421
27. Ortiz-Olvera NX, Gonzalez-Martinez M, Ruiz-Flores LG, Blancas-Valencia JM, Moran-Villota S, Dehesa-Violante M. [Causes of non-cardiac chest pain: multidisciplinary perspective]. *Rev Gastroenterol Mex*. 2007; 72(2):92-99.
28. Bugiardini R, Merz CNB. Angina with "normal" coronary arteries: A changing philosophy. *J Am Med Assoc*. 2005; 293(4):477-484. <https://doi.org/10.1001/jama.293.4.477> PMID:15671433
29. Nakamura Y, Shinozaki N, Hirasawa M, et al. Prevalence of migraine and Raynaud's phenomenon in Japanese patients with vasospastic angina. *Jpn Circ J*. 2000; 64(4):239-242. <https://doi.org/10.1253/jcj.64.239> PMID:10783043
30. Koh KK, Kim SH, Lee KH, et al. Does prevalence of migraine and Raynaud's phenomenon also increase in Korean patients with proven variant angina? *Int J Cardiol*. 1995;51(1):37-46. [https://doi.org/10.1016/0167-5273\(95\)02371-3](https://doi.org/10.1016/0167-5273(95)02371-3)