



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Spo13 prevents premature cohesin cleavage during meiosis

Citation for published version:

Galander, S, Barton, RE, Kelly, DA & Marston, AL 2019, 'Spo13 prevents premature cohesin cleavage during meiosis', Wellcome Open Research , vol. 4. <https://doi.org/10.12688/wellcomeopenres.15066.2>

Digital Object Identifier (DOI):

[10.12688/wellcomeopenres.15066.2](https://doi.org/10.12688/wellcomeopenres.15066.2)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Wellcome Open Research

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





RESEARCH ARTICLE

REVISED **Spo13 prevents premature cohesin cleavage during meiosis [version 2; peer review: 3 approved]**

Stefan Galander , Rachael E. Barton , David A. Kelly , Adèle L. Marston 

Wellcome Centre for Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3BF, UK

v2 **First published:** 13 Feb 2019, 4:29 (<https://doi.org/10.12688/wellcomeopenres.15066.1>)

Latest published: 02 Sep 2019, 4:29 (<https://doi.org/10.12688/wellcomeopenres.15066.2>)

Abstract

Background: Meiosis produces gametes through two successive nuclear divisions, meiosis I and meiosis II. In contrast to mitosis and meiosis II, where sister chromatids are segregated, during meiosis I, homologous chromosomes are segregated. This requires the monopolar attachment of sister kinetochores and the loss of cohesion from chromosome arms, but not centromeres, during meiosis I. The establishment of both sister kinetochore mono-orientation and cohesion protection rely on the budding yeast meiosis I-specific Spo13 protein, the functional homolog of fission yeast Moa1 and mouse MEIKIN.

Methods: Here we investigate the effects of loss of *SPO13* on cohesion during meiosis I using a live-cell imaging approach.

Results: Unlike wild type, cells lacking *SPO13* fail to maintain the meiosis-specific cohesin subunit, Rec8, at centromeres and segregate sister chromatids to opposite poles during anaphase I. We show that the cohesin-destabilizing factor, Wpl1, is not primarily responsible for the loss of cohesion during meiosis I. Instead, premature loss of centromeric cohesin during anaphase I in *spo13Δ* cells relies on separase-dependent cohesin cleavage. Further, cohesin loss in *spo13Δ* anaphase I cells is blocked by forcibly tethering the regulatory subunit of protein phosphatase 2A, Rts1, to Rec8.





Conclusions: Our findings indicate that separase-dependent cleavage of phosphorylated Rec8 causes premature cohesin loss in *spo13Δ* cells.


Keywords

Meiosis, cohesin, Spo13

Open Peer Review

Reviewer Status 

	Invited Reviewers		
	1	2	3
REVISED			
version 2 published 02 Sep 2019	report		
			
version 1 published 13 Feb 2019	? report	 report	 report

- Andreas Hochwagen** , New York University, New York City, USA
- Dean S. Dawson**, Oklahoma Medical Research Foundation, Oklahoma City, USA
- Enrique Martinez-Perez**, Imperial College London, London, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Adèle L. Marston (adele.marston@ed.ac.uk)

Author roles: **Galander S:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Barton RE:** Investigation, Methodology, Writing – Review & Editing; **Kelly DA:** Software; **Marston AL:** Conceptualization, Funding Acquisition, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was funded by Wellcome through PhD studentships to SG [096994] and REB [102316], as well as a Senior Research Fellowship to ALM [107827] and core funding awarded to the Wellcome Centre for Cell Biology [203149].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 Galander S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Galander S, Barton RE, Kelly DA and Marston AL. **Spo13 prevents premature cohesin cleavage during meiosis [version 2; peer review: 3 approved]** Wellcome Open Research 2019, 4:29 (<https://doi.org/10.12688/wellcomeopenres.15066.2>)

First published: 13 Feb 2019, 4:29 (<https://doi.org/10.12688/wellcomeopenres.15066.1>)

REVISED Amendments from Version 1

Our revised manuscript consists of the changes in response to the reviewers and we additionally added a further figure (Figure 3) which presents data adding further support to the conclusion that Spo13 prevents loss of all cohesion during anaphase I, and responds to point 2 raised by reviewer 1 (Hochwagen). This figure includes three new experiments in which we analysed *spo13Δ mad2Δ* cells which undergo two meiotic divisions. Similar to one-division *spo13Δ* meiosis, in two-division *spo13Δ* meiosis, we observed reduced Rec8 at centromeres in anaphase I and frequent segregation of sister chromatids to opposite poles. The result is that *spo13Δ mad2Δ* mutants exhibit profound aneuploidy. We believe that this interesting data confirms the central conclusion of our manuscript which is that Spo13 is important for the retention of cohesion during meiosis I.

Any further responses from the reviewers can be found at the end of the article

Introduction

Sexual reproduction relies on a cell division programme called meiosis. In humans, this is highly error-prone and may give rise to infertility, miscarriage or chromosomal abnormalities such as Down syndrome (reviewed by Hassold & Hunt, 2001). Meiosis consists of two consecutive divisions, where homologous chromosome segregation in meiosis I is followed by mitosis-like sister chromatid segregation in meiosis II. Homologue segregation requires a number of adaptations to the chromosome segregation machinery (Marston & Amon, 2004), including recombination of homologues, mono-orientation of sister kinetochores and the protection of pericentromeric cohesin in meiosis I.

Cohesin is a multi-subunit protein complex made up of the core subunits Smc1, Smc3 and the kleisin α -Scc1 (Losada *et al.*, 1998; Michaelis *et al.*, 1997) as well as the accessory subunits Scc3 (Tóth *et al.*, 1999) and Pds5 (Hartman *et al.*, 2000; Panizza *et al.*, 2000). In mitosis, cohesin resists the spindle forces that pull sister chromatids towards opposite poles, likely by topologically linking sister chromatids (Gruber *et al.*, 2003; Haering *et al.*, 2002). Upon successful bi-orientation, securin (Pds1 in yeast) is ubiquitinated and destroyed by the proteasome, freeing separase (Esp1), which proteolytically cleaves Scc1 and thereby allows chromosome segregation.

Meiotic cohesin contains an alternative kleisin called Rec8 (Buonomo *et al.*, 2000; Watanabe & Nurse, 1999). Rec8 supports a number of meiosis-specific functions of cohesin, particularly during recombination. Rec8 cleavage is dependent on its prior phosphorylation by casein kinase 1 δ (Hrr25), Dbf4-dependent kinase (DDK) Cdc7 (Katis *et al.*, 2010) and, potentially, Polo kinase (Cdc5) (Brar *et al.*, 2006). However, it is currently unclear how these kinases contribute to cohesin removal, with the role of Cdc5 in cohesin cleavage coming under particular scrutiny (Attner *et al.*, 2013; Argüello-Miranda *et al.*, 2017; Brar *et al.*, 2006; Galander *et al.*, 2019; Katis *et al.*, 2010). Hrr25 and Cdc7 are both independently sufficient for cohesin removal at anaphase I, most likely by promoting its cleavage (Katis *et al.*, 2010). Conversely, there is mounting evidence that Cdc5

facilitates cleavage-independent cohesin loss upon prophase exit (Challa *et al.*, 2019; Yu & Koshland, 2005), although a contribution to cleavage has also been argued (Attner *et al.*, 2013; Brar *et al.*, 2006). While cohesin phosphorylation occurs along the length of the chromosome, the pericentromeric adapter protein shugoshin (Sgo1) binds protein phosphatase 2A (PP2A) to dephosphorylate Rec8 in the pericentromere and prevent its cleavage (Katis *et al.*, 2004a; Kitajima *et al.*, 2006; Kitajima *et al.*, 2004; Lee *et al.*, 2008; Marston *et al.*, 2004; Riedel *et al.*, 2006; Tang *et al.*, 2006). In meiosis II, Rec8 becomes deprotected by the action of Hrr25, which is thought to initiate Sgo1 degradation and phosphorylate Rec8 for cleavage (Argüello-Miranda *et al.*, 2017; Jonak *et al.*, 2017).

In mammalian and *Drosophila* mitosis, cohesin is also removed in two steps. First, during prophase, Wapl opens the cohesin ring at the Smc3-Scc1 interface to trigger separase- and cleavage-independent cohesin removal (Buheitel & Stemann, 2013; Sumara *et al.*, 2000; Waizenegger *et al.*, 2000; Warren *et al.*, 2000). A subset of cohesin is resistant to Wapl due to its prior acetylation and association with sororin (Lafont *et al.*, 2010; Nishiyama *et al.*, 2010; Rankin *et al.*, 2005; Rolef Ben-Shahar *et al.*, 2008; Schmitz *et al.*, 2007; Unal *et al.*, 2008). Notably, pericentromeric cohesin is shielded from Wapl during mammalian mitosis by Sgo1-PP2A, which associates with, and dephosphorylates, both cohesin and sororin to prevent cohesin ring opening (Kitajima *et al.*, 2006; Liu *et al.*, 2013b; McGuinness *et al.*, 2005; Yamada *et al.*, 2017). Second, upon sister kinetochore bi-orientation, Sgo1 relocates from the kinetochore to the pericentromeric chromatin, and separase-dependent cohesin cleavage triggers anaphase onset (Liu *et al.*, 2013a; Liu *et al.*, 2013b). A similar Wapl/Rad61-dependent, cleavage-independent cohesin removal pathway has been suggested to occur in budding yeast meiosis. Although condensin, Cdc5 and DDK have been identified as regulators of this pathway (Challa *et al.*, 2016; Challa *et al.*, 2019; Yu & Koshland, 2005), budding yeast lacks an obvious sororin homologue. Thus, the mechanisms of Wapl-mediated cohesin removal in meiosis I are different to those in mammalian and *Drosophila* mitosis.

While previous research has identified key mechanisms governing cohesin protection, a number of additional proteins have been implicated in this process, but their roles remain unclear. Amongst them is the meiosis I-specific Spo13 (Wang *et al.*, 1987). Cells without *SPO13* only undergo a single meiotic division and show a variety of meiotic defects, including failure to mono-orient sister kinetochores in meiosis I and inability to protect cohesin (Katis *et al.*, 2004b; Klapholz & Esposito, 1980; Lee *et al.*, 2004; Shonn *et al.*, 2002). Spo13 is thought to have functional orthologs in both fission yeast (Moa1) and mouse (MEIKIN) (Kim *et al.*, 2015). The unifying feature of these proteins is their interaction with Polo kinases, whose kinetochore recruitment by Spo13, Moa1 and MEIKIN has been proposed to enable mono-orientation and cohesin protection (Galander *et al.*, 2019; Kim *et al.*, 2015; Matos *et al.*, 2008; Miyazaki *et al.*, 2017).

The exact role of Spo13 in cohesin protection is currently unclear. Interestingly, *SPO13* overexpression blocks cohesin

cleavage during mitosis (Lee *et al.*, 2002; Shonn *et al.*, 2002; Varela *et al.*, 2010), suggesting that Spo13 may also influence cohesin cleavage in meiosis, but how it might do so remains unresolved. Although Spo13 was implicated in ensuring the proper pericentromeric localization of Sgo1 (Kiburz *et al.*, 2005), other studies have found no difference in chromosomally associated Sgo1 (Galander *et al.*, 2019; Lee *et al.*, 2004). In fact, it has been suggested that *spo13Δ* cells might retain residual pericentromeric cohesion in meiosis I (Katis *et al.*, 2004b).

Here, we take a live cell imaging approach to re-evaluate the importance of Spo13 for cohesin protection. We show that both cohesin and sister chromatid cohesion are lost upon anaphase I onset in *spo13Δ* cells. Furthermore, we confirm that cohesin removal results from separase-mediated cleavage rather than removal by the prophase pathway. We also provide evidence that PP2A is capable of preventing cohesin cleavage in *spo13Δ* cells.

Results

Pericentromeric cohesin is prematurely lost in *spo13Δ* cells
Previous analyses of fixed cells found that centromeric Rec8 is undetectable or greatly diminished in *spo13Δ* anaphase I cells (Klein *et al.*, 1999; Katis *et al.*, 2004b; Lee *et al.*, 2004).

Further evidence that Spo13 is important for protection of centromeric cohesion came from the analysis of cells lacking the monopolin subunit, Mam1, which biorient, rather than monoorient sister kinetochores, yet fail to segregate sister chromatids due to the persistence of centromeric cohesion. Importantly inactivation of *SPO13* allowed *mam1Δ* cells to segregate sister chromatids during anaphase I (Katis *et al.*, 2004b; Lee *et al.*, 2004). Together, these findings provide evidence that centromeric cohesion is impaired in *spo13Δ* cells. However, it has been argued that residual centromeric cohesin persists after securin destruction in *spo13Δ* cells and prevents timely spindle elongation (Katis *et al.*, 2004b). To clarify the importance of Spo13 in centromeric cohesion, we used live cell imaging of cells progressing through meiosis. We scored the percentage of cells where cohesin (Rec8-GFP) was retained at the pericentromere in anaphase I, as indicated by co-localization with the kinetochore protein Mtw1 (Figure 1A, B). To ensure that observed effects in *spo13Δ* cells were not a consequence of mono-orientation loss, which partially impacts cohesion (Nerusheva *et al.*, 2014), we simultaneously imaged *mam1Δ* cells for comparison. Quantification of pericentromeric Rec8 (Figure 1C) showed that, strikingly, deletion of *SPO13* leads to complete loss of cohesin in anaphase I. This is not due to impaired cohesin loading in early meiosis, since




Figure 1. Cohesin is lost at anaphase I in the absence of *SPO13*. (A) Representative images of Rec8-GFP, Mtw1-tdTomato and Pds1-tdTomato in live sporulating wild-type (AM13716), *spo13Δ* (AM15133), *mam1Δ* (AM15134) and *spo13Δ mam1Δ* (AM15135) cells. Scale bars represent 1 μ m. Arrows indicate pericentromeric cohesin. (B) The number of cells with pericentromeric Rec8-GFP in anaphase I is shown after scoring 50 cells from (A). (C) Rec8-GFP intensity was measured for 50 cells from (A) in the area occupied by the tdTomato-labeled kinetochore protein Mtw1. *** $p < 0.001$ (Welch two-sample t-test). (D) Rec8 loading is unaffected by deletion of *SPO13*. Rec8-3Ha association with the indicated sites was measured in prophase I in wild-type (AM4015), *spo13Δ* (AM15343), *mam1Δ* (AM15342) and *spo13Δ mam1Δ* (AM15344) cells carrying *ndt80Δ* and a no tag control (AM11633). Cells were arrested in prophase by harvesting 5 h after resuspension in sporulation medium and anti-Ha ChIP-qPCR performed. Error bars show standard error of the mean from three independent biological experiments.

prophase I-arrested *spo13Δ* cells have similar levels of Rec8 on centromeres compared to wild type (Figure 1D). We conclude that Spo13 is required for the retention of pericentromeric cohesin in anaphase I.

spo13Δ cells prematurely segregate sister chromatids

To assess sister chromatid cohesion in *spo13Δ* cells, we labelled one copy of chromosome V near the centromere with an array of tet operators (*tetO*), expressed GFP-tagged TetR repressor (Michaelis *et al.*, 1997) and imaged *CEN5*-GFP foci in live meiotic cells. Upon anaphase I entry (as judged by degradation of yeast securin Pds1 (Salah & Nasmyth, 2000)), three different phenotypes may be observed, depending on whether cells successfully mono-orient sister kinetochores and protect pericentromeric cohesin (Figure 2A). In wild-type cells, a single GFP focus segregates to one of the spindle poles (as marked by the spindle pole body component Spc42-tdTomato). Alternatively, in case of defective mono-orientation, split GFP foci stay in close proximity (<2 μm) because sister chromatids are cohered by pericentromeric cohesin. Lastly, in cells lacking both mono-orientation and sister chromatid cohesion, GFP foci split over a greater distance (>2 μm). Note that, using this assay, pericentromeric cohesion loss during

anaphase I can only be readily observed where it is accompanied by sister kinetochore bi-orientation. We subsequently scored the number of cells falling into either of these categories for each of the mutants analysed. This revealed that sister centromeres separate over large (>2 μm) distances in the half of *spo13Δ* anaphase I cells that bi-orient sister kinetochores (Figure 2B), consistent with all cohesion being lost. A small fraction of centromeres in *spo13Δ mam1Δ* cells, which bi-orient almost exclusively, stay in close proximity in the 30-minute time frame measured (Figure 2B), indicating that these cells at least temporarily retain sister chromatid cohesion. However, the loss of cohesion in all *spo13Δ* cells with bi-oriented kinetochores, the near-complete absence of Rec8, and the fact that deletion of *SPO13* permits efficient sister chromatid segregation in most *mam1Δ* cells (Figure 2B) (Katis *et al.*, 2004b; Lee *et al.*, 2004) together confirm that pericentromeric cohesion is predominantly non-functional in *spo13Δ* anaphase I cells.

Restoring the second nuclear division in *spo13Δ* cells does not prevent chromosome missegregation

We reasoned that the chromosome missegregation events seen in *spo13Δ* mutants might be related to the absence of the second nuclear division in these cells. Thus, restoring two




Figure 2. Deletion of *SPO13* permits sister chromosome segregation in anaphase I in *mam1Δ* mutants. (A) Assay for mono-orientation and cohesion defects using heterozygous centromeric fluorescent markers. Representative images are shown. Scale bars represent 1 μm. Images for $\Delta CEN5=0\mu\text{m}$, $\Delta CEN5=0-2\mu\text{m}$ and $\Delta CEN5>2\mu\text{m}$ were taken from wild-type, *mam1Δ* and *spo13Δ* cells, respectively. (B) Frequency of *CEN5* distance categories is shown for the indicated genotypes after live-cell imaging. Wild-type (AM15190), *spo13Δ* (AM15118), *mam1Δ* (AM15119) and *spo13Δ mam1Δ* (AM15120) cells carrying *SPC42-tdTomato*, *PDS1-tdTomato* and heterozygous TetR-GFP foci at *CEN5*, were sporulated for 2.5 h before imaging on a microfluidics plate.

divisions to *spo13Δ* cells by deletion of *MAD2* (Shonn *et al.*, 2002) would be expected to allow accurate chromosome segregation in the absence of Spo13. Our analysis of pericentromeric Rec8-GFP in anaphase I showed that, while pericentromeric cohesin in anaphase I is retained in wild-type and *mad2Δ* strains, it is lost to a similar degree in *spo13Δ* and *spo13Δ mad2Δ* mutants (Figure 3A–C). Intriguingly, *mad2Δ* cells were frequently unable to separate kinetochores in anaphase I, despite successful cleavage of arm cohesin (Figure 3A). While the reasons for this phenotype are unclear, we speculate that unattached kinetochores might persist into anaphase I when *MAD2* is deleted.

To analyse chromosome segregation in more detail, we followed cells carrying chromosomes labelled with Htb1-mCherry and heterozygous *CEN5*-GFP foci through meiosis (Figure 3D). To assess global chromosome segregation, we assayed the chromosomal content of spores by measuring the area occupied by Htb1-mCherry after meiosis II and calculated the ratio of the largest and smallest chromosomal mass in each cell. In wild-type cells, this ratio is close to 1 in most cells (Figure 3E) since all four nuclei are expected to be of similar size. In contrast, *spo13Δ mad2Δ* cells show a large variation in the chromosomal content of nuclei destined for spores (Figure 3E), indicating gross chromosome missegregation. We additionally investigated the segregation of heterozygous *CEN5*-GFP foci in these cells (Figure 3F, G). Similar to *spo13Δ* single mutants, a large proportion of *spo13Δ mad2Δ* double mutant cells split sister chromatids upon the first nuclear division (Figure 3F). Furthermore, 20% of *spo13Δ mad2Δ* cells display *CEN5*-GFP dot(s) in only one out of four spores after meiosis II (Figure 3G). This is largely caused by the absence of Spo13, since *mad2Δ* single mutants display a more modest missegregation phenotype (Figure 3G). Therefore, *spo13Δ mad2Δ* cells fail to accurately segregate chromosomes during both the first and second nuclear divisions. We conclude that the lack of Spo13 causes loss of centromeric cohesion during meiosis I and severe chromosome missegregation even when the second nuclear division is restored.

Sister chromatid cohesion is restored by preventing cohesin cleavage

A cleavage-independent, Rad61/Wpl1-dependent, cohesin removal pathway, similar to that which occurs in mammalian mitosis, operates during prophase I of budding yeast meiosis (Challa *et al.*, 2016; Challa *et al.*, 2019; Yu & Koshland 2005). We considered the possibility that cells lacking Spo13 lose cohesion, not due to its cleavage, but as a result of ectopic Rad61 activity. However, deletion of *RAD61* did not restore cohesion to *spo13Δ* cells (Figure 4A), indicating that a failure to counteract cleavage-independent cohesin removal is not solely responsible for the cohesion defect of cells lacking Spo13.

Next, we assessed whether cohesin cleavage is required for cohesion loss during anaphase I in *spo13Δ* cells. First, we inactivated Esp1 (separase), using the temperature-sensitive *esp1-2* mutant (Buonomo *et al.*, 2000) and followed Rec8-GFP by live cell imaging (Figure 4B–D). As expected, cohesin

remained on chromosomes even after anaphase I onset in both in *esp1-2* and *esp1-2 spo13Δ* cells and, consequently, sister chromatid segregation was largely prevented (Figure 4E).

Additionally, we prevented cohesin cleavage by mutating the separase cleavage site in Rec8 (Rec8-N) (Buonomo *et al.*, 2000). We followed GFP-tagged versions of this Rec8 variant through meiosis in wild- and *spo13Δ* cells (Figure 5A). Similar to *esp1-2* mutants, *rec8-N* prevents cleavage of cohesin along the length of the chromosome in *spo13Δ* cells (Figure 5B) and pericentromeric cohesin intensity is greatly increased (Figure 5C). Furthermore, we find that Rec8-N prevented the segregation of sister chromatids in *spo13Δ* mutants (Figure 5D). We conclude that cohesin cleavage is required for sister chromatid segregation in *spo13Δ* cells.

Interestingly, neither *esp1-2* (Figure 4E) nor Rec8-N (Figure 5D) prevented the splitting of sister centromeres in *spo13Δ* anaphase I, suggesting that pericentromeric cohesin may have been removed independently of cleavage in the absence of Spo13, allowing centromeres to come apart. However, cells lacking the mono-orientation protein Mam1 also split sister centromeres in anaphase I, despite intact pericentromeric cohesin protection (Tóth *et al.*, 2000). This suggests that the presence of uncleaved pericentromeric cohesin in anaphase I cannot prevent the sister centromere splitting resulting from defective mono-orientation in *spo13Δ* cells (Katis *et al.*, 2004b; Lee *et al.*, 2004). Moreover, centromere breathing – the splitting of centromeres in response to spindle tension despite high concentrations of cohesin in the pericentromere – has been observed in pre-anaphase mitotic cells of multiple species (Goshima & Yanagida, 2000; He *et al.*, 2000; Nabeshima *et al.*, 1998; Pearson *et al.*, 2001; Shelby *et al.*, 1996; Tanaka *et al.*, 2000). Thus, the splitting of centromeres in anaphase I in *spo13Δ* in the absence of cohesin cleavage does not confirm cleavage-independent cohesin removal in the pericentromere.

PP2A is functional in the absence of Spo13

Rec8 cleavage during wild-type meiosis relies on its prior phosphorylation (Brar *et al.*, 2006; Katis *et al.*, 2010) which is reversed in the pericentromere by PP2A. We considered the possibility that PP2A function may be impaired in *spo13Δ* cells, rendering it unable to dephosphorylate, and therefore protect, cohesin. We assessed whether tethering PP2A directly to cohesin could prevent Rec8 cleavage in the absence of Spo13. We fused GFP-binding protein (GBP), a nanobody specifically recognising GFP (Rothbauer *et al.*, 2006), to the PP2A regulatory subunit Rts1 to irreversibly tether PP2A to GFP-tagged Rec8. This was sufficient to prevent cohesin removal, both in *pCLB2-SGO1* and *spo13Δ* cells (Figure 6A–C). To further confirm the full functionality of Rts1 in *spo13Δ* cells, we utilised a separase biosensor (Yaakov *et al.*, 2012) where a cleavable Rec8 moiety is fused to GFP and LacI, with the latter allowing targeting of the biosensor to a *lacO* array on chromosome arms (Figure 7A). In wild-type and *spo13Δ* cells, this biosensor appears as a single GFP focus in meiosis I until separase is




Figure 3. Deletion of *MAD2* restores the second nuclear division, but not accurate chromosome segregation to *spo13Δ* mutants.

(A–C) Representative images of Rec8-GFP, Mtw1-tdTomato and Pds1-tdTomato in live sporulating wild-type (AM13716), *spo13Δ* (AM24843), *mad2Δ* (AM24844) and *spo13Δ mad2Δ* (AM24845) cells. Scale bars represent 1 μ m. Arrows indicate pericentromeric cohesin. (B) The number of cells with pericentromeric Rec8-GFP in anaphase I is shown after scoring 50 cells from (A). (C) Rec8-GFP intensity was measured for 50 cells from (A) as described for Figure 1C. *** $p < 0.001$, n.s. = not significant (Welch two-sample t-test). For *spo13Δ mad2Δ* mutants, the analysis in (B) and (C) was performed exclusively for cells that performed two divisions (as judged by the presence of four Mtw1-tdTomato foci after meiosis II). (D–G) Severe chromosome missegregation occurs in *spo13Δ mad2Δ* cells. (D) Representative images of wild-type (AM24848), *spo13Δ* (AM24849), *mad2Δ* (AM25221) and *spo13Δ mad2Δ* (AM25222) cells carrying heterozygous TetR-GFP foci at *CEN5* and *HTB1-mCherry*. Green arrows indicate *CEN5*-GFP segregation outcomes after meiosis I, cyan arrows indicate *CEN5*-GFP segregation outcomes after meiosis II. (E) Spores of *spo13Δ mad2Δ* vary greatly in the amount of nuclear DNA, as estimated by Htb1-mCherry area, thus indicating gross chromosome missegregation. The area occupied by Htb1-mCherry was measured in cells with four (wild type ($n=45$), *mad2Δ* ($n=31$) and *spo13Δ mad2Δ* ($n=33$)), or two (*spo13Δ* ($n=50$)) nuclear masses after meiosis II and variation in chromosomal area estimated by obtaining the ratio of the largest and smallest nuclear mass for each cell. ** $p < 0.01$, n.s. = not significant (Welch two-sample t-test). (F–G) *CEN5* missegregation in *spo13Δ mad2Δ* cells. Segregation of heterozygous *CEN5*-GFP foci was scored in 50 cells after the first (F) and second (G) nuclear division in the indicated strains. For *spo13Δ mad2Δ* mutants, the analysis in (F) and (G) was performed exclusively for cells that performed two divisions (as judged by the presence of four distinct Htb1-mCherry signals after meiosis II). Note that while a large proportion of *spo13Δ mad2Δ* cells end up with *CEN5*-GFP foci in two separate spores after meiosis II (similar to wild type), many of these cells have already segregated sister chromosomes in meiosis I (purple stripes), rather than meiosis II (gray).




Figure 4. Cohesin protection in *spo13Δ* cells is rescued by inhibition of separase, but not by ablation of the prophase pathway. (A) Deletion of *RAD61/WPL1* does not rescue sister chromatid cohesion in *spo13Δ* cells. Categorization of *CEN5*-GFP distances in wild-type (AM15190), *spo13Δ* (AM20146), *rad61Δ* (AM21068) and *spo13Δ rad61Δ* (AM21358) cells carrying *SPC42-tdTomato*, *PDS1-tdTomato* and heterozygous TetR-GFP dots at *CEN5* was carried out as described in Figure 2A. (B–D) Separase activity is required for Rec8 removal in *spo13Δ* mutants. Wild-type (AM13716), *spo13Δ* (AM20033), *esp1-2* (AM20868) and *spo13Δ esp1-2* (AM21949) cells carrying *REC8-GFP*, *MTW1-tdTomato* and *PDS1-tdTomato* were resuspended in sporulation medium at 32°C and grown in flasks for 3h before transferring to a microfluidics plate and imaged at 32°C. (B) The number of cells with the indicated patterns of Rec8-GFP localization in anaphase I was scored for 50 cells per strain. (C) The intensity of pericentromeric Rec8-GFP for the indicated genotypes is shown. The mean of the two maximum intensity values on a straight line connecting both kinetochores in anaphase I (within the first two time points after Pds1-tdTomato degradation) was measured for 50 cells. Error bars represent standard error. *** $p < 0.001$ (Welch two-sample t-test). (D) Representative images are shown. Scale bars represent 1 μm . Arrows indicate pericentromeric cohesin. (E) Inhibition of separase activity restores sister chromatid cohesion to *spo13Δ* mutants. Cohesion was assayed by categorization of *CEN5*-GFP distances as described in Figure 2A. Strains used were wild-type (AM15190), *spo13Δ* (AM20146), *esp1-2* (AM22498) and *spo13Δ esp1-2* (AM22499) cells carrying *SPC42-tdTomato*, *PDS1-tdTomato* and heterozygous TetR-GFP dots at *CEN5*.

activated in anaphase I, causing biosensor cleavage and GFP focus dispersal (Figure 7B, C). Tethering of Rts1 to the biosensor, however, prevents biosensor cleavage (Figure 7B, C). Therefore, our results indicate that PP2A is functional and capable of dephosphorylating cohesin in *spo13Δ* mutants.

Conclusions

The successful protection of pericentromeric cohesin is a key modification to the meiotic chromosome segregation machinery

as it ensures the fidelity of chromosome segregation in meiosis II. Key players in regulating cohesin cleavage are known. The kinases Hrr25 and Cdc7 (and possibly Cdc5) phosphorylate cohesin along the length of the chromosome to promote its cleavage by separase (Attner *et al.*, 2013; Brar *et al.*, 2006; Katis *et al.*, 2010), while pericentromeric Sgo1 recruits the phosphatase PP2A to dephosphorylate Rec8 and thereby protect it (Katis *et al.*, 2004a; Kitajima *et al.*, 2004; Lee *et al.*, 2008; Marston *et al.*, 2004; Riedel *et al.*, 2006; Tang *et al.*, 2006).




Figure 5. Cohesin cleavage is required for loss of sister chromatid cohesion in *spo13Δ* cells. (A–C) Non-cleavable Rec8 blocks efficient removal of cohesin in *spo13Δ* cells. **(A)** Representative images from movies of cells carrying Rec8-GFP, Mtw1-tdTomato, Pds1-tdTomato and with the indicated genotypes are shown. Scale bars represent 1 μm. Arrows indicate pericentromeric Rec8-GFP. **(B)** Frequency of cells with the indicated pattern of Rec8-GFP localization is shown for the indicated genotypes. **(C)** Rec8-GFP intensity was measured for the indicated genotypes as described in Figure 4C. Error bars represent standard error. ** $p < 0.01$, *** $p < 0.001$ (Welch two-sample t-test). Strains used in **(A–C)** were *REC8-GFP* (AM22190), *REC8-GFP spo13Δ* (AM22191), *rec8-N-GFP* (AM22192) and *rec8-N-GFP spo13Δ* (AM22193) cells carrying *MTW1-tdTomato* and *PDS1-tdTomato*. **(D)** Non-cleavable Rec8 prevents sister chromatid segregation in *spo13Δ* mutants. Cohesion functionality was determined for the indicated genotypes by categorization of *CEN5-GFP* distances as described for Figure 2A. Strains were *REC8-3HA* (AM22346), *REC8-3HA spo13Δ* (AM22347), *rec8-N-3HA* (AM22348) and *rec8-N-3HA spo13Δ* (AM22349) and carried *SPC42-tdTomato*, *PDS1-tdTomato* and heterozygous TetR-GFP dots at *CEN5*.

However, the meiosis I-specific Spo13, is also required to retain pericentromeric cohesin in anaphase I (Katis et al., 2004b; Lee et al., 2004; Shonn et al., 2002) but its function is much less well understood. Our study demonstrates that pericentromeric cohesin is prematurely removed in *spo13Δ* cells in a manner that requires cohesin cleavage and phosphorylation. Our recent work indicates that Spo13 achieves this by counteracting the activity of the cohesin kinases, Hrr25 and DDK (Galander et al., 2019). Future work should focus on elucidating how Spo13 elicits its effects on kinase function, and how this might be linked to its functions in both sister kinetochore mono-orientation and meiotic cell cycle control.

Methods

Yeast strains and plasmids

All strains are SK1-derivatives and are listed in Table 1. Plasmids generated in this study are listed in Table 2. Gene deletions, promoter replacements and gene tags were introduced using PCR-based methods (Gauss et al., 2005; Knop et al., 1999; Longtine et al., 1998; Moqtaderi & Struhl, 2008). *pCLB2-CDC20* (Lee & Amon, 2003), *REC8-GFP*, *PDS1-tdTomato* (Matos et al., 2008), *ndt80Δ* (Vincenten et al., 2015),

SPC42-tdTomato (Fox et al., 2017), *REC8-3HA* (Klein et al., 1999), *CEN5-GFP* dots, *mam1Δ::TRP1* (Tóth et al., 2000) and *REC8-N* (Buonomo et al., 2000) were described previously. Separase biosensor constructs (Yaakov et al., 2012) were a kind gift from David Morgan (Departments of Physiology and Biochemistry and Biophysics, UCSF).

Growth conditions

Cells were prepared for sporulation as described by Vincenten et al. (2015).

Chromatin immunoprecipitation

ChIP-qPCR was performed as previously described (Vincenten et al., 2015), using mouse anti-Ha (12CA5, Roche). All parameters and equipment are identical to those described previously, including qPCR mixes and thermocycling conditions. Primers for qPCR analysis are listed in Table 3.

Live cell imaging

Live cell imaging was performed on a DeltaVision Elite system (Applied Precision) connected to an inverted Olympus IX-71 microscope with a 100x UPlanSApo NA 1.4 oil lens. Images




Figure 6. PP2A can prevent cohesin cleavage and sister chromatid segregation in *spo13Δ* cells. (A–C) Cohesin is retained on chromosomes when PP2A^{Rts1} is tethered to Rec8. **(A)** Representative images of Rec8-GFP, Mtw1-dtTomato and Pds1-tdTomato in wild-type (AM13716), *spo13Δ* (AM20033), *pCLB2-SGO1* (AM21315), *RTS1-GBP* (AM21316), *spo13Δ pCLB2-SGO1* (AM21317), *spo13Δ RTS1-GBP* (AM21319), *pCLB2-SGO1 RTS1-GBP* (AM21318) and *spo13Δ pCLB2-SGO1 RTS1-GBP* (AM21320) cells undergoing meiosis. Scale bars represent 1 μ m. Arrows indicate pericentromeric cohesin. **(B)** The number of cells with pericentromeric cohesin in anaphase I was scored for 50 cells per strain. **(C)** Rec8-GFP intensity in anaphase I was measured as described in Figure 2A. ** $p < 0.01$, *** $p < 0.001$ (Welch two-sample t-test).

were taken using a Photometrics Cascade II EMCCD camera. The Deltavision system was controlled using SoftWoRx software, version 5.5. Live-cell imaging for Figure 3 was performed on a Zeiss Axio Observer Z1 (Zeiss UK, Cambridge) equipped with a Hamamatsu Flash 4 sCMOS camera, Prior motorised stage and Zen 2.3 acquisition software.

Cells were imaged at 30°C (unless stated) on an ONIX microfluidic perfusion platform by CellASIC. Cells were pre-grown in culture flasks for ~3 h before transfer to microfluidics plates.

Imaging began about 30 min later with images being acquired every 15 min for 12–15 h. Seven z-stacks were acquired with 0.85 μ m spacing. Image panels were assembled using Image-Pro Premier 3D, version 9.1 (Media Cybernetics). Images were analysed using ImageJ 1.48v (National Institutes of Health). Final image assembly was carried out using Adobe Photoshop CS5.1 and Adobe Illustrator CS5.1. Rec8-GFP intensities were measured using the DV_DotCounter custom plugin for ImageJ (Kelly, 2019a). The plugin applied a Z projection to each colour channel and allowed the user to select a cell of interest. Kinetochores




Figure 7. Fusion of Rts1 to a separase biosensor prevents its cleavage in both wild-type and *spo13Δ* cells. (A) Schematic illustration of the separase biosensor and its Rts1 fusion. (B and C) Wild-type (AM21557) and *spo13Δ* cells (AM21558) carrying a wild-type separase biosensor (*pCUP1-GFP-REC8(110-500)-LacI*) or an Rts1 fused biosensor (*pCUP1-GFP-REC8(110-500)-LacI-RTS1*; wild type: AM21559, *spo13Δ*: AM21800) as well as *lys2::lacOx256* and *PDS1-tdTomato* were sporulated in the presence of 100 nM CuSO_4 for 2.5 h before imaging on a microfluidics plate. (B) Representative images are shown. Scale bars represent 1 μm . (C) Scoring of 50 cells per strain for the presence of GFP foci (uncleaved biosensor) or diffuse GFP signal (cleaved biosensor) within 30 min (two time points) of Pds1 degradation.

in the red channel were identified by Yen *Auto Threshold* (Yen *et al.*, 1995) and their XY central coordinates, mean intensity and area recorded. The coordinates were then used to measure mean intensity in the corresponding location in the green channel, equivalent to pericentromeric Rec8-GFP. In

experiments where pericentromeric cohesin was likely to be found in between kinetochores (which is thought to occur in cells that bi-orient in meiosis I but retain cohesin), the XY coordinates in the red channel were used to generate a line profile between the 2 kinetochores in both colour channels over

Table 1. *Saccharomyces cerevisiae* strains used in this study.

AM strain	Relevant genotype
4015	<i>ndt80Δ::LEU2/ndt80Δ::LEU2</i> <i>REC8-3HA::URA3/REC8-3HA::URA3</i>
11633	<i>ndt80Δ::LEU2/ndt80Δ::LEU2</i>
13716	<i>REC8-GFP::URA3/REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i>
15118	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1"</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i>
15119	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
15120	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
15133	<i>REC8-GFP::URA3/REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i>
15134	<i>REC8-GFP::URA3/REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
15135	<i>REC8-GFP::URA3/REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
15190	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i>
15342	<i>ndt80Δ::LEU2/ndt80Δ::LEU2</i> <i>REC8-3HA::URA3/REC8-3HA::URA3</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
15343	<i>ndt80Δ::LEU2/ndt80Δ::LEU2</i> <i>REC8-3HA::URA3/REC8-3HA::URA3</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i>
15344	<i>ndt80Δ::LEU2/ndt80Δ::LEU2</i> <i>REC8-3HA::URA3/REC8-3HA::URA3</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i> <i>mam1Δ::TRP1/mam1Δ::TRP1</i>
20033	<i>REC8-GFP::URA3/REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i>

AM strain	Relevant genotype
20868	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>esp1-2/esp1-2</i>
21068	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>rad61Δ::KanMX6/rad61Δ::KanMX6</i>
21315	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>sgo1::KanMX6::pCLB2-SGO1/sgo1::KanMX6::pCLB2-SGO1</i>
21316	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>RTS1-GBP::His3MX6/RTS1-GBP::His3MX6</i>
21317	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>sgo1::KanMX6::pCLB2-SGO1/sgo1::KanMX6::pCLB2-SGO1</i>
21318	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>sgo1::KanMX6::pCLB2-SGO1/sgo1::KanMX6::pCLB2-SGO1</i> <i>RTS1-GBP::His3MX6/RTS1-GBP::His3MX6</i>
21319	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>RTS1-GBP::His3MX6/RTS1-GBP::His3MX6</i>
21320	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>sgo1::KanMX6::pCLB2-SGO1/sgo1::KanMX6::pCLB2-SGO1</i> <i>RTS1-GBP::His3MX6/RTS1-GBP::His3MX6</i>
21358	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>rad61Δ::KanMX6/rad61Δ::KanMX6</i>
21557	<i>his3::pCUP1-GFP-REC8(110-500)-LacI::HIS3/his3::pCUP1-GFP-REC8(110-500)-LacI::HIS3</i> <i>lys2::LEU2::lacOx256/lys2::LEU2::lacOx256</i> <i>PDS1-tdTomato-KITRP1/PDS1-tdTomato-KITRP1</i>
21558	<i>his3::pCUP1-GFP-REC8(110-500)-LacI::HIS3/his3::pCUP1-GFP-REC8(110-500)-LacI::HIS3 ></i> <i>lys2::LEU2::lacOx256/lys2::LEU2::lacOx256</i> <i>PDS1-tdTomato-KITRP1/PDS1-tdTomato-KITRP1</i> <i>spo13Δ::hphMX6/spo13Δ::hphMX6</i>
21559	<i>his3::pCUP1-GFP-REC8(110-500)-LacI-RTS1::HIS3/his3::pCUP1-GFP-REC8(110-500)-LacI-RTS1::HIS3</i> <i>lys2::LEU2::lacOx256/lys2::LEU2::lacOx256</i> <i>PDS1-tdTomato-KITRP1/PDS1-tdTomato-KITRP1</i>

AM strain	Relevant genotype
21800	<i>his3::pCUP1-GFP-REC8(110-500)-LacI-RTS1::HIS3/his3::pCUP1-GFP-REC8(110-500)-LacI-RTS1::HIS3 lys2::LEU2::lacOx256/lys2::LEU2::lacOx256</i> <i>PDS1-tdTomato-KITRP1/PDS1-tdTomato-KITRP1</i> <i>spo13Δ::hphMX6/spo13Δ::hphMX6</i>
21949	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>esp1-2/esp1-2</i>
22190	<i>rec8::REC8-GFP::LEU2::KanMX4/rec8::REC8-GFP::LEU2::KanMX4</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i>
22191	<i>rec8::REC8-GFP::LEU2::KanMX4/rec8::REC8-GFP::LEU2::KanMX4</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i>
22192	<i>rec8::rec8-N-GFP::LEU2::KanMX4/rec8::rec8-N-GFP::LEU2::KanMX4</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i>
22193	<i>rec8::rec8-N-GFP::LEU2::KanMX4/rec8::rec8-N-GFP::LEU2::KanMX4</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i>
22346	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>rec8::REC8-3HA::LEU2::KanMX4/rec8::REC8-3HA::LEU2::KanMX4</i>
22347	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>rec8::REC8-3HA::LEU2::KanMX4/rec8::REC8-3HA::LEU2::KanMX4</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i>
22348	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>rec8::rec8-N-3HA::LEU2::KanMX4/rec8::rec8-N-3HA::LEU2::KanMX4</i>
22349	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>rec8::rec8-N-3HA::LEU2::KanMX4/rec8::rec8-N-3HA::LEU2::KanMX4</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i>
22498	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>esp1-2/esp1-2</i>
22499	<i>SPC42-tdTomato::NatMX6/SPC42-tdTomato::NatMX6</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>spo13Δ::HphMX6/spo13Δ::HphMX6</i> <i>esp1-2/esp1-2</i>

AM strain	Relevant genotype
24843	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::hisG/spo13Δ::hisG</i>
24844	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>mad2Δ::KanMX6/mad2Δ::KanMX6</i>
24845	<i>REC8-GFP::URA3/ REC8-GFP::URA3</i> <i>PDS1-tdTomato::KITRP1/PDS1-tdTomato::KITRP1</i> <i>MTW1-tdTomato::NatMX6/MTW1-tdTomato::NatMX6</i> <i>spo13Δ::hisG/spo13Δ::hisG</i> <i>mad2Δ::KanMX6/mad2Δ::KanMX6</i>
24848	<i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>HTB1-mCherry::His3MX6/HTB1-mCherry::His3MX6</i>
24849	<i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>HTB1-mCherry::His3MX6/HTB1-mCherry::His3MX6</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i>
25221	<i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>HTB1-mCherry::His3MX6/HTB1-mCherry::His3MX6</i> <i>mad2Δ::HphMX6/mad2Δ::HphMX6</i>
25222	<i>leu2::pURA3-TetR-GFP::LEU2/leu2::hisG</i> <i>CEN5::tetOx224::HIS3/CEN5</i> <i>HTB1-mCherry::His3MX6/HTB1-mCherry::His3MX6</i> <i>spo13Δ::KanMX6/spo13Δ::KanMX6</i> <i>mad2Δ::HphMX6/mad2Δ::HphMX6</i>

Table 2. Plasmids generated in this study.

Plasmid	Description	Purpose and notes
AMp1317	YIplac128-REC8-GFP	<i>LEU2</i> integration plasmid carrying <i>REC8-GFP</i> .
AMp1368	YIplac128-rec8-N-GFP	<i>LEU2</i> integration plasmid carrying <i>rec8-N-GFP</i> .

Table 3. qPCR primers used in this study. For distances from centromeres, “-” indicates the location is upstream of the centromere, whereas “+” indicates the location is downstream of the centromere.

Chr.	Location	Distance from centromere	Primer pair	Sequence
III	Centromere	+0.25kb	1279	TGTTGATGGGTTTACAATTT
			1280	CTTCAATGATTGCTCTAAATC
IV	Arm	-95kb	782	AGATGAAACTCAGGCTACCA
			783	TGCAACATCGTTAGTTCCTG
IV	Centromere	+0.15kb	794	CCGAGGCTTTCATAGCTTA
			795	ACCGGAAGGAAGAATAAGAA

exactly the same pixels. The two brightest peaks in the line profile of the green channel were calculated to give the maximum intensity value for each. Rec8-GFP intensity was measured in this manner for [Figure 4C](#) and [Figure 5C](#). The plugin used was the custom YeastLineProfiler for ImageJ ([Kelly, 2019b](#)). Chromosomal area in [Figure 3E](#) was measured using a custom ImageJ plugin ([Kelly, 2019c](#)) that identifies the regions of bright fluorescence in the red channel using Yen Auto Threshold and subsequently measures the area of these regions of interest.

An earlier version of this article can be found on bioRxiv (DOI: <https://doi.org/10.1101/488312>)

Data availability

Raw data for scoring imaging experiments and ChIP-qPCR, arranged by figure, is available from OSF. DOI: <https://doi.org/10.17605/OSF.IO/VBKU9> ([Marston, 2019](#)).

Data are available under the terms of the Creative Commons Zero “No rights reserved” data waiver (CC0 1.0 Public domain dedication).

The file size of the raw microscopy movies precludes uploading them to OSF, but are available upon request from adele.marston@ed.ac.uk.

Software availability

Source code for DV_DotCounter is available from: https://github.com/dkelly604/DV_DotCounter.

Archived source code at time of publication: <https://doi.org/10.5281/zenodo.2553081> ([Kelly, 2019a](#)).

Source code for YeastLineProfiler is available from: <https://github.com/dkelly604/YeastLineProfiler>.

Archived source code at time of publication: <https://doi.org/10.5281/zenodo.2560099> ([Kelly, 2019b](#)).

Source code for Size_and_Area is available from: https://github.com/dkelly604/Size_and_Area.

Archived source code at time of publication: <https://doi.org/10.5281/zenodo.3358842> ([Kelly, 2019c](#)).

License: MIT License.

Grant information

This research was funded by Wellcome through PhD studentships to SG [096994] and REB [102316], as well as a Senior Research Fellowship to ALM [107827] and core funding awarded to the Wellcome Centre for Cell Biology [203149].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We are very grateful to Robin Allshire, Angelika Amon, Kevin Hardwick, Heinrich Leonhardt, David Morgan and Ulrich Rothbauer for sharing plasmids. We also thank members of the Marston lab for helpful discussion.

References

- Argüello-Miranda O, Zagoriy I, Mengoli V, *et al.*: **Casein Kinase 1 Coordinates Cohesin Cleavage, Gametogenesis, and Exit from M Phase in Meiosis II.** *Dev Cell.* 2017; **40**(1): 37–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Attner MA, Miller MP, Ee LS, *et al.*: **Polo kinase Cdc5 is a central regulator of meiosis I.** *Proc Natl Acad Sci U S A.* 2013; **110**(35): 14278–14283.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brar GA, Kiburz BM, Zhang Y, *et al.*: **Rec8 phosphorylation and recombination promote the step-wise loss of cohesins in meiosis.** *Nature.* 2006; **441**(7092): 532–536.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Buheitel J, Stemmann O: **Prophase pathway-dependent removal of cohesin from human chromosomes requires opening of the Smc3-Scc1 gate.** *EMBO J.* 2013; **32**(5): 666–676.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buonomo SB, Clyne RK, Fuchs J, *et al.*: **Disjunction of homologous chromosomes in meiosis I depends on proteolytic cleavage of the meiotic cohesin Rec8 by separin.** *Cell.* 2000; **103**(3): 387–398.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Challa K, Fajish VG, Shinohara M, *et al.*: **Meiosis-specific prophase-like pathway controls cleavage-independent release of cohesin by Wapl phosphorylation.** *PLoS Genet.* 2019; **15**(1): e1007851.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challa K, Lee MS, Shinohara M, *et al.*: **Rad61/Wpl1 (Wapl), a cohesin regulator, controls chromosome compaction during meiosis.** *Nucleic Acids Res.* 2016; **44**(7): 3190–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fox C, Zou J, Rappsilber J, *et al.*: **Cdc14 phosphatase directs centrosome re- duplication at the meiosis I to meiosis II transition in budding yeast.** [version 2; referees: 4 approved] *Wellcome Open Res.* 2017; **2**: 2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Galander S, Barton RE, Borek WE, *et al.*: **Reductional Meiosis I Chromosome Segregation Is Established by Coordination of Key Meiotic Kinases.** *Dev Cell.* 2019; **49**(4): 526–541.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gauss R, Trautwein M, Sommer T, *et al.*: **New modules for the repeated internal and N-terminal epitope tagging of genes in *Saccharomyces cerevisiae*.** *Yeast.* 2005; **22**(1): 1–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Goshima G, Yanagida M: **Establishing biorientation occurs with precocious separation of the sister kinetochores, but not the arms, in the early spindle of budding yeast.** *Cell.* 2000; **100**(6): 619–633.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gruber S, Haering CH, Nasmyth K: **Chromosomal cohesin forms a ring.** *Cell.* 2003; **112**(6): 765–777.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Haering CH, Löwe J, Hochwagen A, *et al.*: **Molecular architecture of SMC proteins and the yeast cohesin complex.** *Mol Cell.* 2002; **9**(4): 773–788.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hartman T, Stead K, Koshland D, *et al.*: **Pds5p is an essential chromosomal protein required for both sister chromatid cohesion and condensation in *Saccharomyces cerevisiae*.** *J Cell Biol.* 2000; **151**(3): 613–626.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hassold T, Hunt P: **To err (meiotically) is human: the genesis of human aneuploidy.** *Nat Rev Genet.* 2001; **2**(4): 280–291.
[PubMed Abstract](#) | [Publisher Full Text](#)
- He X, Asthana S, Sorger PK: **Transient sister chromatid separation and elastic deformation of chromosomes during mitosis in budding yeast.** *Cell.* 2000;

- 101(7): 763–775.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jonak K, Zagorij I, Oz T, *et al.*: **APC/C-Cdc20 mediates deprotection of centromeric cohesin at meiosis II in yeast.** *Cell Cycle*. 2017; 16(12): 1145–1152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Katis VL, Gálová M, Rabitsch KP, *et al.*: **Maintenance of cohesin at centromeres after meiosis I in budding yeast requires a kinetochore-associated protein related to MEI-S332.** *Curr Biol*. 2004a; 14(7): 560–572.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Katis VL, Lipp JJ, Imre R, *et al.*: **Rec8 phosphorylation by casein kinase 1 and Cdc7-Dbf4 kinase regulates cohesin cleavage by separate during meiosis.** *Dev Cell*. 2010; 18(3): 397–409.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Katis VL, Matos J, Mori S, *et al.*: **Spo13 facilitates monopolin recruitment to kinetochores and regulates maintenance of centromeric cohesion during yeast meiosis.** *Curr Biol*. 2004b; 14(24): 2183–2196.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kelly DA: **dkelly604/DV_DotCounter: Yeast Spot Counter (Version v1.0).** *Zenodo*. 2019a.
<http://www.doi.org/10.5281/zenodo.2553082>
- Kelly DA: **dkelly604/YeastLineProfiler: YeastLineProfiler_V1.0 (Version V1).** *Zenodo*. 2019b.
<http://www.doi.org/10.5281/zenodo.2560099>
- Kelly DA: **dkelly604/Size_and_Area: Yeast Measure Red dot fluorescence (Version V1).** *Zenodo*. 2019c.
<http://www.doi.org/10.5281/zenodo.3358842>
- Kiburz BM, Reynolds DB, Megee PC, *et al.*: **The core centromere and Sgo1 establish a 50-kb cohesin-protected domain around centromeres during meiosis I.** *Genes Dev*. 2005; 19(24): 3017–3030.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kim J, Ishiguro K, Nambu A, *et al.*: **Meikin is a conserved regulator of meiosis-I-specific kinetochore function.** *Nature*. 2015; 517(7535): 466–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kitajima TS, Kawashima SA, Watanabe Y: **The conserved kinetochore protein shugoshin protects centromeric cohesion during meiosis.** *Nature*. 2004; 427(6974): 510–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kitajima TS, Sakuno T, Ishiguro K, *et al.*: **Shugoshin collaborates with protein phosphatase 2A to protect cohesin.** *Nature*. 2006; 441(7089): 46–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Klapholz S, Esposito RE: **Recombination and chromosome segregation during the single division meiosis in SPO12-1 and SPO13-1 diploids.** *Genetics*. 1980; 96(3): 589–611.
[PubMed Abstract](#) | [Free Full Text](#)
- Klein F, Mahr P, Gálová M, *et al.*: **A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis.** *Cell*. 1999; 98(1): 91–103.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Knop M, Siegers K, Pereira G, *et al.*: **Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines.** *Yeast*. 1999; 15(10B): 963–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lafont AL, Song J, Rankin S: **Sororin cooperates with the acetyltransferase Eco2 to ensure DNA replication-dependent sister chromatid cohesion.** *Proc Natl Acad Sci U S A*. 2010; 107(47): 20364–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee BH, Amon A: **Role of Polo-like kinase CDC5 in programming meiosis I chromosome segregation.** *Science*. 2003; 300(5618): 482–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee BH, Amon A, Prinz S: **Spo13 regulates cohesin cleavage.** *Genes Dev*. 2002; 16(13): 1672–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee BH, Kiburz BM, Amon A: **Spo13 maintains centromeric cohesion and kinetochore coorientation during meiosis I.** *Curr Biol*. 2004; 14(24): 2168–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee J, Kitajima TS, Tanno Y, *et al.*: **Unified mode of centromeric protection by shugoshin in mammalian oocytes and somatic cells.** *Nat Cell Biol*. 2008; 10(1): 42–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu H, Jia L, Yu H: **Phospho-H2A and cohesin specify distinct tension-regulated Sgo1 pools at kinetochores and inner centromeres.** *Curr Biol*. 2013a; 23(19): 1927–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu H, Rankin S, Yu H: **Phosphorylation-enabled binding of SGO1-PP2A to cohesin protects sororin and centromeric cohesion during mitosis.** *Nat Cell Biol*. 2013b; 15(1): 40–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Longtine MS, McKenzie A, Demarini DJ, *et al.*: **Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*.** *Yeast*. 1998; 14(10): 953–961.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Losada A, Hirano M, Hirano T: **Identification of *Xenopus* SMC protein complexes required for sister chromatid cohesion.** *Genes Dev*. 1998; 12(13): 1986–1997.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Marston A: **Spo13 prevents premature cohesin cleavage during meiosis.** 2019.
<http://www.doi.org/10.17605/OSF.IO/VBKU9>
- Marston AL, Amon A: **Meiosis: cell-cycle controls shuffle and deal.** *Nat Rev Mol Cell Biol*. 2004; 5(12): 983–997.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Marston AL, Tham WH, Shah H, *et al.*: **A genome-wide screen identifies genes required for centromeric cohesion.** *Science*. 2004; 303(5662): 1367–1370.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Matos J, Lipp JJ, Bogdanova A, *et al.*: **Dbf4-dependent CDC7 kinase links DNA replication to the segregation of homologous chromosomes in meiosis I.** *Cell*. 2008; 135(4): 662–678.
[PubMed Abstract](#) | [Publisher Full Text](#)
- McGuinness BE, Hirota T, Kudo NR, *et al.*: **Shugoshin prevents dissociation of cohesin from centromeres during mitosis in vertebrate cells.** *PLoS Biol*. 2005; 3(3): e86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Michaelis C, Ciosk R, Nasmyth K: **Cohesins: chromosomal proteins that prevent premature separation of sister chromatids.** *Cell*. 1997; 91(1): 35–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Miyazaki S, Kim J, Yamagishi Y, *et al.*: **Meikin-associated polo-like kinase specifies Bub1 distribution in meiosis I.** *Genes Cells*. 2017; 22(6): 552–567.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Moqtaderi Z, Struhl K: **Expanding the repertoire of plasmids for PCR-mediated epitope tagging in yeast.** *Yeast*. 2008; 25(4): 287–292.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nabeshima K, Nakagawa T, Straight AF, *et al.*: **Dynamics of centromeres during metaphase-anaphase transition in fission yeast: Dis1 is implicated in force balance in metaphase bipolar spindle.** *Mol Biol Cell*. 1998; 9(11): 3211–3225.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nerusheva OO, Galander S, Fernus J, *et al.*: **Tension-dependent removal of pericentromeric shugoshin is an indicator of sister chromosome biorientation.** *Genes Dev*. 2014; 28(12): 1291–1309.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nishiyama T, Ladurner R, Schmitz J, *et al.*: **Sororin mediates sister chromatid cohesion by antagonizing Wapl.** *Cell*. 2010; 143(5): 737–749.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Panizza S, Tanaka T, Hochwagen A, *et al.*: **Pds5 cooperates with cohesin in maintaining sister chromatid cohesion.** *Curr Biol*. 2000; 10(24): 1557–1564.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pearson CG, Maddox PS, Salmon ED, *et al.*: **Budding yeast chromosome structure and dynamics during mitosis.** *J Cell Biol*. 2001; 152(6): 1255–1266.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rankin S, Ayad NG, Kirschner MW: **Sororin, a substrate of the anaphase-promoting complex, is required for sister chromatid cohesion in vertebrates.** *Mol Cell*. 2005; 18(2): 185–200.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Riedel CG, Katis VL, Katou Y, *et al.*: **Protein phosphatase 2A protects centromeric sister chromatid cohesion during meiosis I.** *Nature*. 2006; 441(7089): 53–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rolef Ben-Shahar T, Heeger S, Lehane C, *et al.*: **Eco1-Dependent Cohesin Acetylation During Establishment of Sister Chromatid Cohesion.** *Science*. 2008; 321(5888): 563–566.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rothbauer U, Zolghadr K, Tillib S, *et al.*: **Targeting and tracing antigens in live cells with fluorescent nanobodies.** *Nat Methods*. 2006; 3(11): 887–889.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Salah SM, Nasmyth K: **Destruction of the securin Pds1p occurs at the onset of anaphase during both meiotic divisions in yeast.** *Chromosoma*. 2000; 109(1–2): 27–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schmitz J, Watrin E, Lénárt P, *et al.*: **Sororin is required for stable binding of cohesin to chromatin and for sister chromatid cohesion in interphase.** *Curr Biol*. 2007; 17(7): 630–636.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shelby RD, Hahn KM, Sullivan KF: **Dynamic elastic behavior of alpha-satellite DNA domains visualized in situ in living human cells.** *J Cell Biol*. 1996; 135(3): 545–557.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shonn MA, McCarroll R, Murray AW: **Spo13 protects meiotic cohesin at centromeres in meiosis I.** *Genes Dev*. 2002; 16(13): 1659–1671.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sumara I, Vorlaufer E, Gieffers C, *et al.*: **Characterization of vertebrate cohesin complexes and their regulation in prophase.** *J Cell Biol*. 2000; 151(4): 749–762.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tanaka T, Fuchs J, Loidl J, *et al.*: **Cohesin ensures bipolar attachment of microtubules to sister centromeres and resists their precocious separation.** *Nat Cell Biol*. 2000; 2(8): 492–499.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tang Z, Shu H, Qi W, *et al.*: **PP2A is required for centromeric localization of**

Sgo1 and proper chromosome segregation. *Dev Cell.* 2006; 10(5): 575–585.

[PubMed Abstract](#) | [Publisher Full Text](#)

Tóth A, Ciosk R, Uhlmann F, *et al.*: **Yeast Cohesin complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between sister chromatids during DNA replication.** *Genes Dev.* 1999; 13(3): 320–333.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tóth A, Rabitsch KP, Gálová M, *et al.*: **Functional genomics identifies monopolin: a kinetochore protein required for segregation of homologs during meiosis I.** *Cell.* 2000; 103(7): 1155–1168.

[PubMed Abstract](#) | [Publisher Full Text](#)

Unal E, Heidinger-Pauli JM, Kim W, *et al.*: **A molecular determinant for the establishment of sister chromatid cohesion.** *Science.* 2008; 321(5888): 566–569.

[PubMed Abstract](#) | [Publisher Full Text](#)

Varela E, Schlecht U, Moina A, *et al.*: **Mitotic expression of Spo13 alters M-phase progression and nucleolar localization of Cdc14 in budding yeast.** *Genetics.* 2010; 185(3): 841–854.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Vincenten N, Kuhl LM, Lam I, *et al.*: **The kinetochore prevents centromere-proximal crossover recombination during meiosis.** *eLife.* 2015; 4: pii: e10850.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Waizenegger IC, Hauf S, Meinke A, *et al.*: **Two distinct pathways remove mammalian cohesin from chromosome arms in prophase and from centromeres in anaphase.** *Cell.* 2000; 103(3): 399–410.

[PubMed Abstract](#) | [Publisher Full Text](#)

Wang HT, Frackman S, Kowalysyn J, *et al.*: **Developmental regulation of SPO13, a gene required for separation of homologous chromosomes at meiosis I.** *Mol Cell Biol.* 1987; 7(4): 1425–1435.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Warren WD, Steffensen S, Lin E, *et al.*: **The *Drosophila* RAD21 cohesin persists at the centromere region in mitosis.** *Curr Biol.* 2000; 10(22): 1463–6.

[PubMed Abstract](#) | [Publisher Full Text](#)

Watanabe Y, Nurse P: **Cohesin Rec8 is required for reductional chromosome segregation at meiosis.** *Nature.* 1999; 400(6743): 461–4.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yaakov G, Thorn K, Morgan DO: **Separase Biosensor Reveals that Cohesin Cleavage Timing Depends on Phosphatase PP2A(Cdc55) Regulation.** *Dev Cell.* 2012; 23(1): 124–36.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yamada T, Tahara E, Kanke M, *et al.*: ***Drosophila* Dalmatian combines sororin and shugoshin roles in establishment and protection of cohesion.** *EMBO J.* 2017; 36(11): 1513–1527.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Yen JC, Chang FJ, Chang S: **A new criterion for automatic multilevel thresholding.** *IEEE Trans Image Process.* 1995; 4(3): 370–8.

[PubMed Abstract](#) | [Publisher Full Text](#)

Yu HG, Koshland D: **Chromosome morphogenesis: condensin-dependent cohesin removal during meiosis.** *Cell.* 2005; 123(3): 397–407.

[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:   

Version 2

Reviewer Report 10 September 2019

<https://doi.org/10.21956/wellcomeopenres.16829.r36375>

© 2019 Hochwagen A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Andreas Hochwagen 

Department of Biology, New York University, New York City, NY, USA

The revised manuscript adequately addresses all points raised in my review of the original manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Chromosome biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 19 March 2019

<https://doi.org/10.21956/wellcomeopenres.16436.r34862>

© 2019 Martinez-Perez E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Enrique Martinez-Perez

MRC London Institute of Medical Sciences, Imperial College London, London, UK

Accurate formation of haploid gametes during meiosis requires the step-wise removal of cohesin during the consecutive meiotic divisions. Two different pathways contribute to this process, one mediated by Wapl, which removes cohesin before the onset of the first meiotic division, and a second dependent on the protease separase, which cleaves the kleisin subunit (Rec8) of cohesin at the onset of the meiotic divisions. Phosphorylation of Rec8 by multiple kinases promotes cohesin release by Wapl and cleavage

by separase, therefore the pool of cohesin bound to centromeric regions must be protected from phosphorylation during the first meiotic division to prevent premature loss of cohesion. Galander et al. use *in vivo* imaging to investigate cohesin protection in budding yeast, focusing on Spo13, which role in this process remains poorly understood.

The authors show convincingly that *spo13* mutants display premature loss of Rec8 and sister chromatid cohesion during the first meiotic division, that this premature loss of cohesion requires separase but not Wapl, and that expression of a separase-resistant Rec8 rescues cohesin loss in *spo13* mutants. These results demonstrate an important role for Spo13 in preventing separase-dependent Rec8 removal during meiosis I.

Specific comments:

Figure 4 shows that expression of Rec8-N (separase resistant) prevents loss of Rec8 from pericentromeric regions in *spo13* mutants, but despite this, sister centromeres still show substantial separation in ~50% of the cells. How can sister centromeres achieve this level of separation despite extensive Rec8 binding?

The introduction doesn't mention the Challa et al 2019¹ paper describing the role of Wapl in promoting Rec8 removal before anaphase I. Since distinguishing the contribution of the Wapl and separase pathways to the cohesion defects of *spo13* mutants is a key aspect of the manuscript, mentioning the Challa et al 2019 in the introduction will help the reader.

Non-yeast experts will benefit from a more detailed description of some of the markers used in the study, such as Mtw1.

References

1. Challa K, Fajish V G, Shinohara M, Klein F, Gasser SM, Shinohara A: Meiosis-specific prophase-like pathway controls cleavage-independent release of cohesin by Wapl phosphorylation. *PLoS Genet.* **15** (1): e1007851 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Meiosis

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 05 Aug 2019

Adele Marston, University of Edinburgh, Edinburgh, UK

Figure 4 shows that expression of Rec8-N (separase resistant) prevents loss of Rec8 from pericentromeric regions in spo13 mutants, but despite this, sister centromeres still show substantial separation in ~50% of the cells. How can sister centromeres achieve this level of separation despite extensive Rec8 binding?

Please refer to our comment on the final point made by reviewer 2. Sister kinetochores are bioriented.

The introduction doesn't mention the Challa et al 2019¹ paper describing the role of Wapl in promoting Rec8 removal before anaphase I. Since distinguishing the contribution of the Wapl and separase pathways to the cohesion defects of spo13 mutants is a key aspect of the manuscript, mentioning the Challa et al 2019 in the introduction will help the reader.

We agree and are sorry for this omission. We have now done this.

Non-yeast experts will benefit from a more detailed description of some of the markers used in the study, such as Mtw1.

We have clarified the function of Mtw1 in our updated manuscript and also provided a better explanation of the *mam1Δ* phenotype.

Competing Interests: No competing interests were disclosed.

Reviewer Report 14 March 2019

<https://doi.org/10.21956/wellcomeopenres.16436.r34866>

© 2019 Dawson D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Dean S. Dawson

Program in Cell Cycle and Cancer Biology, Oklahoma Medical Research Foundation, Oklahoma City, OK, USA

Galander and colleagues have revisited the role of Spo13 in protecting or regulating sister chromatid cohesion in meiosis I. Previous work has shown that Spo13 affects the disposition of cohesin at

centromeres but the molecular basis for the modulation of centromeric cohesion by Spo13 has remained unclear. Recent work has shown that there are two pathways for cohesin removal in meiosis in budding yeast. Here the authors test which pathway is impacted by Spo13. The experiments have moved the field forward by using live cell imaging methods to address this question. The results demonstrate that Spo13, at least in part, protects Rec8 at the centromeres from cleavage by separase.

Introduction

Para. 3, line 7. Might be good to adjust the sentence saying Rec8 cleavage is dependent on Cdc5. The next sentence indicates the lack of clarity on this point.

Paragraph 4: It would be worth mentioning the two-step removal of cohesins in budding yeast meiosis here (Yu and Koshland, 2005¹; Challa et al., 2019²), along with referencing the mammalian and fly two-step processes.

Results and Discussion

Page 4, second column, 4 lines from bottom: “withbi”

Fig. 3 E/ Fig. 4 D – spo13 delete allows meiosis I sister centromere separation in esp1 and REC8-N mutants. This shows centromeres are more able to separate in spo13 mutants even without Rec8 cleavage. There are multiple possible explanations for these results. Is it because sister centromeres are more easily bi-oriented in spo13 mutants? Alternatively, could it be that Spo13 also promotes sister centromere cohesion also protects pericentromeric cohesion through a pathway that doesn't involve cleavage? The manuscript would benefit from brief comments from the authors on the implications of these observations.

References

1. Yu HG, Koshland D: Chromosome morphogenesis: condensin-dependent cohesin removal during meiosis. *Cell*. 2005; **123** (3): 397-407 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Challa K, Fajish V G, Shinohara M, Klein F, Gasser SM, Shinohara A: Meiosis-specific prophase-like pathway controls cleavage-independent release of cohesin by Wapl phosphorylation. *PLoS Genet*. **15** (1): e1007851 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.**Reviewer Expertise:** meiotic chromosome biology**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Author Response 05 Aug 2019

Adele Marston, University of Edinburgh, Edinburgh, UK

Para. 3, line 7. Might be good to adjust the sentence saying Rec8 cleavage is dependent on Cdc5. The next sentence indicates the lack of clarity on this point.

Thanks for the suggestion, we have done this.

Paragraph 4: It would be worth mentioning the two-step removal of cohesins in budding yeast meiosis here (Yu and Koshland, 2005¹; Challa et al., 2019²), along with referencing the mammalian and fly two-step processes.

Thanks for pointing out this omission, we have added the suggested text and references.

Results and Discussion

Page 4, second column, 4 lines from bottom: "withbi"

This has been corrected

Fig. 3 E/ Fig. 4 D – spo13 delete allows meiosis I sister centromere separation in esp1 and REC8-N mutants. This shows centromeres are more able to separate in spo13 mutants even without Rec8 cleavage. There are multiple possible explanations for these results. Is it because sister centromeres are more easily bi-oriented in spo13 mutants? Alternatively, could it be that Spo13 also promotes sister centromere cohesion also protects pericentromeric cohesion through a pathway that doesn't involve cleavage? The manuscript would benefit from brief comments from the authors on the implications of these observations.

We believe that this is due to the fact that sister centromeres are bioriented in *spo13* mutants. We have added a few sentences explaining these observations and the previous evidence that shows that defects in monoorientation result in sister centromere splitting even without loss of cohesion.

Competing Interests: No competing interests were disclosed.

Reviewer Report 27 February 2019

<https://doi.org/10.21956/wellcomeopenres.16436.r34863>

© 2019 Hochwagen A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Andreas Hochwagen

Department of Biology, New York University, New York City, NY, USA

This manuscript by Galander and colleagues investigates the function of the Spo13, a central but poorly understood regulator of meiotic chromosome segregation, using live-cell imaging in *S. cerevisiae*. Previous work had shown that Spo13 weakens the protection of centromeric cohesin during meiosis I, but the extent of this weakening had been questioned. The authors show an essentially complete loss of centromeric cohesin by fluorescence microscopy and support this data by monitoring the segregation of sister centromeres. In addition, they propose that this weakening occurs through increased separase activity because of reduced cohesin phosphorylation in the pericentromeric regions. The latter conclusion is complicated by the fact that the phosphorylation experiments do not distinguish between cohesion at centromeres and along chromosome arms.

I suggest the authors address the following points:

1. The data shown in panel 1C is central to the presented conclusions. As this bar graph relies on standard error, please provide statistical analysis including multiple hypothesis testing for this graph.
2. I am confused why there are cells that did not lose cohesion in the mam1 spo13 double mutants (panel 2B). Does this data not indicate that there may be some cohesion remaining in the absence of Spo13?
3. The Rts1-GBP construct will lead to ectopic protection also along chromosome arms. This increased signal is expected lead to an elevated Rec8 fluorescence intensity in the pericentromeric regions given the low spatial resolution of this assay. I think a ChIP experiment comparing Rec8 at arm and pericentromeric sites would be important to exclude the possibility that there is Rts1-independent Rec8 removal in the spo13 mutant. This issue is particularly relevant given a recent paper by Mehta et al¹ (Curr Genet) that came to the conclusion that Spo13 acts independently of Rts1.

Typo:

Page 4, second paragraph: withbi-oriented

References

1. Mehta G, Anbalagan GK, Bharati AP, Gadre P, Ghosh SK: An interplay between Shugoshin and Spo13 for centromeric cohesin protection and sister kinetochore mono-orientation during meiosis I in *Saccharomyces cerevisiae*. *Curr Genet*. 2018; **64** (5): 1141-1152 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Chromosome biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Aug 2019

Adele Marston, University of Edinburgh, Edinburgh, UK

1. We performed statistical analysis using Welch's two sample t-test, which suggests that the observed differences are highly significant.
2. While we agree that this is a possibility, we believe that there are alternative explanations for this observation. Firstly, it could be that, for unknown reasons, the deletion of *MAM1* affects successful segregation of chromosomes in anaphase I in *spo13Δ* cells (which otherwise retain some monopolin at kinetochores (Lee et al. (2004) *Curr Biol* 14, 2168-2182). Indeed, in Figure 2B, as well as additional figures in this manuscript (new Figures 4E and 5D), the fraction of otherwise wild-type *spo13Δ* cells that fail to segregate sister chromatids is minimal (2%). Secondly, the cells that appear to retain cohesin may be in the process of losing cohesin but our imaging could have captured them immediately after Pds1 degradation, so that they have not yet had a chance to segregate chromosomes before we scored them. To further test whether cohesin is retained at centromeres in the absence of Spo13, we analysed cells that also lack Mad2 and which have previously been found to undergo two divisions. This data, shown in a new Figure 3, confirms our previous conclusions that Spo13 is required for cohesin protection.
3. While we agree that Rts1-independent Rec8 removal in the *spo13Δ* mutant is an interesting possibility, we decided not to do this experiment because our opinion is that it would be extremely difficult to interpret due to the difficulty of obtaining comparable cell cycle stages in the different mutants. The experiment would require analysis of cells in anaphase I (or at least past metaphase I) and, due to the metaphase I delay observed in *spo13Δ* cells even when synchronised using the *GAL-NDT80* block release system (Carlile and Amon, 2008), it would be difficult to obtain populations in which similar numbers of cells of each mutant are in this stage. Furthermore, it would be difficult to assess the cell cycle stage, since RTS1-GBP cells would presumably not exhibit standard anaphase spindles due to the failure to segregate chromosomes in anaphase I (compare distances of kinetochores in our imaging at the 15 minute time point in Figure 6A of the updated version of the manuscript). Additionally, *spo13Δ* mutants prematurely exit meiosis after the first division, precluding cell cycle analysis by immunoblot against an anaphase I marker.

However, our main conclusion from these experiments is that PP2A-Rts1 is functional in the absence of Spo13 and we believe are data supports this interpretation.

We also respectfully disagree with the conclusion in Mehta et al. based on the available data. Their conclusion that Spo13 acts independently of Rts1 was derived from data presented in Figure 3 of their paper where they showed that *CEN5-GFP* foci segregate to opposite poles during anaphase I in more *spo13Δ rts1Δ* than *spo13* cells. However, these genotypes are not complete: the methods section and strain table reveal that all *spo13Δ* cells (but none of the other mutations) also carry *mad2Δ* delete to allow the cells to go through two meiotic divisions. Therefore, this experiment is not properly controlled, especially because *mad2* cells themselves are compromised in chromosome segregation (Figure 3 in our current manuscript).

In conclusion, while Rts1-independent Rec8 removal by Spo13 remains a possibility, there is currently no evidence for it.

The typo on Page 4 has been corrected.

Competing Interests: No competing interests were disclosed.