



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Abnormal fatty acid metabolism is a core component of spinal muscular atrophy

Citation for published version:

Deguisse, M-O, Baranello, G, Mastella, C, Beauvais, A, Michaud, J, Leone, A, De Amicis, R, Battezzati, A, Dunham, C, Selby, K, Warman Chardon, J, McMillan, HJ, Huang, Y-T, Courtney, NL, Mole, AJ, Kubinski, S, Claus, P, Murray, LM, Bowerman, M, Gillingwater, TH, Bertoli, S, Parson, SH & Kothary, R 2019, 'Abnormal fatty acid metabolism is a core component of spinal muscular atrophy', *Annals of Clinical and Translational Neurology*, vol. 6, no. 8, pp. 1519-1532. <https://doi.org/10.1002/acn3.50855>

Digital Object Identifier (DOI):

[10.1002/acn3.50855](https://doi.org/10.1002/acn3.50855)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Annals of Clinical and Translational Neurology

Publisher Rights Statement:

This is an open access article under the terms of the Creative Commons AttributionNonCommercialNoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is noncommercial and no modifications or adaptations are made.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.



Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH ARTICLE

Abnormal fatty acid metabolism is a core component of spinal muscular atrophy

Marc-Olivier Deguise^{1,2,3} , Giovanni Baranello^{4,5}, Chiara Mastella⁶, Ariane Beauvais¹, Jean Michaud⁷, Alessandro Leone⁸, Ramona De Amicis⁸, Alberto Battezzati⁸, Christopher Dunham⁹, Kathryn Selby¹⁰, Jodi Warman Chardon^{2,3,11,12,13}, Hugh J. McMillan¹⁴, Yu-Ting Huang^{15,16}, Natalie L. Courtney^{15,16,17}, Alannah J. Mole^{15,16,17}, Sabrina Kubinski^{18,19}, Peter Claus^{18,19}, Lyndsay M. Murray^{15,16,17}, Melissa Bowerman^{20,21,22}, Thomas H. Gillingwater^{15,16}, Simona Bertoli⁸, Simon H. Parson^{15,23} & Rashmi Kothary^{1,2,3,13} 

¹Regenerative Medicine Program, Ottawa Hospital Research Institute, Ottawa, Ontario, Canada

²Department of Cellular and Molecular Medicine, University of Ottawa, Ottawa, Ontario, Canada

³Centre for Neuromuscular Disease, University of Ottawa, Ottawa, Ontario, Canada, K1H 8M5

⁴UO Neurologia dello Sviluppo, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy

⁵The Dubowitz Neuromuscular Centre, NIHR BRC University College London Great Ormond Street Institute of Child Health & Great Ormond Street Hospital, London, United Kingdom

⁶SAPRE-UONPIA, Fondazione IRCCS Cà' Granda Ospedale Maggiore Policlinico, Milan, Italy

⁷Department of Pathology and Laboratory Medicine, Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada

⁸International Center for the Assessment of Nutritional Status (ICANS), Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Milan, Italy

⁹Division of Anatomic Pathology, Children's and Women's Health Centre of B.C., Vancouver, British Columbia, Canada

¹⁰Division of Neurology, Department of Pediatrics, BC Children's Hospital, Vancouver, British Columbia, Canada

¹¹Neuroscience Program, Ottawa Hospital Research Institute, Ottawa, Ontario, Canada

¹²Department of Pediatrics, Children's Hospital of Eastern Ontario, Ottawa, Ontario, Canada

¹³Department of Medicine, University of Ottawa, Ottawa, Ontario, Canada

¹⁴Children's Hospital of Eastern Ontario Research Institute, University of Ottawa, Ottawa, Ontario, Canada

¹⁵Euan MacDonald Centre for Motor Neurone Disease Research, University of Edinburgh, Edinburgh, United Kingdom

¹⁶College of Medicine & Veterinary Medicine, University of Edinburgh, Edinburgh, United Kingdom

¹⁷Centre for Discovery Brain Science, University of Edinburgh, Edinburgh, United Kingdom

¹⁸Institute of Neuroanatomy and Cell Biology, Hannover Medical School, Hannover, Germany

¹⁹Center of Systems Neuroscience, Hannover, Germany

²⁰School of Medicine, Keele University, Staffordshire, United Kingdom

²¹Institute for Science and Technology in Medicine, Stoke-on-Trent, United Kingdom

²²Wolfson Centre for Inherited Neuromuscular Disease, RJA Orthopaedic Hospital, Oswestry, United Kingdom

²³Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom

Correspondence

Rashmi Kothary, Ottawa Hospital Research Institute, 501 Smyth Road, Ottawa, Ontario, Canada K1H 8L6. Tel: (613) 737-8707; Fax: (613) 737-8803; E-mail: rkothary@ohri.ca

Funding Information

RK was supported by Cure SMA/Families of SMA Canada (KOT-1819); Muscular Dystrophy Association (USA) (grant number 575466); Canadian Institutes of Health Research (CIHR) (grant number PJT-156379); and the E-Rare-2 program from the CIHR (grant number ERL-138414). The Italian group (GB, CM, RDA, AB, AL) was funded by the Italian Association of SMA Families (Famiglie SMA, 2015-2016 contribution) and by Fondazione Telethon (Application

Abstract

Objective: Spinal muscular atrophy (SMA) is an inherited neuromuscular disorder leading to paralysis and subsequent death in young children. Initially considered a motor neuron disease, extra-neuronal involvement is increasingly recognized. The primary goal of this study was to investigate alterations in lipid metabolism in SMA patients and mouse models of the disease. **Methods:** We analyzed clinical data collected from a large cohort of pediatric SMA type I–III patients as well as SMA type I liver necropsy data. In parallel, we performed histology, lipid analysis, and transcript profiling in mouse models of SMA. **Results:** We identify an increased susceptibility to developing dyslipidemia in a cohort of 72 SMA patients and liver steatosis in pathological samples. Similarly, fatty acid metabolic abnormalities were present in all SMA mouse models studied. Specifically, *Smn*^{2B/-} mice displayed elevated hepatic triglycerides and dyslipidemia, resembling non-alcoholic fatty liver disease (NAFLD). Interestingly, this phenotype appeared prior to denervation. **Interpretation:** This work highlights metabolic abnormalities as an important feature of SMA, suggesting

GUP15014, 2015, Italy). LMM is supported by grants from Cure SMA (grant number MU1415); Fight SMA; Muscular Dystrophy Association (grant number 417757); Tenovus Scotland (E15/4); and Newlife foundation for disabled children (SG/14-15/08). THG was supported by UK SMA Research Consortium and SMA Europe. SHP was supported by Tenovus (Scotland) and The Euan Macdonald Centre for Research into Motor Neurone Diseases. MB was supported by UK SMA Research Consortium and SMA Angels Charity. The Vanderbilt Mouse Metabolic Phenotyping Center was supported by National Institutes of Health (NIH) grant DK59637. The University of Massachusetts Medical School National Mouse Metabolic Phenotyping Center (MMPC) is supported by NIH grant (2U2C-DK093000). MOD was supported by Frederick Banting and Charles Best CIHR Doctoral Research Award.

Received: 5 July 2019; Accepted: 9 July 2019

***Annals of Clinical and Translational Neurology* 2019; 6(8): 1519–1532**

doi: 10.1002/acn3.50855

Introduction

Spinal muscular atrophy (SMA) is an autosomal recessive disorder characterized primarily by motor neuron death. The incidence of SMA is around 1 in 11,000 live births, and most patients rapidly succumb to their symptoms.¹ More than half of patients with SMA have a severe, infant-onset form of the disease with a median life expectancy of 12 months without supportive treatment. SMA is caused by a mutation or deletion in the ubiquitously expressed *Survival motor neuron 1 (SMN1)* gene,² which produces a protein (SMN) involved in a number of key cellular pathways, including RNA metabolism and splicing, amongst others (reviewed in 3).

SMA has traditionally been considered a motor neuron disease. However, this view has evolved as defects in multiple non-neuronal cell types have been identified.^{4–18} It is unclear whether extra-neuronal components contribute to the clinical picture of SMA (reviewed in 15, 19), but they may become particularly relevant for SMA patients where restoration of SMN protein, largely in the nervous system, has been achieved through therapeutic intervention (e.g. Nusinersen; 20).

Metabolic defects in SMA have been reported previously. Pancreatic defects were observed in mouse models of SMA and in SMA patients.^{11,12,21} These alterations appear to be the cause of abnormal glucose

implementation of nutritional and screening guidelines in patients, as such defects are likely to increase metabolic distress and cardiovascular risk. This study emphasizes the need for a systemic therapeutic approach to ensure maximal benefits for all SMA patients throughout their life.

homeostasis.^{12,21} Furthermore, defects in amino acid metabolism in SMA have been described.²² Lipid metabolism and fatty acid oxidation defects have also been reported in early studies of patients with SMA,^{23,24} where increased esterified carnitine, and reduced β -oxidation capacity are seen.²⁴ There are also three reports of microvesicular steatosis in livers of patients with SMA.^{24–26} The etiology, importance, or generalizability of these findings remain unclear. Most recently, non-neuromuscular phenotypes including metabolic defects were reported prior to their first clinical signs of neuromuscular degeneration in SMA patients.²⁷ As such, standard of care statements have highlighted the need for further research in metabolic status in SMA patients to inform future nutritional guidelines,^{28,29} but strong comprehensive studies are currently still lacking.

The primary goal here is to provide foundational evidence of defects in lipid metabolism in SMA patients and mouse models of SMA. We find an increased propensity of dyslipidemia in SMA patients as well as hepatic fatty deposition. Strikingly, the human findings are reproduced in the *Smn*^{2B/-} mouse model, which develop nonalcoholic fatty liver disease (NAFLD) prior to denervation. Altogether, this work highlights the critical need for investigation of lipid metabolism and the liver in SMA and how this affects the treatment and care of SMA patients in the future.

Methods

Study design

Identification of fatty livers in *Smn*^{2B/-} mice was serendipitous. Upon review of the literature, this sparked a project with the following objectives: Identify whether findings are translated in SMA patients and identify the extent of the fatty acid defect. All objectives were pursued in a simultaneous manner. The data from SMA patients was obtained retrospectively in Italy. Pre-clinical data made use of multiple mouse models of SMA. Serum analysis and lipid quantification were outsourced, and analyses were performed in a blinded fashion. Sample size calculation were not performed for the human data collection as it was retrospective. N number are described in each figure legend. Statistical approach is as described below and in figure captions. Collaboration between laboratories of Kothary and Parson and colleagues occurred midproject, given overlapping results that were converging. Hence, the resulting manuscript offers preclinical data that have been concordant in two independent laboratories.

Patient data

Infant and young SMA patients were recruited from two clinical referral centers for SMA in Italy (UO Neurologia dello Sviluppo, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy and SAPRE-UONPIA, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy). The study protocol was approved by institution ethics review boards (University of Milan #7/16 and Carlo Besta Neurological Institute Foundation #37/2016). All patients with blood work results have genetically confirmed diagnosis of SMA and they were explained benefits and risks of the study, and consented to the study. None of the patients were enrolled in any clinical trials at the time of the blood draw. This study used cut-off values proposed by the National Cholesterol Education Program (NCEP).³⁰ Adult dyslipidemia cut-off values were extracted from the third report of the National Cholesterol Education Program (NCEP) expert panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) and The National Lipid Association recommendations for patient-centered management of dyslipidemia: Part 1 - executive summary.

Liver pathology of human necropsies

Necropsies were obtained at BC Children's Hospital (Protocol #H18-00038). Ten cases of human SMA were retrieved from the pathology files of the British

Columbia's Children Hospital and the Children's Hospital of Eastern Ontario. Given the timeframe of the necropsies (between 1977 and 1996), patients were diagnosed based on clinical presentation and histological features apart from #6 and #7, which were done after 1992 and were genetically confirmed. After review, two cases were not retained: one had associated features of olivo-ponto-cerebellar atrophy; the other because of negative familial genetic studies performed 16 years later and because of the presence of nonmotor changes at histology, in the spinal cord. The autopsies were performed between 1977 and 1996. Slides from the liver were available in all cases. Seven cases were of SMA type I: four females and three males, aged from less than a month to 3 years. One SMA type II case was a female aged 13 years.

Mouse models

The *Smn*^{-/-};SMN2 (Jackson Laboratory), *Smn*^{-/-};SMN2^{+/+};SMNΔ7 and *Smn*^{2B/-} (wild type BL/6J background)³¹ mouse lines were housed at the University of Ottawa Animal Facility and cared for according to the Canadian Council on Animal Care. Experimentation and breeding were performed under protocol OHRI-1948 and OHRI-1927. *Smn*^{+/-} mice were crossed to *Smn*^{2B/2B} mice to obtain *Smn*^{2B/+} and *Smn*^{2B/-} animals. C57BL/6J wild-type mice were bred separately. The Taiwanese *Smn*^{-/-};SMN2 (FVB/N background, FVB.Cg-Smn1^{tm1Hung}Tg(SMN2)2Hung/J from Jackson Laboratory #005058) and *SOD1*^{G93A} mice (B6.Cg-Tg(SOD1*G93A)1Gur/J from Jackson Laboratory #004435) were housed at the Biomedical Sciences Unit, University of Oxford or within Biological Research Resources at the University of Edinburgh. All experiments using mice in the UK were performed in accordance with the licensing procedures authorized by the UK Home Office (Animal Scientific Procedures Act 1986). All tissue for quantitative biochemical analysis were collected at the same time of the day to limit the effect of the circadian rhythm.

Tissue handling and histological analysis

Gross morphology, tissue processing, and staining of animal tissues was as described before.¹³

Lipid quantification and plasma analysis in mice

Tissue lipid analysis for quantification and profiles were performed at the Vanderbilt Mouse Metabolic Phenotyping Center. Lipids were extracted and analyzed as described.^{32,33} Cholesterol and unesterified cholesterol quantification protocol was adapted from 34. Following

decapitation of the mice, blood was collected via capillary using Microcuvette CB 300 K2E coated with K2 EDTA (16.444.100). All the blood collected in this study was sampled ad libitum (i.e. no fasting period) between 9 and 11 AM to limit the effect of the circadian rhythm. Samples were then spun at 2000g for 5 min at room temperature to extract plasma. Lipoproteins analysis was performed at the National Mouse Metabolic Phenotyping Center (MMPC) at the University of Massachusetts Medical School using a Cobas Clinical Chemistry Analyzer (Roche Diagnostics, Indianapolis, IN). Blood glucose was obtained using blood glucose and ketone monitoring system FreeStyle Precision Neo and FreeStyle precision glucose strips.

Gene expression studies

RNA from liver and skeletal muscle was extracted using Qiagen RNeasy Mini kit and reverse transcribed using RT² first strand kit according to manufacturer's protocol. Qiagen microarray fatty acid metabolism (PAMM-007Z) and fatty liver (PAMM-157Z) were used and were analyzed using RT² Profiler PCR Array Data Analysis (<http://saweb2.sabiosciences.com/pcr/arrayanalysis.php>). Automatic selection from Housekeeping group panel was used as a method of normalization.

Statistics

Data are presented as the mean \pm standard error of the mean. A two-sided Student's *t* test was performed using Microsoft Excel or Graphpad Prism 7 to compare the means of data when only two groups were compared (i.e. wild type vs. *Smn*^{2B/-}). One-way ANOVA analysis was used to distinguish differences between more than two groups when multiple comparisons were necessary (i.e. wild type vs. *Smn*^{2B/+} vs. *Smn*^{2B/-}). The post-test used for the ANOVA was Tukey. Significance was set at $P \leq 0.05$ for *, $P \leq 0.01$ for **, $P \leq 0.001$ for ***, and $P \leq 0.0001$ for ****. *N* number for each experiment is as indicated in the figure legends.

Results

SMA patients are at an increased risk of dyslipidemia and fatty liver

We performed lipid profiling on 72 pediatric SMA patients (14 type I, 52 type II, 6 type III – demographics in Table 1). Briefly, the cohort was fairly evenly split between male (54.17%) and female (45.83%). The median age for the whole cohort was 3.8 years, while for males was 3.7 years old and for females was 4 years old. The

median time before their last meal was 5 h. Note that fasting has minimal effect on lipid levels in comparison to nonfasting.³⁵ We focused on total cholesterol (TC), low density lipoprotein (LDL), high density lipoprotein (HDL), non-HDL, and triglycerides to assess abnormalities in fatty acid metabolism in a minimally invasive manner. Over a third (37.5%) of SMA patients, most commonly type I and II, had at least one positive readout out of the five indicative tests for laboratory-defined dyslipidemia (Table 2), in comparison to less than a quarter (20–24%) of the general population in published data sets.³⁶ Furthermore, close to 20% and 13% of SMA patients had more than two or three positive readouts out of the five tests of laboratory-defined dyslipidemia respectively (Table 2). LDL prevalence was doubled in comparison to the general pediatric population.^{36–38} Patients with borderline values made up 61% of SMA patients with at least one indicative lipid result, and 40% would have 3 or more (Table 2). Notably, 3/8 (37.5%) of pediatric SMA liver necropsies revealed steatosis, reminiscent of the proportion of SMA patients showing dyslipidemia (37.5% as well) (Table 3). This is in marked contrast to reported prevalence of NALFD in the pediatric population 2–19 years of age, estimated to be between 2% and 13%.^{39,40} If we limit the age range to 2–4 years, which is more in line with our SMA necropsy cohort, it has been reported that liver steatosis incidence is only 0.7% in the normal pediatric population.⁴⁰

We next assessed whether any associated glucose mis-handling defects may be present, as we had previously identified pancreatic defects.^{11,12} We obtained HbA_{1C} data, a measure of the mean glucose level over the previous 3 months, for 53 of the 72 patients in our cohort. Interestingly, HbA_{1C} trended lower in most SMA patients, with 30 of 53 having an abnormally low readout

Table 1. SMA pediatric patient cohort demographics.

Pediatric cohort	Number	Percentage	Median age (years)	Time before last meal (hours)
Total	72	100	3.8	5
Male	39	54.17	3.7	5
Female	33	45.83	4	5
Type I	14	19.44	3.1	5
Male	5	6.94	2	5
Female	9	12.5	3.2	5
Type II	52	72.22	3.8	5
Male	31	43.05	3.7	5
Female	21	29.17	4.2	5
Type III	6	8.33	6.4	5
Male	3	4.17	6.2	5
Female	3	4.17	6.6	5

Table 2. SMA patients are more susceptible to dyslipidemia than the normal population.

	Criteria	All SMA patients	Type I	Type II	Type III	Normal population*
Abnormal	TC > 200 mg/dL	10/72 (13.89%)	1/14 (7.14%)	9/52 (17.31%)	0/6 (0%)	7.7–10.7% ^{36–38}
	LDL > 130 mg/dL	9/72 (12.5%)	1/14 (7.14%)	7/52 (13.46%)	1/6 (16.67%)	3.2–7.2% ^{36–38}
	HDL < 40 mg/dL	12/72 (16.67%)	1/14 (7.14%)	10/52 (19.23%)	1/6 (16.67%)	4.1–19.3% ^{36,38,43,57}
	TG > 100 mg/dL‡	15/72 (20.83%)	5/14 (35.71%)	7/52 (13.46%)	3/6 (50%)	13.2–22.1% ^{36,38,57}
	Non HDL-cholesterol > 145 mg/dL	10/72 (13.89%)	1/14 (7.14%)	8/52 (15.38%)	1/6 (16.67%)	8.4% ⁴³
	1/5 abnormal dyslipidemia reading	27/72 (37.5%)	6/14 (42.85%)	18/52 (34.62%)	3/6 (50%)	20.2–22.9% ^{36,43}
	2/5 < abnormal dyslipidemia reading	14/72 (19.44%)	2/14 (14.29%)	11/52 (21.15%)	1/6 (16.67%)	5.37% ^{†36}
	3/5 < abnormal dyslipidemia reading	10/72 (13.89%)	1/14 (7.14%)	8/52 (15.38%)	1/6 (16.67%)	–
	HbA1C < 5	30/53 (56.60%)	5/8 (62.5%)	23/41 (56.09%)	2/4 (50%)	–
Borderline	TC > 170 mg/dL	30/72 (41.67%)	5/14 (35.71%)	23/52 (44.23%)	2/6 (33.33%)	–
	LDL > 110 mg/dL	21/72 (29.17%)	2/14 (14.29%)	18/52 (34.62%)	1/6 (16.67%)	–
	HDL < 45 mg/dL	20/72 (27.78%)	5/14 (35.71%)	13/52 (25%)	2/6 (33.33%)	–
	TG > 75 mg/dL§	23/72 (31.94%)	7/14 (50%)	13/52 (25%)	3/6 (50%)	–
	Non HDL-cholesterol > 120 mg/dL	32/72 (44.44%)	6/14 (42.86%)	23/52 (44.23%)	3/6 (50%)	–
	1/5 < borderline dyslipidemia reading	44/72 (61.1%)	11/14 (78.57%)	30/52 (57.69%)	3/6 (50%)	–
	2/5 < borderline dyslipidemia reading	35/72 (48.61%)	6/14 (42.86%)	26/52 (50%)	3/6 (50%)	–
	3/5 < borderline dyslipidemia reading	29/72 (40.28%)	6/14 (42.86%)	20/52 (38.46%)	3/6 (50%)	–

*Note that these values were taken from multiple studies and criteria may have varied and not be identical to the present study.

†Calculated from results in the particular study – see reference.

‡High is defined as >100 for 0–9 years and >130 for 10–19 years of age.

§Borderline is defined as >75 for 0–9 years and >90 for 10–19 years of age.

Table 3. Presence of steatosis in SMA liver necropsies.

Case	Sex	Age	Type	Cause death	Steatosis*
1	F	14 mo	I	Bronchopneumonia	-/**
2	F	7 mo	I	Bronchopneumonia	-
3	M	8 mo	I	Aspiration pneumonia	+++
4	M	<1 mo	I	Aspiration pneumonia	-
5	F	9 mo	I	Respiratory insufficiency	+/**
6	F	14 mo	I	Bronchopneumonia	-
7	M	12 mo	I	Respiratory insufficiency + HIE + Chronic pneumonitis	+/**
8	F	13 y	II	Undetermined	-

HIE, Hypoxic ischemic encephalopathy.

*: -: no steatosis; +: mild panlobular; ++ and +++: moderate and severe, panlobular. In all cases with steatosis (cases 3, 5, and 7), it was of the microvesicular type, predominantly and the periportal regions were more involved than the mid or central regions of the hepatic lobules.

** : Presence of moderate increase of glycogen.

(HbA_{1C} < 5%, normal 5%–6.5%). In fact, most SMA patients had an HbA_{1C} around 5% as the calculated median was 4.94% and calculated mean 4.93% in our cohort (Table 2). Overall, a large subset of SMA patients show clinical test results consistent with considerable metabolic abnormalities, especially dyslipidemia and fatty liver.

Abnormal fatty acid metabolism in SMA mouse models

To investigate whether fatty acid metabolism defects identified above may be related to SMN depletion, we assessed livers of the *Smn*^{2B/-} mouse model at pre-symptomatic

age (postnatal day 4 (P4)) and at symptomatic age (P17–19). The livers from P17–19 *Smn*^{2B/-} mice were paler and displayed microvesicular steatosis (Fig. 1A–B, E–G). The level of triglycerides, the main storage form of fatty acids, was 25-fold higher in livers of P19 *Smn*^{2B/-} mice compared to controls (Fig. 1H). Triglyceride chain length alterations, especially of long chain fatty acids, were noted in P19 *Smn*^{2B/-} mice compared to controls (Fig. 1I–J). Phospholipid, free fatty acid, diglycerides, cholesterol esters, unesterified cholesterol and total cholesterol (Fig. 1K–P) in livers of P19 *Smn*^{2B/-} mice showed alterations, indicating a global misregulation in fatty acid metabolism. By comparison, histology (Fig. 1C and D),




Figure 1. *Smn*^{2B/-} mice have fat accumulation in the liver. Gross morphology (0.75X) and histology (H&E – 40X, Oil Red O – 400X) of *Smn*^{2B/-} livers showing fatty inclusions at P17–19 (A and B, E–G) but not P4 (C and D). Lipid profiling identified elevation of triglycerides at P19 in *Smn*^{2B/-} livers (H), with altered chain length (I–J). Other lipid classes, such as phospholipid, free fatty acids, diglycerides, cholesterol esters, unesterified cholesterol, and total cholesterol, were also misregulated in P19 *Smn*^{2B/-} livers (K–P). P4 lipid levels were unchanged from control (H, K–P). Scale bar: (A–D) 50 μm, (E and F) 10 μm. (N value for each experiment is as follows: N = 5–6 for A–D, 3–5 for E–G, 4 for H–P, unpaired two-sided student’s t-test, *P ≤ 0.05, **P ≤ 0.01, and ***P ≤ 0.001).

and lipid levels (Fig. 1K–P) were unchanged in P4 *Smn*^{2B/-} livers.

Triglyceride levels were determined in three additional mouse models of SMA. Fat accumulation was confirmed in livers from symptomatic P9 *Smn*^{Δ7} mice (*Smn*^{-/-};

SMN2^{+/+};*Smn*^{Δ7}+/+) (Fig. 2). Conversely, more severe mouse models of SMA, such as the “Taiwanese” mice (P9) and the *Smn*^{-/-};*SMN2* mice (P5), showed reduced lipid accumulation in the liver compared to control littermates (Fig. 2A–B), likely due to the reduced life span of




Figure 2. Hepatic triglyceride misregulation is a common feature in different SMA models at symptomatic. Quantification of hepatic triglycerides showed a fivefold reduction in P5 *Smm*^{2/-}; *SMN2* mice (A), a twofold reduction in P9 Taiwanese mice (B), and a threefold increase in P10 *Smm*^{Δ7} mice (C) in comparison to control littermates. Analysis of hepatic triglyceride levels for each SMA mouse model in A–C involved a comparison to their own control (*N* value for each experiment is as follows: *N* = 4–6 for A and B, 9–10 for C, 4–9 for D, unpaired two-sided student's *t*-test, **P* ≤ 0.05 and ***P* ≤ 0.01).

these animals preventing the opportunity for these pathologies to develop (Fig. 2D). A fatty acid pathway-focused PCR array in liver of symptomatic *Smm*^{2B/-} and “Taiwanese” mice, revealed several overlapping alterations, suggesting similar pathogenic etiologies (Fig. 3). Significantly, lipid metabolism defects are present across multiple mouse models of SMA. The severity, stage of disease progression, and genetic background of the mouse models may influence the extent of lipid accumulation in the liver and the pathological presentation.

To better understand the mechanisms underlying the NAFLD in *Smm*^{2B/-} mice, we next examined the progression of fat accumulation in the liver. Interestingly, the liver became progressively fatty starting at P9 (Fig. 4A–E). This raised the possibility that muscle denervation, which may start around this time in *Smm*^{2B/-} mice, could be sufficient to induce liver steatosis. However, *SOD1*^{G93A} mice, a well-established model of amyotrophic lateral sclerosis (ALS) that shows widespread denervation, showed no hepatosteatosis at symptomatic age (20 weeks) (Fig. 4F–I). Nevertheless, we cannot discount the possibility that denervation may yet contribute to a lower metabolic demand by skeletal muscle or through alterations of metabolic pathways.

***Smm*^{2B/-} mice also display dyslipidemia and abnormal fatty acid metabolism in skeletal muscle**

We next assessed whether *Smm*^{2B/-} mice displayed dyslipidemia and fatty acid metabolism abnormalities in other

tissues. Analysis of plasma lipoproteins from P19 *Smm*^{2B/-} mice revealed a significant increase in TC, very low density lipoprotein (VLDL - derived from triglycerides) and LDL, while HDL levels were reduced (Fig. 5A–D). Consequently, the ratios of TC/HDL, a measure of increased cardiovascular risk⁴¹ were significantly elevated (Fig. 5E). In addition, glucose always trended lower and its level plummeted after P7 in *Smm*^{2B/-} mice (Fig. 5F). The clear dyslipidemic profile and low glucose levels align well with our clinical findings in SMA patients. We then determined if the altered lipoprotein in the plasma translated to an increased delivery of lipids to motor neurons and skeletal muscle, two primary targets in SMA pathogenesis. No changes were observed in the spinal cord and *tibialis anterior* muscle of P19 *Smm*^{2B/-} mice (Fig. 5G and H). Given that we had previously identified fatty infiltrates in the skeletal muscle of *Smm*^{2B/-} mice,⁴² we further investigated whether some alterations in lipid metabolism pathways are present despite no change in triglyceride content. A focused fatty acid PCR array showed that a number of genes were misregulated in muscles of these mice (Fig. 5I). These data suggest that fatty acid metabolism is dysregulated in different tissues in SMA mice.

Discussion

Studies from three decades ago had identified potential fatty acid defects with dicarboxylic aciduria, and reduced muscle fatty oxidation, with three subsequent reports of hepatic fatty vacuolization in SMA patients.^{23–26} Despite these initial findings and the recognition for studies in




Figure 3. Commonalities identified in expression of fatty acid metabolism genes between *Taiwanese* and *Smn^{2B/-}* mice. Volcano plot presentation of all changes (1.5X, $P < 0.05$) in a focused fatty acid metabolism PCR array in *Smn^{2B/-}* mice (A) and *Taiwanese* mice (C) identify general downregulation. Changes more than two-fold are represented for *Smn^{2B/-}* (B) and *Taiwanese* (D). Analysis of commonalities between *Smn^{2B/-}* and *Taiwanese* are represented by Venn diagrams, which identify nine genes with similar changes (E), listed in (F). ($N = 4$, for *Smn^{2B/-}* mice, and $N = 3$ for *Taiwanese* mice).

metabolic defects in SMA by consensus statements in the care of SMA patients,^{28,29} minimal research has been done to fill this gap. Indeed, systematic research on fatty acid metabolism in SMA has remained largely unexplored and translational foundational data are lacking. We show that patients with SMA are more prone to develop dyslipidemia as well as liver steatosis than the general pediatric population,^{36–38,40,43} with a large subset showing abnormalities when screened with a common lipid and cholesterol panel and in SMA liver necropsies. In preclinical models, fatty acid abnormalities were identified in four commonly used SMA mouse models. With a focus on the *Smn*^{2B/-} mice, we observe the development of a NAFLD phenotype that does not appear to be caused by the denervation.

In this study, we provide strong evidence for the susceptibility to dyslipidemia and low glycated hemoglobin levels in a large cohort of SMA patients. To our knowledge, this is the first time that such abnormalities are reported in SMA. Current statistics on dyslipidemia in one laboratory-defined measure in otherwise healthy children is estimated to be roughly 20%.^{36,43} Our studies showed that 37% of SMA patients have dyslipidemia. More strikingly, 14% of patients had more than 3 laboratory-defined measures of dyslipidemia, for which prevalence data is sparse in the normal pediatric population but is suspected to be quite low in the absence of familial dyslipidemia. Pathological examination of SMA liver necropsies showed a similar proportion (37.5%) displaying liver steatosis. This is much higher than the prevalence of liver steatosis in 2–4 year old normal children (0.7%).⁴⁰ The proportion of fatty liver in our SMA samples was closer to fatty liver prevalence reported in older obese pediatric population, where prevalence can range from 28–77%.³⁹ Additionally, low HbA_{1C}, observed in the majority of our cohort, has been associated with increased all-cause mortality⁴⁴ and liver disease⁴⁵ in the general population. Given the significant size of our cohort, we suggest that dyslipidemia is an important feature of SMA. Early metabolic studies on SMA patients had focused on urinary organic acids, muscle β -oxidation enzyme function, and plasma acylcarnitine and free fatty acid profiling.^{23,24,26} These tests are rarely used, are not widely available, and their interpretation requires a specialist's advice, hence making them poor choices in the screening and identification of SMA patients with potential metabolic abnormalities. Our study provides a widely accessible manner to identify and monitor fatty acid metabolic abnormalities in SMA patients that can be acted upon with current cholesterol-lowering therapy if needed. While necropsies were used for the identification of fatty liver in this study, this can be easily determined through ultrasonography, which consists of a widely used and

non-invasive imaging modality.⁴⁶ In the upcoming years, screening and preventive treatment of dyslipidemia and fatty liver could be particularly important to limit significant co-morbidities, such as cardiovascular and cerebrovascular disease, in the newly aging demographics of treated SMA patients.

While dyslipidemia and liver steatosis were present in a subset of SMA patients, these phenotypes were present in all *Smn*^{2B/-} mice studied. This is perhaps not surprising as the human population is much more heterogenous than congenic mouse colonies. Nevertheless, the consistency of these findings in human and a preclinical model make it clear that SMN depletion predisposes the organism to fatty acid defects, dyslipidemia, and NAFLD. It is likely that other gene polymorphisms may be required for (or even protect against) the onset and severity of these phenotypes. Similarly, investigation of hepatic fatty acid accumulation in four different mouse models of SMA all revealed abnormalities, following an interesting pattern of presentation. It appears that the severity of disease model and related period of survival critically modulates how the alterations in fatty acid metabolism present, reminiscent of our findings in skeletal muscle.^{5,42} We identified that the most severe mouse models (*Smn*^{-/-}; *SMN2*, and “Taiwanese” mice) show low lipid content in the liver while less severe and slightly longer lived SMA models (*Smn*^{-/-}; *SMN2*^{+/+}; *Smn* Δ 7^{+/+} and *Smn*^{2B/-} mice) display liver steatosis. Even though liver pathology presentation in the “Taiwanese” and *Smn*^{2B/-} mice differs, we identified nine genes in a fatty liver PCR array with similar patterns of misregulation. Their expression profile as a whole were in line with a compensatory mechanism to limit further triglyceride accumulation. This suggests that some of the molecular etiologies are likely the same, but sufficient time (i.e. longer survival) and perhaps some genetic modifiers may be required to develop the fat accumulation.

A common question in non-neuronal findings in SMA is whether it is a consequence of denervation. In other diseases where denervation is present, such as in spinal cord injury (SCI)⁴⁷ and spinal and bulbar muscular atrophy (SBMA),⁴⁸ some reports identify an increased risk for dyslipidemia. However, SCI and SBMA patients are generally older adults and dyslipidemia is well-known to be more prominent with age. Thus, generalizing these results to the SMA population is not obvious. In addition, these reports^{47,48} do not account for prevalence of patients with multiple measures of dyslipidemia, such as in our study. A major limitation of our study was the absence of a denervated patient control group. However, previous studies had shown that fatty acid alterations in SMA patients could not be explained by denervation alone, as denervated control patients did not display these defects.²³




Figure 4. Fat accumulation is first observed between P9 and P11 in *Smn*^{2B/-} mice and denervation is not sufficient to trigger hepatic steatosis. (A and B) Triglycerides and cholesterol esters quantification in livers from *Smn*^{2B/-} mice at different ages. (C–E) Oil Red O staining (400X) additionally showed increased fat at P9. (F–H) H&E staining (40X) of livers of 20-week-old *SOD1*^{G93A} mutant mice, a model of ALS, did not show hepatic fat accumulation in comparison to livers from *Smn*^{2B/-} mice, even though denervation is well-established at this time point. (I) Triglycerides and cholesterol esters quantification showed no difference between mutant *SOD1*^{G93A} and WT controls. Scale bar represents 50 μ m in C and D (10 μ m in the inset), and in F–H. (N value for each experiment is as follows: *N* = 4–6 for A and B, 3 for C–E, and 3–5 for I, unpaired two-sided student's *t*-test, **P* \leq 0.05, ***P* \leq 0.01, ****P* \leq 0.001).

Nevertheless, to overcome this limitation, we turned to preclinical mouse models. We found that denervation, such as in the *SOD1*^{G93A} mouse model of ALS, did not lead to development of liver steatosis. Previous literature in ALS described lipid redistribution rather than accumulation in *SOD1*^{G93A} mice, consistent with our findings.⁴⁹ It should be noted that this model is also associated with increased lipid clearance in the periphery, which could abrogate fatty accumulation brought on by denervation.⁵⁰ Interestingly, we found that the rate of triglyceride accumulation in the liver increases very rapidly with progression of disease in the mouse, which could reflect the overall denervation status of the animal. Indeed, it is possible that denervation plays some role in induction of this phenotype by changing the metabolic demand of the muscle, modulating the molecular metabolism upon denervation, or through other unknown factors.

In simple terms, we believe that NAFLD develops through the imbalance of input (diet, peripheral lipolysis, de novo lipogenesis) and output (export of lipids or usage through β -oxidation). We propose that this phenotype in SMA could be caused by potential perturbation in the pancreas-liver axis (as exemplified by the low blood glucose), potential mitochondrial defects, liver-intrinsic defects, or the interplay of the three. Indeed, it appears that hyperglucagonemia, a characteristic found in *Smn*^{2B/-} mice¹² likely in response to low blood sugar, can lead to peripheral lipolysis of white adipose tissue and increase in circulating lipids.⁵¹ Mitochondrial defects have previously been identified in other cell types as well as in human muscle biopsies^{52–54} and could likely contribute to reduce β -oxidation in hepatocytes. Additionally, the unknown splicing deficiency in the hepatocytes could predispose the cells to fat accumulation. Another possibility, in the context of low glucose, includes severe starvation, which also been linked to NAFLD development.⁵⁵ Nevertheless, the NAFLD phenotype observed in *Smn*^{2B/-} mice occurs prior to overt motor dysfunction that would thus not prevent the mice from obtaining appropriate nutrition. Moreover, in the case of SMA patients, they are generally followed by nutritionists to ensure appropriate caloric intake. Further molecular studies should attempt to dissect the etiologies and identify nutritional or therapeutic strategies to abrogate this co-morbidity in susceptible patients.

The extent of fatty accumulation identified in *Smn*^{2B/-} mice is likely to result in functional consequences. The liver is the metabolic factory of the body, producing plasmatic proteins, processing toxins as well as medications, and regulating glucose, lipid, and amino acid homeostasis. Going forward, the functionality of the liver in SMA will be a particularly important consideration in formulation of new therapeutic for SMA. Indeed, pro-drugs metabolized through the liver may not gain optimal levels while those cleared/processed by the liver might harbor more significant hepatotoxicity. In fact, historically there has been only cursory examination of liver function in SMA. Embryonic lethality and iron overload are the main features of liver restricted *Smn* conditional knockout in mice.⁵⁶ More recently, increased erythropoiesis, megakaryocyte and platelet production, together with mild iron storage abnormalities, were identified in the severe “Taiwanese” mouse model of SMA.¹⁰

Altogether, our clinical studies in SMA patients as well as in preclinical mouse models, provide strong evidence of defects in fatty acid metabolism. The greater predisposition to develop dyslipidemia and fatty liver in SMA patients as well as the identification of NAFLD in *Smn*^{2B/-} mice emphasize that defects in metabolism can lead to added co-morbidities, especially in the new therapeutic era of SMA, where lifespan is extended. Indeed, this work further highlights the importance of establishing currently lacking nutritional guidelines, performing early screening for metabolic defects in treated SMA patients, as well as developing systemic therapeutic strategies that incorporate non-neuronal organs to ensure overall optimal management of SMA.

Acknowledgments

We thank Jocelyn Coté and his student Andr anne Didillon for providing *Smn*^{-/-}; *SMN2*; *SMN17* mice to the Kothary laboratory. We extend our gratitude to Eva Szunyogova, Sabrina Gibeault, My Tran Trung, and Rebecca Yaworski for assistance with experiments. RK was supported by Cure SMA/Families of SMA Canada (KOT-1819); Muscular Dystrophy Association (USA) (grant number 575466); Canadian Institutes of Health Research (CIHR) (grant number PJT-156379); and the E-Rare-2 program from the CIHR (grant




Figure 5. *Smn*^{2B/-} mice display dyslipidemia and abnormal fatty acid metabolism in skeletal muscle, but not in spinal cord. (A–D) Significant upregulation of total cholesterol, VLDL and LDL in the plasma of *Smn*^{2B/-} animals, while HDL levels were significantly lower. (E) Parameters for cardiovascular risks such as TC/HDL were significantly increased for *Smn*^{2B/-} mice. (F) Glucose trend lower early and plummet later in life in *Smn*^{2B/-} mice. (G,H) Every lipid class in the P19 *Smn*^{2B/-} skeletal muscle or spinal cord were at similar levels to WT. (I) Many genes involved in fatty acid metabolism were altered in P19 *Smn*^{2B/-} skeletal muscle in a focused fatty acid PCR array. (N value for each experiment is as follows: N = 10 for A–F, 5 for G, 4 for H–I, unpaired two-sided student’s *t*-test, ***P* ≤ 0.01, ****P* ≤ 0.001 and *****P* ≤ 0.0001).

number ERL-138414). The Italian group (GB, CM, RDA, AB, AL) was funded by the Italian Association of SMA Families (Famiglie SMA, 2015–2016 contribution) and by Fondazione Telethon (Application GUP15014, 2015, Italy). LMM is supported by grants from Cure SMA (grant number MU1415); Fight SMA; Muscular Dystrophy Association (grant number 417757); Tenovus Scotland (E15/4); and Newlife foundation for disabled children (SG/14-15/08). THG was supported by UK SMA Research Consortium and SMA Europe. SHP was supported by Tenovus (Scotland) and The Euan Macdonald Centre for Research into Motor Neurone Diseases. MB was supported by UK SMA Research Consortium and SMA Angels Charity. The Vanderbilt Mouse Metabolic Phenotyping Center was supported by National Institutes of Health (NIH) grant DK59637. The University of Massachusetts Medical School National Mouse Metabolic Phenotyping Center (MMPC) is supported by NIH grant (2U2C-DK093000). MOD was supported by Frederick Banting and Charles Best CIHR Doctoral Research Award.

Author Contribution

MOD, MB, THG, SHP, and RK contributed to the conception and design of the study; MOD, AB, GB, AL, RDA, Alberto B, JWC, HJM, YTH, NLC, AJM, SK, PC, LMM, MB, and SB contributed to the acquisition and analysis of data; CM participated in the enrollment of patients; JM, CD, and KS contributed to the retrieval of pathological specimens and analysis; MOD and RK contributed to the drafting of the text and preparing the figures. GB, HJM, PC, LMM, MB, THG, SB, and SHP read and edited the manuscript.

Conflict of Interest

Marc-Olivier Deguise received honoraria and travel accommodations by Biogen for the SMA Summit 2018

held in Montreal, Canada. Rashmi Kothary and the Ottawa Hospital Research Institute have a licensing agreement with Biogen for the *Smn*^{2B/-} mouse model. All other authors have no competing interests to declare.

References

- Sugarman EA, Nagan N, Zhu H, et al. Pan-ethnic carrier screening and prenatal diagnosis for spinal muscular atrophy: clinical laboratory analysis of >72,400 specimens. *Eur J Hum Genet* 2012;20:27–32.
- Lefebvre S, Burglen L, Reboullet S, et al. Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* 1995;80:155–165.
- Tisdale S, Pellizzoni L. RNA-processing dysfunction in spinal muscular atrophy. In: Sumner CJ, Paushkin S, C. Ko CP, eds. *Spinal muscular atrophy: disease mechanisms and therapy*. pp. 113–131. London: Academic Press, 2017.
- Shafey D, Cote PD, Kothary R. Hypomorphic *Smn* knockdown C2C12 myoblasts reveal intrinsic defects in myoblast fusion and myotube morphology. *Exp Cell Res* 2005;311:49–61.
- Boyer JG, Deguise MO, Murray LM, et al. Myogenic program dysregulation is contributory to disease pathogenesis in spinal muscular atrophy. *Hum Mol Genet* 2014;23:4249–4259.
- Bricceno KV, Martinez T, Leikina E, et al. Survival motor neuron protein deficiency impairs myotube formation by altering myogenic gene expression and focal adhesion dynamics. *Hum Mol Genet* 2014;23:4745–4757.
- Somers E, Lees RD, Hoban K, et al. Vascular defects and spinal cord hypoxia in spinal muscular atrophy. *Ann Neurol* 2016;79:217–230.
- Thomson AK, Somers E, Powis RA, et al. Survival of motor neurone protein is required for normal postnatal development of the spleen. *J Anat* 2017;230:337–346.
- Maxwell GK, Szunyogova E, Shorrock HK, et al. Developmental and degenerative cardiac defects in the Taiwanese mouse model of severe spinal muscular atrophy. *J Anat* 2018;232:965–978.
- Szunyogova E, Zhou H, Maxwell GK, et al. Survival Motor Neuron (SMN) protein is required for normal mouse liver development. *Sci Rep* 2016;6:34635.
- Bowerman M, Michalski JP, Beauvais A, et al. Defects in pancreatic development and glucose metabolism in SMN-depleted mice independent of canonical spinal muscular atrophy neuromuscular pathology. *Hum Mol Genet* 2014;23:3432–3444.
- Bowerman M, Swoboda KJ, Michalski JP, et al. Glucose metabolism and pancreatic defects in spinal muscular atrophy. *Ann Neurol* 2012;72:256–268.
- Deguise MO, De Repentigny Y, McFall E, et al. Immune dysregulation may contribute to disease pathogenesis in spinal muscular atrophy mice. *Hum Mol Genet* 2017;26:801–819.
- Deguise MO, Kothary R. New insights into SMA pathogenesis: immune dysfunction and neuroinflammation. *Ann Clin Transl Neurol* 2017;4:522–530.
- Deguise MO, Patitucci TN, Ebert AD, et al. Contributions of different cell types to Spinal Muscular Atrophy pathogenesis. In: Sumner CJ, Paushkin S, Ko CP, eds. *Spinal muscular atrophy: disease mechanisms and therapy*. pp. 167–181. London: Academic Press, 2017.
- Shababi M, Habibi J, Yang HT, et al. Cardiac defects contribute to the pathology of spinal muscular atrophy models. *Hum Mol Genet* 2010;19:4059–4071.
- Hunter G, Aghamaleky Sarvestany A, Roche SL, et al. SMN-dependent intrinsic defects in Schwann cells in mouse models of spinal muscular atrophy. *Hum Mol Genet* 2014;23:2235–2250.
- McGivern JV, Patitucci TN, Nord JA, et al. Spinal muscular atrophy astrocytes exhibit abnormal calcium regulation and reduced growth factor production. *Glia* 2013;61:1418–1428.
- Hamilton G, Gillingwater TH. Spinal muscular atrophy: going beyond the motor neuron. *Trends Mol Med* 2013;19:40–50.
- Groen EJM, Talbot K, Gillingwater TH. Advances in therapy for spinal muscular atrophy: promises and challenges. *Nat Rev Neurol* 2018;14:214–224.
- Davis RH, Miller EA, Zhang RZ, Swoboda KJ. Responses to fasting and glucose loading in a cohort of well children with spinal muscular atrophy type II. *J Pediatr* 2015;167:1362–1368.e1.
- Walter LM, Deguise MO, Meijboom KE, et al. Interventions targeting glucocorticoid-Kruppel-like factor 15-branched-chain amino acid signaling improve disease phenotypes in spinal muscular atrophy mice. *EBioMedicine* 2018;31:226–242.
- Crawford TO, Sladky JT, Hurko O, et al. Abnormal fatty acid metabolism in childhood spinal muscular atrophy. *Ann Neurol* 1999;45:337–343.
- Tein I, Sloane AE, Donner EJ, et al. Fatty acid oxidation abnormalities in childhood-onset spinal muscular atrophy: primary or secondary defect(s)? *Pediatr Neurol* 1995;12:21–30.
- Zolkipli Z, Sherlock M, Biggar WD, et al. Abnormal fatty acid metabolism in spinal muscular atrophy may predispose to perioperative risks. *Eur J Paediatr Neurol* 2012;16:549–553.
- Kelley RI, Sladky JT. Dicarboxylic aciduria in an infant with spinal muscular atrophy. *Ann Neurol* 1986;20:734–736.
- Lipnick SL, Agniel DM, Aggarwal R, et al. Systemic nature of spinal muscular atrophy revealed by studying insurance claims. *PLoS One* 2019;14:e0213680.

28. Finkel RS, Sejersen T, Mercuri E; Group ESWS. 218th ENMC International Workshop: revisiting the consensus on standards of care in SMA Naarden, The Netherlands, 19–21 February 2016. *Neuromuscul Disord* 2017;27:596–605.
29. Wang CH, Finkel RS, Bertini ES, et al. Consensus statement for standard of care in spinal muscular atrophy. *J Child Neurol* 2007;22:1027–1049.
30. NCEP. National Cholesterol Education Program (NCEP): highlights of the report of the Expert Panel on Blood Cholesterol Levels in Children and Adolescents. *Pediatrics* 1992;89:495–501.
31. Eshraghi M, McFall E, Gibeault S, Kothary R. Effect of genetic background on the phenotype of the *Smn2B*^{-/-} mouse model of spinal muscular atrophy. *Hum Mol Genet* 2016;25:4494–4506.
32. Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 1957;226:497–509.
33. Morrison WR, Smith LM. Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride-methanol. *J Lipid Res* 1964;5:600–608.
34. Rudel LL, Kelley K, Sawyer JK, et al. Dietary monounsaturated fatty acids promote aortic atherosclerosis in LDL receptor-null, human ApoB100-overexpressing transgenic mice. *Arterioscler Thromb Vasc Biol* 1998;18:1818–1827.
35. Langsted A, Nordestgaard BG. Nonfasting versus fasting lipid profile for cardiovascular risk prediction. *Pathology* 2019;51:131–141.
36. Li J, Motsko SP, Goehring EL Jr, et al. Prevalence of pediatric dyslipidemia: comparison of a population-based claims database to national surveys. *Pharmacoepidemiol Drug Saf* 2010;19:1031–1040.
37. Ford ES, Li C, Zhao G, Mokdad AH. Concentrations of low-density lipoprotein cholesterol and total cholesterol among children and adolescents in the United States. *Circulation* 2009;119:1108–1115.
38. Dathan-Stumpf A, Vogel M, Hiemisch A, et al. Pediatric reference data of serum lipids and prevalence of dyslipidemia: results from a population-based cohort in Germany. *Clin Biochem* 2016;49:740–749.
39. Pacifico L, Poggiogalle E, Cantisani V, et al. Pediatric nonalcoholic fatty liver disease: a clinical and laboratory challenge. *World J Hepatol* 2010;2:275–288.
40. Schwimmer JB, Deutsch R, Kahen T, et al. Prevalence of fatty liver in children and adolescents. *Pediatrics* 2006;118:1388–1393.
41. Millan J, Pinto X, Munoz A, et al. Lipoprotein ratios: physiological significance and clinical usefulness in cardiovascular prevention. *Vasc Health Risk Manag* 2009;5:757–765.
42. Deguise MO, Boyer JG, McFall ER, et al. Differential induction of muscle atrophy pathways in two mouse models of spinal muscular atrophy. *Sci Rep* 2016;6:28846.
43. Kit BK, Kuklina E, Carroll MD, et al. Prevalence of and trends in dyslipidemia and blood pressure among US children and adolescents, 1999–2012. *JAMA Pediatr* 2015;169:272–279.
44. Carson AP, Fox CS, McGuire DK, et al. Low hemoglobin A1c and risk of all-cause mortality among US adults without diabetes. *Circ Cardiovasc Qual Outcomes* 2010;3:661–667.
45. Christman AL, Lazo M, Clark JM, Selvin E. Low glycated hemoglobin and liver disease in the U.S. population. *Diabetes Care* 2011;34:2548–2550.
46. Zeng MD, Fan JG, Lu LG, et al. Guidelines for the diagnosis and treatment of nonalcoholic fatty liver diseases. *J Dig Dis* 2008;9:108–112.
47. Koyuncu E, Nakipoglu Yuzev GF, Yenigun D, Ozgirgin N. The analysis of serum lipid levels in patients with spinal cord injury. *J Spinal Cord Med* 2017;40:567–572.
48. Francini-Pesenti F, Querin G, Martini C, et al. Prevalence of metabolic syndrome and non-alcoholic fatty liver disease in a cohort of Italian patients with spinal-bulbar muscular atrophy. *Acta Myol* 2018;37:204–209.
49. Finkelstein A, Kunis G, Seksenyan A, et al. Abnormal changes in NKT cells, the IGF-1 axis, and liver pathology in an animal model of ALS. *PLoS One* 2011;6:e22374.
50. Fergani A, Oudart H, Gonzalez De Aguilar JL, et al. Increased peripheral lipid clearance in an animal model of amyotrophic lateral sclerosis. *J Lipid Res* 2007;48:1571–1580.
51. Campbell JE, Drucker DJ. Islet alpha cells and glucagon-critical regulators of energy homeostasis. *Nat Rev Endocrinol* 2015;11:329–338.
52. Acsadi G, Lee I, Li X, et al. Mitochondrial dysfunction in a neural cell model of spinal muscular atrophy. *J Neurosci Res* 2009;87:2748–2756.
53. Berger A, Mayr JA, Meierhofer D, et al. Severe depletion of mitochondrial DNA in spinal muscular atrophy. *Acta Neuropathol* 2003;105:245–251.
54. Ripolone M, Ronchi D, Violano R, et al. Impaired muscle mitochondrial biogenesis and myogenesis in spinal muscular atrophy. *JAMA Neurol* 2015;72:666–675.
55. Kneeman JM, Misdraji J, Corey KE. Secondary causes of nonalcoholic fatty liver disease. *Therap Adv Gastroenterol* 2012;5:199–207.
56. Vitte JM, Davoult B, Roblot N, et al. Deletion of murine *Smn* exon 7 directed to liver leads to severe defect of liver development associated with iron overload. *Am J Pathol* 2004;165:1731–1741.
57. Johnson WD, Kroon JJ, Greenway FL, et al. Prevalence of risk factors for metabolic syndrome in adolescents: National Health and Nutrition Examination Survey (NHANES), 2001–2006. *Arch Pediatr Adolesc Med* 2009;163:371–377.