

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from The Water Center

Water Center, The

2019

Irrigation Water Quality—A Contemporary Perspective

Arindam Malakar

University of Nebraska - Lincoln, amalakar2@unl.edu

Daniel D. Snow

University of Nebraska at Lincoln, dsnow1@unl.edu

Chittaranjan Ray

University of Nebraska-Lincoln, cray@nebraska.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/watercenterpubs>



Part of the [Environmental Indicators and Impact Assessment Commons](#), [Fresh Water Studies Commons](#), [Hydraulic Engineering Commons](#), [Hydrology Commons](#), [Sustainability Commons](#), and the [Water Resource Management Commons](#)

Malakar, Arindam; Snow, Daniel D.; and Ray, Chittaranjan, "Irrigation Water Quality—A Contemporary Perspective" (2019). *Faculty Publications from The Water Center*. 44.

<https://digitalcommons.unl.edu/watercenterpubs/44>

This Article is brought to you for free and open access by the Water Center, The at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from The Water Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Review

Irrigation Water Quality—A Contemporary Perspective

Arindam Malakar ¹, Daniel D. Snow ^{2,*} and Chittaranjan Ray ³

¹ Nebraska Water Center, part of the Robert B. Daugherty Water for Food Global Institute, 109 Water Sciences Laboratory, University of Nebraska, Lincoln, NE 68583-0844, USA

² School of Natural Resources and Nebraska Water Center, part of the Robert B. Daugherty Water for Food Global Institute, 202 Water Sciences Laboratory, University of Nebraska, Lincoln, NE 68583-0844, USA

³ Nebraska Water Center, part of the Robert B. Daugherty Water for Food Global Institute 2021 Transformation Drive, University of Nebraska, Lincoln, NE 68588-6204, USA

* Correspondence: dsnow1@unl.edu; Tel.: +01-402-472-7539

Received: 29 May 2019; Accepted: 13 July 2019; Published: 17 July 2019



Abstract: In the race to enhance agricultural productivity, irrigation will become more dependent on poorly characterized and virtually unmonitored sources of water. Increased use of irrigation water has led to impaired water and soil quality in many areas. Historically, soil salinization and reduced crop productivity have been the primary focus of irrigation water quality. Recently, there is increasing evidence for the occurrence of geogenic contaminants in water. The appearance of trace elements and an increase in the use of wastewater has highlighted the vulnerability and complexities of the composition of irrigation water and its role in ensuring proper crop growth, and long-term food quality. Analytical capabilities of measuring vanishingly small concentrations of biologically-active organic contaminants, including steroid hormones, plasticizers, pharmaceuticals, and personal care products, in a variety of irrigation water sources provide the means to evaluate uptake and occurrence in crops but do not resolve questions related to food safety or human health effects. Natural and synthetic nanoparticles are now known to occur in many water sources, potentially altering plant growth and food standard. The rapidly changing quality of irrigation water urgently needs closer attention to understand and predict long-term effects on soils and food crops in an increasingly fresh-water stressed world.

Keywords: crop uptake; food quality; geogenic; emerging contaminants; nanomaterials

1. Introduction

Irrigation is the controlled use of multiple water sources in a timely manner for increased or sustained crop production. Irrigation comprises of the water that is applied by an irrigation system during the growing season and also includes water applied during field preparation, pre-irrigation, weed control, harvesting, and for leaching salts from the root zone [1]. In 2015 it was estimated that in the United States irrigation alone accounted for 62% of water usage [1]. Globally, irrigation is the highest consumptive use of freshwater [2]. As the world's population grows, the risk increases that more people will be deprived of adequate food supplies in impoverished areas, particularly those subject to water scarcity [3]. Agricultural production of food needs to increase by an estimated 60% by 2050 to ensure global food security [3] and irrigation will increasingly be called upon to help meet this demand. In the race to enhance agricultural productivity, irrigation will become even more dependent on substandard sources of water. Therefore, it is of utmost importance to access our current state of knowledge and explore the effects of irrigation water quality on crops. This understanding will help ensure adequate crop production to meet increased demand as well as to maintain proper food and soil quality.

Groundwater exploitation (withdrawal for irrigation) can release naturally occurring geogenic contaminants, such as arsenic, from the solid phase to groundwater, while wastewater reuse can

concentrate pesticides, pharmaceuticals and other emerging contaminants in irrigation water [4,5]. Use of untreated wastewater is becoming prevalent in developing countries where around 80–90% of wastewater remains untreated [6]. Polluted municipal, industrial or agricultural water used for irrigation significantly changes soil quality, increases the amount of trace elements in soil and plants, and acts as a source of various pathogens which affects food quality and safety [7,8]. Water of inadequate quality is a potential source of both direct and indirect contamination to food crops [9], and leads to increased contamination of soil and water [10,11]. In addition, the presence of synthetic and natural nanomaterials is beginning to be identified in crops [12–14]. In locations where excess irrigation is practiced, contaminants in soils are leached to the vadose zone, where they can contribute to geogenic contaminant mobilization and potentially increase contaminant levels in local groundwater [15]. Many aspects of water composition, such as hardness and iron content, also affect the suitability of a water source for newer, more efficient spray or drip irrigation techniques. Runoff, return flow, and leaching of irrigation water also contribute to local surface and groundwater contamination [16]. Increased usage of irrigation water has already led to impaired irrigation water and soil quality. Considering the presence of new contaminant types in different water sources (see Figure 1), it is essential to evaluate the impact of these contaminants within the context of modern agriculture. To date, very little research and regulatory attention has been paid to contaminants in irrigation water. Contamination of irrigation water supplies is likely to worsen unless additional efforts (research, guidelines, regulations, treatment methods) are brought to bear on this problem.




Figure 1. Main sources of irrigation water and different types of contaminants present in those sources impacting food, soil, and water quality. Note that surface water and wastewater are subject to similar types of contamination.

This review article looks at previous approaches to define irrigation water quality and compares to a current perspective with respect to impacts on human health. It evaluates the long-term effect and influence of the changing quality of water sources used in agricultural production. Although the article discusses traditional irrigation water quality concerns, such as salinization, it mainly emphasizes contemporary water quality issues like new or emerging contaminants, pathogens, geogenic trace elements and engineered nanomaterials. These contaminants are now widespread in various conventional and unconventional water sources used for modern-day irrigation. The article is organized as a short summary of conventional measures of irrigation water quality followed by a more detailed evaluation of the impact of contemporary irrigation water quality issues on soil and crop quality. Contemporary topics include emerging contaminants with separate sections on pharmaceuticals, antibiotics, steroid hormones, pesticides, cyanotoxins and mycotoxins, biological contaminants bacteria, virus and antibiotic resistance genes, modern inorganic contaminants, such as geogenic trace elements and nanomaterials. The review is summarized by considering the changing quality of water sources used for irrigation, and the need for additional work and improved regulation of irrigation water, especially for food production. The primary focus of this review is to recognize

water quality issues that have a direct or indirect influence on surface soil contamination, crop uptake of contaminants and their potential to impact human health. This article shows a need for modern guidelines, regulations and research to understand the complex nature of irrigation water. Though it is a critically important topic from a human health standpoint, this review does not include an exhaustive discussion of contamination of irrigation water by human pathogens. While wastewater treatment technologies are constantly evolving and can address some of the issues presented here, a review of wastewater quality as a function of treatment technology is beyond the scope of this article. Moreover, treatment approaches are likely to be tailored to sources, and irrigation water sources are highly varied depending on climate, population, industry, crop, and livestock density.

2. Conventional Measures of Irrigation Water Quality

The effect of irrigation water composition on soil properties for crop production has been a focus for the past half century. Previous studies of water quality issues, and the suitability of freshwater sources for irrigation, have primarily been directed toward an understanding of potential problems to soil salinity, fertility and crop growth. For example, early work by the United States Geological Survey (USGS) [17] evaluated groundwater quality in Texas for irrigation and other potentially competing uses. A subsequent report by Schwennesen and Forbes characterized groundwater in San Simon Valley, Arizona and New Mexico, for domestic use and irrigation [18]. Clark reported on the chemical composition of groundwater in the Morgan Hill area of California [19], while Scofield and Headly [20] evaluated water composition with respect to irrigation potential. Most of these early works focused on understanding the impact of water quality on long-term viability of irrigation in arid regions of the United States.

Globally, irrigation water quality was described in Tanzania, Africa, with respect to pH and alkaline and alkaline-earth elements [21]. Taylor et al. reported that irrigation water pH was one of the main factors for wheat growth in Punjab, India [22]. A subsequent work reiterated that alkaline elements such as sodium play a crucial role in continued use of water for irrigation of cropland and quantified the maximum amount that may be tolerated [23]. The effects of soil salinization and trace element composition on crop growth have become more apparent over time. Eaton et al. reported that boron present in water around Hollister, California affected the growth of apricots and prunes [24]. In the subsequent years, the United States Department of Agriculture (USDA) conducted further studies and reported that sodium, boron and electrical conductivity are the best general measures for judging the suitability of water for irrigation [25]. From these studies, it was evident that continuous irrigation with water of marginal quality impacted soil and also affected crop growth [26–31]. In 1967, the American Society for Testing Materials (ASTM) developed a quantitative assessment of irrigation water quality, including new formulas for maximum permissible quantity of chloride and electrical conductivity based on infiltration rate, evapotranspiration rate, irrigation frequency and duration [32]. Traditionally, discussion on irrigation water quality has mainly focused on its effect on soil quality, and how soil quality was predicted to affect crop growth and yield. Color, turbidity, total dissolved solids (TDS), pH, specific conductance, odor and foam characterized the quality of water. Colorless, odorless, foamless water with minimum turbidity, TDS below 1000 mg L⁻¹ at circumneutral pH and specific conductance below 1.5 mmhos/m is generally considered to be of good quality for irrigation purposes [33,34]. A higher TDS is not recommended for most crops as it can impact the salinity of soil and pore water will become highly concentrated when taken up by roots via osmosis. Excessive dissolved solids content, or salinity of irrigation water, has historically been the primary characteristic determining water suitability for irrigation. Salt accumulation in the crop root zone impedes water uptake and can eventually prevent plant growth altogether [34]. Excess salinity from sodium can affect soil structure and water infiltration. The proportion of sodium to calcium and magnesium is the primary factor controlling the hydraulic conductivity of water in soil [33–35]. Sodium is generally expressed as a sodium absorption ratio (SAR) [9]. Long-term irrigation of soils with elevated sodium concentrations relative to calcium and magnesium, bicarbonate, carbonate, and TDS will be limiting soil aggregate formation, which reduces infiltration and makes less water available to crops [34].

Seiler et al., under the National Irrigation Water Quality Program (NIWQP) of the U.S. Department of the Interior (DOI), studied the effect of irrigation-induced contamination of water, soil and biota in the western United States. NIWQP data from the 26 areas under study suggested that degradation of groundwater quality due to irrigation is a common occurrence [11,36]. The study indicated that selenium was the most common contaminant, followed by arsenic, uranium and molybdenum [11,37]. This study also suggested regular co-occurrence of these contaminants. For example, selenium was found to be elevated with uranium, and these contaminants were accumulating in the soils and affecting long-term suitability for crop production. This was one of the first reports to correlate trace element contamination in water sources used for irrigation to soil quality. These findings led to the appreciation of the intricate complexities of irrigation water quality and its role in ensuring proper crop growth and long-term food quality. These studies mainly focused on the impact of water quality on crop productivity and soil quality, while effects to food quality and safety were just beginning to be recognized.

3. Impact of Contemporary Irrigation Water Quality Issues on Soil and Crop Quality

Irrigation water quality has mainly been characterized with respect to effects on crop growth and yield, though an emerging and pressing issue relates to plant uptake and soil enrichment with inorganic and organic contaminants (Figure 2). These “new” issues with respect to irrigation water quality can lead to food quality and safety concerns, as well as affect crop growth and yield [38–41]. Wastewater reuse for irrigation contributes to increasing incidence of organic microcontaminants [42], such as pharmaceuticals and other synthetic organics in soils and crops. Increasing reliance on groundwater also contributes to the probability for elevated concentrations of natural geogenic contaminants such as arsenic and selenium in irrigation water and soils. Understanding the occurrence and fate of these new contaminants in irrigation water sources is paramount in limiting the effects to modern agricultural products [43]. Long term impacts to soil and crop quality (see Figure 2) need to be understood.




Figure 2. A conceptual model of the impact of inadequate quality of irrigation water sources on soil and crop quality.

3.1. Emerging Contaminants: Organic Pollutants

3.1.1. Pharmaceuticals

Traces of pharmaceuticals and personal care products have been identified in a variety of freshwater sources, including drinking water [44], groundwater [45], and surface water [45]. Pharmaceuticals can enter the water system from various sources, including direct disposal and human excretion into sewers leading to elevated concentrations of pharmaceuticals in wastewater [46]. Pharmaceuticals often detected in sewage sludge include non-steroidal anti-inflammatory drugs (NSAIDs), blood thinners, psychiatric drugs, antidiuretics and β -blockers [47–49]. Plant uptake of a wide variety of pharmaceutical groups like NSAIDs, antihistamine, β -blockers, calcium channel blockers, antiepileptics, steroid hormones, antidepressants, antineoplastic agents, anti-itch compounds, x-ray contrast agents, lipid-lowering agents, benzodiazepines, tranquilizers and veterinary drugs from soil and contaminated water has been observed and studied [50–52]. Wu et al. reported that a primary pathway for contamination by pharmaceuticals in food crops is through irrigation water [53,54]. For example, a recent study found traces of carbamazepine, caffeine, lamotrigine, gabapentin and acesulfame in a variety of vegetables grown with treated wastewater in Jordan [55]. Treated wastewater is well known to contain a large variety of pharmaceuticals and personal care products, many of which are known to accumulate in food crops [56,57]. The occurrence of these and other synthetic organic chemicals is likely to increase in water supplies, especially in areas with water scarcity, and irrigation with contaminated water will lead to soil contamination and plant uptake.

3.1.2. Antibiotics

Environmental contamination by antibiotic residues in food production systems is a growing problem worldwide, and the potential implications to proliferation of antibiotic resistance have been the subject of multiple reviews and opinion articles [58–60]. The occurrence of persistent antibiotic residues in various water sources [61,62] is well documented which not only includes municipal [63,64], agricultural [65,66] and hospital sewage [67,68] but also groundwater [69] and surface water [70–72]. The concentration range of antibiotics is generally measured at ng L^{-1} to a few $\mu\text{g L}^{-1}$ in many water sources [73], though concentrated wastewater can have much higher levels [65]. Recent studies have demonstrated that plants can take up antibiotics (like amoxicillin, ketoconazole, lincomycin, oxytetracycline, sulfamethoxazole, sulfonamides, and tetracyclines) [74–76] and antibiotic contaminated irrigation water can play a significant role in the uptake [77]. The environmental fate and transport of antibiotics depend on various physical properties such as water solubility, lipophilicity, volatility and sorption potential [77].

The implications to human health due to the presence of antibiotics in food crops is not clear, but other potential adverse impacts include allergic reactions, disruption of digestive function and chronic toxic effects as a result of prolonged low-level exposure [78–81]. One of the major concerns for the increasing prevalence of antibiotics in the environment is the development and spread of antibiotic-resistant gene and bacteria [82], which is discussed in Section 3.2.2. Clearly, the absence of antibiotic residues in irrigation water cannot be assumed.

3.1.3. Steroids

Land application of livestock manure can contribute to accumulation of steroid hormones [83] and veterinary pharmaceuticals. Very low concentrations of natural steroid hormones, such as estrone, 17α -estradiol, 17β -estradiol, estriol, testosterone, androstenedione and progesterone that occur in animal waste and wastewater have been documented as accumulating in soil [84]. Traces of steroid hormones have also been reported in groundwater [85] and surface water [86], and often in treated municipal wastewater [87–91]. Wastewater treatment plants are known to discharge these hormones into river water and also other recipients [92]. Laboratory experiments have suggested that traces of steroid hormones and pharmaceuticals can be taken up in crops [93] and recent studies of food

crops irrigated with treated municipal water have confirmed this can occur in the field. Further work is needed to understand the significance and impact of these chemicals in the environment and to human health [94], though at present, the reported concentrations are relatively low in comparison to other contaminants.

3.1.4. Agrochemicals

Regular use of pesticides in irrigated crops is also likely to lead to the occurrence of these residues in irrigation water and food crops, especially in regions where regulation and training in proper application of these substances is lacking. Application of large quantities of agrichemicals and improper management can create a substantial effect on the environment. The leaching and runoff of agrochemicals is a potential source of groundwater and surface water contamination [95,96]. The occurrence of agrichemical residues in vegetables has been documented [97] and their uptake by crops is well studied [98,99] and regulated. Leaching of nitrate from fertilizer over application to groundwater below is well reported, and accumulation of reactive nitrogen is also thought to initiate mobilization of other geogenic contaminants [15].

3.1.5. Cyanotoxins and Mycotoxins

Cyanotoxins, which comprise a large range of naturally produced organic compounds, are produced and released by cyanobacteria when they are present in large quantities (blooms) and especially when these organisms die off and decay in surface water. Cyanobacteria also referred to as blue-green algae, naturally occur in all freshwater ecosystems [100]. Warmer temperatures coupled with high nutrient concentrations are thought to favor conditions for algae blooms to form in surface water. Of many different groups of cyanotoxins, hepatotoxic cyclic peptides collectively known as microcystins are the most commonly studied cyanotoxins, which cause a wide range of symptoms in humans [101]. Other studies have also shown that cyanotoxins, which include hepatotoxic microcystins and neurotoxic compounds such as anatoxin-a and beta-N-methylamino-alanine, can make their way to human and animal food chain from contaminated reservoirs [102–105]. A recent review has summarized the extent of the literature investigating the fate in soils, and agricultural crops [103]. It seems quite clear that toxins can accumulate in plants, including food crops and under some conditions can also inhibit plant growth [102,103,106–108]. Though there are many gaps, and only a handful of studies have investigated this route for exposure. There is evidence for human health effects through consumption of plants contaminated with cyanotoxins by irrigation using surface water sources impacted by cyanotoxins.

Mycotoxins are naturally occurring fungal toxins (chemicals), which can cause a variety of adverse health effects to both humans and livestock. A few mycotoxins are known or suspected carcinogens. Fungi do pose potential hazards to human health. However, there were relatively few studies of mycotoxins in water sources until recently. Fungal contamination has been observed in drinking water [109] and recently it was reported that untreated surface water can be breeding place for these fungi, generating mycotoxins [110]. Kolpin et al. led a broad scale study on the occurrence of mycotoxins across streams in the United States (US) [111]. Their study concluded that the ecotoxicological effects from long-term, low-level exposures to mycotoxins are poorly understood and would require further investigation. Mycotoxin uptake in rice has been studied [112] and these chemicals have been reported to be present in various food grains [113,114]. The prevalence of mycotoxins in surface water makes it an important consideration regarding modern water quality of irrigation as mycotoxin health hazards are widely reported.

3.2. Biological Contaminants: Bacteria, Virus and Antibiotic Resistance

3.2.1. Pathogens

Often because of its nutrient content and accessibility, untreated wastewater from municipal and domestic sources containing excessive levels of pathogens is often directly used for irrigation in developing countries [6]. Untreated wastewater generally carries a high pathogen load compared to other irrigation water sources. Risks from pathogen (bacteria, viruses, or protozoan or larger organisms) contamination to irrigation water quality will continue to be a topic of primary concern [115–117], and it is impossible to adequately address this topic in a few paragraphs. Pathogenic microorganisms in irrigation water likely pose the greatest acute risk to human health and will continue to be a concern especially in freshly-eaten produce. Pathogens are biological organisms that may influence modern-day irrigation water quality. Pathogen contamination is generally related to surface water sources, but groundwater may also be under threat if it is recharged with wastewater sources [116]. The complexity of reproducibly measuring microbiological contamination of irrigation water has made monitoring difficult. Several different types of pathogens have been detected in diverse irrigation water sources including bacteria (e.g., *Salmonella* and *Escherichia coli*), protozoa (e.g., *Cryptosporidium* and *Giardia*), as well as viruses (e.g., noroviruses) [118,119]. Irrigation of food crops with surface water clearly has the highest potential for contaminating freshly eaten produce, and this topic has had the greatest research and regulatory effort in recent years.

There have been quite a few comprehensive reviews emphasizing irrigation water as a source of pathogenic microorganisms in fresh produce [120–123]. Between 1973 and 2012, the Centers for Disease Control and Prevention reported 606 leafy-vegetable associated pathogenic outbreaks (norovirus (55% of outbreaks), Shiga toxin-producing *Escherichia coli* (STEC) (18%), and *Salmonella* (11%)), with 20,003 associated illness and 19 deaths [124]. From 2013 to 2017, the number of outbreaks (mainly from norovirus (32%) STEC (23%), and *Salmonella* (32%)) associated with leafy greens and vegetables decreased to 21, with 699 illness and five deaths [125]. In 2018, 272 infections were reported from two outbreaks (*E. Coli*) associated with romaine lettuce resulting in five deaths [126,127], and another multi-state outbreak was linked to parasite *Cyclospora*, which reported 511 cases of infection [128]. However, a 2014 risk-based review conducted in California suggests that recycled water quality criteria, along with proper agricultural management practices do not lead to increased public health risk [129]. In the US, the Center for Produce Safety has published information on the factors that affect the microbiological safety of agricultural water [130]. The Foodborne Disease Outbreak Surveillance System (FDOSS) has an online tool, the National Outbreak Report System (NORS), which keeps track of outbreaks in the United States. Reports of pathogen contamination from inadequately treated wastewater have also been documented in developing countries [8]. The occurrence of pathogens in water used to irrigate food crops is considered a severe problem affecting human health both in both developing and even in developed countries [131]. Groundwater sources are generally considered less vulnerable to contamination by pathogenic microorganisms, while surface water and wastewater have a much higher potential for contamination. Farmers utilizing surface water for food crops, which are consumed raw should follow proper mitigation strategies to control contamination [132]. The method utilized for irrigation has a substantive role in pathogenic contamination of crops. For example, subsurface drip may have the lowest risk as the water is generally applied at the root zone, unlike other methods (e.g., sprinkler irrigation) where the edible portions of crops can come in contact with contaminated water [133]. New and more intensive monitoring approaches and potential disinfection and treatment techniques for surface water used to irrigate food crops are needed to improve food safety [8].

3.2.2. Antibiotic Resistance

The World Health Organization (WHO) has listed antibiotic resistance among today's biggest threats for global health, food safety, and development, as this threatens the ability to treat common infectious diseases [134]. Antibiotic resistome is defined as the sum of all genes directly or indirectly

contributing to antibiotic resistance both in the clinics and the environment [135]. Aquatic ecosystems are regarded as a primary reservoir of antibiotic-resistant bacteria (ARB) [136]. The presence of ARB and their resistance determinants in surface water sources have been well documented and is generally linked to nearby wastewater treatment plant effluent [137–141]. Wastewater treatment plants enrich ARB and their resistance determinants as it favors exchange of antibiotic resistance genes (ARG) among bacteria and selection of resistant strains [142]. In a recent study, it was found that multidrug-resistant (MDR) bacteria were found to be more prevalent in surface waters than in treated wastewater [143].

Irrigation water is one of the major sources for contamination of fresh produce with antibiotic resistance bacteria [58,144,145]. Similar to pathogens, the incidence of ARB contamination is higher when using overhead sprinklers as water can directly come in contact with fresh produce. When fresh produce is consumed raw it can act as an ideal vector for exposure. The diversity of ARB present in fresh produce is significant and can have a severe impact on human health.

3.3. Inorganic Contaminants: Geogenic Source and Nanomaterials

3.3.1. Geogenic Contaminants in Irrigation Water

Selected naturally occurring geogenic contaminants, such as boron, arsenic and selenium, have been the subject of much previous work focused on irrigation water quality [146,147], especially in areas with extensive use of groundwater. With the exception of boron, trace element contaminants were not studied with respect to soil quality, crop productivity and phytotoxicity [9]. Boron is an essential trace element for plant growth, but elevated boron concentrations (>1 mg/L) in irrigation water can cause stunted growth and reduced productivity in sensitive crops such as wheat. The occurrence of geogenic contaminants is a growing contemporary issue because of the potential impact on food quality and human health. Concentrations of geogenic contaminants have likely been increasing over time in a variety of irrigation water sources, often due to increasing agricultural intensification [148,149]. Researchers have reported that groundwater in China has elevated levels of arsenic; this water is used for irrigating feed crops [150]. Similarly, studies in the United States have reported higher levels of uranium [15] and arsenic [151] in its aquifers. India, Bangladesh and Vietnam have widespread arsenic contamination in groundwater used for drinking and irrigation, especially in areas where the use of contaminated water has led to contaminated soils and crops [152–154]. Presently there are no federal guidelines regulating geogenic contaminant levels in irrigation water except in the case of direct wastewater reuse [155,156]. The increasing levels of contaminants in soils and irrigation water are a growing issue across the globe, and there is little work to date regarding strategies to mitigate accumulation in plants and food crops. Contaminant uptake by crops has been well studied and plant uptake has even been used as a remediation method, viz. phytoremediation. However, mitigation strategies for prevention of geogenic contaminant uptake by plants have received scant attention in the literature. Plant uptake and accumulation of specific trace elements may not affect plant growth, but accumulation and consumption may pose hazards to animal or humans.

Table 1 summarizes the estimated ranges of aqueous concentrations of many geogenic elements with respect to guidelines and recommendations of water use. Irrigation water guidelines compiled by the Food and Agriculture Organization (FAO) in 1976 are generally based on the toxic effects to crops and plant growth [157]. FAO has recommended using these values as guidelines for irrigation utilizing groundwater and surface water sources, but not for irrigation with wastewater containing measurable levels of trace elements [157]. Wastewater guidelines set by FAO (Table 1) are equivalent to irrigation water quality recommendations in 1976, though may contradict newer guidelines for special constituents of wastewater [157]. Early recommendations rarely consider uptake of trace elements by crops and the impact on food quality, which may affect human health. Recommended concentrations to ensure consistent food quality are absent, as even low concentrations of geogenic contaminants can impact food quality (Table 1) [158–172]. Continuous use of irrigation water with low concentrations of geogenic contaminants concentrations can result in soil enrichment and affect food crop quality [158–172].

Table 1. Common geogenic trace element contaminants in irrigation water with current regulatory and recommended levels (NA: Not Available, AL: Action level, CCC: Criterion continuous concentration, CMC: Criterion maximum concentration, SDWR: Secondary drinking water regulations). (* Few examples of plant uptake, ** in μg , *** greenhouse experiment concentration was equivalent to natural condition).

Trace Element	EPA Drinking Water Guideline ($\mu\text{g L}^{-1}$) [173]	Regulatory Limits for Wastewater ($\mu\text{g L}^{-1}$) [156,174,175]	Freshwater CCC, CMC Limits for Aquatic Life ($\mu\text{g L}^{-1}$) [176]	Recommended Maximum Concentrations of Trace Elements in Irrigation Waters ($\mu\text{g L}^{-1}$) [157]	Reported Ranges for Trace Elements in Glacier Aquifer System of USA (Includes Wells Used for Agriculture) ($\mu\text{g L}^{-1}$) [177]	Reported Ranges of Water Known to Impact Food and Soil Quality * ($\mu\text{g L}^{-1}$)
Arsenic	10	100	340, 150	100	0.09–340	58.7 [158], 0.2–164 [159]
Boron	NA	NA	NA	1000	NA	3351–16,000 [165–167]
Cadmium	5	10	1.8, 0.72	10	0.018–1	<1–3200 [168]
Cobalt	NA	50	NA	50	0.007–95	0.21–0.81 [169]
Copper	1300 (AL)	200	NA	200	0.126–127	10–133 ** [170]
Chromium (III) or (VI)	100 (Cr)	100 (Cr)	570, 74 (III) 16, 11 (VI)	100 (Cr)	0.4–22 (Cr)	1–46 (VI) [171] ≤ 250 *** [172]
Iron	300 (SDWR)	5000	NA, 1000	5000	3–38,100	-
Lead	15 (AL)	5000	65, 2.5	5000	0.04–9.0	≤ 140 [160]
Lithium	NA	2500	NA	2500	0.040–126	-
Manganese	500 (SDWR)	200	NA	200	0.056–28,200	≤ 100 [160]
Nickel	-	200	470, 52	200	0.035–56	≤ 50 [160]
Selenium	50	20	NA	20	0.173–223	0.12–341 [161]
Silver	100 (SDWR)	NA	3.2, NA	NA	NA	-
Uranium	30	NA	NA	NA	0.009–162	1–1200 [162]
Zinc	5000 (SDWR)	2000	120, 120	2000	0.536–1000	~ 130 [168]

It is likely that irrigation with elevated levels of geogenic contaminants leads to contaminant enrichment in crops. For example, Bundschuh et al. [178] compiled data from different regions of South America known for high occurrences of arsenic in groundwater and surface water. Their study indicated that arsenic concentrations in edible plants and crops were associated with the elevated arsenic concentrations in soil and irrigation waters. The study showed that regions with high arsenic concentrations in surface water and groundwater relate directly to accumulation in plants, fish, livestock meat, milk, and milk products. Their study indicates that there is a need for more rigorous studies in evaluating pathways of arsenic exposure through the food chain in Latin America and other regions.

Factors such as arsenic speciation, type and composition of soil, and plant species also plays a significant role in crop uptake [179]. An interesting aspect of arsenic transformation is where arsenic present in soil may occur as oxidized arsenate (As(V)) but may become reduced to arsenite (As(III)) after uptake in crops [180]. Arsenite is regarded as 25–60 times more toxic to humans than arsenate [181]. Arsenic, by far, is the most studied geogenic contaminant in crops, especially rice, and is included in many review articles detailing its impact of food quality and human health [182–186]. Ongoing research is focused on managing arsenic uptake by crops [154,187,188]. Other geogenic contaminants in groundwater, such as uranium, are less well studied with respect to food contamination.

Elevated levels of selenium may be toxic, though selenium is an essential micronutrient for crops. Selenium is known to accumulate in crops grown on soils with high selenium content [161,189], and selenium enrichment has been reported in soils throughout the United States [190]. In their recent study, Wang et al. [191] found that volatile organic compounds released by plant growth-promoting rhizobacteria increases both selenium and iron uptake. Uranium is another geogenic, potentially toxic contaminant [192] and studies in nutrient culture show its uptake by crops [193,194]. Several studies confirmed that irrigation water contaminated with uranium has an impact on crop quality though less to soil contamination by uranium [195–197]. Lead, mercury [38], chromium [198] and cadmium [199] are also known to accumulate in crops grown on soils with high levels of these contaminants. While there is extensive information in the literature on uptake by plants, most studies have focused on the general type of contaminant, or its accumulation in crops, use in phytoremediation of soils and the pathway of uptake within the crop. Contaminated irrigation water is capable of enriching surface soil with these contaminants [200] and likely enhances availability for uptake by crops [197]. There are few federal guidelines for trace element limits in foods in the United States, other than in arsenic, lead, cadmium, and mercury in wastewater [201], and the evidence suggests there is an urgent need for irrigation water quality standards that include geogenic contaminants.

3.3.2. Engineered Nanomaterials

Earth is rich with natural nanomaterials and it is estimated that thousands of megatons move through the hydrosphere annually [14]. Natural nanoparticles in water can easily pass through conventional membrane filter pore sizes of 0.2- μm and may not be accounted for as nanoparticles [202], which adds to the complexity in understanding their impact on crop health. In the past decade, production and use of engineered nanoparticles have also risen significantly and continues to trend upwards. Nanoparticles are used in a wide variety of contemporary products, ranging from electronics and cosmetics to processed food. This proliferation in use of nanoparticles has paved their way to increasing occurrence in water sources [203,204]. Nanotechnology is also used widely for water treatment for both groundwater and surface water sources [205,206], but its repercussions are still not well understood [207]. In 2018, the occurrence of nanoparticle size plastics (nanoplastics) were critically reviewed with respect to human health and growing global occurrence in freshwater [208,209]. Analytical methods capable of detecting and quantifying nanoparticles in complex aqueous matrices are lacking, increasing the challenge in tracking fate and transport of these particles [207]. In 2005, Oberdörster et al. [210] reviewed the interaction of nanomaterials in regards to human health. Still, the full-scale toxicity of natural or engineered nanomaterials is not well understood in the context of complex biosystems [14]. The size of nanomaterials (100 nm or less in size) is the main factor [14]

which makes studying nanoparticle contamination challenging. Therefore, the paradigm of risk assessment of nanoparticles needs to be reevaluated with respect to the unique challenges involved in monitoring environmental pathways and assessing impacts on human health.

Broadly, engineered nanoparticles can be classified into carbon and metal-based nanoparticles. Nanoparticles derived from carbon (e.g., carbon nanotubes) simultaneously act like particles and high molecular weight organic compounds. Metal oxides or metal nanoparticles must be evaluated for their metal-related chemotoxicity, and also toxicity arising due to their particulate form. These dual behaviors again add to the complexity of studying and assessing risk. Nanotoxicity can elicit significant effects to human health. Nanoparticles can be carcinogenic and may produce reactive species inside the body [211]. Ganguly et al. have recently reviewed the toxicity of nanomaterials, emphasizing major exposure pathways [211].

The nanoparticle life cycle is poorly understood. Therefore, the Environmental Protection Agency (EPA) has allowed limited manufacturing of new materials by administrative orders or new use rules under the Toxic Substances Control Act. These new rules have significantly expanded since their inception in 2010, which is on par with our increase in understanding. However, the use of and occurrence of nanomaterials in irrigation water is not monitored or regulated [206,212].

Nanotechnology usage is vital in modern civilization and will have a substantial impact on the world economy [213]. It is projected that the nanomaterials market will reach 55 billion US dollars by year 2022 [213]. Production of engineered nanoparticles is expected to rise in the coming years, making these contaminants more likely to occur in different water sources, both conventional (surface and/or groundwater) and in treated wastewater. Both carbon-based and metal-based nanoparticles are of concern and it has been reported these particles are persistent in water [214,215]. Nanomaterial transformation in fresh water systems is an active field of research [216]. Nanoparticle assessments have concluded that fine particulate matter occurs in a variety of water sources [217–221], so its impact on irrigation water, accumulation in soil, and potential for crop uptake is of paramount significance.

The growing availability of engineered nanomaterials/particles is relevant to understand how these nanosized particles may impact water and food quality in agriculture. The use of nanomaterials in agriculture is also increasing, and little data is available to understand how occurrence in irrigation water may influence crops [222,223]. There is recent work on uptake of both carbon and metal-based nanoparticles by plants. For example, multi-walled carbon nanotubes have been observed in broccoli, resulting in a positive impact on plant growth [224] but potentially creating health concerns. Carbon nanotubes are also known to act as a carrier for other contaminants (like organochlorine pesticide, etc.) in plants and enhance contaminant translocation [225]. These contrasting effects of carbon nanotubes have been well summarized in a recent review article by Vithanage et al. [226].

Metal oxide nanoparticles, which form a bulk of engineered nanomaterials, have been well studied under the purview of plant uptake [227]. Metal oxides like titanium dioxide, silver oxide, iron oxide, copper oxide and metal nanoparticles have been shown to accumulate in a variety of food crops and have even been detected in commercial produce [228–232]. A recent review by Ma et al. describes studies of nanoparticle uptake by crops and their occurrence in the final produce [40]. Similar to carbon-based nanoparticles, metal or metal oxide-based nanoparticles are also known to be beneficial to plant growth and have been marketed as nanofertilizers [233]. For example, iron oxide nanoparticles have shown to be a potential iron source for peanut crops [234]. Nanoparticles are also known to induce oxidative stress in crops [235]. A recent study by Liu et al. suggested that combinations of nanomaterials might have different impacts on the soil microcosm compared to a single nanomaterial [236]. In actual conditions, mixed nanomaterials will be more prevalent in soil and irrigation water. In addition to engineered nanomaterial, natural nanomaterials of silver are found in groundwater [237]. Life cycle assessment of engineered titanium dioxide nanoparticles showed that their impact is not just limited to crop uptake, but exhibit marine aquatic ecotoxicity and human toxicity [238,239]. Nanoscale is an important factor in ensuring uptake of fertilizer [240,241] but the long-term effects of nanoagrochemicals have yet to be

studied [242]. While nanoagrochemicals may be beneficial to crops [243], they may have undesired effects on the environment and on human health [242].

There are many questions and few answers for understanding the effects of nanosized contaminants in irrigated agriculture. Future studies should be focused on understanding retention times and fate in water, plants and soils, including degradation and transformation rates, and biological effects of different forms. Naturally-formed nanoparticles can occur in irrigation water and in food crops and it is clear this route for exposure needs to be better understood and monitored. Are nanoparticles easily broken down in the environment or are they stable, do they form aggregates, and what is the accumulation in both soil and water sources? If they accumulate, how do they impact soil health and modern irrigation techniques? Nanoparticle occurrence and behavior in irrigation water sources, soils and plants is clearly an emerging area of research.

4. Changing Quality of Water Sources Used for Irrigation

This discussion on the changing composition and quality of different water sources for irrigation—be it conventional sources like surface- or groundwater, or nonconventional source like reclaimed water—signifies an urgent need for increased efforts to monitor the quality of agricultural water [244]. Recently, there have been various efforts to address changing water quality [9] by incorporating measurement of new contaminant types (e.g., arsenic, fluoride) [10,245,246] in water quality guidelines. Still, it is challenging to address the complexity of irrigation water quality covering different forms of geogenic and emerging contaminants. In the coming decades, the concentrations of specific contaminants will likely increase and continue to affect the quality of water sources for irrigation. Impending climate change may make the situation more extreme as drought and water scarcity may concentrate contaminants in water. The increasing prevalence of cyanotoxins may increase due to more intensive agriculture, fertilization and water use. Although there are strict guidelines for contaminant concentrations in drinking water, few guidelines exist for the use of water for any irrigation, including irrigation of food crops. The substantial expense of monitoring, including sampling and laboratory analysis, for “new” recently identified contaminants is a reason to implement a more practical approach for monitoring irrigation water quality, in comparison to the complex and expensive framework currently used for drinking water in developed countries.

Establishing regulations and clear guidelines for irrigation water quality in different countries or individual states of the United States for conventional sources of water is necessary, but not sufficient, to ensure healthy produce for human consumption. Proper food quality can only be insured if water sources are regulated and regularly tested. Testing can also be used to monitor accumulation of contaminants in soil. Moreover, plant tissue should also be checked periodically for contaminant uptake to ensure appropriate produce quality. Presently in the United States, reclaimed wastewater is regulated for irrigation usage [247,248] but almost any other source may be used without restriction. Guidelines are provided for pathogen levels in reclaimed wastewater [174]. There are recommendations for limits of geogenic contaminants in irrigation water in traditional sources [157,249] but there is a complete lack of recommendations or information on new and emerging organic contaminants or nanomaterials. There is a whole range of second and third generation nanomaterials proposed for commercial uses (e.g., nanocomposites and multi-element materials). These new nanomaterials will occur in waste streams and we have little understanding of their fate or toxicity. Moreover, the basis for existing recommendations is questionable, as concentrations far lower than some recommended values have been shown to be biomagnified in crops (see Table 1).

The present review focuses on the changing quality of water used for irrigation, and clearly there are many gaps to be addressed in future research. High priority should be given to research focused on improving our understanding and address the increasing occurrence of geogenic contaminants, pathogens and other biological contaminants in irrigation water, especially as they relate to food crops. Increased wastewater reuse for agricultural purposes will likely increase the occurrence of biologically active organic contaminants such as pharmaceuticals and antibiotics, and the effect on food crops

and human health is also a major research gap. The dynamic nature of the chemical composition of different irrigation water makes it very important that relevant and periodic data is available for both conventional and nonconventional water sources. A more comprehensive set of water quality guidelines needs to be created incorporating our present understanding of the occurrence and effects of emerging contaminants. There should be specific recommendations for monitoring and tests to check irrigation water quality used in agriculture. These recommendations can be soil specific, crop specific and water specific. The availability of new guidelines would help ensure better food quality, as the next generation will not only need larger quantities of irrigation water to feed the growing population, but health concerns may rise as we resort to the use of low-quality water to irrigate food and feed crops.

Author Contributions: A.M. led writing of this article, including extensive literature investigation on irrigation water quality, original draft preparation and sources on geogenic and nanomaterial contaminants. D.D.S. provided conceptualization of the need for a review on this topic, editing and input on literature sources for emerging contaminants. C.R. provided additional input on wastewater contaminants, nanomaterials and additional editing.

Funding: This work is based on research that was partially supported by the Nebraska Agricultural Experiment Station with funding from the Hatch Multistate Research (Accession Number 1011588) through the USDA National Institute of Food and Agriculture.

Acknowledgments: Malakar thanks NET for salary support. The author thank feedback on drafts by Jason White and Sushil Kanel, as well as editorial comments from Erin Haackker and Lacey Bodnar.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Dieter, C.A.; Maupin, M.A.; Caldwell, R.R.; Harris, M.A.; Ivahnenko, T.I.; Lovelace, J.K.; Barber, N.L.; Linsey, K.S. *Estimated Use of Water in the United States in 2015*; U.S. Geological Survey: Reston, VA, USA, 2018.
2. *World Agriculture: Towards 2015/2030 an FAO Perspective*; Bruinsma, J. (Ed.) Earthscan Publications Ltd.: London, UK, 2003; ISBN 9251048355.
3. FAO. How to Feed the World in 2050. *Insights Expert Meet. FAO* **2009**, *2050*, 1–35. [[CrossRef](#)]
4. Sauv , S.; Desrosiers, M. A review of what is an emerging contaminant. *Chem. Cent. J.* **2014**, *8*, 15. [[CrossRef](#)] [[PubMed](#)]
5. National Research Council. *Identifying Future Drinking Water Contaminants*; National Academies Press: Washington, DC, USA, 1999; ISBN 978-0-309-06432-3.
6. WWAP (United Nations World Water Assessment Programme). *The United Nations World Water Development Report 2017. Wastewater: The Untapped Resource*; United Nations Educational, Scientific and Cultural Organization: Paris, France, 22 March 2017.
7. Hass, A.; Mingelgrin, U.; Fine, P. Heavy metals in soils irrigated with wastewater. In *Treated Wastewater in Agriculture: Use and Impacts on the Soil Environment and Crops*; Wiley-Blackwell: Hoboken, NJ, USA, 2010; ISBN 9781405148627.
8. Allende, A.; Monaghan, J. Irrigation water quality for leafy crops: A perspective of risks and potential solutions. *Int. J. Environ. Res. Public Health* **2015**, *12*, 7457–7477. [[CrossRef](#)] [[PubMed](#)]
9. Singh, S.; Ghosh, N.C.; Gurjar, S.; Krishan, G.; Kumar, S.; Berwal, P. Index-based assessment of suitability of water quality for irrigation purpose under Indian conditions. *Environ. Monit. Assess.* **2018**, *190*, 29. [[CrossRef](#)] [[PubMed](#)]
10. Islam, M.A.; Romi , D.; Akber, M.A.; Romi , M. Trace metals accumulation in soil irrigated with polluted water and assessment of human health risk from vegetable consumption in Bangladesh. *Environ. Geochem. Health* **2018**, *40*, 59–85. [[CrossRef](#)] [[PubMed](#)]
11. Seiler, R.L.; Skorupa, J.P.; Naftz, D.L.; Nolan, B.T. *Irrigation-Induced Contamination of Water, Sediment, and Biota in the Western United States—Synthesis of Data from the National Irrigation Water Quality Program*; U.S. Geological Survey Professional Paper 1655; U.S. Geological Survey: Reston, VA, USA, 2003.
12. Servin, A.D.; De la Torre-Roche, R.; Castillo-Michel, H.; Pagano, L.; Hawthorne, J.; Musante, C.; Pignatello, J.; Uchimiya, M.; White, J.C. Exposure of agricultural crops to nanoparticle CeO₂ in biochar-amended soil. *Plant Physiol. Biochem.* **2017**, *110*, 147–157. [[CrossRef](#)] [[PubMed](#)]

13. Zhao, Q.; Ma, C.; White, J.C.; Dhankher, O.P.; Zhang, X.; Zhang, S.; Xing, B. Quantitative evaluation of multi-wall carbon nanotube uptake by terrestrial plants. *Carbon* **2017**, *114*, 661–670. [[CrossRef](#)]
14. Hochella, M.F.; Mogk, D.W.; Ranville, J.; Allen, I.C.; Luther, G.W.; Marr, L.C.; McGrail, B.P.; Murayama, M.; Qafoku, N.P.; Rosso, K.M.; et al. Natural, incidental, and engineered nanomaterials and their impacts on the Earth system. *Science* **2019**, *363*, eaau8299. [[CrossRef](#)] [[PubMed](#)]
15. Nolan, J.; Weber, K.A. Natural Uranium Contamination in Major U.S. Aquifers Linked to Nitrate. *Environ. Sci. Technol. Lett.* **2015**, *2*, 215–220. [[CrossRef](#)]
16. Mateo-Sagasta, J.; Marjani, S.; Turrall, H.; Burke, J. *Water Pollution from Agriculture: A Global Review Executive Summary*; The Food and Agriculture Organization of the United Nations: Rome, Italy; The International Water Management Institute: Colombo, Sri Lanka, 2017.
17. Dexssen, A.; Dole, R.B. *Ground Water in LaSalle and McMullen Counties*; Texas. U.S. Geol. Surv. Water-Supply Paper 375-G; U.S. GPO: Washington, DC, USA, 1916.
18. Schwennesen, A.T.; Forbes, R.H. *Ground Water in San Simon Valley, Arizona and New Mexico*; Water-Supply Paper 425-A; U.S. GPO: Washington, DC, USA, 1917.
19. Clark, W.O. *Ground Water for Irrigation in the Morgan Hill Area, California*; Water-Supply Paper 400-E; U.S. GPO: Washington, DC, USA, 1917.
20. Scofield, C.S.; Headley, F.B. Quality of irrigation water in relation to land reclamation. *J. Agric. Res.* **1921**, *21*, 265–278. [[CrossRef](#)]
21. Sturdy, D.; Calton, W.E.; Milne, G. A chemical survey of the waters of Mount Meru, Tanganyika Territory, especially with regard to their qualities for irrigation. *J. East Afr. Uganda Nat. Hist. Soc.* **1932**, *45–46*, 1–38.
22. Taylor, E.M.; Puri, A.N.; Asghar, A.G. Soil deterioration in the canal-irrigated areas of the Punjab. I. Equilibrium between calcium and sodium ions in base-exchange reactions. *Res. Publ.* **1934**, *4*, 7.
23. Mados, L. The qualifications of irrigation waters. *Mezogazdasagi Kut* **1940**, *12*, 121–131.
24. Eaton, F.M.; McCallum, R.D.; Mayhugh, M.S. *Quality of Irrigation Waters of the Hollister area of California with Special Reference to Boron Content and Its Effect on Apricots and Prunes*; Technical Bulletin; United States Department Agriculture: Washington, DC, USA, 1941; Volume 746, p. 59.
25. Wilcox, L.V. The quality of water for irrigation use. *U.S. Dept. Agr. Tech. Bull.* **1948**, *962*, 40.
26. Pacheco, J.d.I.R.; Lopez-Rubio, F.B. Analysis of waters for agricultural uses. *Inf. Quim. Anal.* **1949**, *3*, 90–96.
27. Thorne, D.W.; Thorne, J.P. Changes in composition of irrigated soils as related to the quality of irrigation waters. *Soil Sci. Soc. Am. Proc.* **1949**, *18*, 92–97. [[CrossRef](#)]
28. Lewis, G.C.; Juve, R.L. Some effects of irrigation-water quality on soil characteristics. *Soil Sci.* **1956**, *81*, 125–137. [[CrossRef](#)]
29. Pearson, H.E.; Huberty, M.R. Response of citrus to irrigation with waters of different chemical characteristics. *Proc. Am. Soc. Hortic. Sci.* **1959**, *73*, 248–256.
30. Babcock, K.L.; Carlson, R.M.; Schulz, R.K.; Overstreet, R. A study of the Effect of irrigation water composition on soil properties. *Hilgardia* **1959**, *29*, 155–170. [[CrossRef](#)]
31. Longenecker, D.E.; Lysterly, P.J. Chemical characteristics of soils of west Texas as affected by irrigation water quality. *Soil Sci.* **1959**, *87*, 207–216. [[CrossRef](#)]
32. Bernstein, L. Quantitative assessment of irrigation water quality. In *Water Quality Criteria*; Bramer, H., Ed.; ASTM International: West Conshohocken, PA, USA, 1967; pp. 51–65.
33. Park, D.M.; White, S.A.; McCarty, L.B.; Menchyk, N.A. *Interpreting Irrigation Water Quality Reports*; CU-14-700; Clemson University Cooperative Extension: Clemson, SC, USA, 2014.
34. Frenkel, H. Reassessment of Water Quality Criteria for Irrigation. *Ecol. Stud. Anal. Synth.* **1984**, *51*, 142–172.
35. Bauder, T.; Waskom, R.; Davis, J.; Sutherland, P. Irrigation water quality criteria. *Crop Ser. Irrig. Fact Sheet* **2007**, *506*, 10–13.
36. Feltz, H.; Engberg, R.; Sylvester, M. Investigations of water quality, bottom sediment, and biota associated with irrigation drainage in the western United States. In *Proceedings of the International Symposium on the Hydrologic Basis for Water Resources Management, Beijing, China, 23–26 October 1990*. Publ. no. 197.
37. Seiler, R.L. Synthesis of data from studies by the national irrigation water-quality program. *J. Am. Water Resour. Assoc.* **1996**, *32*, 1233–1245. [[CrossRef](#)]
38. Tangahu, B.V.; Sheikh Abdullah, S.R.; Basri, H.; Idris, M.; Anuar, N.; Mukhlisin, M. A review on heavy metals (As, Pb, and Hg) uptake by plants through phytoremediation. *Int. J. Chem. Eng.* **2011**, *2011*, 31. [[CrossRef](#)]

39. Intawongse, M.; Dean, J.R. Uptake of heavy metals by vegetable plants grown on contaminated soil and their bioavailability in the human gastrointestinal tract. *Food Addit. Contam.* **2006**, *23*, 36–48. [[CrossRef](#)] [[PubMed](#)]
40. Ma, C.; White, J.C.; Zhao, J.; Zhao, Q.; Xing, B. Uptake of engineered nanoparticles by food crops: Characterization, mechanisms, and implications. *Annu. Rev. Food Sci. Technol.* **2018**, *9*, 129–153. [[CrossRef](#)]
41. Calderón-Preciado, D.; Matamoros, V.; Bayona, J.M. Occurrence and potential crop uptake of emerging contaminants and related compounds in an agricultural irrigation network. *Sci. Total Environ.* **2011**, *412–413*, 14–19. [[CrossRef](#)]
42. Sedlak, D.L.; Gray, J.L.; Pinkston, K.E. Peer reviewed: Understanding microcontaminants in recycled water. *Environ. Sci. Technol.* **2000**, *34*, 508A–515A. [[CrossRef](#)]
43. IOM (Institute of Medicine) and NRC (National Research Council). *A Framework for Assessing Effects of the Food System*; The National Academic Press: Washington, DC, USA, 2015; ISBN 978-0-309-30780-2.
44. World Health Organization. *Pharmaceuticals in Drinking Water*; WHO: Geneva, Switzerland, 2012; ISBN 9789241502085.
45. Balakrishna, K.; Rath, A.; Praveenkumarreddy, Y.; Guruge, K.S.; Subedi, B. A review of the occurrence of pharmaceuticals and personal care products in Indian water bodies. *Ecotoxicol. Environ. Saf.* **2017**, *137*, 113–120. [[CrossRef](#)]
46. Yang, Y.; Ok, Y.S.; Kim, K.H.; Kwon, E.E.; Tsang, Y.F. Occurrences and removal of pharmaceuticals and personal care products (PPCPs) in drinking water and water/sewage treatment plants: A review. *Sci. Total Environ.* **2017**, *596–597*, 303–320. [[CrossRef](#)]
47. Fijalkowski, K.; Rorat, A.; Grobelak, A.; Kacprzak, M.J. The presence of contaminations in sewage sludge—The current situation. *J. Environ. Manag.* **2017**, *203*, 1126–1136. [[CrossRef](#)] [[PubMed](#)]
48. Subedi, B.; Balakrishna, K.; Joshua, D.I.; Kannan, K. Mass loading and removal of pharmaceuticals and personal care products including psychoactives, antihypertensives, and antibiotics in two sewage treatment plants in southern India. *Chemosphere* **2017**, *167*, 429–437. [[CrossRef](#)] [[PubMed](#)]
49. Subedi, B.; Lee, S.; Moon, H.B.; Kannan, K. Emission of artificial sweeteners, select pharmaceuticals, and personal care products through sewage sludge from wastewater treatment plants in Korea. *Environ. Int.* **2014**, *68*, 33–40. [[CrossRef](#)] [[PubMed](#)]
50. Madikizela, L.M.; Ncube, S.; Chimuka, L. Uptake of pharmaceuticals by plants grown under hydroponic conditions and natural occurring plant species: A review. *Sci. Total Environ.* **2018**, *636*, 477–486. [[CrossRef](#)] [[PubMed](#)]
51. Wu, X.; Dodgen, L.K.; Conkle, J.L.; Gan, J. Plant uptake of pharmaceutical and personal care products from recycled water and biosolids: A review. *Sci. Total Environ.* **2015**, *536*, 655–666. [[CrossRef](#)]
52. Tasho, R.P.; Cho, J.Y. Veterinary antibiotics in animal waste, its distribution in soil and uptake by plants: A review. *Sci. Total Environ.* **2016**, *563–564*, 366–376. [[CrossRef](#)] [[PubMed](#)]
53. Wu, X.; Conkle, J.L.; Ernst, F.; Gan, J. Treated wastewater irrigation: Uptake of pharmaceutical and personal care products by common vegetables under field conditions. *Environ. Sci. Technol.* **2014**, *48*, 11286–11293. [[CrossRef](#)]
54. Santiago, S.; Roll, D.M.; Ray, C.; Williams, C.; Moravcik, P.; Knopf, A. Effects of soil moisture depletion on vegetable crop uptake of pharmaceuticals and personal care products (PPCPs). *Environ. Sci. Pollut. Res.* **2016**, *23*, 20257–20268. [[CrossRef](#)]
55. Riemenschneider, C.; Al-Raggad, M.; Moeder, M.; Seiwert, B.; Salameh, E.; Reemtsma, T. Pharmaceuticals, their metabolites, and other polar pollutants in field-grown vegetables irrigated with treated municipal wastewater. *J. Agric. Food Chem.* **2016**, *64*, 5784–5792. [[CrossRef](#)]
56. Colon, B.; Toor, G.S. A review of uptake and translocation of pharmaceuticals and personal care products by food crops irrigated with treated wastewater. *Adv. Agron.* **2016**, *140*, 75–100.
57. Calderón-Preciado, D.; Jiménez-Cartagena, C.; Matamoros, V.; Bayona, J.M. Screening of 47 organic microcontaminants in agricultural irrigation waters and their soil loading. *Water Res.* **2011**, *45*, 221–231. [[CrossRef](#)] [[PubMed](#)]
58. Christou, A.; Agüera, A.; Bayona, J.M.; Cytryn, E.; Fotopoulou, V.; Lambropoulou, D.; Manaia, C.M.; Michael, C.; Revitt, M.; Schröder, P.; et al. The potential implications of reclaimed wastewater reuse for irrigation on the agricultural environment: The knowns and unknowns of the fate of antibiotics and antibiotic resistant bacteria and resistance genes—A review. *Water Res.* **2017**, *123*, 448–467. [[CrossRef](#)] [[PubMed](#)]

59. Williams-Nguyen, J.; Sallach, J.B.; Bartelt-Hunt, S.; Boxall, A.B.; Durso, L.M.; McLain, J.E.; Singer, R.S.; Snow, D.D.; Zilles, J.L. Antibiotics and antibiotic resistance in agroecosystems: State of the science. *J. Environ. Qual.* **2016**, *45*, 394–406. [[CrossRef](#)] [[PubMed](#)]
60. Durso, L.M.; Cook, K.L. Impacts of antibiotic use in agriculture: What are the benefits and risks? *Curr. Opin. Microbiol.* **2014**, *19*, 37–44. [[CrossRef](#)] [[PubMed](#)]
61. Kümmerer, K. Antibiotics in the aquatic environment—A review—Part I. *Chemosphere* **2009**, *75*, 417–434. [[CrossRef](#)] [[PubMed](#)]
62. Kümmerer, K. Antibiotics in the aquatic environment—A review—Part II. *Chemosphere* **2009**, *75*, 435–441. [[CrossRef](#)]
63. Behera, S.K.; Kim, H.W.; Oh, J.E.; Park, H.S. Occurrence and removal of antibiotics, hormones and several other pharmaceuticals in wastewater treatment plants of the largest industrial city of Korea. *Sci. Total Environ.* **2011**, *409*, 4351–4360. [[CrossRef](#)] [[PubMed](#)]
64. Zhou, L.J.; Ying, G.G.; Liu, S.; Zhao, J.L.; Yang, B.; Chen, Z.F.; Lai, H.J. Occurrence and fate of eleven classes of antibiotics in two typical wastewater treatment plants in South China. *Sci. Total Environ.* **2013**, *452–453*, 365–376. [[CrossRef](#)]
65. Aga, D.S.; Lenczewski, M.; Snow, D.; Muurinen, J.; Sallach, J.B.; Wallace, J.S. Challenges in the measurement of antibiotics and in evaluating their impacts in agroecosystems: A critical review. *J. Environ. Qual.* **2016**, *45*, 407–419. [[CrossRef](#)]
66. Burkholder, J.A.; Libra, B.; Weyer, P.; Heathcote, S.; Kolpin, D.; Thorne, P.S.; Wichman, M. Impacts of waste from concentrated animal feeding operations on water quality. *Environ. Health Perspect.* **2007**, *115*, 308–312. [[CrossRef](#)]
67. Chang, X.; Meyer, M.T.; Liu, X.; Zhao, Q.; Chen, H.; Chen, J.A.; Qiu, Z.; Yang, L.; Cao, J.; Shu, W. Determination of antibiotics in sewage from hospitals, nursery and slaughter house, wastewater treatment plant and source water in Chongqing region of Three Gorge Reservoir in China. *Environ. Pollut.* **2010**, *158*, 1444–1450. [[CrossRef](#)] [[PubMed](#)]
68. Duong, H.A.; Pham, N.H.; Nguyen, H.T.; Hoang, T.T.; Pham, H.V.; Pham, V.C.; Berg, M.; Giger, W.; Alder, A.C. Occurrence, fate and antibiotic resistance of fluoroquinolone antibacterials in hospital wastewaters in Hanoi, Vietnam. *Chemosphere* **2008**, *72*, 968–973. [[CrossRef](#)] [[PubMed](#)]
69. Hirsch, R.; Ternes, T.; Haberer, K.; Kratz, K.L. Occurrence of antibiotics in the aquatic environment. *Sci. Total Environ.* **1999**, *225*, 109–118. [[CrossRef](#)]
70. Kolpin, D.W.; Furlong, E.T.; Meyer, M.T.; Thurman, E.M.; Zaugg, S.D.; Barber, L.B.; Buxton, H.T. Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams, 1999–2000: A national reconnaissance. *Environ. Sci. Technol.* **2002**, *36*, 1202–1211. [[CrossRef](#)] [[PubMed](#)]
71. Yan, C.; Yang, Y.; Zhou, J.; Liu, M.; Nie, M.; Shi, H.; Gu, L. Antibiotics in the surface water of the Yangtze Estuary: Occurrence, distribution and risk assessment. *Environ. Pollut.* **2013**, *175*, 22–29. [[CrossRef](#)] [[PubMed](#)]
72. Deng, W.; Li, N.; Zheng, H.; Lin, H. Occurrence and risk assessment of antibiotics in river water in Hong Kong. *Ecotoxicol. Environ. Saf.* **2016**, *125*, 121–127. [[CrossRef](#)] [[PubMed](#)]
73. Zuccato, E.; Castiglioni, S.; Bagnati, R.; Melis, M.; Fanelli, R. Source, occurrence and fate of antibiotics in the Italian aquatic environment. *J. Hazard. Mater.* **2010**, *179*, 1042–1048. [[CrossRef](#)]
74. Ahmed, M.B.M.; Rajapaksha, A.U.; Lim, J.E.; Vu, N.T.; Kim, I.S.; Kang, H.M.; Lee, S.S.; Ok, Y.S. Distribution and accumulative pattern of tetracyclines and sulfonamides in edible vegetables of cucumber, tomato, and lettuce. *J. Agric. Food Chem.* **2015**, *63*, 398–405. [[CrossRef](#)]
75. Chitescu, C.L.; Nicolau, A.I.; Stolker, A.A.M. Uptake of oxytetracycline, sulfamethoxazole and ketoconazole from fertilised soils by plants. *Food Addit. Contam. Part A* **2013**, *30*, 1138–1146. [[CrossRef](#)]
76. Sallach, J.B.; Bartelt-Hunt, S.L.; Snow, D.D.; Li, X.; Hodges, L. Uptake of antibiotics and their toxicity to lettuce following routine irrigation with contaminated water in different soil types. *Environ. Eng. Sci.* **2018**, *35*. [[CrossRef](#)]
77. Azanu, D.; Mortey, C.; Darko, G.; Weisser, J.J.; Styryshave, B.; Abaidoo, R.C. Uptake of antibiotics from irrigation water by plants. *Chemosphere* **2016**, *157*, 107–114. [[CrossRef](#)] [[PubMed](#)]
78. Phillips, I.; Casewell, M.; Cox, T.; De Groot, B.; Friis, C.; Jones, R.; Nightingale, C.; Preston, R.; Waddell, J. Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *J. Antimicrob. Chemother.* **2004**, *53*, 28–52. [[CrossRef](#)] [[PubMed](#)]

79. Kuppusamy, S.; Kakarla, D.; Venkateswarlu, K.; Megharaj, M.; Yoon, Y.E.; Lee, Y.B. Veterinary antibiotics (VAs) contamination as a global agro-ecological issue: A critical view. *Agric. Ecosyst. Environ.* **2018**, *257*, 47–59. [[CrossRef](#)]
80. Bedford, M. Removal of antibiotic growth promoters from poultry diets: Implications and strategies to minimise subsequent problems. *Worlds Poult. Sci. J.* **2000**, *56*, 347–365. [[CrossRef](#)]
81. Schuijt, T.J.; van der Poll, T.; de Vos, W.M.; Wiersinga, W.J. The intestinal microbiota and host immune interactions in the critically ill. *Trends Microbiol.* **2013**, *21*, 221–229. [[CrossRef](#)] [[PubMed](#)]
82. Davies, J.; Davies, D. Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* **2010**, *74*, 417–433. [[CrossRef](#)]
83. Zhang, F.S.; Xie, Y.F.; Li, X.W.; Wang, D.Y.; Yang, L.S.; Nie, Z.Q. Accumulation of steroid hormones in soil and its adjacent aquatic environment from a typical intensive vegetable cultivation of North China. *Sci. Total Environ.* **2015**, *538*, 423–430. [[CrossRef](#)] [[PubMed](#)]
84. Donk, S.v.; Biswas, S.; Kranz, W.; Snow, D.; Bartelt-Hunt, S.; Mader, T.; Shapiro, C.; Shelton, D.; Tarkalson, D.; Zhang, T.; et al. Transport of steroid hormones in the vadose zone after land application of beef cattle manure. *Biol. Syst. Eng. Pap. Publ.* **2013**, *56*, 1327–1338.
85. Bartelt-Hunt, S.; Snow, D.D.; Damon-Powell, T.; Miesbach, D. Occurrence of steroid hormones and antibiotics in shallow groundwater impacted by livestock waste control facilities. *J. Contam. Hydrol.* **2011**, *123*, 94–103. [[CrossRef](#)]
86. Torres, N.H.; Aguiar, M.M.; Ferreira, L.F.R.; Américo, J.H.P.; Machado, Â.M.; Cavalcanti, E.B.; Tornisielo, V.L. Detection of hormones in surface and drinking water in Brazil by LC-ESI-MS/MS and ecotoxicological assessment with *Daphnia magna*. *Environ. Monit. Assess.* **2015**, *187*, 379. [[CrossRef](#)]
87. Pauwels, B.; Noppe, H.; De Brabander, H.; Verstraete, W. Comparison of steroid hormone concentrations in domestic and hospital wastewater treatment plants. *J. Environ. Eng.* **2008**, *134*, 933–936. [[CrossRef](#)]
88. Servos, M.R.; Bennie, D.T.; Burnison, B.K.; Jurkovic, A.; McInnis, R.; Neheli, T.; Schnell, A.; Seto, P.; Smyth, S.A.; Ternes, T.A. Distribution of estrogens, 17 β -estradiol and estrone, in Canadian municipal wastewater treatment plants. *Sci. Total Environ.* **2005**, *336*, 155–170. [[CrossRef](#)] [[PubMed](#)]
89. Andersen, H.; Siegrist, H.; Halling-Sørensen, B.; Ternes, T.A. Fate of estrogens in a municipal sewage treatment plant. *Environ. Sci. Technol.* **2003**, *37*, 4021–4026. [[CrossRef](#)] [[PubMed](#)]
90. Baronti, C.; Curini, R.; D’Ascenzo, G.; Di Corcia, A.; Gentili, A.; Samperi, R. Monitoring natural and synthetic estrogens at activated sludge sewage treatment plants and in a receiving river water. *Environ. Sci. Technol.* **2000**, *34*, 5059–5066. [[CrossRef](#)]
91. Sellin, M.K.; Snow, D.D.; Akerly, D.L.; Kolok, A.S. Estrogenic compounds downstream from three small cities in Eastern Nebraska: Occurrence and biological effect. *J. Am. Water Resour. Assoc.* **2009**, *45*, 14–21. [[CrossRef](#)]
92. Yarahmadi, H.; Duy, S.V.; Hachad, M.; Dorner, S.; Sauvé, S.; Prévost, M. Seasonal variations of steroid hormones released by wastewater treatment plants to river water and sediments: Distribution between particulate and dissolved phases. *Sci. Total Environ.* **2018**, *635*, 144–155. [[CrossRef](#)] [[PubMed](#)]
93. Zheng, W.; Wiles, K.N.; Holm, N.; Deppe, N.A.; Shipley, C.R. Uptake, Translocation, and Accumulation of Pharmaceutical and Hormone Contaminants in Vegetables. In *ACS Symposium Series*; American Chemical Society: Washington, DC, USA, 2014; Volume 1171, pp. 167–181.
94. Adeel, M.; Song, X.; Wang, Y.; Francis, D.; Yang, Y. Environmental impact of estrogens on human, animal and plant life: A critical review. *Environ. Int.* **2017**, *99*, 107–119. [[CrossRef](#)]
95. Rose, S.C.; Carter, A.D. Agrochemical leaching and water contamination. In *Conservation Agriculture*; Springer: Dordrecht, The Netherlands, 2003; pp. 417–424.
96. Jimoh, O.D.; Ayodeji, M.A.; Mohammed, B. Effects of agrochemicals on surface waters and groundwaters in the Tunga-Kawo (Nigeria) irrigation scheme. *Hydrol. Sci. J.* **2003**, *48*, 1013–1023. [[CrossRef](#)]
97. Yu, Y.; Hu, S.; Yang, Y.; Zhao, X.; Xue, J.; Zhang, J.; Gao, S.; Yang, A. Successive monitoring surveys of selected banned and restricted pesticide residues in vegetables from the northwest region of China from 2011 to 2013. *BMC Public Health* **2017**, *18*, 91. [[CrossRef](#)]
98. Retention, uptake, and translocation of agrochemicals in plants. In *ACS Symposium Series*; Myung, K.; Satchivi, N.M.; Kingston, C.K. (Eds.) American Chemical Society: Washington, DC, USA, 2014; Volume 1171, ISBN 0-8412-2972-4.
99. Juraske, R.; Castells, F.; Vijay, A.; Muñoz, P.; Antón, A. Uptake and persistence of pesticides in plants: Measurements and model estimates for imidacloprid after foliar and soil application. *J. Hazard. Mater.* **2009**, *165*, 683–689. [[CrossRef](#)]

100. Elmore, S.A.; Boorman, G.A. Environmental toxicologic pathology and human health. *Haschek Rousseaux's Handb. Toxicol. Pathol.* **2013**, 1029–1049. [[CrossRef](#)]
101. US Protection Agency. *Cyanobacteria and Cyanotoxins: Information for Drinking Water Systems*; EPA-810F11001; USEPA: Washington, DC, USA, 2014.
102. Crush, J.R.; Briggs, L.R.; Sprosen, J.M.; Nichols, S.N. Effect of irrigation with lake water containing microcystins on microcystin content and growth of ryegrass, clover, rape, and lettuce. *Environ. Toxicol.* **2008**, *23*, 246–252. [[CrossRef](#)] [[PubMed](#)]
103. Corbel, S.; Mouglin, C.; Bouaïcha, N. Cyanobacterial toxins: Modes of actions, fate in aquatic and soil ecosystems, phytotoxicity and bioaccumulation in agricultural crops. *Chemosphere* **2014**, *96*, 1–15. [[CrossRef](#)] [[PubMed](#)]
104. Loftin, K.A.; Graham, J.L.; Hilborn, E.D.; Lehmann, S.C.; Meyer, M.T.; Dietze, J.E.; Griffith, C.B. Cyanotoxins in inland lakes of the United States: Occurrence and potential recreational health risks in the EPA National Lakes Assessment 2007. *Harmful Algae* **2016**, *56*, 77–90. [[CrossRef](#)] [[PubMed](#)]
105. Al-Sammak, M.A.; Hoagland, K.D.; Cassada, D.; Snow, D.D. Co-occurrence of the cyanotoxins BMAA, DABA and anatoxin-a in Nebraska reservoirs, fish, and aquatic plants. *Toxins* **2014**, *6*, 488–508. [[CrossRef](#)] [[PubMed](#)]
106. Miller, A.; Russell, C. Food crops irrigated with cyanobacteria-contaminated water: An emerging public health issue in Canada. *Environ. Heal. Rev.* **2017**, *60*, 58–63. [[CrossRef](#)]
107. Saqrane, S.; Oudra, B. CyanoHAB occurrence and water irrigation cyanotoxin contamination: Ecological impacts and potential health risks. *Toxins* **2009**, *1*, 113–122. [[CrossRef](#)] [[PubMed](#)]
108. Abeysiriwardena, N.M.; Gascoigne, S.J.L.; Anandappa, A. Algal bloom expansion increases cyanotoxin risk in food. *Yale J. Biol. Med.* **2018**, *91*, 129–142. [[PubMed](#)]
109. Al-Gabr, H.M.; Zheng, T.; Yu, X. Fungi contamination of drinking water. *Rev. Environ. Contam. Toxicol.* **2014**, *228*, 121–139. [[PubMed](#)]
110. Oliveira, B.R.; Mata, A.T.; Ferreira, J.P.; Barreto Crespo, M.T.; Pereira, V.J.; Bronze, M.R. Production of mycotoxins by filamentous fungi in untreated surface water. *Environ. Sci. Pollut. Res.* **2018**, *25*, 17519–17528. [[CrossRef](#)]
111. Kolpin, D.W.; Hoerger, C.C.; Meyer, M.T.; Wettstein, F.E.; Hubbard, L.E.; Bucheli, T.D. Phytoestrogens and mycotoxins in Iowa streams: An examination of underinvestigated compounds in agricultural basins. *J. Environ. Qual.* **2010**, *39*, 2089–2099. [[CrossRef](#)] [[PubMed](#)]
112. Rao, G.J.; Govindaraju, G.; Sivasithamparam, N.; Shanmugasundaram, E.R.B. Uptake, translocation and persistence of mycotoxins in rice seedlings. *Plant Soil* **1982**, *66*, 121–123. [[CrossRef](#)]
113. Mohammad-Hasani, F.; Mirlohi, M.; Mosharraf, L.; Hasanzade, A. Occurrence of aflatoxins in wheat flour specified for sangak bread and its reduction through fermentation and baking. *Qual. Assur. Saf. Crop. Foods* **2016**, *8*, 1–8. [[CrossRef](#)]
114. Tola, M.; Kebede, B. Occurrence, importance and control of mycotoxins: A review. *Cogent Food Agric.* **2016**. [[CrossRef](#)]
115. Tanaka, H.; Asano, T.; Schroeder, E.D.; Tchobanoglous, G. Estimating the safety of wastewater reclamation and reuse using enteric virus monitoring data. *Water Environ. Res.* **1998**, *70*, 39–51. [[CrossRef](#)]
116. Asano, T.; Cotruvo, J.A. Groundwater recharge with reclaimed municipal wastewater: Health and regulatory considerations. *Water Res.* **2004**, *38*, 1941–1951. [[CrossRef](#)] [[PubMed](#)]
117. Lothrop, N.; Bright, K.R.; Sexton, J.; Pearce-Walker, J.; Reynolds, K.A.; Verhougstraete, M.P. Optimal strategies for monitoring irrigation water quality. *Agric. Water Manag.* **2018**, *199*, 86–92. [[CrossRef](#)]
118. Jongman, M.; Chidamba, L.; Korsten, L. Bacterial biomes and potential human pathogens in irrigation water and leafy greens from different production systems described using pyrosequencing. *J. Appl. Microbiol.* **2017**, *123*, 1043–1053. [[CrossRef](#)]
119. Truchado, P.; Hernandez, N.; Gil, M.I.; Ivanek, R.; Allende, A. Correlation between *E. coli* levels and the presence of foodborne pathogens in surface irrigation water: Establishment of a sampling program. *Water Res.* **2018**, *128*, 226–233. [[CrossRef](#)]
120. Steele, M.; Odumeru, J. Irrigation water as source of foodborne pathogens on fruit and vegetables. *J. Food Prot.* **2004**, *67*, 2839–2849. [[CrossRef](#)]
121. Pachepsky, Y.; Shelton, D.R.; McLain, J.E.T.; Patel, J.; Mandrell, R.E. Irrigation waters as a source of pathogenic microorganisms in produce: A review. *Adv. Agron.* **2011**, *113*, 75–141. [[CrossRef](#)]

122. Park, S.; Szonyi, B.; Gautam, R.; Nightingale, K.; Anciso, J.; Ivanek, R. Risk factors for microbial contamination in fruits and vegetables at the preharvest level: A systematic review. *J. Food Prot.* **2012**, *75*, 2055–2081. [[CrossRef](#)] [[PubMed](#)]
123. Jongman, M.; Korsten, L. Irrigation water quality and microbial safety of leafy greens in different vegetable production systems: A review. *Food Rev. Int.* **2018**, *34*, 308–328. [[CrossRef](#)]
124. Herman, K.M.; Hall, A.J.; Gould, L.H. Outbreaks attributed to fresh leafy vegetables, United States, 1973–2012. *Epidemiol. Infect.* **2015**, *143*, 3011–3021. [[CrossRef](#)] [[PubMed](#)]
125. National Outbreak Reporting System (NORS) Dashboard | CDC. Available online: <https://wwwn.cdc.gov/norsdashboard/> (accessed on 25 March 2019).
126. Multistate Outbreak of *E. coli* O157:H7 Infections Linked to Romaine Lettuce (Final Update) | Investigation Notice: Multistate Outbreak of *E. coli* O157:H7 Infections April 2018 | *E. coli* | CDC. Available online: <https://www.cdc.gov/ecoli/2018/o157h7-04-18/index.html> (accessed on 25 March 2019).
127. Outbreak of *E. coli* Infections Linked to Romaine Lettuce | *E. coli* Infections Linked to Romaine Lettuce | November 2018 | *E. coli* | CDC. Available online: <https://www.cdc.gov/ecoli/2018/o157h7-11-18/index.html> (accessed on 25 March 2019).
128. FDA Investigation of a Multistate Outbreak of Cyclospora Illnesses Linked to Fresh Express Salad Mix Served at McDonald’s Ends | FDA. Available online: <https://www.fda.gov/food/outbreaks-foodborne-illness/fda-investigation-multistate-outbreak-cyclospora-illnesses-linked-fresh-express-salad-mix-served#Cyclospora> (accessed on 3 July 2019).
129. Olivieri, A.W.; Seto, E.; Cooper, R.C.; Cahn, M.D.; Colford, J.; Crook, J.; Debroux, J.-F.; Mandrell, R.; Suslow, T.; Tchobanoglous, G.; et al. Risk-based review of California’s water-recycling criteria for agricultural irrigation. *J. Environ. Eng.* **2014**, *140*, 04014015. [[CrossRef](#)]
130. Leaman, S.; Gorny, J.; Wetherington, D.; Belkris, H. *Agricultural Water: Five Year Research Review*; Center for Produce Safety: Davis, CA, USA, 2014.
131. U.S. EPA. *Regulations Governing Agricultural Use of Municipal Wastewater and Sludge*; National Academy Press: Washington, DC, USA, 1996; ISBN 0309054796.
132. Jones, L.A.; Worobo, R.W.; Smart, C.D. UV light inactivation of human and plant pathogens in unfiltered surface irrigation water. *Appl. Environ. Microbiol.* **2014**, *80*, 849–854. [[CrossRef](#)] [[PubMed](#)]
133. Uyttendaele, M.; Jaykus, L.A.; Amoah, P.; Chiodini, A.; Cunliffe, D.; Jacxsens, L.; Holvoet, K.; Korsten, L.; Lau, M.; McClure, P.; et al. Microbial hazards in irrigation water: Standards, norms, and testing to manage use of water in fresh produce primary production. *Compr. Rev. Food Sci. Food Saf.* **2015**, *14*, 336–356. [[CrossRef](#)]
134. World Health Organization World Health Organization (WHO): Antibiotic Resistance—Fact Sheet. Available online: <https://www.who.int/en/news-room/fact-sheets/detail/antibiotic-resistance> (accessed on 26 June 2019).
135. Perry, J.A.; Wright, G.D. The antibiotic resistance “mobilome”: Searching for the link between environment and clinic. *Front. Microbiol.* **2013**, *4*, 138. [[CrossRef](#)]
136. Marti, E.; Variatza, E.; Balcazar, J.L. The role of aquatic ecosystems as reservoirs of antibiotic resistance. *Trends Microbiol.* **2014**, *22*, 36–41. [[CrossRef](#)]
137. Bergeron, S.; Brown, R.; Homer, J.; Rehage, S.; Boopathy, R. Presence of antibiotic resistance genes in different salinity gradients of freshwater to saltwater marshes in southeast Louisiana, USA. *Int. Biodeterior. Biodegrad.* **2016**, *113*, 80–87. [[CrossRef](#)]
138. Pepper, I.; Brooks, J.P.; Gerba, C.P. Antibiotic resistant bacteria in municipal wastes: Is there reason for concern? *Environ. Sci. Technol.* **2018**, *52*, 3949–3959. [[CrossRef](#)] [[PubMed](#)]
139. Zhang, X.X.; Zhang, T.; Fang, H.H.P. Antibiotic resistance genes in water environment. *Appl. Microbiol. Biotechnol.* **2009**, *82*, 397–414. [[CrossRef](#)] [[PubMed](#)]
140. Fahrenfeld, N.; Ma, Y.; O’Brien, M.; Pruden, A. Reclaimed water as a reservoir of antibiotic resistance genes: Distribution system and irrigation implications. *Front. Microbiol.* **2013**, *4*, 130. [[CrossRef](#)] [[PubMed](#)]
141. Aslan, A.; Cole, Z.; Bhattacharya, A.; Oyibo, O.; Aslan, A.; Cole, Z.; Bhattacharya, A.; Oyibo, O. Presence of antibiotic-resistant *Escherichia coli* in wastewater treatment plant effluents utilized as water reuse for irrigation. *Water* **2018**, *10*, 805. [[CrossRef](#)]
142. Gekenidis, M.-T.; Qi, W.; Hummerjohann, J.; Zbinden, R.; Walsh, F.; Drissner, D. Antibiotic-resistant indicator bacteria in irrigation water: High prevalence of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli*. *PLoS ONE* **2018**, *13*, e0207857. [[CrossRef](#)] [[PubMed](#)]

143. Farkas, A.; Bocoş, B.; Butiuc-Keul, A. Antibiotic resistance and intI1 carriage in waterborne enterobacteriaceae. *Water Air Soil Pollut.* **2016**, *227*, 251. [[CrossRef](#)]
144. Olaimat, A.N.; Holley, R.A. Factors influencing the microbial safety of fresh produce: A review. *Food Microbiol.* **2012**, *32*, 1–19. [[CrossRef](#)]
145. Vital, P.G.; Zara, E.S.; Paraoan, C.E.M.; Dimasupil, M.A.Z.; Abello, J.J.M.; Santos, I.T.G.; Rivera, W.L. Antibiotic resistance and extended-spectrum beta-lactamase production of escherichia coli isolated from irrigationwaters in selected urban farms in Metro Manila, Philippines. *Water* **2018**, *10*, 548. [[CrossRef](#)]
146. Fipps, G. *Irrigation Water Quality Standards and Salinity Management Strategies*; Texas Agriculture Extension Service 7-96 Rev edition; Texas A&M University System: College Station, TX, USA, 1996.
147. Stoner, J.D. *Water-Quality Indices for Specific Water Uses*; Circular 770; United States Department of the Interior, Geological Survey: Arlington, VA, USA, 1978. [[CrossRef](#)]
148. Ayotte, J.D.; Gronberg, J.A.M.; Apodaca, L.E. *Trace Elements and Radon in Groundwater across the United States, 1992–2003*; Scientific Investigations Report 2011–5059; U.S. Geological Survey: Reston, VA, USA, 2011; Volume i–xi, pp. 1–115.
149. Welch, A.H.; Westjohn, D.B.; Helsel, D.R.; Wanty, R.B. Arsenic in ground water of the United States: Occurrence and geochemistry. *Ground Water* **2000**, *38*, 589–604. [[CrossRef](#)]
150. Rodríguez-Lado, L.; Sun, G.; Berg, M.; Zhang, Q.; Xue, H.; Zheng, Q.; Johnson, C.A. Groundwater arsenic contamination throughout China. *Science* **2013**, *341*, 866–868. [[CrossRef](#)]
151. Selck, B.J.; Carling, G.T.; Kirby, S.M.; Hansen, N.C.; Bickmore, B.R.; Tingey, D.G.; Rey, K.; Wallace, J.; Jordan, J.L. Investigating anthropogenic and geogenic sources of groundwater contamination in a semi-arid alluvial basin, Goshen Valley, UT, USA. *Water Air Soil Pollut.* **2018**, *229*, 186. [[CrossRef](#)]
152. Signes-Pastor, A.J.; Mitra, K.; Sarkhel, S.; Hobbes, M.; Burló, F.; De Groot, W.T.; Carbonell-Barrachina, A.A. Arsenic speciation in food and estimation of the dietary intake of inorganic arsenic in a rural village of West Bengal, India. *J. Agric. Food Chem.* **2008**, *56*, 9469–9474. [[CrossRef](#)] [[PubMed](#)]
153. Erban, L.E.; Gorelick, S.M.; Fendorf, S. Arsenic in the multi-aquifer system of the Mekong Delta, Vietnam: Analysis of large-scale spatial trends and controlling factors. *Environ. Sci. Technol.* **2014**, *48*, 6081–6088. [[CrossRef](#)] [[PubMed](#)]
154. Huhmann, B.; Harvey, C.F.; Uddin, A.; Choudhury, I.; Ahmed, K.M.; Duxbury, J.M.; Ellis, T.; van Geen, A. Inversion of high-arsenic soil for improved rice yield in Bangladesh. *Environ. Sci. Technol.* **2019**, *53*, 3410–3418. [[CrossRef](#)] [[PubMed](#)]
155. Pick, T. *Assessing Water Quality for Human Consumption, Agriculture, and Aquatic Life Uses*; United States Department of Agriculture: Washington, DC, USA, 2011.
156. U.S. EPA. *Guidelines for Water Reuse 2012*; US Agency for International Development: Washington, DC, USA, 2012; p. 643.
157. Ayers, R.S.; Westcot, D.W. *Water Quality for Agriculture*; Food and Agricultural Organization, United Nations: Rome, Italy, 1976.
158. Kandakji, T.; Udeigwe, T.K.; Dixon, R.; Li, L. Groundwater-induced alterations in elemental concentration and interactions in semi-arid soils of the Southern High Plains, USA. *Environ. Monit. Assess.* **2015**, *187*, 665. [[CrossRef](#)] [[PubMed](#)]
159. Scanlon, B.R.; Nicot, J.P.; Reedy, R.C.; Kurtzman, D.; Mukherjee, A.; Nordstrom, D.K. Elevated naturally occurring arsenic in a semiarid oxidizing system, Southern High Plains aquifer, Texas, USA. *Appl. Geochem.* **2009**, *24*, 2061–2071. [[CrossRef](#)]
160. Malan, M.; Müller, F.; Cyster, L.; Raitt, L.; Aalbers, J. Heavy metals in the irrigation water, soils and vegetables in the Philippi horticultural area in the Western Cape Province of South Africa. *Environ. Monit. Assess.* **2015**, *187*, 1–8. [[CrossRef](#)] [[PubMed](#)]
161. Gupta, M.; Gupta, S. An overview of selenium uptake, metabolism, and toxicity in Plants. *Front. Plant Sci.* **2017**, *7*, 1–14. [[CrossRef](#)]
162. Hakonson-Hayes, A.C.; Fresquez, P.R.; Whicker, F.W. Assessing potential risks from exposure to natural uranium in well water. *J. Environ. Radioact.* **2002**, *59*, 29–40. [[CrossRef](#)]
163. Islam, S.M.A.; Fukushi, K.; Yamamoto, K. Contamination of agricultural soil by arsenic containing irrigation water in Bangladesh: Overview of status and a proposal for novel biological remediation. *WIT Trans. Biomed. Heal.* **2006**, *6*, 295–316. [[CrossRef](#)]

164. Jeambrun, M.; Pourcelot, L.; Mercat, C.; Boulet, B.; Pelt, E.; Chabaux, F.; Cagnat, X.; Gauthier-Lafaye, F. Potential sources affecting the activity concentrations of ²³⁸U, ²³⁵U, ²³²Th and some decay products in lettuce and wheat samples. *J. Environ. Monit.* **2012**, *14*, 2902–2912. [[CrossRef](#)] [[PubMed](#)]
165. Bañuelos, G.S.; Ajwa, H.A.; Caceres, L.; Dyer, D. Germination responses and boron accumulation in germplasm from Chile and the United States grown with boron-enriched water. *Ecotoxicol. Environ. Saf.* **1999**, *43*, 62–67. [[CrossRef](#)] [[PubMed](#)]
166. Rhoades, J.D.; Bingham, F.T.; Letey, J.; Hoffman, G.J.; Dedrick, A.R.; Pinter, P.J.; Replogle, J.A. Use of saline drainage water for irrigation: Imperial Valley study. *Agric. Water Manag.* **1989**, *16*, 25–36. [[CrossRef](#)]
167. Hopkins, B.G.; Horneck, D.A.; Stevens, R.G.; Ellsworth, J.W.; Sullivan, D.M. *Managing Irrigation Water Quality for Crop Production in the Pacific Northwest*; PNW597-E; USDA: Washington, DC, USA, 2007. Available online: <https://catalog.extension.oregonstate.edu/sites/catalog/files/project/pdf/pnw597.pdf> (accessed on 19 March 2019).
168. Hem, J.D. Chemistry and occurrence of cadmium and zinc in surface water and groundwater cadmium is reported of compounds in rice. *Water Resour. Res.* **1972**, *8*, 661–679. [[CrossRef](#)]
169. Alexakis, D. Assessment of water quality in the Messolonghi-Etoliko and Neochorio region (West Greece) using hydrochemical and statistical analysis methods. *Environ. Monit. Assess.* **2011**, *182*, 397–413. [[CrossRef](#)] [[PubMed](#)]
170. Irmak, S. Copper correlation of irrigation water, soils and plants in the Cukurova Region of Turkey. *Int. J. Soil Sci.* **2009**, *4*, 46–56. [[CrossRef](#)]
171. Manning, A.H.; Mills, C.T.; Morrison, J.M.; Ball, L.B. Insights into controls on hexavalent chromium in groundwater provided by environmental tracers, Sacramento Valley, California, USA. *Appl. Geochem.* **2015**, *62*, 186–199. [[CrossRef](#)]
172. Stasinou, S.; Zabetakis, I. The uptake of nickel and chromium from irrigation water by potatoes, carrots and onions. *Ecotoxicol. Environ. Saf.* **2013**, *91*, 122–128. [[CrossRef](#)]
173. USEPA. *2018 Edition of the Drinking Water Standards and Health Advisories*; USEPA: Washington, DC, USA, 2018.
174. Gurel, M.; Iskender, G.; Ovez, S.; Arslan-Alaton, I.; Tanik, A.; Orhon, D. A global overview of treated wastewater guidelines and standards for agricultural reuse. *Fresenius Environ. Bull.* **2007**, *16*, 590–595.
175. Pescod, M.B. *Wastewater Treatment and Use in Agriculture*; Food and Agricultural Organization, United Nations: Rome, Italy, 1992; ISBN 9251031355.
176. US EPA. National Recommended Water Quality Criteria—Aquatic Life Criteria Table. Available online: <https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table> (accessed on 27 December 2018).
177. Groschen, G.E.; Arnold, T.L.; Morrow, W.S.; Warner, K.L. *Occurrence and Distribution of Iron, Manganese, and Selected Trace elements in Ground Water in the Glacial Aquifer System of the Northern United States*; U.S. Geological Survey Scientific Investigations Report 2009–5006; USGS: Reston, VA, USA, 2008.
178. Bundschuh, J.; Nath, B.; Bhattacharya, P.; Liu, C.W.; Armienta, M.A.; Moreno López, M.V.; Lopez, D.L.; Jean, J.S.; Cornejo, L.; Lauer Macedo, L.F.; et al. Arsenic in the human food chain: The Latin American perspective. *Sci. Total Environ.* **2012**, *429*, 92–106. [[CrossRef](#)]
179. Meharg, A.A.; Rahman, M. Arsenic contamination of Bangladesh paddy field soils: Implications for rice contribution to arsenic consumption. *Environ. Sci. Technol.* **2003**, *37*, 229–234. [[CrossRef](#)] [[PubMed](#)]
180. Finnegan, P.M.; Chen, W. Arsenic toxicity: The effects on plant metabolism. *Front. Physiol.* **2012**, *3*, 182. [[CrossRef](#)] [[PubMed](#)]
181. Kim, J.Y.; Davis, A.P.; Kim, K.W. Stabilization of available arsenic in highly contaminated mine tailings using iron. *Environ. Sci. Technol.* **2003**, *37*, 189–195. [[CrossRef](#)] [[PubMed](#)]
182. Bakhat, H.F.; Zia, Z.; Fahad, S.; Abbas, S.; Hammad, H.M.; Shahzad, A.N.; Abbas, F.; Alharby, H.; Shahid, M. Arsenic uptake, accumulation and toxicity in rice plants: Possible remedies for its detoxification: A review. *Environ. Sci. Pollut. Res.* **2017**, *24*, 9142–9158. [[CrossRef](#)] [[PubMed](#)]
183. Arslan, B.; Djamgoz, M.B.A.; Akün, E. ARSENIC: A review on exposure pathways, accumulation, mobility and transmission into the human food chain. *Rev. Environ. Contam. Toxicol.* **2017**, *243*, 27–51. [[PubMed](#)]
184. Zhao, F.J.; Ma, J.F.; Meharg, A.A.; McGrath, S.P. Arsenic uptake and metabolism in plants. *New Phytol.* **2009**, *181*, 777–794. [[CrossRef](#)] [[PubMed](#)]
185. Zhao, F.-J.; McGrath, S.P.; Meharg, A.A. Arsenic as a food chain contaminant: Mechanisms of plant uptake and metabolism and mitigation strategies. *Annu. Rev. Plant Biol.* **2010**, *61*, 535–559. [[CrossRef](#)] [[PubMed](#)]

186. Chakraborty, S.; Alam, M.O.; Bhattacharya, T.; Singh, Y.N. Arsenic accumulation in food crops: A potential threat in Bengal Delta Plain. *Water Qual. Expo. Heal.* **2014**, *6*, 233–246. [[CrossRef](#)]
187. Brammer, H. Mitigation of arsenic contamination in irrigated paddy soils in South and South-East Asia. *Environ. Int.* **2009**, *35*, 856–863. [[CrossRef](#)]
188. Polizzotto, M.L.; Birgand, F.; Badruzzaman, A.B.M.; Ali, M.A. Amending irrigation channels with jute-mesh structures to decrease arsenic loading to rice fields in Bangladesh. *Ecol. Eng.* **2015**, *74*, 101–106. [[CrossRef](#)]
189. Winkel, L.H.E.; Johnson, C.A.; Lenz, M.; Grundl, T.; Leupin, O.X.; Amini, M.; Charlet, L. Environmental selenium research: From microscopic processes to global understanding. *Environ. Sci. Technol.* **2012**, *46*, 571–579. [[CrossRef](#)] [[PubMed](#)]
190. USGS. *Geochemical and Mineralogical Data for Soils of the Conterminous United States*; USGS: Reston, VA, USA, 2014.
191. Wang, J.; Zhou, C.; Xiao, X.; Xie, Y.; Zhu, L.; Ma, Z. Enhanced iron and selenium uptake in plants by volatile emissions of *Bacillus amyloliquefaciens* (BF06). *Appl. Sci.* **2017**, *7*, 85. [[CrossRef](#)]
192. Mitchell, N.; Pérez-Sánchez, D.; Thorne, M.C. A review of the behaviour of U-238 series radionuclides in soils and plants. *J. Radiol. Prot.* **2013**, *33*, R17–R48. [[CrossRef](#)] [[PubMed](#)]
193. Soudek, P.; Petrova, T.; Benesova, D.; Dvorakova, M.; Vanek, T. Uranium uptake by hydroponically cultivated crop plants. *J. Environ. Radioact.* **2011**, *102*, 598–604. [[CrossRef](#)] [[PubMed](#)]
194. Boghi, A.; Roose, T.; Kirk, G.J.D. A model of uranium uptake by plant roots allowing for root-induced changes in the soil. *Environ. Sci. Technol.* **2018**, *52*, 3536–3545. [[CrossRef](#)] [[PubMed](#)]
195. Hayes, A.C.; Fresquez, P.R.; Whicker, W.F. *Uranium Uptake Study, Nambe, New Mexico: Source Document*; Los Alamos National Laboratory: Los Alamos, NM, USA, 2000.
196. Neves, O.; Abreu, M.M. Are uranium-contaminated soil and irrigation water a risk for human vegetables consumers? A study case with *Solanum tuberosum* L., *Phaseolus vulgaris* L. and *Lactuca sativa* L. *Ecotoxicology* **2009**, *18*, 1130–1136. [[CrossRef](#)] [[PubMed](#)]
197. Neves, M.O.; Abreu, M.M.; Figueiredo, V. Uranium in vegetable foodstuffs: Should residents near the Cunha Baixa uranium mine site (Central Northern Portugal) be concerned? *Environ. Geochem. Health* **2012**, *34*, 181–189. [[CrossRef](#)]
198. Gomes, M.A.d.C.; Hauser-Davis, R.A.; Suzuki, M.S.; Vitória, A.P. Plant chromium uptake and transport, physiological effects and recent advances in molecular investigations. *Ecotoxicol. Environ. Saf.* **2017**, *140*, 55–64. [[CrossRef](#)]
199. Song, Y.; Jin, L.; Wang, X. Cadmium absorption and transportation pathways in plants. *Int. J. Phytoremediation* **2017**, *19*, 133–141. [[CrossRef](#)]
200. Amrhein, C.; Mosher, P.A.; Brown, A.D. The effects of redox on Mo, U, B, V, and As solubility in evaporation pond soils. *Soil Sci.* **1993**, *155*, 249–255. [[CrossRef](#)]
201. Metals-U.S. Food & Drug Administration. Available online: <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/default.htm> (accessed on 18 February 2019).
202. Lapworth, D.J.; Stolpe, B.; Williams, P.J.; Gooddy, D.C.; Lead, J.R. Characterization of suboxic groundwater colloids using a multi-method approach. *Environ. Sci. Technol.* **2013**, *47*, 2554–2561. [[CrossRef](#)] [[PubMed](#)]
203. Praetorius, A.; Scheringer, M.; Hungerbühler, K. Development of environmental fate models for engineered nanoparticles—A case study of TiO₂ nanoparticles in the Rhine River. *Environ. Sci. Technol.* **2012**, *46*, 6705–6713. [[CrossRef](#)] [[PubMed](#)]
204. González-Gálvez, D.; Janer, G.; Vilar, G.; Vilchez, A.; Vázquez-Campos, S. *The Life Cycle of Engineered Nanoparticles*; Springer: Cham, Switzerland, 2017; pp. 41–69.
205. Thomé, A.; Reddy, K.R.; Reginatto, C.; Cecchin, I. Review of nanotechnology for soil and groundwater remediation: Brazilian perspectives. *Water Air Soil Pollut.* **2015**, *226*, 121. [[CrossRef](#)]
206. Gehrke, I.; Geiser, A.; Somborn-Schulz, A. Innovations in nanotechnology for water treatment. *Nanotechnol. Sci. Appl.* **2015**, *8*, 1–17. [[CrossRef](#)] [[PubMed](#)]
207. Troester, M.; Brauch, H.-J.; Hofmann, T. Vulnerability of drinking water supplies to engineered nanoparticles. *Water Res.* **2016**, *96*, 255–279. [[CrossRef](#)] [[PubMed](#)]
208. Alimi, O.S.; Farnier Budarz, J.; Hernandez, L.M.; Tufenkji, N. Microplastics and nanoplastics in aquatic environments: Aggregation, deposition, and enhanced contaminant transport. *Environ. Sci. Technol.* **2018**, *52*, 1704–1724. [[CrossRef](#)] [[PubMed](#)]
209. Lehner, R.; Weder, C.; Petri-Fink, A.; Rothen-Rutishauser, B. Emergence of nanoplastic in the environment and possible impact on human health. *Environ. Sci. Technol.* **2019**, *53*, 1748–1765. [[CrossRef](#)] [[PubMed](#)]

210. Oberdörster, G.; Oberdörster, E.; Oberdörster, J. Nanotoxicology: An emerging discipline evolving from studies of ultrafine particles. *Environ. Health Perspect.* **2005**, *113*, 823–839. [[CrossRef](#)]
211. Ganguly, P.; Breen, A.; Pillai, S.C. Toxicity of nanomaterials: Exposure, pathways, assessment, and recent advances. *ACS Biomater. Sci. Eng.* **2018**, *4*, 2237–2275. [[CrossRef](#)]
212. Environmental Protection Agency. *Technical Fact Sheet—Nanomaterials*; USEPA: Washington, DC, USA, 2017. Available online: https://www.epa.gov/sites/production/files/2014-03/documents/ffrofactsheet_emergingcontaminant_nanomaterials_jan2014_final.pdf (accessed on 2 February 2019).
213. Inshakova, E.; Inshakov, O. World market for nanomaterials: Structure and trends. *MATEC Web Conf.* **2017**. [[CrossRef](#)]
214. Hyung, H.; Kim, J.-H. Dispersion of C60 in natural water and removal by conventional drinking water treatment processes. *Water Res.* **2009**, *43*, 2463–2470. [[CrossRef](#)] [[PubMed](#)]
215. Zhang, Y.; Chen, Y.; Westerhoff, P.; Hristovski, K.; Crittenden, J.C. Stability of commercial metal oxide nanoparticles in water. *Water Res.* **2008**, *42*, 2204–2212. [[CrossRef](#)] [[PubMed](#)]
216. Hedberg, J.; Blomberg, E.; Odnevall Wallinder, I. In the search for nonspecific effects of dissolution of metallic nanoparticles at freshwater-like conditions: A critical review. *Environ. Sci. Technol.* **2019**, *53*, 4030–4044. [[CrossRef](#)] [[PubMed](#)]
217. Gottschalk, F.; Sonderer, T.; Scholz, R.W.; Nowack, B. Modeled environmental concentrations of engineered nanomaterials (TiO₂, ZnO, Ag, CNT, Fullerenes) for different regions. *Environ. Sci. Technol.* **2009**, *43*, 9216–9222. [[CrossRef](#)] [[PubMed](#)]
218. Blaser, S.A.; Scheringer, M.; MacLeod, M.; Hungerbühler, K. Estimation of cumulative aquatic exposure and risk due to silver: Contribution of nano-functionalized plastics and textiles. *Sci. Total Environ.* **2008**, *390*, 396–409. [[CrossRef](#)] [[PubMed](#)]
219. Brar, S.K.; Verma, M.; Tyagi, R.D.; Surampalli, R.Y. Engineered nanoparticles in wastewater and wastewater sludge—Evidence and impacts. *Waste Manag.* **2010**, *30*, 504–520. [[CrossRef](#)] [[PubMed](#)]
220. Baalousha, M.; Yang, Y.; Vance, M.E.; Colman, B.P.; McNeal, S.; Xu, J.; Blaszcak, J.; Steele, M.; Bernhardt, E.; Hochella, M.F. Outdoor urban nanomaterials: The emergence of a new, integrated, and critical field of study. *Sci. Total Environ.* **2016**, *557–558*, 740–753. [[CrossRef](#)] [[PubMed](#)]
221. Min Park, C.; Hoon Chu, K.; Her, N.; Jang, M.; Baalousha, M.; Heo, J.; Yoon, Y. Occurrence and removal of engineered nanoparticles in drinking water treatment and wastewater treatment processes. *Sep. Purif. Rev.* **2017**, *46*, 255–272. [[CrossRef](#)]
222. Kaphle, A.; Navya, P.N.; Umapathi, A.; Daima, H.K. Nanomaterials for agriculture, food and environment: Applications, toxicity and regulation. *Environ. Chem. Lett.* **2018**, *16*, 43–58. [[CrossRef](#)]
223. Pourzahedi, L.; Pandorf, M.; Ravikumar, D.; Zimmerman, J.B.; Seager, T.P.; Theis, T.L.; Westerhoff, P.; Gilbertson, L.M.; Lowry, G.V. Life cycle considerations of nano-enabled agrochemicals: Are today's tools up to the task? *Environ. Sci. Nano* **2018**, *5*, 1057–1069. [[CrossRef](#)]
224. Martínez-Ballesta, M.C.; Zapata, L.; Chalbi, N.; Carvajal, M. Multiwalled carbon nanotubes enter broccoli cells enhancing growth and water uptake of plants exposed to salinity. *J. Nanobiotechnol.* **2016**, *14*, 42. [[CrossRef](#)] [[PubMed](#)]
225. Chen, G.; Qiu, J.; Liu, Y.; Jiang, R.; Cai, S.; Liu, Y.; Zhu, F.; Zeng, F.; Luan, T.; Ouyang, G. Carbon nanotubes act as contaminant carriers and translocate within plants. *Sci. Rep.* **2015**, *5*, 15682. [[CrossRef](#)] [[PubMed](#)]
226. Vithanage, M.; Seneviratne, M.; Ahmad, M.; Sarkar, B.; Ok, Y.S. Contrasting effects of engineered carbon nanotubes on plants: A review. *Environ. Geochem. Health* **2017**, *39*, 1421–1439. [[CrossRef](#)] [[PubMed](#)]
227. Lv, J.; Christie, P.; Zhang, S. Uptake, translocation, and transformation of metal-based nanoparticles in plants: Recent advances and methodological challenges. *Environ. Sci. Nano* **2019**, *6*, 41–59. [[CrossRef](#)]
228. Larue, C.; Khodja, H.; Herlin-Boime, N.; Brisset, F.; Flank, A.M.; Fayard, B.; Chaillou, S.; Carrière, M. Investigation of titanium dioxide nanoparticles toxicity and uptake by plants. *J. Phys. Conf. Ser.* **2011**, *304*, 012057. [[CrossRef](#)]
229. Siddiqi, K.S.; Husen, A. Plant response to engineered metal oxide nanoparticles. *Nanoscale Res. Lett.* **2017**, *12*, 92. [[CrossRef](#)] [[PubMed](#)]
230. Rastogi, A.; Zivcak, M.; Sytar, O.; Kalaji, H.M.; He, X.; Mbarki, S.; Brestic, M. Impact of metal and metal oxide nanoparticles on plant: A critical review. *Front. Chem.* **2017**, *5*, 78. [[CrossRef](#)]
231. Zhu, H.; Han, J.; Xiao, J.Q.; Jin, Y. Uptake, translocation, and accumulation of manufactured iron oxide nanoparticles by pumpkin plants. *J. Environ. Monit.* **2008**, *10*, 713–717. [[CrossRef](#)]

232. Tripathi, D.K.; Tripathi, A.; Shweta; Singh, S.; Singh, Y.; Vishwakarma, K.; Yadav, G.; Sharma, S.; Singh, V.K.; Mishra, R.K.; et al. Uptake, Accumulation and toxicity of silver nanoparticle in autotrophic plants, and heterotrophic microbes: A concentric review. *Front. Microbiol.* **2017**, *8*, 7. [[CrossRef](#)]
233. Dimkpa, C.O.; Bindraban, P.S. Nanofertilizers: New products for the industry? *J. Agric. Food Chem.* **2018**, *66*, 6462–6473. [[CrossRef](#)]
234. Rui, M.; Ma, C.; Hao, Y.; Guo, J.; Rui, Y.; Tang, X.; Zhao, Q.; Fan, X.; Zhang, Z.; Hou, T.; et al. Iron oxide nanoparticles as a potential iron fertilizer for peanut (*Arachis hypogaea*). *Front. Plant Sci.* **2016**, *7*, 815. [[CrossRef](#)] [[PubMed](#)]
235. Motyka, O.; Štrbová, K.; Olšovská, E.; Seidlerová, J. Influence of Nano-ZnO Exposure to Plants on L-Ascorbic Acid Levels: Indication of Nanoparticle-Induced Oxidative Stress. *J. Nanosci. Nanotechnol.* **2019**, *19*, 3019–3023. [[CrossRef](#)] [[PubMed](#)]
236. Liu, J.; Williams, P.C.; Goodson, B.M.; Geisler-Lee, J.; Fakharifar, M.; Gemeinhardt, M.E. TiO₂ nanoparticles in irrigation water mitigate impacts of aged Ag nanoparticles on soil microorganisms, Arabidopsis thaliana plants, and Eisenia fetida earthworms. *Environ. Res.* **2019**, *172*, 202–215. [[CrossRef](#)] [[PubMed](#)]
237. Hu, G.; Cao, J. Occurrence and significance of natural ore-related Ag nanoparticles in groundwater systems. *Chem. Geol.* **2019**, *515*, 9–21. [[CrossRef](#)]
238. Céspedes, C.; Yeo, M.-K. Life cycle assessment of a celery paddy macrocosm exposed to manufactured Nano-TiO₂. *Toxicol. Environ. Health Sci.* **2018**, *10*, 288–296. [[CrossRef](#)]
239. Wu, F.; Zhou, Z.; Hicks, A.L. Life cycle impact of titanium dioxide nanoparticle synthesis through physical, chemical, and biological routes. *Environ. Sci. Technol.* **2019**, *53*, 4078–4087. [[CrossRef](#)] [[PubMed](#)]
240. Solanki, P.; Bhargava, A.; Chhipa, H.; Jain, N.; Panwar, J. Nano-fertilizers and their smart delivery system. In *Nanotechnologies in Food and Agriculture*; Springer International Publishing: Cham, Switzerland, 2015; pp. 81–101.
241. Rai, M.; Ribeiro, C.; Mattoso, L.; Duran, N. Nanotechnologies in food and agriculture. *Nanotechnol. Food Agric.* **2015**, 1–347. [[CrossRef](#)]
242. Kah, M.; Kookana, R.S.; Gogos, A.; Bucheli, T.D. A critical evaluation of nanopesticides and nanofertilizers against their conventional analogues. *Nat. Nanotechnol.* **2018**, *13*, 677–684. [[CrossRef](#)]
243. White, J.C.; Gardea-Torresdey, J. Achieving food security through the very small. *Nat. Nanotechnol.* **2018**, *13*, 627–629. [[CrossRef](#)]
244. Jackson, R.B.; Carpenter, S.R.; Dahm, C.N.; McKnight, D.M.; Naiman, R.J.; Postel, S.L.; Running, S.W. Water in a changing world. *Ecol. Appl.* **2001**, *11*, 1027–1045. [[CrossRef](#)]
245. RamyaPriya, R.; Elango, L. Evaluation of geogenic and anthropogenic impacts on spatio-temporal variation in quality of surface water and groundwater along Cauvery River, India. *Environ. Earth Sci.* **2018**, *77*, 1–17. [[CrossRef](#)]
246. Etheridge, A.B.; MacCoy, D.E.; Weakland, R.J. *Water-Quality and Biological Conditions in Selected Tributaries of the Lower Boise River, Southwestern Idaho, Water Years 2009–12 Scientific Investigations Report 2014–5132*; U.S. Geological Survey: Reston, VA, USA, 2014; pp. 1–70.
247. *Use of Reclaimed Water and Sludge in Food Crop Production*; National Academies Press: Washington, DC, USA, 1996; ISBN 978-0-309-05479-9.
248. Crook, J.; Surampalli, R.Y. Water reclamation and reuse criteria in the U.S. *Water Sci. Technol.* **1996**, *33*, 451–462. [[CrossRef](#)]
249. Bortolini, L.; Maucieri, C.; Borin, M. A tool for the evaluation of irrigation water quality in the arid and semi-arid regions. *Agronomy* **2018**, *8*, 23. [[CrossRef](#)]

