



Complete Genome Sequences of *Mycoplasma anatis*, *M. anseris*, and *M. cloacale* Type Strains

Dénes Gróznér,^a Barbara Forró,^a Kinga Mária Sulyok,^a Szilvia Marton,^a Zsuzsa Kreizinger,^a Krisztián Bányai,^a Miklós Gyuranecz^{a,b}

^aInstitute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest, Hungary

^bDepartment of Microbiology and Infectious Diseases, University of Veterinary Medicine, Budapest, Hungary

ABSTRACT *Mycoplasma anatis*, *M. anseris*, and *M. cloacale* are pathogens of waterfowl. Airsacculitis, nervous disease, and reproductive disorders are the main symptoms in the affected flocks. Here, we report the complete genome sequences of the *M. anatis* (NCTC 10156), *M. anseris* (ATCC 49234), and *M. cloacale* (NCTC 10199) type strains.

Mycoplasma anatis, *M. anseris*, and *M. cloacale* are waterfowl-pathogenic bacteria. *M. anatis* may cause serious nervous symptoms under stress conditions in ducks (1), while airsacculitis, peritonitis, and the increase of embryo lethality were described after experimental inoculation of the pathogen (2). The type strain was isolated from a duck with sinusitis (3). *M. anseris* causes airsacculitis, peritonitis, and embryo lethality (4) and probably has a role in cloaca and phallus inflammation of ganders (5). The type strain was isolated from a flock with a history of phallus inflammation (6). The *M. cloacale* type strain was isolated from a turkey (7), but this species could be isolated from ducks and geese as well (8, 9). Egg infertility is the most common symptom caused by this agent in waterfowl (10). All three of these *Mycoplasma* strains can be transmitted vertically (4, 11, 12). The coexistence of waterfowl-pathogenic mycoplasmas has been described (13).

The *M. anatis* (NCTC 10156), *M. anseris* (ATCC 49234), and *M. cloacale* (NCTC 10199) type strains were purchased directly from the repositories. Cells were grown in Oxoid *Mycoplasma* broth medium (pH 7.8) (Thermo Fisher Scientific, Inc., Waltham, MA) supplemented with 0.5% (wt/vol) sodium pyruvate, 0.5% (wt/vol) glucose, and 0.005% (wt/vol) phenol red and were incubated at 37°C. DNA was extracted with the QIAamp DNA minikit (Qiagen, Inc., Hilden, Germany). DNA libraries were prepared with the Nextera mate pair library preparation kit (Illumina, Inc., San Diego, CA). Two genome sequencing runs were performed on an Illumina NextSeq 500 instrument for each strain, generating 2 × 150-bp (300 cycles) and 2 × 75-bp (150 cycles) mate pair reads. NxTrim software (14) was used to trim the junction adapters from all the raw mate pair reads, generating shorter paired-end reads as well. First, contigs were generated per strain from the paired-end output data by the SPAdes Genome Assembler 3.11 (15) with the assembly-only option. Then, the paired-end contigs and the trimmed mate pair output data were assembled with the same option, generating the draft genomes. Trimmed reads (mate pair and paired-end) were control mapped to the draft *de novo* genome and curated with Geneious 9.1.8 software (16). Circularization of the contigs was performed by primer pairs and PCR assays specific for the contigs' ends (data not shown), and the PCR products were sequenced on the ABI Prism 3100 (Applied Biosystems, Foster City, CA) automated DNA sequencer. The NCBI Prokaryotic Genome Annotation Pipeline (17) online service was used to annotate the genomes. The rRNA

Received 10 July 2018 Accepted 4

September 2018 Published 27 September 2018

Citation Gróznér D, Forró B, Sulyok KM, Marton S, Kreizinger Z, Bányai K, Gyuranecz M. 2018. Complete genome sequences of *Mycoplasma anatis*, *M. anseris*, and *M. cloacale* type strains. *Microbiol Resour Announc* 7:e00939-18. <https://doi.org/10.1128/MRA.00939-18>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2018 Gróznér et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Miklós Gyuranecz, mgyuranecz@gmail.com.

D.G. and B.F. contributed equally to this work.

TABLE 1 Genome information and GenBank accession numbers of *Mycoplasma anatis*, *M. anseris*, and *M. cloacale* type strains^a

Strain	GenBank accession no.	SRA no.	Size (bp)	Coverage (×)	G+C content (%)	No. of CDS ^b	No. of rRNAs	No. of tRNAs	No. of ncRNAs ^c
<i>M. anatis</i> (NCTC 10156)	CP030141	SRP155810	956,093	292	26.7	791	6	33	1
<i>M. anseris</i> (ATCC 49234)	CP030140	SRP155813	750,010	1,833	26.4	617	6	32	2
<i>M. cloacale</i> (NCTC 10199)	CP030103	SRP155814	659,552	1,439	27.0	541	4	31	2

^aThe number of tmRNAs was 1 and the number of regulatory elements was 1 for each strain listed.

^bCDS, coding sequences.

^cncRNAs, noncoding RNAs.

and the tRNA genes and the transfer-messenger RNA (tmRNA) genes were verified by RNAmmer (18) and ARAGORN (19), respectively.

The total genome sizes and information concerning the strains are detailed in Table 1. We hope that the presented complete and circularized genomes will improve research of the waterfowl-pathogenic *Mycoplasma* species.

Data availability. The annotated genome sequences were deposited in GenBank, and the raw read data are available in the Sequence Read Archive. The accession numbers are listed in Table 1.

ACKNOWLEDGMENTS

This work was supported by the Lendület (Momentum) program (LP2012-22) of the Hungarian Academy of Sciences. M.G., S.M., and Z.K. were supported by the Bolyai János Fellowship of the Hungarian Academy of Sciences. M.G. was supported by the Bolyai+ Fellowship (ÚNKP-18-4) of the New National Excellence Program of the Ministry of Human Capacities. The funders had no role in study design, analysis and interpretation, decision to publish, or preparation of the manuscript.

REFERENCES

- Ivanics E, Glávits R, Takács G, Molnár É, Bitay Z, Meder M. 1988. An outbreak of *Mycoplasma anatis* infection associated with nervous symptoms in large-scale duck flocks. *J Vet Med B* 35:368–378. <https://doi.org/10.1111/j.1439-0450.1988.tb00509.x>.
- Stipkovits L. 1979. The pathogenicity of avian mycoplasmas. *Zentralbl Bakteriol Orig A* 245:171–183.
- Roberts DH. 1964. The isolation of an influenza virus and a *Mycoplasma* associated with duck sinusitis. *Vet Rec* 76:470–473.
- Stipkovits L, Kempf I. 1996. Mycoplasmoses in poultry. *Rev Sci Tech* 15:1495–1525. <https://doi.org/10.20506/rst.15.4.986>.
- Stipkovits L, Szathmary S. 2012. *Mycoplasma* infection of ducks and geese. *Poult Sci* 91:2812–2819. <https://doi.org/10.3382/ps.2012-02310>.
- Bradbury JM, Jordan FTW, Shimizu T, Stipkovits L, Varga Z. 1988. *Mycoplasma anseris* sp. nov. found in geese. *Int J Syst Bacteriol* 38:74–76. <https://doi.org/10.1099/00207713-38-1-74>.
- Bradbury JM, Forrest M. 1984. *Mycoplasma cloacale*, a new species isolated from a turkey. *Int J Syst Evol Microbiol* 34:389–392. <https://doi.org/10.1099/00207713-34-4-389>.
- Bencina D, Dorner D, Tadina T. 1987. *Mycoplasma* species isolated from six avian species. *Avian Pathol* 16:653–664. <https://doi.org/10.1080/03079458708436413>.
- Stipkovits L, Varga Z, Dobos-Kovács M, Sántha M. 1984. Biochemical and serological examination of some *Mycoplasma* strains of goose origin. *Acta Vet Hung* 32:117–125.
- Stipkovits L, Varga Z, Czifra G, Dobos-Kovács M. 1986. Occurrence of mycoplasmas in geese affected with inflammation of the cloaca and phallus. *Avian Pathol* 15:289–299. <https://doi.org/10.1080/03079458608436289>.
- Bencina D, Tadina T, Dorner D. 1988. Natural infection of ducks with *Mycoplasma synoviae* and *Mycoplasma gallisepticum* and *Mycoplasma* egg transmission. *Avian Pathol* 17:441–449. <https://doi.org/10.1080/03079458808436462>.
- Samuel AMD, Goldberg DR, Thomas CB, Sharp P. 1995. Effects of *Mycoplasma anatis* and cold stress on hatching success and growth of mallard ducklings. *J Wildl Dis* 31:172–178. <https://doi.org/10.7589/0090-3558-31.2.172>.
- Hinz K-H, Pfützer H, Behr K-P. 1994. Isolation of mycoplasmas from clinically healthy adult breeding geese in Germany. *Zentralbl Veterinarmed B* 41:145–147. <https://doi.org/10.1111/j.1439-0450.1994.tb00217.x>.
- O'Connell J, Schulz-Trieglaff O, Carlson E, Hims MM, Gormley NA, Cox AJ. 2015. NxTrim: optimized trimming of Illumina mate pair reads. *Bioinformatics* 31:2035–2037. <https://doi.org/10.1093/bioinformatics/btv057>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
- Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Lagesen K, Hallin P, Rødland EA, Staerfeldt H-H, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <https://doi.org/10.1093/nar/gkm160>.
- Laslet D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.