

Phytopathol. Mediterr. (2008) 47, 214–222

Effect of pea cultivar, pathogen isolate, inoculum concentration and leaf wetness duration on *Ascochyta* blight caused by *Mycosphaerella pinodes*

BENALI SETTI¹, MOHAMED BENCHEIKH¹, JAMEL HENNI² and CLAIRE NEEMA³

¹Institut de Biologie, Université de Chlef, BP151, 02000 Chlef, Algérie

²Institut des Sciences, Université d'Es Senia, Oran, Algérie

³UMR de Pathologie Végétale, INRA/INA-PG/Université Paris VI,
16 rue Claude Bernard, 75231 Paris Cedex, France

Summary. The effect of host leaf wetness duration, *Mycosphaerella pinodes* inoculum concentration and pathogen isolate on the latent period and the incubation period of the pathogen or disease severity were quantified on pea (*Pisum sativum* L.). Seedlings of two widely grown pea cultivars, Onward and Merveille de Kelvedon, respectively susceptible and moderately resistant to *M. pinodes* were subjected to six leaf wetness durations of 6, 12, 24, 48 and 72 h, and inoculated with five inoculum concentrations, 2.5×10^3 , 4×10^4 , 3.5×10^5 , 4×10^6 , and 5.2×10^7 , in order to determine whether the cultivars reacted differently to *M. pinodes* isolates inoculated under identical conditions. Increasing the duration of leaf wetness and inoculum concentration caused significant ($P < 0.001$) increases in disease severity within each cultivar. Both the incubation period and the latent period decreased with increasing conidial concentration and leaf wetness duration. Generally, the cv. Onward had a significantly shorter incubation period, and latent period and higher disease severity than cv. Merveille de Kelvedon. Isolates differed in aggressiveness at higher levels of leaf wetness (48 h) duration and of inoculum concentration (4×10^6), but there was no significant interaction between isolates and leaf wetness duration, or between isolates and inoculum concentration. The optimum levels for obtaining a consistent infection and for readily separating the susceptible and the partially resistant cultivars were a leaf wetness of 48 h and an inoculum concentration of 4×10^6 . The study also showed that continuous leaf wetness for 48 h was a threshold for application of fungicides to control the fungus in the susceptible cultivar.

Key words: disease severity, incubation period, *Pisum sativum*.

Introduction

In Algeria, little information is available on the factors affecting the disease severity (DS) and the yield losses caused by *Ascochyta* blight from *Mycosphaerella pinodes* (Berk & Blox) Vesterg on peas (*Pisum sativum* L.) (Bouznad, 1998). According to Van der Plank's theorem, quantitative information on how environmental factors affect specific disease components is useful to determine the quantitative impact that plant genotypes with partial resistance

have on the rate of disease progress in the field. *Ascochyta* blight from *M. pinodes* has become more common in the recent years because of the large amount of inoculum left in the field, which is then spread by wind and rain splash.

The pea cultivars widely grown in Algeria are not all totally resistant to *M. pinodes*, but no information is available on how these cultivars react to epidemiological factors such as temperature, leaf wetness duration (LWD) and inoculum concentration (IC). Typically, the first symptoms of *M. pinodes* occur on the plant stems from where they progress to other parts of the plants (Raynal, 1977; Bouznad, 1988; Maufras, 1996; Xue *et al.*, 1997; Tivoli, 1999; Bretag, 2006; Tivoli and Banniza, 2007). Symptoms consist of lesions surrounded by a yellowish chlorotic ring

Corresponding author: B. Setti
E-mail: setiben@yahoo.fr

which evolves into dark-brown spots lacking distinct margins on the leaves, and to reddish spots with a necrotic centre on the pods (Champion *et al.*, 1984; Pederson and Morrall, 1994; Maufras, 1996; Vloutoglou and Kalogerakis, 2000). Allard *et al.* (1993), Tivoli and Lemarchand (1998) stated that *M. pinodes* is strongly correlated with temperature and LWD. Singh and Reddy (1990) suggested that temperature below than 6°C impedes the infection process. Whereas, temperatures between 18 and 21°C and a LWD of 12 h or more favour infection initiation.

Previous studies also report that moisture affects all stages of the disease process. Hence Shaner (1981), Royle and Butler (1986) and Van Ginkel and Scharen (1988), suggested that moisture parameters should be investigated as a means to predict the likelihood of the infection of airborne spores.

Fitt *et al.* (1989, 1998), Huber and Gillespie (1992), also reported that free water on the leaves affects sporulation. On the other hand, Makowski (1993), Tivoli *et al.* (1997) and Rogger *et al.* (1999) stated that IC had a strong impact on the life cycle of *M. pinodes* and hence on all the components of *Ascochyta* blight. The aim of the present work was to examine the conditions favouring infection, including LWD and IC, and that enable the susceptible cultivars to be separated from the moderately resistant cultivars. This is expected to lead to an accurate timing of fungicide application, compared with a strict calendar approach which rarely takes the cultivar susceptibility into account.

Materials and methods

Plant material

Pea plants cv. Onward and cv. Merveille de Kelvedon (MK) were used in all experiments. The two cultivars are the most commonly cultivated varieties in Algeria, the first one being highly susceptible and the second one moderately resistant to the blight.

Seeds of each cultivar were sown in 20-cm diameter pots (10 seeds per pot) containing an unsterilized soil/compost mixture. The seedlings were thinned to five per pot. The plants were maintained in a glasshouse. Three replicates were used for each combination.

Fungal material and inoculum production

Two isolates of *M. pinodes*, md0202 and tn0203 were used in the study. The isolates came from two localities in the Chellif region and presented

respectively a low and a high score of aggressiveness on 'Onward'.

Strains were raised on potato dextrose agar (PDA) for 10 days at 21°C. Conidia from 10 day-old cultures were collected by adding 10 ml of sterile distilled water to dislodge the spores. The spore suspension was filtered through two layers of cheesecloth to remove mycelium and agar fragments. The spore concentration was determined using a haemocytometer. The conidial suspension was diluted with sterile distilled water to the concentration required for each experiment.

Leaf wetness duration

Two week old pea seedlings of 'Onward' and 'MK' were sprayed to run-off with a conidial suspension of 4×10^6 spores ml⁻¹. The seedlings were then subjected to LWD of 6, 12, 24, 48 or 72 h. To maintain the wetness, the seedlings were covered with transparent polyethylene bags sprayed inside with distilled water. Unbagged seedlings were taken not to have been exposed to leaf wetness. At the end of the LWD, seedlings were uncovered and kept in a glasshouse at temperature from 15 to 25°C.

Inoculum concentration

The IC effect was investigated on 15 day-old (three-leaf stage) pea plants of 'Onward' and 'MK'. Plants were inoculated by spraying to runoff with spore suspensions containing 2.5×10^3 , 4×10^4 , 3.5×10^5 , 4×10^6 , and 5.2×10^7 spores ml⁻¹. Control plants were sprayed with sterile distilled water. Suspensions were applied with a spray atomiser with an adjustable nozzle to form a high density of fine droplets on the aerial parts of the plants. Immediately after inoculation, the plants were covered for 48 h with transparent polyethylene bags sprayed inside with distilled water. After the incubation period (IP), the plants were uncovered and kept in a glasshouse at temperature from 15 to 25°C.

Disease assessment

Mycosphaerella pinodes infection on the leaves was recorded 21 days after inoculation using a 0–5 disease scale following to Tivoli *et al.* (1996), where 0, no lesion; 1, a few scattered flecks; 2, numerous flecks; 3, 10–15% leaf area necrotic and appearance of flecks; 4, 50% of leaf area covered by lesions; 5, 75–100% of leaf area dehydrated or necrotic. To

determine the IP and latent period (LP), plants were inspected daily for up to 20 days.

The IP was defined as the period (in days) from host inoculation to the appearance of the first symptoms on the leaves, and the LP as the time (in days) from inoculation to the first formation of pycnidia in the lesions. Each lesion with pycnidia on leaves was recorded with the aid of a hand lens (10×).

Statistical analysis

Data analyzed were subjected to analysis of variance (ANOVA) considering the effect of factors individually and the interaction of factors using the R Statistics software (Version 2.5.0). ANOVA was carried out to detect the effect of IC, LWD, isolate and host susceptibility on IP, LP and DS. Data for IC were transformed to the natural logarithmic scale before analysis.

Results

Inoculum concentration effects

Analysis of variance detected that IC, cultivar and isolate had significant effects on IP, LP and DS ($P < 0.001$) (Table 1). No significant interactions were found between either isolate × IC, or between cultivar × isolate × IC (Table 1).

tn0203 inoculated on Onward at an IC of 5.2×10^7 spores ml^{-1} and the lowest DS (mean: 1.0, SD: 0.25) was found when isolate md0202 was inoculated on MK at an IC of 2.5×10^3 spores ml^{-1} (Fig. 1a).

The IP was also affected by the IC. The shortest IP (mean: 3.3 days, SD: 0.52) was found when tn0203 was inoculated on 'Onward' with an IC of 5.2×10^7 and the longest (mean: 8 days, SD: 0.86) occurred when md0202 was inoculated on 'Onward' with an IC of 2.5×10^3 (Fig. 1b).

Similarly, the LP was shortest when the IC was highest. The LP for md0202 inoculated on 'Onward' ranged from 11 days (SD: 0.47) respectively for the highest IC (5.2×10^7) to 18 days (SD: 1.12) for the lowest IC (2.5×10^3). When md0202 was inoculated on 'MK', the LP ranged from 10.5 (SD: 0.45) days for the highest IC to 18.6 (SD: 0.63) days for the lowest IC (Fig. 1c).

Leaf wetness effects

Analysis of variance detected that LWD, cultivar and isolates all had a significant effect on DS, IP and LP ($P < 0.05$) (Table 2). For DS and IP, there were also significant differences between cultivars × LWD. The interaction between the cultivar × isolate was also significant for DS and LP. However, there was no interaction between isolate × LWD or between isolate × cultivar × LWD (Table 2).

Table 1. Analysis of variance on the effect of cultivar, isolate and inoculum concentration on some epidemiological components of *Ascochyta* blight evaluated on three- leaf stage pea seedlings.

Source	Disease severity ^a				Incubation period ^a				Latent period ^a			
	DF	MS	F	P>F	DF	MS	F	P>F	DF	MS	F	P>F
Cultivar	1	25.741	278.2865	<0.001	1	3.267	21.7778	<0.001	1	3.75	8.0357	<0.01
IC	4	10.019	108.3171	<0.001	4	29.358	195.7222	<0.001	4	122.15	261.7500	<0.001
Isolate	1	2.774	29.9838	<0.001	1	8.067	53.7778	<0.001	1	6.02	12.8929	<0.001
Cultivar × IC	4	0.655	7.0793	<0.001	4	2.808	18.7222	<0.001	4	0.25	0.5357	<0.05
Cultivar × isolate	1	0.840	9.0829	<0.01	1	0.267	1.7778	<0.05	1	2.82	6.0357	
Isolate × IC	4	0.192	2.0739		4	0.525	3.5000		4	0.18	0.3929	
Cultivar × isolate × IC	4	0.109	1.1820		4	0.308	2.0556		4	0.82	1.7500	

^a DF, degree of freedom; MS, mean square; F, F ratio; P, probability level.

Symptoms were more severe with higher IC on both the sensitive 'Onward' and the moderately resistant 'MK' (Fig. 1a–c). Dead plants, particularly from 'Onward' were found at the higher ICs of 4×10^6 and 5.2×10^7 spores ml^{-1} . No blight symptoms were seen in the control plants. The highest DS (mean: 4.9, SD: 0.45) occurred with the isolate

The DS increased with the increasing LWD (Fig. 2a–c). The highest DS (mean: 4.90, SD: 0.69) was recorded at a LWD of 72 h when tn0203 was inoculated on 'Onward', and the lowest DS was 1.0 (SD: 0.19) at a LWD of 6 h with md0202 inoculated on 'Onward'. When there was no leaf wetting, the DS was generally lower than 1 (Fig. 2a).




Fig. 1. Effect of log inoculum concentration in *Mycosphaerella pinodes* isolates md0202 and tn0203 inoculated on pea cv. Onward and Merveille de Kelvedon: a) disease severity; b) latent period; c) incubation period.

Table 2. Analysis of variance on the effect of cultivar, isolate and leaf wetness duration on some epidemiological components of *Ascochyta* blight evaluated on three- leaf stage pea seedlings.

Source	Disease severity ^a				Incubation period ^a				Latent period ^a			
	DF	MS	F	P>F	DF	MS	F	P>F	DF	MS	F	P>F
Cultivar	1	2.840	27.4115	<0.001	1	8.681	69.4444	<0.001	1	1.68	5.2609	<0.05
LWD	5	25.740	248.4335	<0.001	5	26.792	214.3333	<0.001	5	658.79	412.4609	<0.001
Isolate	1	0.867	8.3660	<0.01	1	0.681	5.4444	<0.05	1	5.01	15.6957	<0.001
Cultivar × LWD	5	0.769	7.4233	<0.001	5	2.414	19.3111	<0.001	5	3.57	2.2348	
Cultivar × isolate	1	0.661	6.3820	<0.05	1	0.014	0.1111		1	7.35	23.0000	<0.001
Isolate × LWD	5	0.023	0.2233	<0.001	5	0.081	0.6444		5	0.24	0.1478	
Cultivar × isolate × LWD	5	0.134	1.2957	<0.001	5	0.014	0.1111		5	1.90	1.1913	

^a, see Table 1.

The IP of *M. pinodes* was also affected by the LWD. It was shortest (mean: 3 days, SD: 0.27) for the isolate tn0203 inoculated on 'Onward' at a LWD of 72 h. At a LWD of 6 h, on the other hand, the IP ranged from 6.5 to 7 days (SD: 0.47) (Fig. 2b).

The LP ranged from 11 to 20 days with md0202 inoculated on 'Onward' and from 10 to 19 days with tn0203 inoculated on 'Onward'. The shortest LP (10 days) was with tn0203 inoculated on 'Onward'. There was only slight difference in LP between a LWD of 48 h and a LWD of 72 h (Fig. 2c).

Cultivar and isolate effect

The two cultivars differed in their DS for all ICs (Fig. 1a) and for all LWDS (Fig. 2a). The highest DS (mean: 4.9, SD: 0.69) was achieved with the tn0203 inoculated on 'Onward'. The two cultivars were equally resistant at low levels of both IC and LWD. Hence, with an IC of 3.5×10^5 or less and with a LWD of up to 24h the two cultivars exhibited the same level of resistance. At higher IC and LWD, however, the cv. Onward had a higher DS (>4.1) than the cv. MK (DS<3.56).

The IP and the LP differed significantly between cultivars, the lowest IP (mean: 3 days, SD: 0.27) and the lowest LP (mean: 10.33 days, SD: 0.63) were found with tn0203 inoculated on Onward (Fig. 1). The interaction between cultivar and LWD was also significant for both DS and LP.

Effect of isolates

Significant differences were found between the

isolates for DS, IP and LP (Tables 1, 2), but not for the IC. On the other hand, isolate and cultivar interacted significantly for DS with both IC and LWD (Table 2). But this effect of isolate aggressiveness with any of the epidemiological components tested was not visible for IC concentrations below 3.5×10^5 or with a LWD of 24 h. At higher IC and longer LWD, however, their effect became more pronounced. The highest DS (4.9) was found with the isolate tn0203 on 'Onward' followed by isolate md0202 also on 'Onward' (Fig. 1a).

Discussion

Leaf wetness effects

Leaf wetness duration had a significant effect on DS, IP and LP. The wetness of leaf in general is an important factor enabling the fungal pathogens to infect the aerial parts of plants. When the temperature is favourable and the moisture requirements for a pathogen on a susceptible host are met for a sufficient length of time, an epidemic occurs. Weather moisture is frequently used as an indicator of the likelihood of an epidemic (Royle and Butler, 1986).

LWD of 6 h was sufficient to initiate symptoms formation on leaves of both the susceptible and the moderately resistant cultivars but with this duration, the reaction of the two cultivars were similar when the LWD increased, however, the DS also increased up to a duration of 48 h, beyond which the DS did not increase further. Most foliar fungi can infect the leaves of a plant only while the leaves are wet. But the optimal LWD varies depending on




Fig. 2. Effect of leaf wetness duration on: a) disease severity, b) latent period, c) incubation period of *Mycosphaerella pinodes* isolates md0202 and tn0203 inoculated on pea cv. Onward and Merveille de Kelvedon.

the fungus group (Trapero-Casas and Kaiser 1992, Pederson and Morrall, 1994; Gilles *et al.*, 2000). Many previous studies have reported that severe disease was obtained with a LWD of at least 48 h (Shew *et al.*, 1988; Davis and Fitt, 1994; Scott *et al.*, 1995; Rogger *et al.*, 1999).

The IP on the other hand was shortest with a LWD of 48 and 72 h and longest with no LWD or a LWD of 6 h on both the susceptible and the partially resistant cultivar. The IP was shorter on 'Onward' than on 'MK' and the IP on both cultivars decreased as LWD increased. The LP is the interval between the moment of infection and the appearance of the first pycnidium, when the infected plant becomes itself infectious. The LP was longest when the LWD had been shortest. The LP for different species of *Ascochyta* blight has been reported in earlier studies. It was 5 days for *A. rabiei* on chickpea (Trapero-Casas and Kaiser, 1992), 6 days for *A. fabae* f. sp. *lentis* on lentil (Pederson and Morrall, 1994) and 8 to 10 days for *A. fabae* on fabae bean (Wallen and Galway, 1977). Our study found that on pea with an optimal LWD and IC, the LP was of 11 to 12 days on 'Onward'. The long LP could be explained in part as caused by fluctuations in the temperature that had occurred in the uncontrolled glasshouse during the experimental period. These temperatures ranged from 15 to 25°C and could have disturbed and delayed pycnidia production.

Lovell *et al.* (2004) found that pycnidia development was strongly influenced by environmental conditions. This relation has already been reported by other studies (Eden *et al.*, 1996; Webb *et al.*, 1997; Hartman *et al.*, 1999; Buloviene and Surviliene, 2006).

Effect of IC and of isolate

The IC clearly affected the DS, the IP and the LP. The DS increased when IC went up. An IC of 5.2×10^7 caused the greatest DS on the two cultivars, 4.9 on 'Onward' and 3.53 on 'MK'. Plant death in the cv. Onward was also noted with higher ICs particularly 4×10^6 and 5.2×10^7 ml⁻¹. Makowski (1993) found that increasing DS increased the risk of plant death, particularly at the juvenile stages. There was a positive correlation between IC and DS. Such a positive correlation between IC and DS was also demonstrated for other *Mycosphaerella* spp. (Scott *et al.*, 1985; Shew *et al.*, 1988; Rogger *et al.*, 1999). In our study the IP and LP of *M. pinodes* decreased significantly as its IC increased. According to Peder-

son and Morrall (1994), the LP is strongly affected by the IC and the pycnidial development was lower when the IC was lower. Furthermore, Hildebrand and Sutton (1984) reported that a high IC shortened both the IP and the LP. These experiments on IC revealed significant differences between the two isolates. The more aggressive isolate caused the highest DS with the shortest IP and LP.

The study found that there were differences between the isolates. The more aggressive isolate caused a greater DS after a short IP and LP. However, no significant differences were found in the interaction between the isolates and LWD or between isolates and IC.

Both LWD and IC were important for the epidemiology of *M. pinodes*. The pea cv. Onward and MK reacted differently with both a higher IC and longer LWD. At IC of 3×10^5 spores ml⁻¹ and a LWD of 24 h, the two cultivars both exhibited the equal level of moderate resistance to the isolates. But at IC of 4×10^6 spores ml⁻¹ and an LWD of 48 h and longer, the cv. Onward gave susceptible reaction. It was also found that an IC of 4×10^6 spores ml⁻¹, and a LWD of 48 h, together produced a consistent infection that enabled the susceptible and moderately resistant cultivars to be readily separated. Overall, under controlled conditions, an appropriate IC and LWD are very important to evaluate the resistance of pea cultivars to *M. pinodes*. The study also found that continuous leaf wetness of 48 h was the threshold duration for the application of fungicides on susceptible cultivars to control of this disease. Furthermore, the present study highlights the importance of taking account of the resistance of the plant when devising a fungicide spraying program, in order to keep the amount of fungicide sprayed to a minimum.

Literature cited

- Allard D., D. Bill and C. Tourand, 1993. *L'antracnose du Pois. Revue Bibliographique et Synthèse*. Ed. Scientifiques. INRA Rennes, France, 5–24.
- Bailey J.A., 1969. Phytoalexin production by leaves of *Pisum sativum* in relation to senescence. *Annals of Applied Biology* 64, 315–324.
- Bouznad Z., 1988. *Contribution à la Connaissance du Genre Ascochyta chez les Légumineuses en Algérie. L'étude Biologique, Ultrastructurale et Cytochimique, des Relations Hôtes-pathogènes*. PhD Thesis, Université Pierre et Marie Curie, Paris, France, 123 pp.
- Bretag T.W., P.J. Keane and T.V. Price, 2006. The epidemiology of ascochyta blight in field peas: a review. *Australian*

- Journal of Agricultural Research* 57, 883–902.
- Bryne J.M., M.K. Hausbeck, C. Meloche, and A.M. Jarosz, 1998. Influence of dew period and temperature on foliar infection of greenhouse grown tomato by *Colletotrichum coccodes*. *Plant Disease* 82, 639–664.
- Buloviene V. and E. Survilienė, 2006. Effect of environmental conditions and inoculum concentration on sporulation of *Peronospora destructor*. *Agronomy Research* 4, 147–150.
- Chakraborty S., D. Rateliff, and F.J. McKay, 1990. Anthracnose of *Stylosanthes scabra*: effect of leaf surface on disease severity. *Plant Disease* 74, 379–384.
- Champion R., R. Bourdind and G. Berthier, 1984. *L'antracnose du Pois, Détermination des Agents Responsables et Divers Fongicides en Traitements*. INRA, Paris, France, 650 pp.
- Clulow S., P. Mathen and P.G. Lewis, 1991. Genetical analysis of resistance to *Mycosphaerella* in pea seedlings. *Euphytica* 58, 183–189.
- Davis H. and B.D.L. Fitt, 1994. Effect of temperature and leaf wetness on the latent period of *Rhynchosporium secalis* (leaf blotch) on leaves of winter barley. *Journal of Phytopathology* 140, 269–279.
- Eden M.A., R.A. Hill, R. Beresford and A. Stewart, 1996. The influence of inoculum concentration, relative humidity, and temperature on infection of greenhouse tomatoes by *Botrytis cinerea*. *Plant Pathology* 45, 795–806.
- Fitt B.D.L., K.J. Doughty, T. Gilles, P. Gladders, J.M. Steed, H. Su and K.G. Sutherland, 1998. Methods for assessment of light leaf spot (*Pyrenopeziza brassicae*) on winter oilseed rape (*Brassica napus*) in the UK. *Annals of Applied Biology* 133, 329–341.
- Gilles T., B.D.L. Fitt, R. Kennedy, S.J. Welham and M.J. Jeger, 2000. Effects of temperatures and wetness duration on conidial infection, latent period and asexual sporulation of *Pyrenopeziza brassicae* on leaves of oil seed rape. *Plant Pathology* 49, 498–508.
- Graham J.H., T.E. Devine, and C.H. Hanson, 1976. Occurrence and interaction of three species of *Colletotrichum* on alfalfa in the mid-atlantic United States. *Phytopathology* 66, 538–541.
- Hare W.W. and J.C. Walker, 1945. Ascochyta diseases of canning pea. *Wisconsin Research Bulletin* 150, 1–8.
- Hartman J.R., L. Paris, and P. Bautreais, 1999. Effect of leaf wetness duration, temperature, and conidial inoculum dose on apple scab infections. *Plant Disease* 8, 531–534.
- Hassani A.A.E., 1981. *Modalités d'Expression de la Résistance d'un Cultivar de Pois Chiche (Cicer arietinum L.) à Ascochyta rabiei (Pass) Lab*. PhD thesis, Université de Rennes, Rennes, France, 41 pp.
- Heath M.C. and R.K.S. Wood, 1971. Role of inhibitors of fungal growth in the limitation of leaf spots caused by *Ascochyta pisi* and *Mycosphaerella pinodes*. *Annals of Botany* 35, 475–491.
- Hildebrand P.D. and J.C. Sutton, 1984. Relationships of temperature, moisture, and inoculum density to the infection cycle of *Peronospora destructor*. *Canadian Journal of Plant Pathology* 6, 127–134.
- Jhorar O.P., D.R. Butter and S.S. Mathauda, 1998. Effect of leaf wetness duration, relative humidity, light and dark on infection and sporulation by *Didymella rabiei* on chickpea. *Plant Pathology* 47, 586–594.
- Jones A.L., 1986. Role of wet periods in predicting foliar diseases. In: *Plant Disease Epidemiology; Population Dynamics and Management. Vol 1*. (K.J. Leonard, W.E. Fry, ed.), Macmillan Publishing, New York, NY, USA, 87–100.
- Klienbaum D.G. and L.L. Kupper, 1978. *Applied Regression Analysis And Other Multivariate Methods*. Duxbury Press, North Scituate, MA, USA, 556 pp.
- Lovell D.J., T. Hunter, S.J. Powers, S.R. Parker and F. VanDenBosch, 2004. Effect of temperature on latent period of Septoria leaf blotch on winter wheat under outdoor conditions. *Plant Pathology* 53, 170–181.
- Maddock S.E., D.S. Ingram and L. Hubber, 1981. Studies of survival and longevity on the light leaf spot pathogen on brassicas, *Pyrenopeziza brassicae*. *Transactions of the British Mycological Society* 77, 153–159.
- Makowski R.M.D., 1993. Effect of inoculum concentration, temperature, dew period and plant growth stage on disease of round leaved mallow and velvet leaf by *Colletotrichum gloeosporioides* f. sp. *malvae*. *Phytopathology* 83, 1229–1234.
- Manandhar J.B., G.L. Hartman and J.B. Sinclair, 1988. Soybean germplasm evaluation for resistance to *Colletotrichum truncatum*. *Plant Disease* 72, 56–59.
- Maufras J.Y., 1996. L'antracnose, maladie du pois. *Perspectives Agricoles* 212, 6–17.
- Pederson E.A. and R.A.A. Morrall, 1994. Effects of cultivars, leaf wetness duration, temperature, and growth stage on infection and development of Ascochyta blight on lentil. *Phytopathology* 84, 1024–1030.
- Potts A.D., 1990. *Expression and Genetics of Resistance to Septoria tritici in Wheat*. PhD thesis, University of Saskatchewan, Saskatoon, Saskatchewan, Canada.
- Raynal G., 1977. *Maladies Cryptogamiques des Légumineuses, Alimentaires, Fourragères et Industrielles*. Ed. INAP-G, Paris, France, 18 p.
- Reddy M.V. and K.B. Singh, 1990. Relationship between temperature, relative humidity and Ascochyta blight development in winter sown chickpea in Syria. *Phytopathologia Mediterranea* 29, 159–162.
- Rogger C., B. Tivoli and L. Hubber, 1999. Effects of interrupted wet periods and different temperatures on the development of Ascochyta blight caused by *Mycosphaerella pinodes* on pea (*Pisum sativum* L.) seedlings. *Plant Pathology* 48, 10–18.
- Royle D.J. and D.R. Butler, 1986. Epidemiological significance of liquid water in crop canopies and its role in disease forecasting. In: *Water Fungi and Plants*. (P.G. Ayres, L. Boddy, ed.), Cambridge University Press, Cambridge, UK, 139–156.
- Salamati S. and H.A. Magnus, 1997. Leaf blotch severity on spring barley infected by isolates of *Rhynchosporium secalis* under different temperature and humidity regimes. *Plant Pathology* 46, 939–945.
- Scott P.R., P.W. Benediks, H.G. Jones and M.A. Ford, 1985. Some effects of canopy structure and microclimate on

- infection of tall and short wheat by *Septoria nodorum*. *Plant Pathology* 34, 578–593.
- Shaner G., 1981. Effect of environment on fungal leaf blights of small grains. *Annual Review of Phytopathology* 19, 273–296.
- Shearer B.J. and J.C. Zadoks, 1972. The latent period of *Septoria nodorum* in wheat. 1- The effect of temperature and moisture treatments under controlled conditions. *Netherlands Journal of Plant Pathology* 78, 231–241.
- Shew B.B., M.K. Beute and J.C. Wynne, 1988. Effects of temperature and relative humidity on expression of resistance to *Cercosporidium personatum* in peanut. *Phytopathology* 78, 493–498.
- Singh G., 1990. Identification and designation of physiological races of *Ascochyta rabiei* in India. *Indian Phytopathology* 43, 48–54.
- Tivoli B., 1999. L'antracnose du pois. Mieux évaluer sa nuisibilité pour mieux raisonner la protection de la culture. *Phytoma* 512, 16–20.
- Tivoli B. and S. Banniza, 2007. Comparative epidemiology of *Ascochyta* blights on grain legumes. *European Journal of Plant Pathology* 119, 59–76.
- Tivoli B., C. Béasse, E. Lemarchand and E. Masson, 1996. Effects of *Ascochyta* blight (*Mycosphaerella pinodes*) on yield components of single pea (*Pisum sativum* L) plants under field conditions. *Annals of Applied Biology* 129, 207–216.
- Tivoli B., H. Halila and A. Porta Puglia, 1997. Les anthracoses des légumineuses à grosses graines dans les pays du pourtour méditerranéen. In: *Les Légumineuses Alimentaires Méditerranéennes*. INRA ed., Paris, France, pp, 81–102.
- Trapero-Casas A. and W.J. Kaiser, 1992. Influence of temperature, wetness period, plant age, and inoculum concentration on infection and development of *Ascochyta* blight of chickpea. *Phytopathology* 82, 589–596.
- Van Ginkel M. and A.L. Scharen, 1988. Host pathogen relationships of wheat and *Septoria tritici*. *Phytopathology* 78, 762–766.
- Voloutoglou I. and S.N. Kalogerakis, 2000. Effects of inoculum concentration, wetness duration and plant age on development of early blight (*Alternaria solani*) and shedding of leaves in tomato plants. *Plant Pathology* 49, 339–345.
- Wallen V.R., 1974. Influence of three *Ascochyta* diseases of peas on plant development and yield. *Canadian Plant Disease Survey, Disease Highlights* 54, 86–90.
- Webb D.H. and F.W.J.R. Nutter, 1997. Effect of leaf wetness duration and temperature on infection efficiency, latent period, and rate of pustule appearance of rust in alfalfa. *Phytopathology* 87, 946–950.
- Xue A.G., T.D. Warkentin and E.O. Kenaschuk, 1997. Effects of timing of inoculation with *Mycosphaerella pinodes* on yield and seed infection of field pea. *Canadian Journal of Plant Science* 78, 685–690.

Accepted for publication: September 29, 2008